

Stillbirths: Recall to action in high-income countries

1.1 Authors

Vicki Flenady ^{PhD}, Aleena M Wojcieszek ^{BPsySci}, Philippa Middleton ^{PhD}, David Ellwood ^{PhD}, Jan Jaap Erwich ^{PhD}, Michael Coory ^{PhD}, T Yee Khong ^{MD}, Robert M Silver ^{PhD}, Gordon Smith ^{DSc}, Frances M Boyle ^{PhD}, Joy E Lawn ^{PhD}, Hannah Blencowe ^{MRCPC}, Susannah Hopkins Leisher ^{MA}, Mechthild M Gross ^{PhD}, Dell Horey ^{PhD}, Lynn Farrales ^{MD}, Frank Bloomfield ^{PhD}, Lesley McCowan ^{MD}, Stephanie J Brown ^{PhD}, K S Joseph ^{PhD}, Jennifer Zeitlin ^{DSc}, Hanna E Reinebrant ^{PhD}, Claudia Ravaldi ^{MD}, Alfredo Vannacci ^{PhD}, Jillian Cassidy, Paul Cassidy ^{MPhil}, Cindy Farquhar ^{MD}, Euan Wallace ^{PhD}, Dimitrios Siassakos ^{MD}, Alexander EP Heazell ^{PhD}, Claire Storey ^{BA}, Lynn Sadler ^{MPH}, Scott Petersen ^{MBBS}, J Frederik Frøen ^{PhD}, Robert L Goldenberg ^{MD}

1.2 The Lancet Ending Preventable Stillbirths study group

J Frederik Frøen, Joy E Lawn, Hannah Blencowe, Alexander Heazell, Vicki Flenady, Luc de Bernis, Mary Kinney, Susannah Hopkins Leisher

1.3 Author affiliations

Mater Research Institute - The University of Queensland (MRI-UQ), Brisbane, Australia (V Flenady, A M Wojcieszek, S Hopkins Leisher, D Horey, H Reinebrant, S Petersen); International Stillbirth Alliance (V Flenady, A M Wojcieszek, P Middleton, D Ellwood, J Erwich, M Coory, T Y Khong, R M Silver, F M Boyle, S Hopkins Leisher, H Reinebrant, L Farrales, L McCowan, C Ravaldi, A Vannacci, J Cassidy, P Cassidy, E Wallace, D Sisassakos, A Heazell, C Storey); Women's & Children's Health Research Institute (WCHRI), The University of Adelaide, Adelaide, Australia (P Middleton); Griffith University & Gold Coast University Hospital, Gold Coast, Australia (D Ellwood); The University of Groningen, University Medical Center Groningen, Groningen, The Netherlands (J Erwich); Murdoch Childrens Research Institute, Melbourne, Australia (M Coory); SA Pathology, University of Adelaide, Adelaide, Australia (T Y Khong); University of Utah Health Sciences Center, United States (R M Silver); NIHR Biomedical Research Centre & Cambridge University, United Kingdom (G Smith); The University of Queensland, Brisbane, Australia (F M Boyle); London School of Hygiene and Tropical Medicine, London, United Kingdom (J E Lawn, H Blencowe); Hannover Medical School, Hannover, Germany; Zurich University of Applied Sciences, Institute for Midwifery, Winterthur, Switzerland (M M Gross); La Trobe University, Melbourne, Australia (D Horey); Still Life Canada: Stillbirth and Neonatal Death Education, Research and Support Society, Canada (L Farrales); Liggins Institute, The University of Auckland, Auckland, New Zealand (F Bloomfield); The University of Auckland, Auckland, New Zealand

(L McCowan, C Farquhar, L Sadler); Murdoch Childrens Research Institute and General Practice and Primary Health Care Academic Centre, The University of Melbourne, Parkville, Australia, S J Brown); University of British Columbia, Vancouver, Canada (L Farrales, K S Joseph); INSERM, Obstetrical, Perinatal and Paediatric Epidemiology Research Team, Centre for Epidemiology and Biostatistics (U1153), Paris-Descartes University, Paris, France (J Zeitlin); CiaoLapo Onlus - Charity for High-Risk Pregnancies and Perinatal Grief Support, Italy (C Ravaldi; A Vannacci); Department of Neurosciences, Psychology, Drug Research and Child Health, University of Florence, Italy (A Vannacci); Umamanita, Spain (J Cassidy, P Cassidy); Monash University, Melbourne Australia (E Wallace); University of Bristol; Southmead Hospital, Bristol, UK (D Siassakos); University of Manchester, UK; St. Mary's Hospital, Central Manchester University Hospitals NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, (A Heazell); Mater Health Services, Brisbane, Australia (S Petersen); Department of International Public Health, Norwegian Institute of Public Health, Oslo, and Center for Intervention Science for Maternal and Child Health, University of Bergen, Norway (J F Frøen); Department of Obstetrics and Gynecology, Columbia University, New York, USA (RL Goldenberg)

1.4 The Lancet stillbirths in high-income countries investigator group

Bia Bichara (London School of Hygiene & Tropical Medicine, London, UK); Beatrice Blondel (National Institute of Health and Medical Research (INSERM), Paris, France); Sheelagh Bonham (Healthcare Pricing Office (HPO), Ireland); Stephanie Bradley; Joanne Cacciatore (Arizona State University, Tempe, USA); Andre Carvalho (Sands Australia); Paul Corcoran (National Perinatal Epidemiology Centre, University College Cork, Ireland); Vicki Culling (Sands Wellington-Hutt Valley, New Zealand); Jane Dahlstrom (Australian National University, Canberra, Australia; International Stillbirth Alliance); Soo Downe (ReaCH group, University of Central Lancashire, UK); Christine East (Monash University & Monash Health, Melbourne, Australia; International Stillbirth Alliance); Mairie Cregan (Feillecain, Ireland; International Stillbirth Alliance); Robin Cronin (The University of Auckland, Auckland, New Zealand); Francine de Montigny (Université du Québec en Outaouais, Gatineau, Canada); Sabine de Wall (Hannover Medical School, Hannover, Germany); Elizabeth Draper (MBRRACE-UK, Dept of Health Sciences, University of Leicester, Leicester, UK); Wes Duke (U.S. Department of Health and Human Services, Washington, USA; International Stillbirth Alliance); Alison Ellis (The University of Bristol, Bristol, UK); Jane P Fox (Mater Research Institute - The University of Queensland (MRI-UQ), Brisbane, Australia); Ruth C Fretts (Harvard Vanguard Medical Associates, Wellesley, USA; International Stillbirth Alliance); Stephanie Fukui (SIDS Family Association, Japan); Glenn Gardener (Mater Health Services, Brisbane, Australia; International Stillbirth Alliance); Kristen Gibbons (Mater Research Institute - The University of Queensland (MRI-UQ), Brisbane, Australia); Katherine J Gold

(Department of Family Medicine and Department of Obstetrics, University of Michigan, Ann Arbor, USA; International Stillbirth Alliance); Sanne Gordijn (The University of Groningen, University Medical Center Groningen, Groningen, The Netherlands; International Stillbirth Alliance); Adrienne Gordon (The University of Sydney, Sydney, Australia); Grace Guyon (Alberta Health Services Perinatal Program, Alberta Canada); Ibinabo Ibiebele (Mater Research Institute - The University of Queensland (MRI-UQ), Brisbane, Australia); Belinda Jennings (International Stillbirth Alliance); Alison Kent (Australian National University Medical School & Centenary Hospital for Women and Children, Canberra, Australia; International Stillbirth Alliance); Sue Kildea (Mater Health Service, Mater Research Institute, University of Queensland (MRI-UQ) & School of Nursing, Midwifery and Social Work, UQ); Fleurisca Korteweg (The University of Groningen, Martini Hospital, Groningen, The Netherlands; International Stillbirth Alliance); Tina Lavender; Rohan Lourie (Mater Health Services, Brisbane, Australia); Brad Manktelow (MBRRACE-UK, Dept of Health Sciences, University of Leicester, Leicester, UK); Vicki Masson (The University of Auckland, Auckland, New Zealand); Elizabeth McClure (Research Triangle Institute, NC, USA; International Stillbirth Alliance); Sarah Meaney (National Perinatal Epidemiology Centre, University College Cork, Ireland); Kelly Merchant (Bears of Hope, Australia); Susan McDonald (La Trobe University & Mercy Hospital for Women, Melbourne, Australia; International Stillbirth Alliance); Margaret Murphy (University College Cork, Ireland; International Stillbirth Alliance); Jeremy Oats (University of Melbourne, Melbourne, Australia; International Stillbirth Alliance); Karin Pettersson (Karolinska University Hospital, Stockholm, Sweden; International Stillbirth Alliance); Ingela Rådestad (Sophiahemmet University, Stockholm, Sweden; International Stillbirth Alliance); Leanne Raven (SIDS and Kids, Australia); Uma Reddy (Eunice Kennedy Shriver National Institute of Child Health and Human Development, Rockville, USA); Jessica Ruidiaz (Fundación Era en Abril; International Stillbirth Alliance); Anne Schirmann (The University of Queensland, Brisbane, Australia); Janet Scott (SANDS UK; International Stillbirth Alliance); Lucy Smith (MBRRACE-UK, Dept of Health Sciences, University of Leicester, Leicester, UK); William Stones (The University of St Andrews, Scotland, UK; FIGO); Scotland, UK); Anna Usynina (Northern State Medical University, Arkhangelsk, Russia); Susanne van Gogh (Zurich University of Applied Sciences, Institute for Midwifery, Winterthur, Switzerland); Susan P Walker (The University of Melbourne, Melbourne, Australia; International Stillbirth Alliance); Patricia Wilson (Mater Health Services, Brisbane, Australia; International Stillbirth Alliance).

1.5 Correspondence to

Associate Professor Vicki Flenady, Mater Research Institute - The University of Queensland (MRI-UQ), Level 3 Aubigny Place, Mater Health Services, South Brisbane, QLD, Australia

vicki.flenady@mater.uq.edu.au

1.6 Abstract

Variation in stillbirth rates across high-income countries (HIC) and large equity gaps within HIC persist. If all HIC achieved stillbirth rates equal to the best performing countries, 19,439 late gestation stillbirths could have been avoided in 2015. Unexplained stillbirths remain high and can be addressed through improvements in data collection, investigation and classification, and better understanding of causal pathways. Substandard care contributes to 20-30% of all stillbirths, and the contribution is even higher for late-gestation intrapartum stillbirths. National perinatal mortality audit programmes need to be implemented across all HIC. Reducing stigma and fatalism around stillbirth and improving bereavement care are also clear, ongoing priorities for action. In HIC, a child has twice the risk of being stillborn due to adverse socioeconomic circumstances. Community- and country-level programs to improve health among disadvantaged families are needed to address these inequities.

1.7 Key words

Stillbirth; Perinatal mortality; High-income countries; Quality of care; Implementation; Perinatal audit; Classification; Bereavement care

1.8 Key messages

- **Late gestation stillbirth rates vary** across HIC from 1.3 to 8.8/1000 births, showing that further reduction in stillbirths is possible. Setting and monitoring targets in all HIC are important for reducing preventable stillbirths.
- **Socially marginalised and disadvantaged women** often have twice or more the risk of stillbirth. **Social determinants** of maternal and fetal wellbeing should be monitored in all HIC, and addressed through education and alleviation of poverty, as well as better access to health care, especially timely, culturally appropriate antenatal care.
- **Stigma and fatalism** continue to exacerbate trauma for families and impede progress in stillbirth prevention. Stronger parent and care provider partnerships are needed to dispel misperceptions and negative attitudes that persist in communities.
- It is the responsibility of all countries to implement high quality **national perinatal mortality audit** that translates into improvements in quality of care. Key performance indicators of quality maternity care should be measured and reported, with the aim of eliminating **substandard antepartum and intrapartum care** - too often present when a stillbirth occurs.
- **Bereavement care** commonly fails to meet the needs of parents, often with devastating consequences. Immediate bereavement care should be provided by appropriately trained health care professionals with sensitive and seamless transition to community support services in all settings.
- **Poor quality data on stillbirths is a major problem across HIC.** Access to high quality investigation into the causes of stillbirth, including **autopsy and placental histopathology** by a skilled perinatal pathologist, should be made available to all parents following stillbirth. Consensus on a **classification system** for stillbirth, which addresses the contribution of placental pathology, and **standard a definition for reporting stillbirths** that makes it possible to report comparable early and late stillbirth rates across HIC are needed.
- Future research must focus on **stillbirth prediction**, understanding **placental pathways** to stillbirth and causal pathways to **unexplained stillbirth**. Effective strategies to reduce the prevalence of **obesity and smoking** in women of reproductive ages are needed. Understanding pathways leading to **early stillbirth** and **spontaneous preterm birth** at early gestation is also important to pursue.

1.9 Introduction

As stated across this Series, stillbirth rate is a key indicator of women's health and quality of care in pregnancy and childbirth.^{1,2} While high income country (HIC) rates are relatively low compared with low- and middle-income countries (LMIC), stillbirth is a major health burden with rates over double neonatal mortality rates,³ and often equal to all deaths in the first year.⁴ Neonatal death rates continue to fall,⁴ whereas stillbirth rates remain steady, and have increased in some regions.⁵ The death of any child is a tragedy for families, often with profound, long-lasting psychosocial and economic impact.⁶ The 2011 Lancet Stillbirth Series (LSS) drew attention to the slow progress in reducing rates across HIC and highlighted prevention.⁷ In this fourth paper of the Lancet's Ending Preventable Stillbirths series, we summarise the current status of stillbirths in HIC, and present strategies to maintain momentum in reducing deaths and meeting parents' needs when their child is stillborn.

1.10 Methods

We used the range of methods described in Panel 1 with further details provided in the webappendix. We also assessed HIC stillbirth rates and annual rate reductions (ARR) from 2000 to 2015 (Blencowe and colleagues).⁸

1.11 Stillbirth rates – Is progress good enough?

The latest global estimates⁸ show an average stillbirth rate (using the 28 weeks' gestation definition) across 49 HIC of 3.5 per 1000 total births. Country-specific rates varied widely from 1.3 to 8.8 (Figure 1). The ARR from 2000 to 2015 varied, with nine countries demonstrating ARRs of <1%, and five countries >4%. Cross-country comparisons are hindered by data capture issues including reporting practices for termination of pregnancy after the gestational age threshold, variation in data capture mechanisms including the use of cross-linkages between birth and death certificate data and birth registry data, and variation in definitions for reporting.⁸ Use of the WHO recommended lower gestational age limit of 28 weeks likely reduces the influence of these issues on reported rates. Regardless of data capture issues, real epidemiological variation in rates is present,⁹ and shows

further reduction is possible. Stillbirth rates for disadvantaged groups are around double those of greater advantage,^{3, 10-12} with recent evidence indicating these gaps can be narrowed.¹⁰ While intrapartum stillbirths comprise a small proportion (<10%) across HIC, variability in this indicator is also evident, as presented by Lawn and colleagues.⁹ Nonetheless, it is important that countries monitor and understand their own temporal trends rather than assess performance based on rankings with other countries.

As shown in Figure 1, 6 of 49 countries (12%) demonstrated third trimester rates of 2.0/1000 births or lower, showing that this is achievable. If all countries had achieved a stillbirth rate ≤ 2.0 , in 2015, 19,439 late gestation stillbirths could have been avoided.

1.11.1 Early gestation stillbirth

Depending on definition, 35%¹³ to 50%¹⁴ of HIC stillbirths occur below the WHO recommended cut-off for international comparison of 1000 g (or 28 weeks).¹³ Due to variability in definitions (e.g. inclusion of terminations), comparisons of early gestation stillbirth rates between and even within countries are difficult. When crude stillbirth rates were compared, Sweden ranked 3rd and Australia 28th in a comparison of 28 HICs, but Sweden dropped to 10th and Australia improved to 11th when rank was based on stillbirths ≥ 1000 g.¹⁵ Under-reporting of stillbirths <28 weeks is also evident in some regions.³ Ascertainment may be influenced by perceptions of viability.⁸ Despite these difficulties, stillbirth rates <28 weeks' in HIC are not falling, and some increases are evident.^{5, 13, 14, 16} In Canada, pregnancy terminations for congenital anomalies at 20-23 weeks gestation explain the increasing stillbirth rate.⁵ In the USA, spontaneous preterm birth at early gestations made a significant contribution to higher stillbirth rates in black non-Hispanic women.¹⁷ Thus, in HIC, stillbirths at <28 weeks comprise an important component of all adverse pregnancy outcomes, particularly among some racial/ethnic groups.

1.11.2 Perceptions of stillbirth – Are they holding back progress?

Fatalism and stigma around stillbirth persist in HIC, both across communities and in the healthcare workforce (as reported in the LSS).¹⁸ In the International Stillbirth Alliance (ISA) surveys (see Panel 1

for methods), 2 in 3 respondents felt their community believed that most stillbirths are not preventable (Figure 2). Around 1 in 2 parents felt their community believed that “*parents should not talk about their stillborn baby because it makes people feel uncomfortable*”. One parent said “*...many women told me that my son's death was likely ‘nature taking care of mistakes’*”. Perceptions and actions that denigrate grief, dismiss the significance of a stillborn child, or support notions that a child was “never supposed to live”, are harmful to bereaved parents and devalue efforts towards prevention.¹⁸ Parent organisations provide powerful mechanisms to challenge stigma and fatalism around stillbirth (“Parents bringing about change”, webappendix).

1.12 Factors leading to stillbirth

1.12.1 Risk factors - Are we sufficiently aware?

Risk factors including demographic and lifestyle factors and medical/pregnancy conditions were reported in the LSS⁷. In the ISA survey of care providers, we asked respondents to select 10 out of 23 risk factors and associated conditions they believed posed the highest risk. We present the survey rankings alongside the adjusted odds ratio (aOR) as reported in the LSS¹⁹ and recent systematic reviews (Figure 1, webappendix).²⁰⁻²² Perceptions were generally consistent with the evidence, but care providers underestimated of the risks of advanced maternal age (>35 years), IVF, and multiple gestations, and overestimated of the risk of preeclampsia, smoking and substance misuse.

Risks associated with maternal obesity were also underestimated. In the survey of community members, 72% (n=1113) rated community awareness of overweight and obesity as very low to moderate (Figure 2, webappendix). With increasing prevalence of overweight and obesity, interventions to increase the number of women beginning pregnancy with a normal body weight are critically important to improving pregnancy outcomes and longer-term health. Modelling of a large Canadian cohort indicates a 10% decrease in pregnancy BMI could decrease stillbirth risk by 10%.²³ However, targeting behavioural change alone fails to recognise the complexity of this problem, resulting in ineffective interventions and added stigma for women who are overweight.²⁴ Antenatal

care that “problematizes” women by focusing on weight rather than a healthy pregnancy can produce feelings of embarrassment, guilt and shame,^{25, 26} leading women to avoid or delay care.

1.12.2 Classification of causes – Where do we need to focus?

Despite the call in the LSS for a uniform approach to the definition and classification of stillbirths,²⁷ a recent systematic review of causes of stillbirth globally²⁸ showed continued use of disparate approaches across HIC (Table 4, webappendix), rendering interpretation difficult. Despite this, placental pathologies were clearly important, accounting for around 50% of stillbirths in systems designed to capture them. Wide variation was shown in capture and definition of these pathologies, consistent with a recent review.²⁹ The contribution of other important factors varied widely; congenital abnormalities ranged from 6-33%; infection 2-22%; and spontaneous preterm birth/preterm ruptured membranes (PROM) 1-14%. In one high quality study, spontaneous preterm birth/PROM was a key factor in 41% of stillbirths <28 weeks.¹⁷

Studies using hierarchical approaches showed higher proportions of FGR³⁰ and congenital abnormalities,^{11, 14} depending on the system used. The categories “Other unspecified” and “Unexplained” showed the widest variation and highest proportions; up to 76% and 53% respectively.

1.12.3 Diagnostic tests for finding the cause of stillbirth

The evidence for many routinely performed stillbirth investigations is limited. While ongoing studies in the Netherlands and Australia will help to address this, the value of placental histopathology, autopsy, and genetic analysis is clear.^{31, 32} Nonetheless, in the ISA parent survey (n=3503), almost one quarter reported not being counselled or given information about autopsy. Failure to offer autopsy denies parents a chance to understand the cause of their baby’s death, increases the proportion of unexplained stillbirths, and hinders the effectiveness of subsequent audit. A critical shortage of perinatal pathologists also hampers efforts.^{33, 34} Such a shortage was shown in our surveys, where only 26% of care providers reported that autopsies were performed or supervised by perinatal or paediatric pathologists. Resources continue to be diverted away from perinatal

pathology,^{33, 35} despite stillbirths and neonatal deaths outnumbering deaths from cancer.^{9, 36} In our survey of care providers, only 33% reported that autopsy was routinely performed upon consent (Figure 3, webappendix). Parental consent and cost were frequently cited barriers to investigations (Figure 4, webappendix), despite evidence that identifying the cause of stillbirth may reduce costs in subsequent pregnancies. As stated by Heazell and colleagues,⁶ the cost of care for subsequent pregnancies following stillbirth with an assigned cause is less than for women whose stillbirths were of unknown cause.³⁷

Changes in diagnostic testing may lead to revisions of causes of stillbirths. Chromosomal microarray is now preferred to karyotype since it overcomes the problem of non-viable tissue. Microarrays also identify abnormalities, such as microdeletions and microduplications, that are not identified by karyotyping.³⁸ However, adoption of diagnostic advances is slow, with 30% of care providers from the ISA survey unsure how frequently microarray was performed; only 4% indicated that microarray was routine. Whichever test is used, it is important to have a perinatal pathology service to determine phenotype to assess the significance of newly described genetic variations.^{39, 40}

1.12.4 Addressing data quality in causes of stillbirth

Classification of cause of death in stillbirth needs to be standardised, especially with regard to placental pathology. Collective agreement of definitions of placental lesions and their significance is also needed. Although the same lesions may be seen in stillbirths and in livebirths, high quality studies suggest that specific placental lesions are significantly more common in cases of stillbirth.⁴¹ This is also true for other “causes” of stillbirth. In a recent review, not a single classification systems met the criteria of a quality system.⁴² Development of the WHO’s International Classification of Diseases Perinatal Mortality (ICD-PM), presented in a commentary to this series,⁴³ aims to address this. While HIC are likely to continue with detailed classification based on sophisticated diagnostics not accessible in LMIC, approaches must be consistent with the ICD-PM system, and HIC must reach consensus on such a system. Underpinning accurate data on causes is the availability of thorough

history and diagnostic testing. Standardised perinatal death datasets are essential. Such datasets are in place nationally in The UK, Ireland and New Zealand, and under pilot-testing in Australia.

1.13 Understanding and tackling disparities in stillbirth risk

Disparities reflect larger systems of structural inequality, including racism and systematic inequities in opportunities and power.⁴⁴ Consequently, health disparities reflect social and political determinants rather than biological origin.^{45, 46} Within HIC, stillbirth rates for disadvantaged groups are often double those of more advantaged groups, and are likely to be underestimated, as health disparities are often only measured by comparing the most advantaged with the least advantaged.⁴⁷ The relationship between stillbirth and social disadvantage is complex, with probable links across preconception, pregnancy pathways and risk factors (Figure 3).

1.13.1 Access to, and quality of, antenatal and maternity care

Disadvantaged women are less likely to receive adequate antenatal care.⁴⁸ Access to, and quality of antenatal care also differs by populations served,⁴⁹ and amongst ethnicities.⁵⁰ There are clear and specific circumstances where differential access or uptake of services contributes to disparities. These include antenatal diagnosis and pregnancy termination for congenital anomalies; timely diagnosis and treatment of preeclampsia; and labour induction for post-term pregnancy. Rural-urban differences in access to services are also likely to contribute, especially among vulnerable populations in remote areas.^{51, 52} Institutionalised racism is commonly reported by women accessing antenatal care.⁵³

1.13.2 Beyond health care delivery

Health disparities are only partly explained by disparities in maternity care.⁴⁷ Complex social determinants, termed “causes of the causes”, include: poverty; experiences of discrimination; incarceration; addiction; chronic stress; inadequate education, child care, employment, transportation, and living conditions.⁵⁴ Intimate partner violence,⁵⁵ mental health issues,⁵⁶ and the cumulative impact of stressful “life events”⁵⁷ are also hidden but potent risks. In a systematic review

of nearly 1,000,000 births in The UK from 1993-2005, stillbirth rates were 1.5-2 fold greater between the highest and lowest area deprivation quintiles.^{58, 59}

In a recent study, women migrating to HIC had stillbirth rates double host country averages, particularly when their country of birth was a humanitarian source country.⁶⁰ Adjusted for age, parity, socioeconomic status and BMI, these differences disappeared,⁶⁰ suggesting ethnicity may not always be a pathway to increased stillbirth. However, disparities in pregnancy outcomes continue between women of differing racial or ethnic background accessing the same healthcare services.⁶¹⁻⁶³ Stillbirth rates for women of South Asian and African origin giving birth in Europe or Australia are 2-3 times higher than those of Caucasian women.⁶¹⁻⁶³ The challenge is to understand why, and how we can manage the excess risk of stillbirth correlated with ethnicity within routine and comprehensive antenatal care.

1.13.3 Addressing disparity

To understand and address socioeconomic disparities in stillbirth, it is critical that all HIC monitor and report SES in vital statistics.¹² Maternal education is one relevant and feasible indicator for within-country and cross-country comparisons.¹² Across 19 European countries with a median population attributable risk of 26% (IQR 16 to 31), Zeitlin and colleagues showed 1,606 out of 6,447 stillbirths would not have occurred in 2010 if rates for all women were the same as for women with post-secondary education. School completion for pregnant women could therefore have a substantial impact on reducing disparities. Structural issues such as housing, employment and food security policies must also be addressed.

Antenatal care, home visiting services and financing of contraceptive services are examples of interventions with capacity to address reproductive health strategies⁶⁴ and therefore to prevent stillbirths. Universal service platforms should be supplemented with efforts to engage vulnerable populations, including outreach strategies and transportation to health services. In 2011 we nominated quality, accessible, culturally responsive and appropriate preconception care among priorities to reduce disparity.⁷ Improving preconception care remains an enormous challenge.

Innovative community programs addressing refugee maternal and child health inequalities⁶⁵ and antenatal care programs involving partnerships between midwives and Aboriginal health workers are underway in Australia.^{66, 67} Universally, women whose first language differs from dominant national languages should be offered care and information in their own language,⁶⁸ with choice of interpreter gender.⁶⁵ The US Affordable Care Act will provide services such as free contraception, screening for infections and alcohol and smoking cessation programs.⁶⁹

1.14 Quality of care

1.14.1 Uptake of interventions in stillbirth prevention

Through the ISA survey of care providers we assessed uptake and perceived barriers to implementation of LSS recommended interventions in stillbirth prevention (Figures 5 and 6, webappendix). Only 60% of respondents said they always provided *smoking cessation advice*. The most frequently cited barriers were lack of time and/or resources (20%) and acceptance to women (35%). Only 36% said they always provided *culturally-appropriate care*, with lack of time and/or resources the most common barrier (17%). *Serial fundal height measurements* were performed not at all or only sometimes by 14%, with 10% identifying lack of evidence as a barrier. Most providers said they always or mostly used *early ultrasound assessment of gestational age* (83%), with cost and acceptance to women cited as barriers by around 12%. *Screening for gestational diabetes at 28 weeks* was always or mostly performed by 77%, with 12% citing acceptance by women as a barrier. Use of *Doppler velocimetry in high-risk pregnancies* had reasonably high usage (68%), with lack of evidence (11%) and cost (9%) the most frequently cited barriers. The least commonly used intervention (used always or mostly by 43%) was *low dose aspirin for high-risk pregnancies*, with 13% reporting lack of evidence as a barrier. Low dose aspirin and heparins have been used to improve placental function and decrease stillbirth, but a high number needed to treat (aspirin) and uncertain efficacy has prevented widespread adoption of these interventions.⁷⁰ While evidence is limited, preconception care is a potentially valuable intervention in stillbirth prevention, yet only 28% of care

providers said that *preconception care for women with risk factors* was performed mostly/always in their facilities.

1.14.2 Antenatal and bereavement care - Information, communication and support

In addition to clinical care, quality maternity care incorporates interpersonal and emotional aspects of care.⁷¹ Since stillbirth is an indicator of quality care, women's experiences of care around stillbirth can be considered an indicator of quality of care processes. Just as actions can be taken to prevent stillbirth, actions can be taken to prevent adverse psychosocial outcomes following stillbirth and, in both instances, suboptimal interpersonal care can undermine even the best clinical care and produce harm (see Heazell and colleagues).⁶ The ISA surveys asked parents and care providers eight questions designed to capture components of quality care consistently identified as important to women (webappendix). The data (Figure 7, webappendix) show that care providers viewed various aspects of care more positively than bereaved parents. At least 4 in 5 providers (83-95%), but only 3 in 5 parents (54-70%), considered these aspects of quality care to be present always or most of the time. Not spending enough time with parents in antepartum care was a point of agreement for parents and care providers. Critically, more than one-third of parents believed their concerns were not taken seriously or felt not listened to, either before or after their baby was stillborn.

Ratings of information-provision and parental involvement in decision-making after stillbirth were lower for parents and care providers alike compared with before stillbirth, underscoring the challenge of providing quality bereavement care. Parents' views of the comprehensibility of information were also less positive; barely half felt the time spent with care providers was adequate. Many of the questions parents had at this time could be readily answered and procedures to answer such questions would be easy to implement (Panel 1, webappendix). Missed opportunities to answer parents' questions might be avoided by measures that recognise parents' need to know more about their child.

These survey findings corroborate the results of a systematic review on parents' and care providers' experiences of bereavement care.⁷² Care providers were found to "hide" behind ritualising guidelines

and checklists; they were frequently not trained to expect and manage parents' reactions and individual needs. In our survey of care providers, only 23% reported being satisfied with training opportunities in bereavement care at their facility, and 30% had no opportunities. As found in the meta-analysis, care providers urgently need emotional, knowledge and system-based support, and training in verbal and non-verbal communication skills.

1.14.3 Addressing quality of care

Stillbirth prevention requires emphasis on quality maternity care that is respectful of a woman's rights and tailored to her needs.⁷¹ Quality can be improved through better communication and information-provision, and timely delivery of evidence-based interventions. Quality bereavement care must also be emphasised, with greater access to training a crucial first step. Maternity units must decide whether this is best accomplished via training and certification of competencies for all staff, or whether to assign the role of bereavement support to a dedicated group. Access to clinical practice guidelines is imperative, but active implementation and evaluation are required.⁷³ The UK provides an extensive range of national clinical guidelines acknowledging every aspect of the key messages of the LSS HIC paper but, for other HIC with a high stillbirth burden, such resources are far less comprehensive (webappendix). Publically available reports of maternal satisfaction with care and other indicators of women's maternity care experiences should be developed, as done in Queensland, Australia,⁷⁴ New Zealand⁷⁵ and across The UK.⁷⁶ Audit and feedback and benchmarking programs that include explicit targets for change and suggestions for how change can be achieved are also effective.⁷⁷

1.14.4 Perinatal mortality audit - Why and how?

Perinatal audit has been described as: "The systematic, critical analysis of the quality of perinatal care, including the procedures used for diagnosis and treatment, the use of resources and the resultant outcome and quality of life for women and their babies".⁷⁸ Audits in the Netherlands,⁷⁹ The UK⁸⁰ and New Zealand¹¹ show substandard care factors are present in an unacceptably high proportion of cases (20-30%, and up to 60% for intrapartum stillbirths). In New Zealand, stillbirth

rates at term have declined over the seven years since national perinatal audit began¹¹ (Figure 8, webappendix). This decline was specifically attributed to a reduction in stillbirth at 37-40 weeks' gestation and ≥ 41 weeks' gestation. Despite their value, few countries have implemented national-level perinatal audit programs (webappendix). Norway, which originally introduced it in 1984, has now abandoned this practice. Among the care providers we surveyed, only 37% reported that their facility conducts regular perinatal audit meetings; these were most commonly held only monthly (34%) or quarterly (26%), and used case discussion only as opposed to formal audit methodology (61% vs 12%).

There is a clear need for greater focus on effective, sustainable implementation of perinatal audit to ensure health services identify areas of suboptimal care. Establishing perinatal mortality audit requires both service- and ministerial-level support, and quarantined time for multidisciplinary team engagement. Other critical components include an agreed set of definitions, adoption of a formal audit methodology, and appropriate indicators for monitoring and evaluation, and effective data systems (Panel 2). Innovative e- and m-health solutions currently being implemented hold promise,⁸¹ as well as structured education programs around institutional perinatal mortality audit and classification such as the IMPROVE program.⁸²

1.15 Antenatal screening and interventions to prevent stillbirth – Are we getting closer?

1.15.1 Early delivery

Routine induction of labour (IOL) at term and post-term reduces the risk of perinatal mortality and caesarean birth.⁸³ However, birth prior to 39 weeks increases the risk of morbidity⁸⁴ and is associated with increased risk of long-term mortality.⁸⁵ Therefore, prior to 39 weeks, early delivery requires balancing any reduction in stillbirth risk against these risks, and should only be considered in the presence of significant risk for maternal and neonatal complications. When IOL is undertaken, adequate information-provision to women is essential, as women have reported not being aware of

the risks of induction or the implications for future pregnancies until after the induction had been performed.⁸⁶

The ARRIVE trial⁸⁷ in the US comparing elective IOL at 39 weeks with expectant management among singleton uncomplicated term pregnancies may help to clarify the risks and benefits of term induction.

1.15.2 Ultrasonic and biochemical prediction of stillbirth risk

Multiple pathophysiological processes result in stillbirth, making it difficult to predict.⁸⁸ A test is likely to perform poorly when assessed against all stillbirths, but possibly more specific for a given cause of stillbirth. A systematic review of biomarker and ultrasonic tests found none of 16 single, or 5 combined, tests performed well as predictors of stillbirth.⁸⁹ However, stillbirth attributed to placental dysfunctional disorders was moderately to strongly associated (positive likelihood ratios between 5 and 15) with low first trimester pregnancy associated plasma protein A (PAPP-A), and abnormal uterine artery Doppler velocimetry in the second trimester. More studies are needed to determine whether closer vigilance or any treatment is effective to prevent stillbirth in this increased-risk group.

1.15.3 Routine late pregnancy ultrasound to screen for fetal growth restriction

Of the estimated 30-50% of stillbirths related to FGR; most are undetected and many occur in women lacking risk factors.⁹⁰ Ultrasonic fetal biometry is widely used in high risk pregnancies as a means of detecting FGR, thus universal ultrasound is one potential approach to screening low-risk women.⁹⁰ However, high-quality evidence on the diagnostic effectiveness of ultrasound is lacking.⁹¹ A recent prospective cohort study found universal scanning was associated with approximately 3-fold increase in the detection of small for gestational age (SGA) (20-57%).⁹² Further, SGA fetuses with reduced abdominal circumference growth velocity were at increased risk of morbidity, whereas SGA fetuses with normal growth velocity were not. This study confirms universal ultrasound is effective in identifying FGR. However, the costs and potential adverse iatrogenic consequences of implementing such an intervention require consideration.⁹³

1.15.4 Challenges in gaining high quality evidence for screening

Sample size calculations demonstrate that even if a screening test has a positive likelihood ratio of 10 and was coupled with an intervention that reduced stillbirth by 50%, a study of screening and intervention would still require approximately 130,000 women to be adequately powered (Figure 9, webappendix). Possible approaches to address this problem are the inclusion of stillbirth as part of a composite outcome and the use of study designs with randomisation at the hospital level, including cluster randomised controlled trials (RCTs) or stepped wedge RCTs,⁸⁸ such as the trials of fetal movement awareness interventions mentioned below.^{94, 95}

1.15.5 Promising antenatal interventions?

Raising awareness of decreased fetal movements (DFM) may aid stillbirth prevention via timely detection and reporting, though concerns exist over the potential to increase anxiety and health service utilisation.⁹⁶ Two large-scale trials of fetal movement awareness interventions are ongoing in Australasia,⁹⁵ and Ireland and The UK.⁹⁴ In a large non-randomised study, an educational program of standardised measurement of fundal height, plotting on customised charts and referral protocols, has also been associated with reduced stillbirth.⁹⁷ Data from RCTs are required to confirm or refute these findings.⁹⁸ The adverse impact of supine sleep position in late pregnancy has recently been highlighted as a potentially modifiable risk factor.⁹⁹⁻¹⁰² Although these findings are biologically plausible, results from further studies in The UK¹⁰³ and New Zealand¹⁰⁰ study are awaited.

1.16 The research agenda

The LSS identified 30 questions derived from opinions of professionals and researchers. Research priority setting methods have since developed to include patient and public views. As part of the ISA surveys, over 7,000 parents, care providers and community members provided stillbirth action and research priorities. While the ISA project is ongoing, preliminary data indicate agreement with the LSS and a recent UK project.¹⁰⁴ Major topics included: stillbirth prevention by application of current tests and development of novel investigations with optimal timing of delivery, understanding

placental pathways in stillbirth and the causes of unexplained stillbirth, optimal bereavement care, and subsequent pregnancy care.

Perinatal mortality audit programs, interventions to reduce the prevalence of overweight and obesity, and initiatives to increase the coverage of smoking cessation programs in pregnancy are also priorities. With static rates of stillbirths <28 weeks across HIC, and with spontaneous preterm labour and/or preterm ROM a major contributor, ongoing efforts in prediction and prevention of preterm birth are important in stillbirth prevention. Strengthening collaborations between researchers and parents to address priorities using similar protocols is key in addressing stillbirths in HIC.

1.17 Conclusions

Stillbirth remains a major public health problem in HIC and reductions in rates have not matched those for neonatal mortality. Variation and socio-economic disparities in stillbirth rates, suboptimal uptake of interventions, low proportions of stillbirths attributed to congenital abnormality and high proportions classified as unexplained, and the contribution of substandard care factors suggest stillbirths are not inevitable, and that further reduction in HIC is possible. Ending preventable stillbirths in HIC can be achieved through improvements in the health status of women, through improvements in quality of maternity care, and by reducing social inequities. High quality perinatal mortality audit informed by thorough investigation is attainable in all HIC and holds the key to (relatively) rapid reductions in stillbirth rates.

The death of a child before birth is a tragedy for families, and stigma and fatalism must be eliminated to optimise bereavement care and to reduce the number of these deaths.

1.18 Contributions

V Flenady was responsible for overall development and writing of the manuscript; A Wojcieszek assisted with development and writing of the manuscript and coordination of author contributions; D Ellwood and P Middleton assisted in editing; V Flenady, A Wojcieszek, D Ellwood, F Boyle, D Horey, A Vanacci, C Ravaldi, J Cassidy, P Cassidy, J F Frøen, L Farrales, M Gross, P Middleton, J J Erwich, G Smith, T Y Khong, R M Silver, S Leisher, L McCowan, D Siassakos, A Heazell and S Petersen contributed to the development and dissemination of the web-based surveys; A Wojcieszek coordinated the dissemination of surveys and S Leisher and A Wojcieszek coordinated the survey translations; A Wojcieszek lead the quantitative survey data analyses with V Flenady, M Coory; F Boyle, D Horey, undertook qualitative survey data analysis; J E Lawn and H Blencowe conducted data analyses for stillbirth rates and risk factors; T Y Khong, R M Silver, S Leisher, J J Erwich developed the diagnostic test section; D Ellwood developed the risk factor and interventions sections; V Flenady, M Coory, and H Reinebrant undertook the causes of stillbirth analysis; J J Erwich, C Farquar, L Sadler developed the perinatal mortality audit section; P Middleton, L Farrales, S Brown, K S Joseph, J Zeitlin, E Wallace developed the disparity section; D Siassakos, A Heazell, C Storey contributed a meta-synthesis on experiences of bereavement care; G Smith and L McCowan assisted to develop the antenatal screening and interventions to prevent stillbirth section; F Bloomfield, G Smith, J F Frøen, and R Goldenberg contributed to framing and drafting of the manuscript. All authors read and approved the final version of the manuscript.

1.19 Conflicts of interest

HB received grants from Saving Newborn Lives, Save the Children, during the conduct of the study. SB received salary support via a Future Fellowship from the Australian Research Council, during the conduct of the study. LS received personal fees from Health Quality and Safety Commission NZ, during the conduct of the study. DS received grants from Sands, outside the submitted work and is a member of the Executive Committee of the Stillbirth Clinical Study Group, UK (RCOG & Sands), the Department of Health Stillbirth task-and-finish groups, and the PROMPT Maternity Foundation. GS received non-financial support from GE, personal fees and non-financial support from Roche, grants and personal fees from GSK, outside the submitted work. All remaining authors have nothing to disclose.

1.20 Acknowledgments

We sincerely thank every parent, care provider and community member who gave their time to complete the ISA web-based surveys. We thank the International Stillbirth Alliance (ISA) Scientific Network, the International Confederation of Midwives (ICM), the International Federation of

35 Gynecology and Obstetrics (FIGO), ISA member organisations, and further national organisations that
36 supported the dissemination of surveys. Finally, we thank Translators Without Borders for assisting
37 with survey translations.

38

39 **1.21 Funding**

40 Mater Research Institute – The University of Queensland provided infrastructure and funding for the
41 research team to enable this work to be undertaken. The Canadian Research Chair in Psychosocial
42 Family Health provided funding for revision of the translation of the French web-based survey of care
43 providers.

1.22 References

- 45 1. de Bernis L, Kinney M, Stones W, et al. Stillbirths: Ending preventable deaths by 2030. *The*
46 *Lancet (Submitted) 2015.*
- 47 2. Frøen JF, Friberg IK, Lawn JE, et al. Stillbirths: Progress and unfinished business. *The Lancet*
48 *(Submitted) 2015.*
- 49 3. Bradley NM, Lucy KS, Alun Evans T, et al. Perinatal mortality surveillance report UK - Perinatal
50 deaths for births from January to December 2013. Leicester: Department of Health Sciences
51 University of Leicester, 2015.
- 52 4. Gregory EC, MacDorman MF. Fetal and Perinatal Mortality: United States, 2013. *Natl Vital*
53 *Stat Rep 2015; 64(8): 1-24.*
- 54 5. Joseph KS, Kinniburgh B, Hutcheon JA, et al. Determinants of increases in stillbirth rates from
55 2000 to 2010. *CMAJ 2013; 185(8): E345-51.*
- 56 6. Heazell AE, Siassakos D, Blencowe H, et al. Stillbirths: Why Invest? *The Lancet (In Press) 2015.*
- 57 7. Flenady V, Middleton P, Smith GC, et al. Stillbirths: the way forward in high-income countries.
58 *Lancet 2011; 377: 1703-17.*
- 59 8. Blencowe H, Lawn J. National, regional, and worldwide estimates of stillbirth rates in 2015
60 with trends since 1995: a systematic analysis. *Lancet Global Health (Submitted) 2015.*
- 61 9. Lawn JE, Blencowe H, Waiswa P, et al. Stillbirths: Data for accelerating progress towards
62 2030. *The Lancet (In Press) 2015.*
- 63 10. Ibiebele I, Coory M, Boyle FM, et al. Stillbirth rates among indigenous and non-indigenous
64 women in Queensland, Australia: is the gap closing? *BJOG: An International Journal of Obstetrics &*
65 *Gynaecology 2014: n/a-n/a.*
- 66 11. PMMRC. Ninth annual report of the Perinatal and Maternal Mortality Review Committee:
67 Reporting mortality 2013. Wellington: Health Quality & Safety Commission, 2015.
- 68 12. Zeitlin J, Mortensen L, Prunet C, et al. Socioeconomic inequalities in stillbirth rates in Europe:
69 measuring the gap using routine data from the Euro-Peristat Project. *BMC Pregnancy Childbirth*
70 *(Submitted to Ending Preventable Stillbirths Supplement) 2015.*
- 71 13. Mohangoo AD, Buitendijk SE, Szamotulska K, et al. Gestational age patterns of fetal and
72 neonatal mortality in Europe: results from the Euro-Peristat project. *PLoS One 2011; 6(11): e24727.*
- 73 14. Hilder L, Li Z, Zeki R, et al. Stillbirths in Australia 1991-2009. Perinatal statistics. Series no. 29.
74 Cat. no. PER 63. Canberra: AIHW, 2014.
- 75 15. Joseph KS, Liu S, Rouleau J, et al. Influence of definition based versus pragmatic birth
76 registration on international comparisons of perinatal and infant mortality: population based
77 retrospective study. *BMJ 2012; 344: e746.*
- 78 16. MacDorman M, Kirmeyer S. Fetal and perinatal mortality, United States, 2005, *Natl Vital Stat*
79 *Rep Natl Vital Stat Rep 2009; 57(8): 1-20.*
- 80 17. Bukowski R, Carpenter M, Conway D, et al. Causes of death among stillbirths. *Jama 2011;*
81 *306(22): 2459-68.*
- 82 18. Froen JF, Cacciatore J, McClure EM, et al. Stillbirths: why they matter. *Lancet 2011;*
83 *377(9774): 1353-66.*
- 84 19. Flenady V, Koopmans L, Middleton P, et al. Major risk factors for stillbirth in high-income
85 countries: a systematic review and meta-analysis. *The Lancet 2011; 377(9774): 1331-40.*
- 86 20. Marufu TC, Ahankari A, Coleman T, et al. Maternal smoking and the risk of still birth:
87 systematic review and meta-analysis. *BMC Public Health 2015; 15: 239.*
- 88 21. Mondal D, Galloway TS, Bailey TC, et al. Elevated risk of stillbirth in males: systematic review
89 and meta-analysis of more than 30 million births. *BMC Med 2014; 12: 220.*
- 90 22. Ovesen PG, Jensen DM, Damm P, et al. Maternal and neonatal outcomes in pregnancies
91 complicated by gestational diabetes. a nation-wide study. *J Matern Fetal Neonatal Medicine 2015: 1-*
92 *5.*

- 93 23. Schummers L, Hutcheon JA, Bodnar LM, et al. Risk of adverse pregnancy outcomes by
94 prepregnancy body mass index: a population-based study to inform prepregnancy weight loss
95 counseling. *Obstet Gynecol* 2015; **125**(1): 133-43.
- 96 24. Ramos Salas X. The ineffectiveness and unintended consequences of the public health war on
97 obesity. *Can J Public Health* 2015; **106**(2): e79-81.
- 98 25. Mulherin K, Miller YD, Barlow FK, et al. Weight stigma in maternity care: women's
99 experiences and care providers' attitudes. *BMC Pregnancy Childbirth* 2013; **13**: 19.
- 100 26. Smith D, Lavender T. The maternity experience for women with a body mass
101 index ≥ 30 kg/m²: a meta-synthesis. *BJOG: An International Journal of Obstetrics & Gynaecology*
102 2011; **118**(7): 779-89.
- 103 27. Froen JF, Pinar H, Flenady V, et al. Causes of death and associated conditions (Codac) - a
104 utilitarian approach to the classification of perinatal deaths. *BMC Pregnancy and Childbirth* 2009;
105 **9**(1): 22.
- 106 28. Reinebrant HE, Teoh Z, Leisher SH, et al. Global causes of stillbirth: A systematic review. *BMC*
107 *Pregnancy Childbirth (Submitted to Ending Preventable Stillbirths Supplement)* 2015.
- 108 29. Ptacek I, Sebire NJ, Man JA, et al. Systematic review of placental pathology reported in
109 association with stillbirth. *Placenta* 2014; **35**(8): 552-62.
- 110 30. Ego A, Zeitlin J, Batailler P, et al. Stillbirth classification in population-based data and role of
111 fetal growth restriction: the example of RECODE. *BMC Pregnancy Childbirth* 2013; **13**: 182.
- 112 31. Korteweg FJ, Erwich JJ, Timmer A, et al. Evaluation of 1025 fetal deaths: proposed diagnostic
113 workup. *Am J Obstet Gynecol* 2012; **206**(1): 53 e1- e12.
- 114 32. Korteweg FJ, Bouman K, Erwich JJ, et al. Cytogenetic analysis after evaluation of 750 fetal
115 deaths: proposal for diagnostic workup. *Obstet Gynecol* 2008; **111**(4): 865-74.
- 116 33. Colgan TJ, Geldenhuys L. The Practice of Pathology in Canada: Decreasing Pathologist Supply
117 and Uncertain Outcomes. *Arch Pathol Lab Med* 2012; **136**: 90-4.
- 118 34. Robboy SJ, Weintraub S, Horvath AE, et al. Pathologist workforce in the United States:
119 Development of a predictive model to examine factors influencing supply. *Archives of Pathology &*
120 *Laboratory Medicine* 2013; **137**(12): 1723-32.
- 121 35. PMMRC. Perinatal and Paediatric Pathology Service Provision in New Zealand 2008: Perinatal
122 and Maternal Mortality Review Committee, 2008.
- 123 36. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide:
124 Sources, methods and major patterns in GLOBOCAN 2012. *International Journal of Cancer* 2015;
125 **136**(5): E359-E86.
- 126 37. Mistry H, Heazell AE, Vincent O, et al. A structured review and exploration of the healthcare
127 costs associated with stillbirth and a subsequent pregnancy in England and Wales. *BMC pregnancy*
128 *and childbirth* 2013; **13**: 236.
- 129 38. Reddy UM, Page GP, Saade GR, et al. Karyotype versus Microarray Testing for Genetic
130 Abnormalities after Stillbirth. *New England Journal of Medicine* 2012; **367**(23): 2185-93.
- 131 39. Kooper A, Faas B, Feenstra I, et al. Best diagnostic approach for the genetic evaluation of
132 fetuses after intrauterine death in first, second or third trimester: QF-PCR, karyotyping and/or
133 genome wide SNP array analysis. *Molecular Cytogenetics* 2014; **7**(1): 6.
- 134 40. Vrijenhoek T, Kraaijeveld K, Elferink M, et al. Next-generation sequencing-based genome
135 diagnostics across clinical genetics centers: implementation choices and their effects. *Eur J Hum*
136 *Genet* 2015.
- 137 41. Pinar H, Goldenberg RL, Koch MA, et al. Placental findings in singleton stillbirths. *Obstetrics*
138 *and gynecology* 2014; **123**(2 Pt 1): 325-36.
- 139 42. Leisher SH, Teoh Z, Reinebrant HE, et al. Classification systems for causes of stillbirth and
140 neonatal death, 2009-2014: An assessment of alignment with characteristics for an effective global
141 system. *BMC Pregnancy Childbirth (Submitted to Ending Preventable Stillbirths Supplement)* 2015.
- 142 43. Allanson E, Tunçalp Ö, Chou D, et al. Perinatal mortality: WHO Application of ICD 10 to
143 perinatal deaths. *The Lancet (Submitted)* 2015.

- 144 44. Dehlendorf C, Harris LH, Weitz TA. Disparities in abortion rates: a public health approach. *Am J Public Health* 2013; **103**(10): 1772-9.
- 145
- 146 45. Bryant AS, Worjolah A, Caughey AB, et al. Racial/ethnic disparities in obstetric outcomes and care: prevalence and determinants. *Am J Obstet Gynecol* 2010; **202**(4): 335-43.
- 147
- 148 46. Dehlendorf C, Bryant AS, Huddleston HG, et al. Health disparities: definitions and measurements. *Am J Obstet Gynecol* 2010; **202**(3): 212-3.
- 149
- 150 47. Braveman P. Health disparities and health equity: concepts and measurement. *Annu Rev Public Health* 2006; **27**: 167-94.
- 151
- 152 48. Cha S, Masho SW. Intimate partner violence and utilization of prenatal care in the United States. *Journal of Interpersonal Violence* 2013.
- 153
- 154 49. Creanga AA, Bateman BT, Mhyre JM, et al. Performance of racial and ethnic minority-serving hospitals on delivery-related indicators. *Am J Obstet Gynecol* 2014; **211**(6): 647 e1-16.
- 155
- 156 50. Brett KM, Schoendorf KC, Kiely JL. Differences between black and white women in the use of prenatal care technologies. *American Journal of Obstetrics and Gynecology* 1994; **170**(1): 41-6.
- 157
- 158 51. Gilbert NL, Auger N, Tjepkema M. Stillbirth and infant mortality in Aboriginal communities in Quebec. *Health Rep* 2015; **26**(2): 3-8.
- 159
- 160 52. Luo ZC, Wilkins R. Degree of rural isolation and birth outcomes. *Paediatr Perinat Epidemiol* 2008; **22**(4): 341-9.
- 161
- 162 53. Salm Ward T, Mazul M, Ngui E, et al. "You learn to go last": Perceptions of prenatal care experiences among African-American women with limited incomes. *Maternal and Child Health Journal* 2013; **17**(10): 1753-9.
- 163
- 164 54. Marmot M. BMA presidency acceptance speech: fighting the alligators of health inequalities; 2010.
- 165
- 166 55. Mogos MF, Araya WN, Masho SW, et al. The fetomaternal health cost of intimate partner violence among delivery-related discharges in the United States, 2002-2009. *Journal of Interpersonal Violence* 2014.
- 167
- 168 56. King-Hele S, Webb RT, Mortensen PB, et al. Risk of stillbirth and neonatal death linked with maternal mental illness: a national cohort study. *Arch Dis Child Fetal Neonatal Ed* 2009; **94**(2): F105-10.
- 169
- 170 57. Hogue CJ, Parker CB, Willinger M, et al. A population-based case-control study of stillbirth: the relationship of significant life events to the racial disparity for African Americans. *Am J Epidemiol* 2013; **177**(8): 755-67.
- 171
- 172 58. Seaton SE, Field DJ, Draper ES, et al. Socioeconomic inequalities in the rate of stillbirths by cause: a population-based study. *BMJ Open* 2012; **2**(3).
- 173
- 174 59. Weightman AL, Morgan HE, Shepherd MA, et al. Social inequality and infant health in the UK: systematic review and meta-analyses. *BMJ Open* 2012; **2**(3).
- 175
- 176 60. Gibson-Helm ME, Teede HJ, Cheng IH, et al. Maternal health and pregnancy outcomes comparing migrant women born in humanitarian and nonhumanitarian source countries: a retrospective, observational study. *Birth* 2015; **42**(2): 116-24.
- 177
- 178 61. Khalil A, Rezende J, Akolekar R, et al. Maternal racial origin and adverse pregnancy outcome: a cohort study. *Ultrasound Obstet Gynecol* 2013; **41**(3): 278-85.
- 179
- 180 62. Ravelli AC, Tromp M, Eskes M, et al. Ethnic differences in stillbirth and early neonatal mortality in The Netherlands. *J Epidemiol Community Health* 2011; **65**(8): 696-701.
- 181
- 182 63. Drysdale H, Ranasinha S, Kendall A, et al. Ethnicity and the risk of late-pregnancy stillbirth. *Med J Aust* 2012; **197**(5): 278-81.
- 183
- 184 64. Mehta P. Addressing reproductive health disparities as a healthcare management priority: pursuing equity in the era of the Affordable Care Act. *Current Opinion in Obstetrics and Gynecology* 2014; **26**(6): 531-8.
- 185
- 186 65. Yelland J, Riggs E, Szwarc J, et al. Bridging the Gap: using an interrupted time series design to evaluate systems reform addressing refugee maternal and child health inequalities. *Implement Sci* 2015; **10**(1): 62.
- 187
- 188
- 189
- 190
- 191
- 192
- 193
- 194

- 195 66. Brown SJ, Weetra D, Glover K, et al. Improving Aboriginal women's experiences of antenatal
196 care: findings from the Aboriginal families study in South Australia. *Birth* 2015; **42**(1): 27-37.
- 197 67. Middleton PF, Brown SJ, Bubner TK, et al. The SA Aboriginal Family Birthing Program – a
198 success story. *PSANZ: Melbourne* April 2015.
- 199 68. Boerleider A, Wiegers T, Mannien J, et al. Factors affecting the use of prenatal care by non-
200 western women in industrialized western countries: a systematic review. *BMC Pregnancy and*
201 *Childbirth* 2013; **13**(1): 81.
- 202 69. Adams EK, Markowitz S, Dietz PM, et al. Expansion of Medicaid covered smoking cessation
203 services: maternal smoking and birth outcomes. *Medicare Medicaid Res Rev* 2013; **3**(3).
- 204 70. Duley L, Henderson-Smart DJ, Meher S, et al. Antiplatelet agents for preventing pre-
205 eclampsia and its complications. *Cochrane Database Syst Rev* 2007; (2): CD004659.
- 206 71. Renfrew MJ, McFadden A, Bastos MH, et al. Midwifery and quality care: findings from a new
207 evidence-informed framework for maternal and newborn care. *The Lancet* 2014; **384**(9948): 1129-45.
- 208 72. Ellis A, Chebsey C, Storey C, et al. Systematic review to understand and improve care after
209 stillbirth: a review of parents and healthcare professionals experiences. *BMC Pregnancy Childbirth*
210 *(Submitted to Ending Preventable Stillbirths Supplement)* 2015.
- 211 73. Michie S, van Stralen M, West R. The behaviour change wheel: A new method for
212 characterising and designing behaviour change interventions. *Implementation Science* 2011; **6**(1): 42.
- 213 74. Prosser SJ, Miller YD, Armanasco A, et al. Findings from the Having a Baby in Queensland
214 Survey, 2012. Brisbane: Queensland Centre for Mothers & Babies, The University of Queensland,
215 2013.
- 216 75. New Zealand Ministry of Health. Maternity Consumer Surveys 2011. Wellington: Ministry of
217 Health, 2012.
- 218 76. Redshaw M, Rowe R, Henderson J. Listening to Parents after stillbirth or the death of their
219 baby after birth. Oxford: National Perinatal Epidemiology Unit, 2014.
- 220 77. Ivers N, Jamtvedt G, Flottorp S, et al. Audit and feedback: effects on professional practice and
221 healthcare outcomes. *Cochrane Database of Systematic Reviews*, 2012.
222 <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD000259.pub3/abstract> (accessed.
- 223 78. Dunn PM, McIlwaine G. Perinatal audit: a report produced for the European Association of
224 Perinatal Medicine. New York, NY: Parthenon; 1996.
- 225 79. Eskes M, Waelput AJM, Erwich JJ, et al. Term perinatal mortality audit in the Netherlands
226 2010-2012: a population based cohort study. *BMJ Open* 2014; **4**: e005652.
- 227 80. West Midlands Perinatal Institute. Confidential enquiry into intrapartum related deaths: NHS
228 Perinatal Institute, 2010.
- 229 81. Frøen JF, Flenady V, Myhre S, et al. Electronic health registries for mothers and their children.
230 *Lancet Global Health (Invited submission in conjunction with Ending Preventable Stillbirth series)*
231 2015.
- 232 82. Gardiner P, Kent AL, Rodriguez V, et al. IMproving Perinatal Mortality Review and Outcomes
233 Via Education; An international educational program for health care professionals on best practice
234 around the time of a perinatal death. *BMC Pregnancy Childbirth (Submitted to Ending Preventable*
235 *Stillbirths Supplement)* 2015.
- 236 83. Gulmezoglu AM, Crowther CA, Middleton P, et al. Induction of labour for improving birth
237 outcomes for women at or beyond term. *Cochrane Database of Systematic Reviews* 2012; (6).
- 238 84. Spong CY, Mercer BM, D'Alton M, et al. Timing of indicated late-preterm and early-term
239 birth. *Obstetrics & Gynecology* 2011; **118**(2, Part 1): 323-33.
- 240 85. Crump C, Sundquist K, Winkleby MA, et al. Early-term birth (37-38 weeks) and mortality in
241 young adulthood. *Epidemiology* 2013; **24**(2): 270-6.
- 242 86. Moore JE, Low LK, Titler MG, et al. Moving toward patient-centered care: Women's decisions,
243 perceptions, and experiences of the induction of labor process. *Birth* 2014; **41**(2): 138-46.
- 244 87. Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD).
245 Induction in nulliparous women at 39 weeks to prevent adverse outcomes: a randomized controlled

- trial. In: ClinicalTrials.gov [Internet] [cited 28th April 2015]; Bethesda (MD): National Library of Medicine (US)]. Available from: <https://clinicaltrials.gov/ct2/show/NCT01990612>: NCT01990612.
88. Hussey MA, Hughes JP. Design and analysis of stepped wedge cluster randomized trials. *Contemporary Clinical Trials* 2007; **28**(2): 182-91.
89. Conde-Agudelo A, Bird S, Kennedy SH, et al. First- and second-trimester tests to predict stillbirth in unselected pregnant women: a systematic review and meta-analysis. *BJOG* 2015; **122**(1): 41-55.
90. Rodger MA, Carrier M, Le Gal G, et al. Meta-analysis of low-molecular-weight heparin to prevent recurrent placenta-mediated pregnancy complications. *Blood* 2014; **123**(6): 822-8.
91. National Collaborating Centre for Women's and Children's Health. NICE Guideline: Antenatal care. London: RCOG Press; 2008.
92. Sovio U, Smith G, Dacey A, et al. 151: Screening for fetal growth restriction (FGR) using universal third trimester ultrasonography: a prospective cohort study of 3,977 nulliparous women. *American Journal of Obstetrics & Gynecology*; **212**(1): S92.
93. Monier I, Blondel B, Ego A, et al. Poor effectiveness of antenatal detection of fetal growth restriction and consequences for obstetric management and neonatal outcomes: a French national study. *BJOG* 2015; **122**(4): 518-27.
94. Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD). Promoting awareness fetal movements to reduce fetal mortality stillbirth, a stepped-wedge cluster randomised trial. (AFFIRM). In: ClinicalTrials.gov [Internet] [cited 30th April 2015]; Available from: <https://www.clinicaltrials.gov/ct2/show/NCT01777022>: NCT01777022.
95. National Health and Medical Research Council of Australia. My Baby's Movements: a stepped wedge cluster randomised controlled trial to raise maternal awareness of fetal movements during pregnancy. In: Australian and New Zealand clinical trials registry [Internet] [cited 30th April 2015]; Available from: <https://www.anzctr.org.au/Trial/Registration/TrialReview.aspx?id=365867>: ACTRN12614000291684.
96. Mangesi L, Hofmeyr GJ, Smith V. Fetal movement counting for assessment of fetal wellbeing. *Cochrane Database of Systematic Reviews* 2007, Issue 1 Art No: CD004909 DOI: 101002/14651858.
97. Gardosi J, Giddings S, Clifford S, et al. Association between reduced stillbirth rates in England and regional uptake of accreditation training in customised fetal growth assessment. *BMJ Open* 2013; **3**: e003942.
98. Carberry Angela E, Gordon A, Bond Diana M, et al. Customised versus population-based growth charts as a screening tool for detecting small for gestational age infants in low-risk pregnant women. *Cochrane Database of Systematic Reviews* 2014, Issue 5 Art No: CD008549 DOI: 101002/14651858.
99. Gordon A, Raynes-Greenow C, Bond D, et al. Sleep Position, Fetal Growth Restriction, and Late-Pregnancy Stillbirth. *Obstetrics & Gynecology* 2015; **125**(2): 347-55.
100. McCowan LME, Thompson JMD, Cronin R, et al. Supine sleep position in late pregnancy is associated with increased risk of late stillbirth. The International Conference on Stillbirth, SIDS and Baby Survival 2014 Amsterdam, 18-21 September; 2014.
101. Owusu JT, Anderson FJ, Coleman J, et al. Association of maternal sleep practices with pre-eclampsia, low birth weight, and stillbirth among Ghanaian women. *International Journal of Gynecology & Obstetrics* 2013; **121**(3): 261-5.
102. Stacey T, Thompson JMD, Mitchell EA, et al. Antenatal care, identification of suboptimal fetal growth and risk of late stillbirth: Findings from the Auckland Stillbirth Study. *Australian and New Zealand Journal of Obstetrics and Gynaecology* 2012; **52**(3): 242-7.
103. Platts J, Mitchell EA, Stacey T, et al. The Midland and North of England Stillbirth Study (MiNESS). *BMC Pregnancy and Childbirth* 2014; **14**(1): 171.
104. Heazell AEP, Whitworth MK, Whitcombe J, et al. Research Priorities for Stillbirth: Process Overview and Results from the UK Stillbirth Priority Setting Partnership. *Ultrasound in Obstetrics & Gynecology* 2015.

- 297 105. MacDorman MF. Race and ethnic disparities in fetal mortality, preterm birth, and infant
298 mortality in the United States: an overview. *Semin Perinatol* 2011; **35**(4): 200-8.
299 106. Zhang S, Cardarelli K, Shim R, et al. Racial disparities in economic and clinical outcomes of
300 pregnancy among medicaid recipients. *Maternal and Child Health Journal* 2013; **17**(8): 1518-25.
301 107. Auger N, Park AL, Zoungrana H, et al. Rates of stillbirth by gestational age and cause in Inuit
302 and First Nations populations in Quebec. *CMAJ* 2013; **185**(6): E256-62.

303

Panels

Panel 1. Methods

ISA web-based surveys

We developed three web-based, multilanguage surveys of bereaved parents, care providers and general community members to assess practices around stillbirth prevention, awareness of stillbirth risk factors, quality of antepartum and bereavement care, uptake of stillbirth investigations, audit and classification of stillbirths and more. A mix of categorical items, open-ended items, ranking items and rating scales were included. Surveys were disseminated chiefly via the International Stillbirth Alliance (ISA) member organisations and additional relevant professional societies between December 2014 and February 2015. Surveys were available in English, Dutch, German, Italian, Spanish and Portuguese. The survey of care providers was also available in French and Japanese.

In total, 6,636 responses were received across 32 HIC. Quantitative data were analysed in SPSS and were weighted to account for uneven distribution of responses across countries (see webappendix). Qualitative data were sorted in NVivo.

Surveys were approved by the Mater Health Services Human Research Ethics Committee, within the guidelines of the Australian National Statement on Ethical Conduct in Human Research and the University of British Columbia Office Of Research Ethics.

Stillbirth rates in HIC / analysis of avoidable deaths

The number of potentially preventable stillbirths across all HIC for the year 2015 was calculated from stillbirth rates and total births in Blencowe and colleagues⁸ by subtracting the anticipated numbers of stillbirths applying the 2015 stillbirth rates from the numbers derived using a rate of 2/1000 births for all countries with a rate above 2/1000.

Summary of disparities in HIC

Socioeconomic disparities in stillbirth rates in HIC were investigated via narrative review, by searching for papers on health inequities and social disadvantage as they relate to stillbirth, by snowballing from those papers, and from citations made to the LSS HIC paper.⁷ We also contacted experts in the field for details of programs addressing disparities in stillbirth and ways of reducing stillbirths in HIC.

335 **Review of national policies and clinical practice guidelines**

336 *National policies on perinatal mortality audit*

337 We searched for policies national policies on perinatal mortality audit across the top 36 HIC
338 according to number of annual stillbirths (equating to 99% of the known stillbirth burden in HIC).
339 Perinatal audit was defined as per Dunn and McIlwaine as “The systematic, critical analysis of the
340 quality of perinatal care, including the procedures used for diagnosis and treatment, the use of
341 resources and the resultant outcome and quality of life for women and their babies”.⁷⁸ Specifically,
342 we searched for policies for national perinatal data collection that is coupled with mandatory in-
343 depth review of care by a multidisciplinary team following a stillbirth (see webappendix for further
344 search details). We also consulted a local key informant to confirm findings and gain further
345 information.

346 *National clinical practice guidelines*


347 National clinical practice guidelines and recommendations addressing stillbirth prevention and
348 investigations were identified across the top five developed countries according to number of annual
349 stillbirths (Russian Federation; United States, Japan, France, The United Kingdom). Guidelines of
350 interest included those addressing the key messages of the 2011 LSS HIC paper,⁷ namely overweight
351 and obesity, alcohol and substance use, smoking cessation, training of health professionals to
352 provide care to disadvantaged pregnant women, and stillbirth investigations protocols to assess
353 cause of death. A structured search was conducted specific for each country using a customised list
354 of organisational websites providing national health care guidelines (see webappendix for further
355 search details). We also consulted a local key informant.

356 **Panel 2. Essential steps to establish perinatal mortality audit at the national level**

- 357 • Execute an information-plan to ensure it is known that stillbirths are not inevitable and that
358 many stillbirths can be prevented by increasing quality of care, particularly around term.
- 359 • Obtain support and budget from national bodies, including ministries of health and
360 professional colleges.
- 361 • Develop a national network to coordinate the data collection and identify missing cases
362 through a check system using births and death certificate; and to lead timely reporting and
363 analysis.
- 364 • Establish a national multidisciplinary leadership/steering group to drive the process, agree on
365 national priorities, develop and monitor formal audit-methodology, establish consistent and
366 robust definitions, ensure consistency across jurisdictions, and ensure perinatal audit
367 remains on the national agenda while it remains relevant.
- 368 • Identify clinical champion(s) at service delivery level.
- 369 • Develop a system for clinical and process data collection, preferably web-based, to be
370 completed by the clinical staff.
- 371 • Ensure that the underlying philosophy of data collection is based on shared ownership of the
372 data to optimise data quality. This includes the ability of units to access their own data in a
373 format that they can use for their own surveillance and perinatal mortality reviews.
- 374 • Allocate human resources to support local or regional audit initiatives.
- 375 • Conduct local review of perinatal cases with multidisciplinary teams that have allocated time.
- 376 • Develop a method to provide useful, automatically-generated feedback to clinicians and
377 facilities of the suggested improvements to support local quality and audit processes.
- 378 • Conduct effective monitoring and evaluation of the audit program with relevant and feasible
379 performance indicators.
- 380 • Implement processes to ensure that disclosure of information cannot be used for disciplinary
381 action.
- 382 • Ensure there is support/funding to support implementation of recommendations.

Figures


Figure 1. Current stillbirth rates and reductions since 2000 in HIC



ARR: Annual rate reduction *Countries with <5000 annual births

Figure 2. Survey data on perceptions of stillbirth in high-income countries

“In the community I live/work, people/my colleagues generally think that:”



**Full statement: “Parents should not talk about their stillborn baby because it makes people feel uncomfortable”*

Figure 3. Disparity pathways for stillbirth




Figure depicts complex relationships and associations for social disparity in stillbirth rates. Stillbirth rates can be double for African-American women;^{105, 106} Aboriginal and Indigenous women in Canada, Australia and New Zealand;^{10, 51, 107} women who have migrated, especially those coming as humanitarian entrants/asylum seekers or refugees;^{60, 107} low income women;⁵⁸ women with low educational achievement;¹² and teenage women under 15 years.¹⁰⁵