

## CELL SCIENCE AT A GLANCE

## Subcellular mRNA localisation at a glance

Richard M. Parton<sup>1,\*</sup>, Alexander Davidson<sup>1,\*</sup>, Ilan Davis<sup>1</sup> and Timothy T. Weil<sup>2,‡</sup>

## ABSTRACT

mRNA localisation coupled to translational regulation provides an important means of dictating when and where proteins function in a variety of model systems. This mechanism is particularly relevant in polarised or migrating cells. Although many of the models for how this

<sup>1</sup>Department of Biochemistry, University of Oxford, South Parks Road, Oxford OX1 3QU, UK. <sup>2</sup>Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK.


\*These authors contributed equally to this work

‡Author for correspondence (tw419@cam.ac.uk)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

is achieved were first proposed over 20 years ago, some of the molecular details are still poorly understood. Nevertheless, advanced imaging, biochemical and computational approaches have started to shed light on the cis-acting localisation signals and trans-acting factors that dictate the final destination of localised transcripts. In this Cell Science at a Glance article and accompanying poster, we provide an overview of mRNA localisation, from transcription to degradation, focusing on the microtubule-dependent active transport and anchoring mechanism, which we will use to explain the general paradigm. However, it is clear that there are diverse ways in which mRNAs become localised and target protein expression, and we highlight some of the similarities and differences between these mechanisms.

**KEY WORDS:** Cell Biology, Development, Polarity, mRNA



## Introduction

mRNA localisation is a common and conserved means of targeting proteins to their site of function, and is important in a diverse range of cellular and developmental functions (reviewed in Medioni et al., 2012) with clear links to human diseases (Jeibmann et al., 2009).

Throughout their journey, mRNA transcripts never exist alone; they bind to a number of proteins to form mRNA–protein complexes, the ribonucleoprotein (RNP) complexes, (Box 1) (reviewed in Kato and Nakamura, 2012). Furthermore, as RNP complexes travel through the cell, there is evidence for dynamic remodeling of these complexes by different trans-acting protein factors that interact with regulatory cis-acting mRNA sequences (Otero et al., 2001). These interactions dictate the fate of mRNAs, so an understanding of these interactions is crucial for explaining the precise spatio-temporal control of localisation and translation. These biological processes have proven challenging to study because of the diversity of proteins and mRNA sequences involved. However, there has recently been significant progress owing to innovative experimental approaches (Box 2).

In this Cell Science at a Glance article and accompanying poster, we focus on localised coding mRNAs in polarised or migrating cells. Using a mechanism of ‘active transport and anchoring’ as a basis for comparison, we follow the fate of mRNA from nuclear export, particle formation (see Box 1), transport, anchoring, translation, to degradation (see poster), highlighting some of the recent exciting findings.

## Nuclear export

mRNAs start their journey in the nucleus where they are transcribed from DNA (Shandilya and Roberts, 2012). Nascent transcripts are spliced, typically co-transcriptionally, to generate the coding mRNA in a process that removes introns, followed by the deposition of the exon–junction complex (EJC) and other components necessary for export and localisation (Baurén and Wieslander, 1994; Natalizio and Wente, 2013; Saulière et al., 2012). At this point, the emerging mRNAs associate, for the first time, with protein co-factors (Box 1) to form RNP complexes (Holt and Bullock, 2009; Natalizio and Wente, 2013).

A growing theme that emerges from the recent literature is that, even at this early stage, the association of localising mRNAs with proteins in the nucleus can dictate their future fate. First, recent work on the EJC suggests that the formation of RNP multimers through interaction with other EJC complexes and with serine- and arginine-rich (SR) proteins plays a role in packaging mRNAs, and in preparing them for nuclear export (Singh et al., 2012). More generally, a host of protein associations is initiated co-transcriptionally in the nucleus and appears to change continually throughout the different stages of mRNA localisation (Marchand et al., 2012; Rodríguez-Navarro and Hurt, 2011; Tran et al., 2007; Trcek et al., 2010). The formation of RNP complexes is directed by specific regions (cis-acting domains), which typically comprise double-stranded secondary structures including hairpins, stem loops and bulges (Amrute-Nayak and Bullock, 2012; Hamilton et al., 2007; Patel et al., 2012; Wilhelm and Vale, 1993). Second, alternative splicing can redefine or shuffle these cis-acting sequences to generate RNP complexes with varying trans-acting protein composition that dictates differential fates of mRNAs (Horne-Badovinac and Bilder, 2008; Trcek et al., 2010). Third, direct interactions of RNP complexes with the nuclear pore complex (NPC) can be essential for the formation of an export- and localisation-competent RNP complex. For example in

## Box 1. Particle formation

Biochemical and genetic studies have shown how, for localising mRNAs, particular proteins determine each step of their journey (reviewed in Kato and Nakamura, 2012). Although a diverse array of proteins are involved, these trans-acting factors can be grouped into several classes (see poster).

Understanding the interaction between cis- and trans-acting components, RNP complex composition and remodeling are crucial for understanding the molecular mechanisms underlying mRNA localisation (Otero et al., 2001). Assembly of the multiple proteins required for regulating the translational state of the mRNA has been shown to involve numerous sites of weak interaction and involve modification of the mRNA folding (Chao et al., 2010; St Johnston, 2005). One of the intriguing questions is how the composition of the RNP complexes confers the specific fates of different mRNAs. Although there is evidence that different mRNAs with the same destination are packaged and transported together in the same RNP complex (Lange et al., 2008), it was hypothesised that particles with distinct destinations can be distinguished by certain trans-acting factors that are specific for the particular destination (Cha et al., 2001). It has been difficult to confirm this hypothesis because many mRNA binding proteins associate with a wide range of different transcripts; for example, the highly conserved double-stranded RNA (dsRNA)-binding protein Staufen (Stau) associates with both *oskar* (*osk*) and *bicoid* (*bcd*) mRNA in *Drosophila* oocytes (Ephrussi et al., 1991; St Johnston et al., 1989). This notion is further complicated by evidence from biochemical analysis such as the ‘atlas’ of mammalian mRNA-interacting proteins, which has revealed the enormous diversity of trans-acting factors (Castello et al., 2012).

These complex data suggest an alternative view of the specificity of RNP complexes, in that their flexibility in regulating transcripts is based on an unique complement of overlapping cofactors for each mRNA species (Castello et al., 2012; Castello et al., 2013; Kato and Nakamura, 2012; McDermott et al., 2012). A major remaining question is what precise combination of factors is required for the correct localization and translational regulation of each mRNA species, and how this is modified during the life cycle of the transcripts.

*Saccharomyces cerevisiae*, mutants for the nucleoporin protein Nup60p, fail to export *ASH1* mRNA from the nucleus and transcripts that are exported do not localise correctly (Powrie et al., 2011).

Once mRNAs are spliced, RNP complexes become export competent through the addition of further co-factors and diffuse through the interchromatin spaces in the nucleoplasm to the NPC (Mor et al., 2010). At the NPC, a conserved and highly complicated structure, they are exported through interactions with co-factors such as nuclear RNA export factor 1 and NTF2-related export protein 1 (NXF1 and NXT1, respectively) and the transcription export complex (TREX), which are essential for nuclear export (Natalizio and Wente, 2013; Rodríguez-Navarro and Hurt, 2011; Valkov et al., 2012). In *Drosophila* embryos and rat myoblasts, the export path has been shown to be random (Politz et al., 2003; Wilkie and Davis, 2001). However, the exact routes taken by RNP complexes remains an area of debate and may indeed be polarised with respect to the cell. Recent breakthroughs achieved by using ‘super registration’ fluorescence microscopy approaches, which allow a direct visualisation of mRNA export in transgenic mouse cell lines (Grünwald and Singer, 2010), show that not all nuclear pores are equally active at

## Box 2. Imaging techniques to study mRNA localisation

Early studies of mRNA localisation were reliant upon *in situ* hybridisation in fixed material, a slow and laborious method. Recent advances in fluorescent *in situ* hybridisation (FISH) now allow the rapid labeling of single mRNA molecules with high sensitivity and selectivity (Buxbaum et al., 2014; Little et al., 2011). FISH has also been applied to live cells (Santangelo et al., 2009). However, despite these improvements, *in situ* hybridisation remains limited in its application to certain tissues. An alternative approach to label mRNA in live cells is through injection of *in vitro* synthesized RNA that incorporates fluorescent dyes. By using this technique, it has been possible to rapidly screen a range of mutated RNAs to determine the crucial cis-acting sequences that determine mRNA localisation in *Drosophila* embryos and oocytes (Bullock and Ish-Horowicz, 2001; Van De Bor et al., 2005).

A breakthrough in live imaging that also emerges as one of the most useful approaches for *in vivo* study of mRNA localisation is the MS2 bacteriophage RNA stem loop bound by MS2 coat protein fusion to a fluorescent protein (MS2-MCP system). It was originally developed in yeast (Bertrand et al., 1998) but has since been extended to several organisms, including *Drosophila* (Forrest et al., 2003), *Mus musculus* (Park et al., 2014), *Xenopus* (Gagnon et al., 2013) and plants (Hamada et al., 2003). It has also been used to detect nascent mRNA as it is transcribed (Hocine et al., 2013; Yunger et al., 2013). Improvements are directed towards maximising the detection of mRNAs, such as by increasing the number and optimisation of MS2 loops (Lionnet et al., 2011). An alternative construct uses the bacteriophage PP7 and also involves the interaction between a stem loop and a viral coat protein (Wu et al., 2012). Using the MS2 and PP7 systems in combination might allow the tracking of two different species of mRNAs simultaneously and could provide insights into how transcripts segregate.

However, one limitation of the MS2 and PP7 systems is the potential lack of contrast caused by unbound MCP. This may be overcome by targeting MCP to the nucleus (Bertrand et al., 1998), although this would compromise the ability to follow mRNAs in and around the nucleus. Recently, a combination of MS2 and PP7 has been described that drives a highly specific split-fluorescent protein complementation on the mRNA and allows for high-contrast labeling of individual transcripts (Hocine et al., 2013).

Furthermore, different tissues present different challenges for imaging approaches, which could be addressed by using advances in imaging techniques. For instance, fast, sensitive widefield microscopy is useful when imaging fast-moving mRNA particles (Zimyanin et al., 2008), whereas when imaging within thick samples, such as *Drosophila* embryos, multi-photon microscopy can be useful (Sinsimer et al., 2013). Total internal reflection microscopy (TIRF) is ideal for *in vitro* applications because of its high axial resolution (Amrute-Nayak and Bullock, 2012), whereas 3D structured illumination microscopy (3D-SIM) gives axial and lateral super-resolution while being flexible enough to be applied when investigating conventionally prepared materials (Weil et al., 2010a).

the same time. Furthermore, mRNAs were shown to move bidirectionally through the pores, with docking at and release from the nuclear pore being the rate-limiting steps, rather than translocation through the channel. There is also evidence that translocation of a large RNP complex through a channel of limited diameter is facilitated by RNA helicases, which are essential for nuclear export and mRNA quality control (reviewed

in Valkov et al., 2012), and can specifically interact with nucleoporins and promote mRNA remodeling (Montpetit et al., 2011).

Interestingly, export through the NPC is not the only path to the cytoplasm. Recently, an alternative mechanism has been discovered in the *Drosophila* neuromuscular junction where large RNP complexes exit the nucleus through vesicular budding (Jokhi et al., 2013; Speese et al., 2012), a process thought to be utilized by some DNA viruses. Regardless of the underlying mechanism, once the mRNA has arrived at the cytoplasmic face of the nucleus, it is ready for the next phase of its journey.

## Transport and anchoring

Early studies of mRNA transport in the cytoplasm focused on the overall localisation of transcripts through the visualisation of endogenous mRNA by *in situ* hybridisation (Jeffery et al., 1983; Lawrence and Singer, 1986; Weil et al., 2010a). The subsequent development of the MS2 bacteriophage RNA stem loop bound by MS2 coat protein fusion to a fluorescent protein (MS2-MCP system) and recent imaging advances (see Box 2) combined with the use of mutants that affect transacting protein factors, motors and the cytoskeleton, and also drug treatments, showed that different mRNAs localise by means of different mechanisms (Bertrand et al., 1998; Forrest et al., 2003; Jaramillo et al., 2008; Lerit and Gavis, 2011; Medioni et al., 2012; Takatori et al., 2010). For example, in late *Drosophila* oogenesis, *bicoid* (*bcd*) is localised by active transport and *nanos* (*nos*) is localised by means of diffusion and trapping (Forrest et al., 2003; Weil et al., 2006). Biochemical experiments have supported and expanded our understanding of the trans-acting factors that are required for the regulation of these processes (McDermott et al., 2012; Müller et al., 2011; Snee et al., 2005). The classic view of RNP complexes is that they are composed of higher order mRNA complexes. This certainly appears to be the case for *oskar* (*osk*) mRNA, where multiple mRNA particles associate with each other to form large transport particles or granules (Chekulaeva et al., 2006; Kato and Nakamura, 2012). However, recent research following individual mRNAs indicates that oligomerisation is not obligatory for transport (Amrute-Nayak and Bullock, 2012). In this section, we focus primarily on the current understanding of mRNA localisation along microtubules by means of active motor-dependent transport in polarised cells. This is exemplified by the classic axis-determining mRNAs in *Drosophila* that move on microtubule tracks, with kinesin transporting *osk* mRNA in the plus-end direction and dynein driving *gurken* (*grk*) mRNA transport to the microtubule minus end (MacDougall et al., 2003; Zimyanin et al., 2008).

An essential requirement for localisation is the linkage of the correct motor to the cargo and the connection of this complex to the cytoskeleton. Bicaudal-D (BicD) and Egalitarian (Egl) are two linkers with key roles in axis determination in *Drosophila* oocytes and embryos (Mach and Lehmann, 1997; Bullock and Ish-Horowicz, 2001; Navarro et al., 2004). Egl directly binds to different cis-acting elements that mediate mRNA localisation and BicD regulates the linkage of mRNA cargo to dynein. A recent study provides insight into the molecular mechanism of motor protein recruitment by solving the crystal structure of the cargo-binding domain of BicD (Liu et al., 2013), which shows that cargo binds to the homotypic domain of BicD and releases the auto inhibition of its heterotypic coiled-coil domain, thus allowing this domain to bind to dynein.

Work *in vitro* has suggested that the number of motors that are recruited to an RNP complex can be controlled by cis-acting localisation elements in the mRNA (Amrute-Nayak and Bullock, 2012). Similarly, recruitment of multiple copies of the class V myosin Myo4 by She2 has been shown to enhance transport of *ASH1* mRNA along actin in *S. cerevisiae* (Chung and Takizawa, 2010). Additional levels of regulating transport efficiency also exist, for instance mediated by proteins such as Pat1, which interacts with cargo adaptors to regulate the motility of kinesin heavy chain on microtubules and is necessary for *osk* mRNA localisation (Loiseau et al., 2010). The microtubule-associated protein (MAP) ensconsin interacts with microtubules and kinesin-1 to increase efficiency of motor recruitment (Sung et al., 2008). This is different to typical MAPs, which affect cytoskeletal stability and organisation.

The classic model of mRNA localisation poses that it is underpinned by a highly polarised cytoskeletal network (Clark et al., 1994). This model predicts that mRNAs exhibit a concerted directional motion. However, research on the dynamics of RNP complexes often shows a non-uniform rather than continuous, processive movement (Sinsimer et al., 2013). One explanation for this is that the dynein motor is capable of reversing direction (Gross, 2004). Another explanation comes from recent *in vivo* and *in vitro* analysis that has shown that individual mRNA cargos can undergo bidirectional movement owing to transport by multiple motors. In the case of *Vg1* mRNA in frogs, there are different phases of transport during localisation. An initial highly unidirectional dynein-dependent phase is followed by multiple rounds of bidirectional transport for which kinesin-1, kinesin-2 and dynein are required, before transcripts are anchored at the vegetal cortex (Gagnon et al., 2013).

In *Drosophila* oocytes, live imaging of *osk* mRNA and visualisation of microtubules using green fluorescent protein (GFP) tagged to end-binding protein 1 (EB1) revealed an apparently random organisation with subtle bias towards the posterior, rather than the expected highly polarised cytoskeletal network (Parton et al., 2011; Zimyanin et al., 2008). Additional control of transport can be mediated through the regulation of the properties of the tracks themselves (Gardner et al., 2011), such as regulating the extension and catastrophic collapse of microtubules.

Once the mRNA has reached its destination, anchoring is a common mechanism for maintaining mRNA localisation. For example *grk* mRNA is anchored by dynein at the *Drosophila* oocyte dorsal anterior corner, whereas *nos* mRNA that is diffusing in the ooplasm is trapped by actin at the posterior pole (Delanoue et al., 2007; Forrest et al., 2003). In *Xenopus*, *Vg1* mRNA is maintained at the vegetal pole of the oocyte by the actin cytoskeleton (Yisraeli et al., 1990).

Actin has also been implicated in the organisation and function of the microtubule cytoskeleton and has been demonstrated to be of particular importance for the localisation of *bicoid* mRNA (Weil et al., 2010b) and in organising a polarised microtubule cytoskeleton in the *Drosophila* oocyte (Dahlgard et al., 2007). Actin can also function as a track on which cargoes can be transported. Here, actin not only acts in the short-range localisation of mRNAs, as is the case for *ASH1* mRNA in yeast (Bertrand et al., 1998), but also in long-range vesicle transport in mouse oocytes (Schuh et al., 2011; reviewed in St Johnston, 2005). Together, these examples demonstrate that there is no clear universal mechanism for mRNA transport. The future challenge is to understand to what extent the differential localisation of mRNA species is

achieved through distinct transport mechanisms that operate in parallel.

### Translation

For effective localised protein expression, mRNAs are kept translationally silent during transport and are activated for translation when they are anchored at their destination. Cis-acting mRNA sequences are almost certainly responsible for both the translational repression of mRNAs and their activation by directing their interaction with trans-acting proteins. Several strategies have been proposed, including the binding of factors that block or mask the interaction of translational activators, either by sequestering these translation activation factors or by occupying their interaction sites on the mRNA. These factors could either physically restrict the access of the translational machinery or act by regulating the length of the poly(A)-tail on the transcripts, whose extension is known to precede translational activation (Rosenthal et al., 1983).

One of the best-studied mechanisms of translational repression involves the eukaryotic translation initiation factor 4E (eIF4E) pathway (Jackson et al., 2010). In this repression pathway, mRNAs are prevented from initiating translation that requires the recruitment of the 40S ribosomal subunit through assembly of eIF4F (eIF4G+eIF4E and eIF4A) at the 5' cap. Translational repressors can bind directly to the mRNA, thus masking the sites for eIF4E binding. For example, the *Drosophila* protein Bruno interacts with Bruno-response elements on mRNA, thus blocking initiation of translation (Kim-Ha et al., 1995). Alternatively, factors such as the *Drosophila* ovarian protein Cup can bind eIF4E and prevent it from accessing mRNA, thereby inhibiting initiation of translation (Chekulaeva et al., 2006; Nakamura et al., 2004; Piccioni et al., 2005; Wilhelm et al., 2003). Overexpression of eIF4E has been shown to lead to autism-like behavior in mice (Santini et al., 2013), and HIV-1 was shown to be able to maintain virus-specific protein synthesis when eIF4E is downregulated (Sharma et al., 2012). This demonstrates that eIF4E-mediated translational repression is important in a range of cells and circumstances.

A further well-characterised means of controlling the level of proteins synthesis from mRNA is through polyadenylation or deadenylation. In frogs and flies, there are many examples of cytoplasmic polyadenylation elements (CPEs) that reside in the 3' UTR of mRNA, which are bound by the CPE-binding protein (CPEB) to control translation (Chang et al., 1999; Christerson and McKearin, 1994; Hake and Richter, 1994; Radford et al., 2008). Recently, work in the hippocampus of mice has shown that mutants in which the translational repressor of poly(A)-binding protein (PABP) has been knocked out show an increased translation of  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase II alpha (*Camk2a*) mRNA, a factor involved in many signalling cascades that are regulated by  $\text{Ca}^{2+}$ . Translational activation through the release of PABP-dependent repression has been demonstrated following electrode stimulation and is important for synaptic plasticity and learning (Khoutorsky et al., 2013).

Mouse models have also highlighted the complexity of translational regulation and revealed new means by which kinase activity regulates mRNA translation. For instance, the mouse protein kinase R (PKR)-like endoplasmic reticulum kinase (PERK, also known as EIF2AK3) has been shown to have a key role in brain function by phosphorylating eIF2 $\alpha$  (also known as EIF2A), a key regulator of translational activity (Trinh et al., 2012). A unique level of intricacy has been shown to be provided

by the interaction of fragile X mental retardation 1 (FMRP, also known as FMR1) protein and the human topoisomerase 3-beta-1 (Top3 $\beta$ , also known as TOP3B) (Xu et al., 2013), as this complex appears to regulate multiple mRNAs in neurons and – in Top3 $\beta$  mutants – there is a reduction in the expression of genes within the neuromuscular junction (NMJ) that are important for neural function. In addition to FMRP, other RNA-binding proteins found in the *Drosophila* nervous system, such as Syncrip (Syp) and Staufen (Stau), have distinct roles in regulation mRNA translation in other stages of development (McDermott et al., 2012; Barbee et al., 2006).

Although non-coding RNAs are thought to primarily have a role in mRNA degradation (see following section), an additional role is emerging in translational regulation. In flies, a reversible mechanism for regulating gene expression through the pathway via microRNA (miRNA) and argonaute 2 (AGO2) (Muddashetty et al., 2011) involves the phosphorylation of *Drosophila* FMR1 at the synapse. The direct targeting of precursor miRNA – a long piece of double-stranded RNA (dsRNA) from which mature miRNAs are generated – to neuronal dendrites has been shown to be mediated by the DEAH-box helicase 36 (DHX36), suggesting that the localisation of precursor miRNA is an important plasticity mechanism (Bicker et al., 2013). Interestingly, an miRNA-independent mechanism for the recruitment of Argonaute-1 (AGO1) to nanos (*nos*) mRNA in the *Drosophila* early embryo has been described that acts through the smaug protein, suggesting that there are different mechanisms of translational regulation that most likely work in concert (Pinder and Smibert, 2013).

The recent characterisation of protein components that are involved in translational repression of mRNAs has revealed that the subcellular organisation of protein complexes contributes to efficient translational regulation (Balagopal and Parker, 2009). Processing bodies (P-bodies) are known locations of mRNA translational control and degradation and have, therefore, been referred to as hubs for RNA metabolism in yeast (Aizer et al., 2008; reviewed in Balagopal and Parker, 2009). It has recently been shown that they are also important in regulating developmentally relevant transcripts. We have also shown that it is important where exactly in P-bodies mRNAs localise because the inside of electron-dense P-bodies does not support translation, whereas their localisation at the P-body edge allows translation to occur (Weil et al., 2012). Other work from yeast suggests that P-bodies are also the place where mRNAs terminate their journey through the cell (Aizer et al., 2008; Brengues et al., 2005).

## Degradation

The tight regulation of mRNAs in both space and time ultimately requires mRNA degradation. Although there are several pathways of degradation, the most common is through 5' to 3' exonuclease activity, following deadenylation and decapping (reviewed in Decker and Parker, 2012). In yeast, this process is linked with cytoplasmic P-bodies, which are associated with translationally repressed transcripts (Teixeira et al., 2005). These are distinct from stress granules, related cytoplasmic foci that have a similar composition and are likely to share functions. Stress granules assemble under physiological conditions when translation is stalled rather than repressed (Buchan and Parker, 2009). Interestingly, recent research indicates that mRNA can be degraded in a 5' to 3' fashion as translation is occurring (Hu et al., 2009), suggesting that the removal of ribosomes and the activity of exonucleases are not necessarily sequential events.

In *Drosophila*, P-body proteins have been found in neurons and oocytes, and have been suggested to be important for regulating transcripts (Barbee et al., 2006; Weil et al., 2012). In the case of *grk* mRNA, these bodies may be involved in regulating the level of transcripts and act as a dosage and temporal control, although degradation was not explicitly demonstrated in our study (Weil et al., 2012). An additional mechanism of mRNA degradation that is relevant for the control of spatially and temporally constrained transcripts is miRNA-mediated degradation. In the case of the maternal to zygotic transition in zebrafish and *Drosophila* embryos, maternal mRNAs are cleared from the embryo by miRNA-mediated deadenylation (Bushati et al., 2008; Giraldez et al., 2006).

Another mechanism of mRNA regulation is at the level of the DNA sequence, which can affect mRNA levels through the control of decay rates. Recent evidence from yeast shows that DNA promoter elements can affect the decay kinetics of their respective mRNAs after nuclear export (Bregman et al., 2011).

The specific mechanisms that regulate the degradation of mRNA transcripts after their final localisation have yet to be extensively studied and, thus, the role of P-bodies as sites of degradation remains to be established, mainly owing to the difficulty in characterising these labile bodies and in observing their interactions with mRNA.

## Perspectives

New imaging technologies, bioinformatics and biochemistry have facilitated major advances in our understanding of the composition, motility and translational regulation of RNP complexes. One successful example of these applications is in the analysis of RNP complex composition of nascent transcripts as they are being processed during splicing in the nucleus and at the NPC.

However, many important questions in the field remain unanswered, such as what is the extent of RNP complex remodeling in the cytoplasm and how is it regulated? How prevalent is the function of small non-coding RNA in the translational regulation of mRNAs? Do the numerous regulatory mechanisms facilitate the localisation of diverse transcripts or are they redundant mechanisms to protect crucial biological processes? The challenge will be to place the biochemical information with regard to the components that are involved in the context of *in vivo* data to fully understand the molecular mechanism of mRNA localisation with respect to cell and tissue function.

## Acknowledgements

We are grateful to Russell Hamilton and David Ish-Horowicz for critical reading of the manuscript.

## Competing interests

The authors declare no competing interests.

## Funding

This work was supported by a Wellcome Trust Senior Research Fellowship to I.D. supporting R.M.P. [grant number: 096144], a studentship from the Wellcome Trust to A.D. [grant number: 097304], the University of Cambridge, ISSF to T.T.W. [grant number 097814]. Deposited in PMC for immediate release.

## Cell science at a glance

A high-resolution version of the poster is available for downloading in the online version of this article at [jcs.biologists.org](http://jcs.biologists.org). Individual poster panels are available as JPEG files at <http://jcs.biologists.org/lookup/suppl/doi:10.1242/jcs.114272/-DC1>.

## References

- Aizer, A., Brody, Y., Ler, L. W., Sonenberg, N., Singer, R. H. and Shav-Tal, Y. (2008). The dynamics of mammalian P body transport, assembly, and disassembly in vivo. *Mol. Biol. Cell* **19**, 4154–4166.
- Amrute-Nayak, M. and Bullock, S. L. (2012). Single-molecule assays reveal that RNA localization signals regulate dynein-dynactin copy number on individual transcript cargoes. *Nat. Cell Biol.* **14**, 416–423.
- Balagopal, V. and Parker, R. (2009). Polyosomes, P bodies and stress granules: states and fates of eukaryotic mRNAs. *Curr. Opin. Cell Biol.* **21**, 403–408.
- Barbee, S. A., Estes, P. S., Cziko, A.-M., Hillebrand, J., Luedeman, R. A., Coller, J. M., Johnson, N., Howlett, I. C., Geng, C., Ueda, R. et al. (2006). Staufen- and FMRP-containing neuronal RNPs are structurally and functionally related to somatic P bodies. *Neuron* **52**, 997–1009.
- Baurén, G. and Wieslander, L. (1994). Splicing of Balbiani ring 1 gene pre-mRNA occurs simultaneously with transcription. *Cell* **76**, 183–192.
- Bertrand, E., Chartrand, P., Schaefer, M., Shenoy, S. M., Singer, R. H. and Long, R. M. (1998). Localization of ASH1 mRNA particles in living yeast. *Mol. Cell* **2**, 437–445.
- Bicker, S., Khudayberdiev, S., Weiß, K., Zocher, K., Baumeister, S. and Schrott, G. (2013). The DEAH-box helicase DHX36 mediates dendritic localization of the neuronal precursor-microRNA-134. *Genes Dev.* **27**, 991–996.
- Bregman, A., Avraham-Kelbert, M., Barkai, O., Duek, L., Guterman, A. and Choder, M. (2011). Promoter elements regulate cytoplasmic mRNA decay. *Cell* **147**, 1473–1483.
- Brengues, M., Teixeira, D. and Parker, R. (2005). Movement of eukaryotic mRNAs between polyosomes and cytoplasmic processing bodies. *Science* **310**, 486–489.
- Buchan, J. R. and Parker, R. (2009). Eukaryotic stress granules: the ins and outs of translation. *Mol. Cell* **36**, 932–941.
- Bullock, S. L. and Ish-Horowicz, D. (2001). Conserved signals and machinery for RNA transport in Drosophila oogenesis and embryogenesis. *Nature* **414**, 611–616.
- Bushati, N., Stark, A., Brennecke, J. and Cohen, S. M. (2008). Temporal reciprocity of miRNAs and their targets during the maternal-to-zygotic transition in Drosophila. *Curr. Biol.* **18**, 501–506.
- Buxbaum, A. R., Wu, B. and Singer, R. H. (2014). Single β-actin mRNA detection in neurons reveals a mechanism for regulating its translatability. *Science* **343**, 419–422.
- Castello, A., Fischer, B., Eichelbaum, K., Horos, R., Beckmann, B. M., Strein, C., Davey, N. E., Humphreys, D. T., Preiss, T., Steinmetz, L. M. et al. (2012). Insights into RNA biology from an atlas of mammalian mRNA-binding proteins. *Cell* **149**, 1393–1406.
- Castello, A., Horos, R., Strein, C., Fischer, B., Eichelbaum, K., Steinmetz, L. M., Krijgsfeld, J. and Hentze, M. W. (2013). System-wide identification of RNA-binding proteins by interactome capture. *Nat. Protoc.* **8**, 491–500.
- Cha, B. J., Koppetsch, B. S. and Theurkauf, W. E. (2001). In vivo analysis of Drosophila bicoid mRNA localization reveals a novel microtubule-dependent axis specification pathway. *Cell* **106**, 35–46.
- Chang, J. S., Tan, L. and Schedl, P. (1999). The Drosophila CPEB homolog, orb, is required for oskar protein expression in oocytes. *Dev. Biol.* **215**, 91–106.
- Chao, J. A., Patskovsky, Y., Patel, V., Levy, M., Almo, S. C. and Singer, R. H. (2010). ZBP1 recognition of beta-actin zipcode induces RNA looping. *Genes Dev.* **24**, 148–158.
- Chekulaeva, M., Hentze, M. W. and Ephrussi, A. (2006). Bruno acts as a dual repressor of oskar translation, promoting mRNA oligomerization and formation of silencing particles. *Cell* **124**, 521–533.
- Christerson, L. B. and McKearin, D. M. (1994). orb is required for anteroposterior and dorsoventral patterning during Drosophila oogenesis. *Genes Dev.* **8**, 614–628.
- Chung, S. and Takizawa, P. A. (2010). Multiple Myo4 motors enhance ASH1 mRNA transport in Saccharomyces cerevisiae. *J. Cell Biol.* **189**, 755–767.
- Clark, I., Giniger, E., Ruohola-Baker, H., Jan, L. Y. and Jan, Y. N. (1994). Transient posterior localization of a kinesin fusion protein reflects anteroposterior polarity of the Drosophila oocyte. *Curr. Biol.* **4**, 289–300.
- Dahlgaard, K., Raposo, A. A. S. F., Niccoli, T. and St Johnston, D. (2007). Capu and Spire assemble a cytoplasmic actin mesh that maintains microtubule organization in the Drosophila oocyte. *Dev. Cell* **13**, 539–553.
- Decker, C. J. and Parker, R. (2012). P-bodies and stress granules: possible roles in the control of translation and mRNA degradation. *Cold Spring Harb. Perspect. Biol.* **4**, a012286.
- Delanoue, R., Herpers, B., Soetaert, J., Davis, I. and Rabouille, C. (2007). Drosophila Squid/hnRNP helps Dynein switch from a Gurken mRNA transport motor to an ultrastructural static anchor in sponge bodies. *Dev. Cell* **13**, 523–538.
- Ephrussi, A., Dickinson, L. K. and Lehmann, R. (1991). Oskar organizes the germ plasm and directs localization of the posterior determinant nanos. *Cell* **66**, 37–50.
- Forrest, K. M. and Gavis, E. R. (2003). Live imaging of endogenous RNA reveals a diffusion and entrapment mechanism for nanos mRNA localization in Drosophila. *Curr. Biol.* **13**, 1159–1168.
- Gagnon, J. A., Kreiling, J. A., Powrie, E. A., Wood, T. R. and Mowry, K. L. (2013). Directional transport is mediated by a Dynein-dependent step in an RNA localization pathway. *PLoS Biol.* **11**, e1001551.
- Gardner, M. K., Zanic, M., Gell, C., Bormuth, V. and Howard, J. (2011). Depolymerizing kinesins Kip3 and MCAK shape cellular microtubule architecture by differential control of catastrophe. *Cell* **147**, 1092–1103.
- Giraldez, A. J., Mishima, Y., Rihel, J., Grocock, R. J., Van Dongen, S., Inoue, K., Enright, A. J. and Schier, A. F. (2006). Zebrafish MiR-430 promotes deadenylation and clearance of maternal mRNAs. *Science* **312**, 75–79.
- Gross, S. P. (2004). Hither and yon: a review of bi-directional microtubule-based transport. *Phys. Biol.* **1**, R1–R11.
- Grünwald, D. and Singer, R. H. (2010). In vivo imaging of labelled endogenous β-actin mRNA during nucleocytoplasmic transport. *Nature* **467**, 604–607.
- Hake, L. E. and Richter, J. D. (1994). CPEB is a specificity factor that mediates cytoplasmic polyadenylation during Xenopus oocyte maturation. *Cell* **79**, 617–627.
- Hamada, S., Ishiyama, K., Choi, S.-B., Wang, C., Singh, S., Kawai, N., Franceschi, V. R. and Okita, T. W. (2003). The transport of prolamine RNAs to prolamine protein bodies in living rice endosperm cells. *Plant Cell* **15**, 2253–2264.
- Hamilton, R. S. and Davis, I. (2007). RNA localization signals: deciphering the message with bioinformatics. *Semin. Cell Dev. Biol.* **18**, 178–185.
- Hocine, S., Raymond, P., Zenklusen, D., Chao, J. A. and Singer, R. H. (2013). Single-molecule analysis of gene expression using two-color RNA labeling in live yeast. *Nat. Methods* **10**, 119–121.
- Holt, C. E. and Bullock, S. L. (2009). Subcellular mRNA localization in animal cells and why it matters. *Science* **326**, 1212–1216.
- Horne-Badovinac, S. and Bilder, D. (2008). Dynein regulates epithelial polarity and the apical localization of stardust A mRNA. *PLoS Genet.* **4**, e8.
- Hu, W., Sweet, T. J., Chamnongpol, S., Baker, K. E. and Coller, J. (2009). Co-translational mRNA decay in *Saccharomyces cerevisiae*. *Nature* **461**, 225–229.
- Jackson, R. J., Hellen, C. U. T. and Pestova, T. V. (2010). The mechanism of eukaryotic translation initiation and principles of its regulation. *Nat. Rev. Mol. Cell Biol.* **11**, 113–127.
- Jaramillo, A. M., Weil, T. T., Goodhouse, J., Gavis, E. R. and Schupbach, T. (2008). The dynamics of fluorescently labeled endogenous Gurken mRNA in Drosophila. *J. Cell Sci.* **121**, 887–894.
- Jeffery, W. R., Tomlinson, C. R. and Brodeur, R. D. (1983). Localization of actin messenger RNA during early ascidian development. *Dev. Biol.* **99**, 408–417.
- Jeibmann, A. and Paulus, W. (2009). Drosophila melanogaster as a model organism of brain diseases. *Int. J. Mol. Sci.* **10**, 407–440.
- Jokhi, V., Ashley, J., Nunnari, J., Noma, A., Ito, N., Wakabayashi-Ito, N., Moore, M. J. and Budnik, V. (2013). Torsin mediates primary envelopment of large ribonucleoprotein granules at the nuclear envelope. *Cell Rep.* **3**, 988–995.
- Kato, Y. and Nakamura, A. (2012). Roles of cytoplasmic RNP granules in intracellular RNA localization and translational control in the Drosophila oocyte. *Dev. Growth Differ.* **54**, 19–31.
- Khoutorsky, A., Yanagiya, A., Gkogkas, C. G., Fabian, M. R., Prager-Khoutorsky, M., Cao, R., Gamache, K., Bouthiette, F., Parsyan, A., Sorge, R. E. et al. (2013). Control of synaptic plasticity and memory via suppression of poly(A)-binding protein. *Neuron* **78**, 298–311.
- Kim-Ha, J., Kerr, K. and Macdonald, P. M. (1995). Translational regulation of oskar mRNA by bruno, an ovarian RNA-binding protein, is essential. *Cell* **81**, 403–412.
- Lange, S., Katayama, Y., Schmid, M., Burkacky, O., Bräuchle, C., Lamb, D. C. and Jansen, R.-P. (2008). Simultaneous transport of different localized mRNA species revealed by live-cell imaging. *Traffic* **9**, 1256–1267.
- Lawrence, J. B. and Singer, R. H. (1986). Intracellular localization of messenger RNAs for cytoskeletal proteins. *Cell* **45**, 407–415.
- Lerit, D. A. and Gavis, E. R. (2011). Transport of germ plasm on astral microtubules directs germ cell development in Drosophila. *Curr. Biol.* **21**, 439–448.
- Lionnet, T., Czaplinski, K., Darzacq, X., Shav-Tal, Y., Wells, A. L., Chao, J. A., Park, H. Y., de Turris, V., Lopez-Jones, M. and Singer, R. H. (2011). A transgenic mouse for in vivo detection of endogenous labeled mRNA. *Nat. Methods* **8**, 165–170.
- Little, S. C., Tkacik, G., Kneeland, T. B., Wieschaus, E. F. and Gregor, T. (2011). The formation of the Bicoid morphogen gradient requires protein movement from anteriorly localized mRNA. *PLoS Biol.* **9**, e1000596.
- Liu, Y., Salter, H. K., Holding, A. N., Johnson, C. M., Stephens, E., Lukavsky, P. J., Walshaw, J. and Bullock, S. L. (2013). Bicaudal-D uses a parallel, homodimeric coiled coil with heterotypic registry to coordinate recruitment of cargos to dynein. *Genes Dev.* **27**, 1233–1246.
- Loiseau, P., Davies, T., Williams, L. S., Mishima, M. and Palacios, I. M. (2010). Drosophila PAT1 is required for Kinesin-1 to transport cargo and to maximize its motility. *Development* **137**, 2763–2772.
- MacDougall, N., Clark, A., MacDougall, E. and Davis, I. (2003). Drosophila Gurken (TGFα) mRNA localizes as particles that move within the oocyte in two dynein-dependent steps. *Dev. Cell* **4**, 307–319.
- Mach, J. M. and Lehmann, R. (1997). An Egalitarian-Bicaudal-D complex is essential for oocyte specification and axis determination in Drosophila. *Genes Dev.* **11**, 423–435.
- Marchand, V., Gaspar, I. and Ephrussi, A. (2012). An intracellular transmission control protocol: assembly and transport of ribonucleoprotein complexes. *Curr. Opin. Cell Biol.* **24**, 202–210.
- McDermott, S. M., Meignin, C., Rappaport, J. and Davis, I. (2012). Drosophila Syncpr binds the Gurken mRNA localisation signal and regulates localised transcripts during axis specification. *Biol. Open* **1**, 488–497.
- Medioni, C., Mowry, K. and Besse, F. (2012). Principles and roles of mRNA localization in animal development. *Development* **139**, 3263–3276.

- Montpetit, B., Thomsen, N. D., Helmke, K. J., Seeliger, M. A., Berger, J. M. and Weis, K.** (2011). A conserved mechanism of DEAD-box ATPase activation by nucleoporins and InsP6 in mRNA export. *Nature* **472**, 238–242.
- Mor, A., Suliman, S., Ben-Yishay, R., Yunger, S., Brody, Y. and Shav-Tal, Y.** (2010). Dynamics of single mRNP nucleocytoplasmic transport and export through the nuclear pore in living cells. *Nat. Cell Biol.* **12**, 543–552.
- Muddashetty, R. S., Nalavadi, V. C., Gross, C., Yao, X., Xing, L., Laur, O., Warren, S. T. and Bassell, G. J.** (2011). Reversible inhibition of PSD-95 mRNA translation by miR-125a, FMRP phosphorylation, and mGluR signaling. *Mol. Cell* **42**, 673–688.
- Müller, M., Heym, R. G., Mayer, A., Kramer, K., Schmid, M., Cramer, P., Urlaub, H., Jansen, R.-P. and Niessing, D.** (2011). A cytoplasmic complex mediates specific mRNA recognition and localization in yeast. *PLoS Biol.* **9**, e1000611.
- Nakamura, A., Sato, K. and Hanyu-Nakamura, K.** (2004). Drosophila cup is an eIF4E binding protein that associates with Bruno and regulates oskar mRNA translation in oogenesis. *Dev. Cell* **6**, 69–78.
- Natalizio, B. J. and Wente, S. R.** (2013). Postage for the messenger: designating routes for nuclear mRNA export. *Trends Cell Biol.* **23**, 365–373.
- Navarro, C., Puthalakath, H., Adams, J. M., Strasser, A. and Lehmann, R.** (2004). Egalitarian binds dynein light chain to establish oocyte polarity and maintain oocyte fate. *Nat. Cell Biol.* **6**, 427–435.
- Otero, L. J., Devaux, A. and Standart, N.** (2001). A 250-nucleotide UA-rich element in the 3' untranslated region of Xenopus laevis Vg1 mRNA represses translation both in vivo and in vitro. *RNA* **7**, 1753–1767.
- Park, H. Y., Lim, H., Yoon, Y. J., Follenzi, A., Nwokafor, C., Lopez-Jones, M., Meng, X. and Singer, R. H.** (2014). Visualization of dynamics of single endogenous mRNA labeled in live mouse. *Science* **343**, 422–424.
- Parton, R. M., Hamilton, R. S., Ball, G., Yang, L., Cullen, C. F., Lu, W., Ohkura, H. and Davis, I.** (2011). A PAR-1-dependent orientation gradient of dynamic microtubules directs posterior cargo transport in the Drosophila oocyte. *J. Cell Biol.* **194**, 121–135.
- Patel, V. L., Mitra, S., Harris, R., Buxbaum, A. R., Lionnet, T., Brenowitz, M., Girvin, M., Levy, M., Almo, S. C., Singer, R. H. et al.** (2012). Spatial arrangement of an RNA zipcode identifies mRNAs under post-transcriptional control. *Genes Dev.* **26**, 43–53.
- Piccioni, F., Zappavigna, V. and Verrotti, A. C.** (2005). A cup full of functions. *RNA Biol.* **2**, 125–128.
- Pinder, B. D. and Smibert, C. A.** (2013). microRNA-independent recruitment of Argonaute 1 to nanos mRNA through the Smaug RNA-binding protein. *EMBO Rep.* **14**, 80–86.
- Politz, J. C., Tuft, R. A. and Pederson, T.** (2003). Diffusion-based transport of nascent ribosomes in the nucleus. *Mol. Biol. Cell* **14**, 4805–4812.
- Powrie, E. A., Zenklusen, D. and Singer, R. H.** (2011). A nucleoporin, Nup60p, affects the nuclear and cytoplasmic localization of ASH1 mRNA in *S. cerevisiae*. *RNA* **17**, 134–144.
- Radford, H. E., Meijer, H. A. and de Moor, C. H.** (2008). Translational control by cytoplasmic polyadenylation in Xenopus oocytes. *Biochim. Biophys. Acta* **1779**, 217–229.
- Rodríguez-Navarro, S. and Hurt, E.** (2011). Linking gene regulation to mRNA production and export. *Curr. Opin. Cell Biol.* **23**, 302–309.
- Rosenthal, E. T., Tansey, T. R. and Ruderman, J. V.** (1983). Sequence-specific adenylations and deadenylations accompany changes in the translation of maternal messenger RNA after fertilization of *Spisula* oocytes. *J. Mol. Biol.* **166**, 309–327.
- Santangelo, P. J., Lifland, A. W., Curt, P., Sasaki, Y., Bassell, G. J., Lindquist, M. E. and Crowe, J. E., Jr** (2009). Single molecule-sensitive probes for imaging RNA in live cells. *Nat. Methods* **6**, 347–349.
- Santini, E., Huynh, T. N., MacAskill, A. F., Carter, A. G., Pierre, P., Ruggero, D., Kaphzan, H. and Klann, E.** (2013). Exaggerated translation causes synaptic and behavioural aberrations associated with autism. *Nature* **493**, 411–415.
- Saulière, J., Murigneux, V., Wang, Z., Marquenet, E., Barbosa, I., Le Tonquéze, O., Audic, Y., Paillard, L., Roest Crollius, H. and Le Hir, H.** (2012). CLIP-seq of eIF4AIII reveals transcriptome-wide mapping of the human exon junction complex. *Nat. Struct. Mol. Biol.* **19**, 1124–1131.
- Schuh, M.** (2011). An actin-dependent mechanism for long-range vesicle transport. *Nat. Cell Biol.* **13**, 1431–1436.
- Shandilya, J. and Roberts, S. G. E.** (2012). The transcription cycle in eukaryotes: from productive initiation to RNA polymerase II recycling. *Biochim. Biophys. Acta* **1819**, 391–400.
- Sharma, A., Yilmaz, A., Marsh, K., Cochrane, A. and Boris-Lawrie, K.** (2012). Thriving under stress: selective translation of HIV-1 structural protein mRNA during Vpr-mediated impairment of eIF4E translation activity. *PLoS Pathog.* **8**, e1002612.
- Singh, G., Kucukural, A., Cenik, C., Leszyk, J. D., Shaffer, S. A., Weng, Z. and Moore, M. J.** (2012). The cellular EJC interactome reveals higher-order mRNP structure and an EJC-SR protein nexus. *Cell* **151**, 750–764.
- Sinsimer, K. S., Lee, J. J., Thibierge, S. Y. and Gavis, E. R.** (2013). Germ plasm anchoring is a dynamic state that requires persistent trafficking. *Cell Rep.* **5**, 1169–1177.
- Snee, M. J., Arn, E. A., Bullock, S. L. and Macdonald, P. M.** (2005). Recognition of the bcd mRNA localization signal in Drosophila embryos and ovaries. *Mol. Cell. Biol.* **25**, 1501–1510.
- Speese, S. D., Ashley, J., Jokhi, V., Nunnari, J., Barria, R., Li, Y., Ataman, B., Koon, A., Chang, Y.-T., Li, Q. et al.** (2012). Nuclear envelope budding enables large ribonucleoprotein particle export during synaptic Wnt signaling. *Cell* **149**, 832–846.
- St Johnston, D.** (2005). Moving messages: the intracellular localization of mRNAs. *Nat. Rev. Mol. Cell Biol.* **6**, 363–375.
- St Johnston, D., Driever, W., Berleth, T., Richstein, S. and Nüsslein-Volhard, C.** (1989). Multiple steps in the localization of bicoid RNA to the anterior pole of the Drosophila oocyte. *Development* **107 Suppl.**, 13–19.
- Sung, H.-H., Telley, I. A., Papadaki, P., Ephrussi, A., Surrey, T. and Rørtø, P.** (2008). Drosophila ensconsin promotes productive recruitment of Kinesin-1 to microtubules. *Dev. Cell* **15**, 866–876.
- Takatori, N., Kumano, G., Saiga, H. and Nishida, H.** (2010). Segregation of germ layer fates by nuclear migration-dependent localization of Not mRNA. *Dev. Cell* **19**, 589–598.
- Teixeira, D., Sheth, U., Valencia-Sánchez, M. A., Brengues, M. and Parker, R.** (2005). Processing bodies require RNA for assembly and contain nontranslating mRNAs. *RNA* **11**, 371–382.
- Tran, E. J., Zhou, Y., Corbett, A. H. and Wente, S. R.** (2007). The DEAD-box protein Dbp5 controls mRNA export by triggering specific RNA:protein remodeling events. *Mol. Cell* **28**, 850–859.
- Trcek, T. and Singer, R. H.** (2010). The cytoplasmic fate of an mRNP is determined cotranscriptionally: exception or rule? *Genes Dev.* **24**, 1827–1831.
- Trinh, M. A., Kaphzan, H., Wek, R. C., Pierre, P., Cavener, D. R. and Klann, E.** (2012). Brain-specific disruption of the eIF2 $\alpha$  kinase PERK decreases ATF4 expression and impairs behavioral flexibility. *Cell Rep.* **1**, 676–688.
- Valkov, E., Dean, J. C., Jani, D., Kuhlmann, S. I. and Stewart, M.** (2012). Structural basis for the assembly and disassembly of mRNA nuclear export complexes. *Biochim. Biophys. Acta* **1819**, 578–592.
- Van De Bor, V., Hartswood, E., Jones, C., Finnegan, D. and Davis, I.** (2005). Gurken and the I factor retrotransposon RNAs share common localization signals and machinery. *Dev. Cell* **9**, 51–62.
- Weil, T. T., Forrest, K. M. and Gavis, E. R.** (2006). Localization of bicoid mRNA in late oocytes is maintained by continual active transport. *Dev. Cell* **11**, 251–262.
- Weil, T. T., Parton, R. M. and Davis, I.** (2010a). Making the message clear: visualizing mRNA localization. *Trends Cell Biol.* **20**, 380–390.
- Weil, T. T., Xanthakis, D., Parton, R., Dobbie, I., Rabouille, C., Gavis, E. R. and Davis, I.** (2010b). Distinguishing direct from indirect roles for bicoid mRNA localization factors. *Development* **137**, 169–176.
- Weil, T. T., Parton, R. M., Herpers, B., Soetaert, J., Veenendaal, T., Xanthakis, D., Dobbie, I. M., Halstead, J. M., Hayashi, R., Rabouille, C. et al.** (2012). Drosophila patterning is established by differential association of mRNAs with P bodies. *Nat. Cell Biol.* **14**, 1305–1315.
- Wilhelm, J. E. and Vale, R. D.** (1993). RNA on the move: the mRNA localization pathway. *J. Cell Biol.* **123**, 269–274.
- Wilhelm, J. E., Hilton, M., Amos, Q. and Henzel, W. J.** (2003). Cup is an eIF4E binding protein required for both the translational repression of oskar and the recruitment of Barentsz. *J. Cell Biol.* **163**, 1197–1204.
- Wilkie, G. S. and Davis, I.** (2001). Drosophila wingless and pair-rule transcripts localize apically by dynein-mediated transport of RNA particles. *Cell* **105**, 209–219.
- Wu, B., Chao, J. A. and Singer, R. H.** (2012). Fluorescence fluctuation spectroscopy enables quantitative imaging of single mRNAs in living cells. *Biophys. J.* **102**, 2936–2944.
- Xu, D., Shen, W., Guo, R., Xue, Y., Peng, W., Sima, J., Yang, J., Sharov, A., Srikanthan, S., Yang, J. et al.** (2013). Top3 $\beta$  is an RNA topoisomerase that works with fragile X syndrome protein to promote synapse formation. *Nat. Neurosci.* **16**, 1238–1247.
- Yisraeli, J. K., Sokol, S. and Melton, D. A.** (1990). A two-step model for the localization of maternal mRNA in Xenopus oocytes: involvement of microtubules and microfilaments in the translocation and anchoring of Vg1 mRNA. *Development* **108**, 289–298.
- Yunger, S., Rosenfeld, L., Garini, Y. and Shav-Tal, Y.** (2013). Quantifying the transcriptional output of single alleles in single living mammalian cells. *Nat. Protoc.* **8**, 393–408.
- Zimyanin, V. L., Belya, K., Pecreaux, J., Gilchrist, M. J., Clark, A., Davis, I. and St Johnston, D.** (2008). In vivo imaging of oskar mRNA transport reveals the mechanism of posterior localization. *Cell* **134**, 843–853.