#### Macroautophagy without LC3 conjugation?

Research Highlight to: "The ATG conjugation systems are important for degradation of the inner autophagosomal membrane" by Tsuboyama *et al., Science* 2016, DOI: 10.1126/science.aaf6136

Maurizio Renna<sup>1</sup>, David C. Rubinsztein<sup>1</sup>

<sup>1</sup>Department of Medical Genetics, University of Cambridge, Cambridge Institute for Medical Research, Wellcome Trust/MRC Building, Addenbrooke's Hospital, Hills Road, Cambridge, CB2 2XY, UK.

\* Address correspondence to: David C. Rubinsztein Phone: +44 (0)1223 762608, Fax: +44 (0)1223 331206 E-mail: <u>dcr1000@hermes.cam.ac.uk</u>

A recent study makes the surprising observation that autophagosomes can still form in the absence of the core conjugation machinery. Furthermore, while such autophagosomes can fuse with lysosomes, their degradation is delayed, and this is associated with delayed destruction of the inner autophagosomal double membrane, highlighting a new role for proteins thought to act exclusively in the formation of autophagosomes in late stages of the autophagic itinerary within autolysosomes.

In the last decade, the study of the mechanisms regulating the intertwined connections between the autophagic biogenesis and lysosomal compartment has gained increasing interest, and has been stimulated by the therapeutic and clinical relevance of this domain<sup>1-3</sup>. During macroautophagy (hereafter referred to as autophagy), cytoplasmic contents, including damaged organelles, pathogens and aggregate-prone proteins, are sequestered into a double-membrane structure called autophagosome, which eventually fuses with the lysosome to become an autolysosome, where the degradation occurs<sup>4</sup>. Lysosomal enzymes selectively degrade the inner (but not the outer) autophagosomal membrane (IAM), which finally allows the degradation of the enclosed autophagic content.

The formation of autophagosomes is enabled by more than 30 proteins. Many of these are so-called ATG proteins. Some of these, including ATG3, ATG5, ATG7, ATG10 and ATG12, regulate the conjugation of the ubiquitin-like ATG8 protein family members, such as LC3B, to autophagic precursor membranes enriched in phophatidylethanolamine. These conjugation events are thought to facilitate the extension and closure of the edges of autophagic precursor structure to allow the formation of completed, sealed autophagosomes<sup>5</sup>.

In a recent report, Mizushima and collaborators describe how autophagosomelike structures can be generated even in the absence of critical components of the canonical ATG conjugation systems, such as ATG3, ATG5 and ATG7<sup>6</sup>. Strikingly, even though these proteins are required for sealing of autophagosomes, these ATGs proteins do not appear to be essential for autolysosome formation but are important for efficient degradation of the inner autophagosomal membrane.

In order to further characterize the autophagosome maturation step, the authors relied on the autophagosomal SNARE syntaxin17 (STX17), whose hairpin-type tail-anchor is essential for targeting and driving the fusion of mature autophagosomes to the endosomal/lysosomal compartment<sup>7</sup>. Using live imaging, the authors were able to identify and discriminate between four steps during autophagosome maturation, namely: STX17 recruitment, lysosomal fusion, IAM degradation and STX17 release. Of particular interest, the STX17 recruitment induces an elliptical-to-spherical transition, which is likely coincident with the fission between the outer and inner autophagosomal membranes (Figure 1).

As expected, the rate of successful autophagosome formation was substantially reduced in ATG3 knockout (KO) cells, confirming that the ATG conjugation systems are indeed important for the late stages of autophagosome formation<sup>8-9</sup>. Consistent with these observation, autophagic substrate degradation was dramatically impaired in such cells. Nevertheless, the STX17-positive autophagosome-like structures in ATG3 KO cells were positive for SNAP29 and capable of undergoing acidification and co-localizing with the late endosome/lysosome marker, LAMP1. Thus, the fusion between autophagosome-like structures and lysosomes occur almost normally in ATG3 KO, as opposed to WT cells, although at a lower rate. Interestingly, their experiments suggested that IAM degradation is significantly delayed in ATG conjugation-deficient cells. This phenomenon was confirmed by immunoelectron microscopy and was not due to a lysosomal defect. Their data also suggested that the collapse of the IAM triggers the immediate dissociation of STX17 from the autolysosomes. In addition, they observed that a higher proportion of the STX17-positive autophagosome-like structures were elliptic in ATG3 KO cells, as opposed to those in WT cells. These results also suggest that, despite the STX17 recruitment, the ultimate fission step is somehow impaired in ATG non-competent cells (Figure 1).

To summarise, the function of ATG proteins had been investigated previously almost exclusively in the context of autophagosome formation. These novel findings reveal a functional role of ATG proteins involved in ATG8 family conjugation in late stages of autophagy after autophagosome-lysosome fusion. The ability to form some type of autophagic structures in the absence of these proteins is consistent with previous observations that a small proportion of autophagosome-like structures can mature into autolysosomes in ATG conjugation-deficient cells. This may explain, at least in part, some of the phenotypic differences and severities existing among different autophagy knock- out mouse models, depending whether they affect ATG8 conjugation or affect alternative or earlier steps of the process<sup>10-12</sup>.

### ACKNOWLEDGEMENTS

We are grateful to the Wellcome Trust (Principal Research Fellowship to DCR

(095317/Z/11/Z)) and a Strategic Grant to Cambridge Institute for Medical Research (100140/Z/12/Z) for funding.

# FIGURE LEGEND

# FIGURE 1 Inner Autophagosomal Membrane (IAM) route to lysosomal degradation depends on ATG proteins.

The function of ATG proteins has been mainly investigated in the context of autophagosome formation. In normal conditions, (WT) inner autophagosomal membrane (IAM) is sensitive to lysosomal degradation. While autophagic activity is strongly suppressed in ATG conjugation-deficient cells (exemplified by ATG3 KO in this Figure), autophagosome-like structures can still form and acquire STX17. These syntaxin17-positive structures are able to fuse with lysosomes. Nevertheless, degradation of the IAM is significantly impaired. These findings reveal a novel function of ATGs with regard to autophagosome maturation steps.

## REFERENCES

1. Mizushima N, Levine B, Cuervo AM, Klionsky DJ. Autophagy fights disease through cellular self-digestion. *Nature* 2008; **451**(7182): 1069-75 doi: 10.1038/nature06639.

2. Ravikumar B, Sarkar S, Davies JE, Futter M, Garcia-Arencibia M, Green-Thompson ZW, Jimenez-Sanchez M, Korolchuk VI, Lichtenberg M, Luo S, Massey DC, Menzies FM, Moreau K, Narayanan U, Renna M, Siddiqi FH, Underwood BR, Winslow AR, Rubinsztein DC. Regulation of mammalian autophagy in physiology and pathophysiology. *Physiol. Rev.* 2010; **90**(4), 1383-1435

3. Mizushima N, Komatsu M. Autophagy: Renovation of cells and tissues. *Cell* 2011; **147**: 728–741 doi:10.1016/j.cell.2011.10.026 pmid:22078875

4. Bento CF, Renna M, Ashkenazi A, Vicinanza M, Ghislat G, Puri C, Menzies FM, Rubinsztein DC. Mammalian Autophagy – how does it work? *ANNUAL REVIEW OF BIOCHEMISTRY* 2016; 85(1) DOI: 10.1146/annurevbiochem-060815-014556

5. Ktistakis NT, Tooze SA. Digesting the Expanding Mechanisms of Autophagy. *Trends Cell Biol.* 2016; **26**(8): 624-35 doi: 10.1016/j.tcb.2016.03.006.

6. Tsuboyama K, Koyama-Honda I, Sakamaki Y, Koike M, Morishita H, Mizushima N. The ATG conjugation systems are important for degradation of the inner autophagosomal membrane. *Science* 2016; DOI: 10.1126/science.aaf6136

7. Itakura E, Kishi-Itakura C, Mizushima N. The hairpin-type tail-anchored SNARE syntaxin 17 targets to autophagosomes for fusion with endosomes/lysosomes. *Cell* 2012; **151**: 256–1269

8. Fujita N, Hayashi-Nishino M, Fukumoto H, Omori H, Yamamoto A, Noda T, Yoshimori T. An Atg4B mutant hampers the lipidation of LC3 paralogues and causes defects in autophagosome closure. *Mol. Biol. Cell* 2008; **19**: 4651–4659 doi:10.1091/mbc.E08-03-0312 pmid:18768752

9. Noda T, Fujita N, Yoshimori T. The late stages of autophagy: How does the end begin? *Cell Death Differ.* 2009; **16**: 984-990 doi:10.1038/cdd.2009.54 pmid:19424283

10. Komatsu M, Waguri S, Ueno T, Iwata J, Murata S, Tanida I, Ezaki J, Mizushima N, Ohsumi Y, Uchiyama Y, Kominami E, Tanaka K, Chiba T, Impairment of starvation-induced and constitutive autophagy in Atg7-deficient mice. *J. Cell Biol.* 2005; **169**:425–434

11. Malhotra R, Warne JP, Salas E, Xu AW, Debnath J. Loss of Atg12, but not Atg5, in pro-opiomelanocortin neurons exacerbates diet-induced obesity.

### Autophagy 2015; 11: 145–154

12. Saitoh T, Fujita N, Jang MH, Uematsu S, Yang BG, Satoh T, Omori H, Noda T, Yamamoto N, Komatsu M, Tanaka K, Kawai T, Tsujimura T, Takeuchi O, Yoshimori T, Akira S. Loss of the autophagy protein Atg16L1 enhances endotoxin-induced IL-1β production. *Nature* 2008; **456**: 264–268



