

Review Article

How Supraphysiological Oxygen Levels in Standard Cell Culture Affect Oxygen-Consuming Reactions

Jeffrey A. Stuart ¹, Joao Fonseca,¹ Fereshteh Moradi,¹ Cassandra Cunningham,¹ Bishoy Seliman,¹ Cydney R. Worsfold,¹ Sarah Dolan,¹ John Abando,¹ and Lucas A. Maddalena²

¹Department of Biological Sciences, Brock University, St. Catharines, ON, Canada L2S 3A1

²MRC Cancer Research Centre, University of Cambridge, Hutchison/MRC Research Centre, Box 197, Cambridge Biomedical Campus, Cambridge CB2 0XZ, UK

Correspondence should be addressed to Jeffrey A. Stuart; jstuart@brocku.ca

Received 3 May 2018; Accepted 1 August 2018; Published 30 September 2018

Academic Editor: Eric E. Kelley

Copyright © 2018 Jeffrey A. Stuart et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Most mammalian tissue cells experience oxygen partial pressures *in vivo* equivalent to 1–6% O₂ (i.e., physioxia). In standard cell culture, however, headspace O₂ levels are usually not actively regulated and under these conditions are ~18%. This drives hyperoxia in cell culture media that can affect a wide variety of cellular activities and may compromise the ability of *in vitro* models to reproduce *in vivo* biology. Here, we review and discuss some specific O₂-consuming organelles and enzymes, including mitochondria, NADPH oxidases, the transplasma membrane redox system, nitric oxide synthases, xanthine oxidase, and monoamine oxidase with respect to their sensitivities to O₂ levels. Many of these produce reactive oxygen and/or nitrogen species (ROS/RNS) as either primary end products or byproducts and are acutely sensitive to O₂ levels in the range from 1% to 18%. Interestingly, many of them are also transcriptional targets of hypoxia-inducible factors (HIFs) and chronic cell growth at physioxia versus 18% O₂ may alter their expression. Aquaporins, which facilitate hydrogen peroxide diffusion into and out of cells, are also regulated by HIFs, indicating that O₂ levels may affect intercellular communication via hydrogen peroxide. The O₂ sensitivities of these important activities emphasize the importance of maintaining physioxia in culture.

1. Introduction

Mammalian cells are typically cultured under hyperoxic conditions. While most cells experience oxygen levels of 1–6% *in vivo* (physioxia; Table 1), almost all mammalian cell culture is done in humidified atmospheric air at 37°C with CO₂ added to 5%. Although the headspace O₂ level is usually not measured, it is 18–19% under these conditions due to displacement of O₂ by water vapour and CO₂. When O₂ levels used in cell culture experiments are measured and reported, these are virtually always those of headspace gas and not media. Particularly in metabolically active cells growing in high density, the pericellular media O₂ levels that cells experience may be substantially lower than headspace O₂ levels [1–3], since O₂ is continually removed from media by mitochondrial respiration and by other O₂-consuming cellular activities.

O₂ participates in many metabolic reactions, some of which are sensitive to O₂ levels in the range between physioxia and 18%. Therefore, mammalian cells should be cultured in physiologically relevant O₂ levels. Where this is not done, it is important to be aware of the potential consequences on cellular functions. One potential consequence of elevated O₂ in cell culture media is increased cellular production of reactive oxygen (ROS) and nitrogen (RNS) species. This probably contributes to the observed effects of high O₂ levels on cell senescence, differentiation, and apoptosis, amongst a wide range of other less well-characterized effects ([4–6]; Fehrer et al. 2007).

There is a growing appreciation of the role of O₂ levels in cell biology (e.g., [7, 8]). However, there is a limited mechanistic understanding of how supraphysiological O₂ levels influence specific O₂-dependent processes. In this review,

TABLE 1: Average oxygen levels measured in human tissues and cells *in vivo*.

Tissue/compartment	pO ₂ (%)	Reference
Brain	4.4 ± 0.3	[66]
Brain glioma cells (intracellular)	4.5 ± 0.5	
Skin	1.1–4.6	[66, 67]
Liver	5.4 ± 0.7	[66, 68]
Liver (mitochondrial)	3–4	[69]
Skeletal muscle	3.8 ± 0.2	[70]
Skeletal muscle (intracellular)	3.4–4.8	Lanza et al. 2010
Kidney	9.5 ± 2.6	[66]
Kidney (intracellular)	6.6–7.9	[71]
Bone marrow	1.3–2	[72, 73]

we consider the O₂ sensitivity of some of the important O₂-consuming organelles and enzymes in mammalian cells in the range between physioxia and 18% O₂ (standard cell culture conditions). We use the Michaelis-Menten constant (K_m(O₂)) for each O₂-consuming enzyme (where this value is available) as a convenient way to compare O₂ sensitivity across a broad range of different organelles and enzymes. All O₂ and K_m(O₂) values from published works have been converted to % O₂ at 37°C, since % O₂ is the unit of measurement used in most descriptions of cell culture experiments. In general, higher K_m(O₂) values are indicative of enzymes that will be more sensitive to the differences between *in vivo* and *in vitro* O₂ levels. We address the physiological role(s) of the metabolite(s) produced from these O₂-consuming reactions and the O₂ sensitivity over the range from physiological to 18% O₂ of that reaction. We further summarize the interesting observation that many of the O₂-consuming and ROS/RNS-producing enzymes are positively regulated by hypoxia, in some instances specifically by hypoxia-inducible factor-1 (HIF-1).

1.1. Oxygen Limitation of Mitochondrial Respiration in Cell Culture. An important goal of maintaining higher O₂ levels in culture is to ensure that mitochondrial respiration is not limited by O₂ availability. Some of the most comprehensive and physiologically relevant data on the O₂ levels required to sustain maximal mitochondrial respiration rates have been provided by Hoffmann et al. (2009), who measured these values for isolated liver mitochondria while systematically varying O₂ concentrations. State 4 respiration of complex I or complex II substrates (glutamate/malate or succinate, respectively) or of palmitoyl carnitine is near maximal at ~1% O₂ (Hoffman 2009). Marcinek et al. (2003) showed that respiration in skeletal muscle is not O₂ limited until O₂ falls below ~0.5%, which is similar to the observation by Gnaiger (2001) for isolated rat liver mitochondria. To understand how this relates to O₂ sufficiency in cell culture, we can compare these values to the levels of O₂ present in media immediately outside of cells (pericellular O₂) or within the cytosol (Table 2).

Pericellular O₂ values for adherent cell monolayers can be measured in real time using a variety of approaches,

TABLE 2: Intracellular O₂ levels under various cell culture conditions.

Cell line and conditions	Extracellular O ₂ (%)	Intracellular O ₂ (%)	References
Confluent, differentiated PC12 cell monolayer	20	~15	
	10	~7	[74]
	6	~1.5	
	20	~17	
HeLa	10	8–9	[75]
	5	2–5	
Undifferentiated PC12 cells	20	~15	
	8	1–2	[76]
Mouse embryonic fibroblasts	20	6–8	[76]
	20	~14	
Mouse embryonic fibroblasts	9	2–3	[9]
	5	~0.5	

including Luxcel and Seahorse platforms (Agilent, USA). We use PreSens OxyDish (PreSens Precision Sensing GMBH, Germany) which has O₂-sensitive fluorescent probes impregnated into the tissue culture dish plastic. For most cell lines seeded at typical densities and maintained in a 5% CO₂ incubator at 37°C without O₂ control, headspace O₂~18% and pericellular O₂ levels are close to this [3]. This is far in excess of what is needed to support maximal mitochondrial respiration rates. When headspace O₂ is maintained at more physiologically relevant levels, however, pericellular and intracellular O₂ levels may become significantly lower, particularly if media changes and/or mixing are infrequent.

Intracellular O₂ levels can be measured in cultured cells using a variety of fluorescent probes (Zhdanov et al. 2012; [9]). Measurements using these tools show that, at the higher levels of O₂ (~20%) typical of standard cell culture, intracellular O₂ levels range from 14–17% depending upon cell type, medium composition (specifically whether fuel source promotes reliance on oxidative phosphorylation), and seeding density (Table 2). Conditions promoting faster rates of O₂ consumption lower these values, but they are generally hyperoxic regardless. At more physiologically relevant headspace O₂ levels of 5–9%, intracellular O₂ levels under typical culture conditions range from 0.5% to 5%. Thus, under standard cell culture conditions, intracellular O₂ levels are typically at least 10 times higher than what is required to sustain maximal mitochondrial respiration rates based on the V_{max} values reported by Hoffman et al. (2009) and others. On the other hand, cell culture at 5–6% headspace O₂ results in intracellular O₂ levels that may be low enough to limit respiration when cells are at high density or growing in respiration-promoting media.

Although a higher than physiological headspace O₂ level in tissue culture helps to ensure that mitochondrial respiration is not oxygen limited, it may have the unintended consequence of stimulating the production of ROS and RNS from various enzymes that are widely expressed in common cell lines. Increases in ROS/RNS with concomitant effects on

redox-sensitive signaling events and potentially oxidative macromolecular damage should be expected under these conditions, and indeed, these are observed. It is therefore important to understand the relationship between oxygen levels and the rates of ROS/RNS production in cultured cells. In mammalian cells, ROS/RNS are produced by a wide range of organelles and enzymes, including mitochondria, NADPH oxidase (Nox), nitric oxide synthase (NOS), monoamine oxidase (MAO), xanthine oxidase/oxidoreductase (XO/XOR), lipoxygenase (LOX), cyclooxygenase (COX), heme oxygenase (HOX), and the transplasma membrane redox system (tPMRS). Here, we discuss how the hyperoxia of standard cell culture is expected to affect the activities and/or expression of all these organelles and enzymes. Our list of oxygen-dependent enzymes is not exhaustive, and we have omitted some oxygen-metabolizing enzymes for which we could not readily find data regarding oxygen sensitivity of reaction rates.

1.2. Oxygen Concentration and Mitochondrial ROS Production. Superoxide/ H_2O_2 production as a byproduct of oxidative phosphorylation has been well studied in isolated mitochondria, and many sites of production, albeit under generally nonphysiological conditions, have been identified (Figure 1). The specific sites of mitochondrial superoxide production have been reviewed recently (e.g., [10]) and will not be detailed here. Superoxide produced at various sites within mitochondria is released into either the matrix or the IMS side of the inner membrane. H_2O_2 arising from superoxide within the mitochondrial matrix can diffuse out of the matrix.

Although mitochondria are often stated to be responsible for the majority of cellular ROS production, this has not been demonstrated [11] and indeed seems unlikely to be universally true given that the total cell volume occupied by mitochondria varies from a few percent in low-metabolic rate cells to as much as 30% in cardiomyocytes [12]. Similarly, the relative levels of other ROS and RNS producers like Nox and NOS vary greatly between cell types and physiological condition. Therefore, while it may be true that mitochondria are the most important sites of ROS production in *some* cell types, they may not be in others. Nonetheless, it is important to consider the sensitivity of mitochondrial ROS production to the oxygen levels prevailing in cells in culture.

Hoffman et al. (2007; 2009) provided detailed measurements and calculations of H_2O_2 production (originating as superoxide) from isolated liver mitochondria respiring in state 4 on various substrates at 37°C (Table 3). Measurements were made over a range of O_2 levels, with and without various respiratory poisons, allowing the calculation of $K_m(O_2)$ values for H_2O_2 production associated with different respiratory substrates and ETC sites. Notably, of the sites contributing to mitochondrial H_2O_2 production in cultured cells, which is likely primarily the complex I FMN site and the complex III Q_o site, both are saturated at the same low O_2 levels as respiration. Note that ETFQOR is probably not a major contributor to mitochondrial H_2O_2 production under standard cell culture conditions because fatty acids are not included as fuels in most commercial media. The

physiological relevance of complex I backflow is established only in ischemia/reperfusion [13], and these would be expected to occur in normal cell culture only as hypoxia-hyperoxia transitions during media changes or passaging. Therefore, based on the above observations, the rate of mitochondrial ROS production is likely not O_2 -limited under most standard cell culture conditions or at head-space O_2 levels as low as 5%, providing that intracellular O_2 remains above 0.5%. Thus, little difference is expected in mitochondrial ROS production between 5% (physioxia) and 18% (standard cell culture) O_2 .

1.3. NADPH Oxidases. There are seven members of the family of nicotinamide adenine dinucleotide phosphate (NADPH) oxidases (Nox1–5 and Duox 1 and 2), several of which (e.g., Nox1, 2, and 4) are quite widely distributed amongst mammalian tissues and cell lines [14]. Noxs are multisubunit membrane-spanning enzymes that transport electrons from NAD(P)H across biological membranes [15]. Although the Nox enzymes localize to a variety of cellular membranes, all can be found in the plasma membrane, where they transport electrons to external O_2 , thus leading to the production of superoxide in the extracellular space [14]. Superoxide produced outside the cell and dismutated to H_2O_2 can diffuse back into the cell of origin or into neighbouring cells.

Importantly, recent work indicates that Nox4 differs from other Nox isoforms in several important ways. Firstly, Nox4 localizes (at least in some cells and/or some physiological conditions) to mitochondria [16] where its activity appears to interact with respiratory complex I and ATP [17, 18]. Nox4 activity is not dependent upon interactions with accessory subunits. Furthermore, Nox4 preferentially produces H_2O_2 rather than superoxide [19]. The rate of Nox4 H_2O_2 production is also very sensitive to O_2 levels in the range between physioxia and 18%.

There is surprisingly little $K_m(O_2)$ data for any of the Nox isoforms. Furthermore, there is some doubt regarding the validity of some assays of Nox activity using isolated membrane fractions [20, 21]. Nonetheless, the available data suggest that the activities of Nox1, Nox2, and Nox4 are all sensitive to O_2 levels in the range from 5% to 18% (Table 4). Nox4, with a $K_m(O_2)$ value~18%, is particularly sensitive over this range; Nox4 activity triples between 3% and 12% O_2 [19]. Direct measurements with cultured PC3 and C2C12 cells indicate a substantial contribution of Nox1 and/or Nox4 to H_2O_2 production in live cells (measured as Amplex Red oxidation), particularly at 18% O_2 , since this value is strongly inhibited by GKT138731 (a Nox1/4 inhibitor; Figure 2). Similarly, in Nox1/2/4 triple knockout mouse dermal fibroblasts, H_2O_2 production at 18% O_2 is only about 1/10 to that in wild-type fibroblasts, and, at 5% O_2 , H_2O_2 production is undetectable (Figure 2). Thus, although somewhat limited, available evidence indicates that Nox isoforms produce H_2O_2 at rates that are strongly dependent upon O_2 levels. This probably underlies observations such as the Nox4 contribution to cellular senescence in primary cell lines, since these studies have been performed at 18% O_2 . Whether Nox4 plays the same roles *in vivo*, where O_2 levels are several times lower, is not clear but must be considered.




FIGURE 1: Sites of H_2O_2 production in the mitochondrial electron transport chain. These include 2-oxoacid dehydrogenase, complex I (site I_F), complex I (site I_Q), complex II (site II_F), complex III (site III_{Q0}), the ETFQOR system (site E_F), glycerol 3-phosphate dehydrogenase (site G_q), and dihydroorotate dehydrogenase (site D_q). Complex I: O00217 (*Homo sapiens*); complex II: Q0QF01 (*Sus scrofa*); complex III: P23004 (*Bos taurus*); complex IV: P68530 (*Bos taurus*); ATP synthase O00217 (*Bos taurus*); cytochrome c: P99999 (*Homo sapiens*); glycerol 3-phosphate: P04406 (*Homo sapiens*); dihydroorotate dehydrogenase: ID3H (*Homo sapiens*); 2-oxoacid dehydrogenase: 2DNE (*Homo sapiens*); ETFQOR: 2GMH (*Sus scrofa*). Data based on Wong et al. [10]. Structures retrieved from <https://www.rcsb.org/>.

TABLE 3: $K_m(\text{O}_2)$ and V_{\max} values for H_2O_2 production from isolated liver mitochondria respiring on various substrates.

Substrates	$K_m(\text{O}_2)$ (%)	H_2O_2 production ($\text{pmol min}^{-1} \cdot \text{mg protein}^{-1}$)
Glutamate + malate with malonate	0.025	250
Succinate (– rotenone)	0.179	330
Succinate (+ rotenone)	0.070	105
Glutamate, malate, and succinate	0.050	330
Palmitoylcarnitine (– rotenone)	0.100	290
Palmitoylcarnitine (+ rotenone)	0.398	250
<i>Sites</i>		
Complex III Q_o site	0.199	150
Complex I FMN site	0.019	170
Complex I electron backflow	0.090	135
ETFQOR	0.498	200

All values are from Hoffman et al. 2009 and measured using an Amplex Red/horseradish peroxidase assay in the presence of superoxide dismutase at 37°C .

1.4. Transplasma Membrane Redox System. The transplasma membrane electron transfer (tPMET) system is a ubiquitous system for transferring electrons from cytosolic NAD(P)H

outside of the cell, similar to the Noxs [22]. The core components of the tPMET system are the NAD(P)H-quinone oxidoreductase (NQO1), NADH-cytochrome b5 reductase (CytB5red), coenzyme Q_{10} , and the ecto-NADH oxidase disulfide thiol exchanger (ENox). Both NQO1 and CytB5red are distributed in multiple intracellular localizations, including in association with the plasma membrane.

The tPMET system makes significant contributions to cellular O_2 consumption in many common cell lines (e.g., Jurkat, RAW264.7, and pancreatic beta cells; see [23, 24]). The tPMET system is highly upregulated in respiration-deficient (ρ^0) cells, which accumulate coenzyme Q_{10} in the plasma membrane [25]. In ρ^0 cells, the tPMRS may be the predominant site of O_2 consumption [24].

A variety of extracellular terminal electron acceptors are possible, including O_2 , which can undergo single-electron reduction to produce superoxide. Purified mammalian CytB5red produces superoxide directly [26], though its overexpression has beneficial effects in some specific contexts [27, 28]. NQO1 is associated with an intracellular antioxidant function (e.g., [29]) but contributes to reduction of extracellular electron acceptors, presumably including O_2 , in pancreatic beta cells [23]. ENox proteins include several isoforms—the age-related arNox may be the most relevant in terms of ROS production in cell culture. This isoform produces superoxide and becomes more highly expressed in the tissues of aged mammals and in late-passage or senescent cells in culture [30].

TABLE 4: $K_m(O_2)$ values for various O_2 -consuming enzymes.

Enzyme	Product	$K_m(O_2)$ (%)	Details	Reference
Nox2	O_2^-	3.5		[77]
Nox2	O_2^-	3.1		[78]
Nox2	O_2^-	2-3		[19]
Nox2	O_2^-	2-3		[79]
Nox4	H_2O_2	18		[78]
Nox4	H_2O_2	18		[19]
Nox4	H_2O_2	16–20		[79]
nNOS	NO	2.3	Purified bovine enzyme	[45]
nNOS	NO/ O_2^-	2.2	Partially uncoupled rat enzyme	[42, 43]
nNOS	O_2^-	3.4	Fully uncoupled rat enzyme	
nNOS	NO	15.7	Purified rat enzyme	[47]
nNOS	NO	39.8	Purified rat enzyme	[80]
nNOS	NO	28.5	Kinetic model based on rat enzyme data	[81, 82]
eNOS	NO	0.8	Purified bovine enzyme	[45]
eNOS	NO	0.3	Kinetic model based on rat enzyme data	[82, 83]
iNOS	NO	0.63 ± 0.09	Purified bovine enzyme	[45]
iNOS	NO	11.0	Isolated enzyme assay	[84]
iNOS	NO	10.6	Kinetic model	[82]
NOS	NO	3.1–10.8	Unspecified isoform	[48, 85]
XO	O_2^-/H_2O_2	2.2–6.8	Isolated from bovine milk	Fridovich et al. 1962, 1964
MAO	H_2O_2	3.4–28	Mammalian enzymes; various substrates	[86–88]; reviewed in [57]
HO	CO, iron, and bilirubin	<1.5	Unspecified isoform from chicken liver	[59]
LOX	H_2O_2	1–2.6	Unspecified isoform	[65]
COX-1	Various	0.4–3.1	Arachidonic acid substrate	[63, 89–91]
COX-2	Various	1.3–1.5	Arachidonic acid substrate	[63, 65]
PHD	Modified HIF	41–46	Various substrates	[92]
FIH	Modified HIF	4–12	Various substrates	[92]

Nox: NADPH oxidase; NOS: nitric oxide synthase; XO: xanthine oxidase; MAO: monoamine oxidase; HO: heme oxygenase; LOX: lipoxygenase; COX: cyclooxygenase; PHD: prolyl hydroxylase; FIH: factor inhibiting HIF-1.

The tPMET system has only been measured under standard cell culture O_2 (18%), and we are not aware of any data regarding the sensitivity of superoxide production to O_2 concentration for the system as a whole or for individual components of the system. Nonetheless, consideration should be given to the possibility that the tPMRS is O_2 sensitive in the 5–18% range and that O_2 -dependent changes in its activity could affect cell physiology.

1.5. Role of Aquaporins in Transmembrane H_2O_2 and Gas Diffusion. The tPMRS and all Nox isoforms (a portion of all Nox isoforms localizes to the plasma membrane) can produce either superoxide or H_2O_2 on the extracellular side of the plasma membrane. Here, these ROS may react with membranes or membrane-bound proteins facing the extracellular space on the originating cell or on neighbouring cells. Alternatively, H_2O_2 , either directly produced or resulting from superoxide dismutation, may cross cell membranes to exert intracellular effects. The H_2O_2 permeability of phospholipid bilayers is limited; indeed, H_2O_2 is less membrane permeant than H_2O [31]. However, H_2O_2 rapidly equilibrates across cell membranes via aquaporins (AQPs). Structural studies

suggest that all water-transporting AQPs may facilitate some degree of H_2O_2 movement across cellular membranes [32]. However, empirical data is less equivocal, suggesting that hAQP3 and hAQP8 are particularly good H_2O_2 transporters [31].

Interestingly, O_2 levels modulate the expression of several AQPs. AQP1 and AQP3 mRNA and protein levels are both increased at 1% versus 20% O_2 ([33]; Hoogewijs et al. 2016), though AQP5 and AQP9 expressions are reduced at lower O_2 levels ([34]; Castro-Parodi et al. 2013). There is also some evidence that hypoxia can affect AQP8 permeability directly via posttranslational modifications [35]. Recent studies ([36]; Zwiazek et al. 2017) also suggest that some AQP isoforms facilitate the transport of O_2 and therefore may facilitate O_2 uptake by cells at low O_2 levels. There is also strong evidence that AQP1 and AQP4 can transport other important ROS/RNS, such as NO [37, 38]. Taken together, these observations indicate that media O_2 levels are likely to modulate the H_2O_2 , NO, and O_2 permeability of cell membranes via their effects on AQP expression and/or activity. This makes it more difficult to understand the detailed kinetics of how ROS/RNS and gases traverse cellular membranes




FIGURE 2: Cells produce more H₂O₂ at 18% versus 5% O₂, but this is prevented or ameliorated by selective NADPH oxidase 1/4 inhibitor GKT137831. (a) C2C12 mouse myoblasts, (b) PC-3 human prostate cancer cells, and (c) wild-type and NOX1/2/4 triple knockout mouse dermal fibroblast cell lines were grown at 18% O₂ in a humidified 37°C CO₂ incubator, at 5%, and assayed at either 5% or 18% O₂ as in Maddalena et al. [3]. H₂O₂ efflux from cells was measured using Amplex Red reagent (10-acetyl-3,7-dihydroxyphenoxazine). Krebs-Ringer buffer (KRB) was used for the assays which consisted of 135 mM NaCl, 5 mM KCl, 1 mM MgSO₄, 0.4 mM K₂HPO₄, 20 mM HEPES, 5.5 mM glucose, and 10% fetal bovine serum. During the experiment, cells were incubated in KRB-contained 50 mM Amplex Red reagent and 0.1 units/mL horseradish peroxidase enzyme in the presence of 5 μM GKT 137831 or vehicle control. Data were analysed using two-tailed *t*-tests. Bars represent the mean ± SEM from at least five independent experiments. **p* < 0.05.

in vivo when measurements are done *in vitro* under nonphysiological O₂ conditions.

1.6. Nitric Oxide Synthases. NOS are a family of enzymes responsible for the production of NO and L-citrulline from L-arginine and O₂ [39, 40]. Three isoforms of NOS have been identified: endothelial NOS (eNOS), neuronal NOS (nNOS), and inducible NOS (iNOS) [41]. These three NOS isoforms share approximately 50% amino acid identity [41], and all three are widely expressed in mammalian tissues and cell lines. Under some conditions, including transient anoxia/reoxygenation, NOS catalyzes an “uncoupled” reaction producing superoxide instead of NO (Stuehr et al. 1991; [42, 43]; reviewed in [44]).

The first detailed study of the O₂ sensitivity of the NOS reaction was by Rengasamy and Johns [45] who calculated a K_m(O₂) value of 0.63 to 2.31% O₂ for NOS isolated from bovine brain and aortic endothelium as well as RAW 264.7 mouse macrophages. Subsequently, K_m(O₂) values have been published for various tissues and cell lines, as well as for specific purified NOS isoforms (Table 4). Although these values vary widely, presumably due to differences in experimental conditions, O₂ affinities of nNOS and iNOS are found relatively consistently to be within a range that is sensitive to

changes in O₂ levels between physioxia and 18% O₂. Cell lines with high levels of these two isoforms will therefore produce more NO under standard cell culture conditions than *in vivo*. Cell culture models of ischemia/reperfusion injury typically employ a period of near anoxia followed by return to “normoxia” where the latter is 18% O₂. NOS activity can be uncoupled under these conditions, and the rate of superoxide production upon reoxygenation at 18% O₂ likely exceeds that which would occur upon a return to physioxia *in vivo*.

NO can also be produced by the reduction of nitrite catalyzed by several enzymes including XO/XOR [46]. This reaction is promoted at lower O₂ levels and makes it difficult to predict the effect of O₂ levels on NO production rates. Thus, the relationship between cellular NO synthesis rates and O₂ levels will depend on the relative abundance of different NOS isoforms, which have different sensitivities to O₂.

In addition to its effects on NO synthesis, O₂ levels influence NO metabolism ([47, 48]; reviewed in [49]). NO is metabolized in cells by poorly characterized pathways. However, it is known that the rate of NO metabolism by cells is faster at higher O₂ levels [50]. Thus, higher O₂ levels will affect the rate of NO production from nNOS and iNOS while simultaneously increasing the rate of their metabolism.

Again, it is difficult to predict how this will affect steady-state NO levels in all cells. In activated RAW 264.7 cells in culture, the maximum rates of NO production were observed at 8% O₂ [50], though the relationship between O₂ and NO levels in other cell types is unknown. The O₂-dependent rate of NO metabolism will in turn affect its diffusional distance and therefore alter the subset of proteins modified in the cell of origin or neighbouring cells. NO participates in many regulatory posttranslational modifications of key proteins, including those driving epigenetic modifications. Given these diverse influences of NO on cellular functions and the effects of O₂ levels on synthesis, diffusion distance, and metabolism of NO, there is clear potential to generate nonphysiologically relevant results at nonphysiological O₂ levels.

1.7. Other Oxidases. XO and MAO are two widely expressed cellular oxidases whose products include superoxide and H₂O₂. XO and its precursor XOR are expressed in mammalian cells, where they localize to the cytosol and the external face of the plasma membrane [51]. Both XO and XOR can catalyze the oxidation of purines, producing superoxide radical and H₂O₂ [52]. Under some conditions, XO may also catalyze NO production from nitrites and nitrates, a reaction that is favoured at lower O₂ [46]. However, the levels of these latter compounds are relatively low in culture media and this activity may therefore be of minor importance in mammalian cell culture. The K_m(O₂) value of bovine XO has been reported in the range of 3–6% O₂ (Table 4). Interestingly, the relative production of superoxide versus H₂O₂ by XO is also sensitive to O₂ levels within the range found in cell culture (1–21%), such that at O₂ levels below 5% O₂, H₂O₂ formation is promoted [53]. In addition, high O₂ levels promote posttranslational modifications of XOR that decrease its specific activity [54, 55]. Thus, O₂ levels in culture could affect the relative rates of ROS production by XO/XOR as well as the relative amounts of superoxide versus H₂O₂ produced.

MAO occurs as two isoforms, MAO-A and MAO-B. Both are widely distributed in mammalian tissues [56]. They localize to the outer mitochondrial membrane where they catalyze the degradation of biogenic and dietary monoamines such as norepinephrine, dopamine, tyramine, serotonin, phenylethylamine, and benzylamine in a reaction producing H₂O₂. The reported K_m(O₂) values of both isoforms vary widely (Table 4) due to a complicated reaction mechanism in which O₂ affinity is strongly affected by monoamine concentrations (reviewed in [57]). MAO enzyme activities are nonetheless predicted to be sensitive to O₂ over the range of 5–18%. Because the H₂O₂ produced by MAOs is near the mitochondrial compartment, aberrant mitochondrial and cytosolic redox signaling and/or macromolecular damage may be caused by MAO enzymes in the O₂ conditions typical of standard cell culture.

1.8. Oxygenases. Oxygenases catalyze the incorporation of oxygen into an organic substrate. Heme oxygenase (HOX), lipoxygenase (LOX), and cyclooxygenase (COX) are all widely expressed in mammalian tissues and cell lines. HOX is important in the process of heme degradation, while both

COX and LOX assist in the breakdown of arachidonic acid via two separate pathways.

HOX is localized in the endoplasmic reticulum (reviewed in [58]). There are two isoforms, HOX-1 and HOX-2, sharing an amino acid sequence identity of >45%. While HOX-2 is constitutively expressed, HOX-1 is induced by endogenous and exogenous stressors such as heavy metals, pharmacological agents, inflammatory mediators, UV light, and oxidative stress. HOX enzymes oxidize heme to produce carbon monoxide, Fe²⁺, and biliverdin. HOX purified from chicken liver has maximal activity at O₂ levels as low as 1.5% ([59]; Table 4), indicating that this enzyme is likely O₂ saturated even in cells cultured in physioxia.

COX is a dioxygenase that catalyzes the first step of arachidonic acid or linoleic acid breakdown leading to the production of prostanoid derivatives (reviewed in [60]). COX enzymes modulate cell growth and signaling pathways and are implicated in cancer progression [61]. There are two isoforms, COX-1 and COX-2, with an amino acid sequence homology of ~60%. COX enzymes localize to the endoplasmic reticulum membrane and nuclear envelope. These enzymes catalyze two-step reactions. In the first step, cyclooxygenase activity using molecular oxygen produces the hydroperoxide prostaglandin G₂ intermediate. Peroxidase activity is then observed resulting in the formation of prostaglandin H₂. The K_m(O₂) value of COX-1-catalyzed arachidonic acid oxidation varies between 0.4% and 3.1% O₂, while the K_m(O₂) value of COX-2 is somewhat lower (Table 4). Activities of COX-1 and COX-2 saturate at around 10–20% O₂ [62, 63]. Thus, these enzymes are also O₂ sensitive in the range of interest, and as their products modulate various cellular activities including cell growth and differentiation [61], it is possible that their increased activities at 18% O₂ affect studies of cell physiology in culture.

LOX catalyzes the deoxygenation of polyunsaturated fatty acids, producing a variety of oxygen and lipid radicals under some conditions (reviewed in [60]). There are six isoforms of LOX that are widely distributed in mammalian tissues and cells. LOX enzymes produce ROS and lipid radical species that participate in intracellular signaling pathways. O₂ levels affect the lipoxygenase reaction in complex ways [64, 65], but the measured K_m(O₂) values are 1–2.6% O₂ (Table 4), making reaction rates O₂ sensitive in the relevant range.

1.9. Transcriptional Regulation of ROS/RNS-Producing Enzymes. In addition to acute effects on the activities of enzyme-catalyzed reactions, O₂ levels in cell culture may affect the expression of various O₂-consuming enzymes and organelles. HIF-1 and HIF-2 are heterodimeric transcription factors whose activities are regulated by O₂ via the degradation of the α -subunits (HIF-1 α , HIF-2 α). This reaction is catalyzed by prolyl hydroxylase (PHD), which uses O₂ and 2-oxoglutarate to hydroxylate the HIF- α subunits leading to their subsequent ubiquitination and degradation. A second reaction, hydroxylation of an asparagine in HIF-1 α catalyzed by the factor inhibiting HIF-1 (FIH), inhibits HIF-1 transcriptional activity. Together, O₂ regulates the levels and activity of HIF-1 and/or HIF-2. Both hydroxylation reactions




FIGURE 3: Reduced levels of NADPH oxidases 1 and 4 at 18% versus 5% O₂. (a) Representative Western blots showing Nox1 and β-tubulin in PC-3 cells or Nox4 and β-tubulin in C2C12 cells, at 5% and 18% O₂. (b, c) Average Nox1 signal (b) or Nox4 signal (c) standardized to β-tubulin. Total cellular proteins were extracted by treating cells with NP-40 buffer (150 mM NaCl, 1% NP-40, 50 mM Tris-HCl pH 8.0). Total protein (15 μg per sample) was resolved on 10% SDS-PAGE, transferred to a polyvinylidene difluoride membrane, and probed for Nox1 or Nox4. β-Tubulin was used as an internal loading and transfer control. All antibodies were purchased from Novus Biologicals; Nox1 (NBPI-31546), Nox4 (NB110-5885), β-tubulin (NB600-936). Data were analysed using two-tailed Student's *t*-tests. Bars represent means ± SEM from at least five independent experiments. **p* < 0.05. The identities of bands at intermediate molecular weight are unknown.

are O₂ sensitive within the physioxia to 18% O₂ range. Mammalian PHD isoforms have a K_m(O₂) value ~25% O₂ and FIH K_m(O₂) value ~11% (Table 4). In addition, PHD activities may be modulated by ROS/RNS produced by many of the enzymes discussed above at higher rates under standard culture conditions.

Hundreds of genes are transcriptionally regulated by HIFs, and interestingly, most of the ROS/RNS-producing enzymes discussed above are amongst them. In our experiments, the relative levels of Nox1 and Nox4 are much higher at 5% compared to 18% O₂ (Figure 3). Indeed, Nox4, all three NOS isoforms, LOX, COX, MAO, and HOX are all transcriptional targets of HIFs (Table 5). In addition, several AQP are HIF regulated, indicating that the H₂O₂ permeability of cellular membranes is likely different at physioxia versus 18% O₂. It is important to note that virtually, all data regarding the HIF regulation of these enzymes has been collected using 18% O₂ as “normoxia” and 1% O₂ as hypoxia but similar results are expected for 5% versus 18% comparison.

In terms of the impact on cell culture experiments, the effects of O₂ on the specific activities (based on K_m(O₂) values) versus levels of O₂-consuming proteins (based on transcriptional upregulation) would tend to oppose each other, though our measurement of higher rates of cellular ROS production at 18% O₂ versus 5% O₂ suggests that they do not cancel. It is unclear to what extent the combined effects of media O₂ on acute metabolic flux through the various O₂-consuming pathways and the transcriptional regulation via HIF-1/2 activation of their component proteins, would impact cell physiology. Indeed, it seems impossible to predict how the hyperoxia of cell culture will impact the numerous O₂-sensitive, O₂-consuming, metabolic reactions overall given these concomitant changes in specific activity and expression.

1.10. Conclusions. Under standard cell culture conditions, media O₂ levels of typically ~18% are hyperoxic with respect to the 1–6% experienced by most mammalian cells *in vivo*

TABLE 5: Hypoxia effects on the expression of some O₂-consuming enzymes.

Enzyme	Effect	Reference
Nox1	Transcriptional stimulation by HIF-1	Goyal et al. 2004
Nox4	Transcriptional stimulation	[93]
nNOS	Transcriptional stimulation	[94]
eNOS	Transcriptional stimulation Stabilization of mRNA	[95] [96]
iNOS	Transcriptional stimulation by HIF-1	[97–99]
XO	Transcriptional stimulation	[54]
MAO	Transcriptional stimulation	[100]
HO	Transcriptional stimulation	[101]
LOX	Transcriptional stimulation	[102]
COX	Transcriptional stimulation	Demasi et al. 2004;
AQP1	Transcriptional stimulation	[103]
AQP3	Transcriptional stimulation	Hoogewijs et al. 2016
AQP5	Transcriptional repression	[34]
AQP9	Unclear mechanism	Castro-Parodi et al. 2013

Nox: NADPH oxidase; NOS: nitric oxide synthase; XO: xanthine oxidase; MAO: monoamine oxidase; HO: heme oxygenase; LOX: lipoxygenase; COX: cyclooxygenase; AQP: aquaporin.

(Table 1). This disparity has important consequences for the accurate modeling of *in vivo* cell physiology, so it is important to have a comprehensive understanding of how O₂ affects specific ROS/RNS-producing processes. Although rates of ROS production from mitochondrial respiration are relatively insensitive to increases in O₂ levels from 5% to 18% O₂, many widespread O₂-consuming cellular enzymes are very sensitive in this same range. In particular, the activities of Nox4, nNOS, eNOS, and both MAO isoforms appear to be strongly induced at higher O₂ levels. These enzymes will thus produce substantially more ROS/RNS, potentially affecting the states of intracellular signaling pathways and the downstream events they are regulating. This is unlikely to be an appropriate starting point upon which to impose further stresses and expect a “normal” physiological response that mimics an *in vivo* context.

In addition to acute effects of O₂ on flux through specific metabolic pathways, there is evidence that chronic exposure to the hyperoxia of cell culture will also affect the expression of Nox, NOS, MAO, and other O₂-consuming enzymes (Table 5). Presumably, this is a homeostatic mechanism for tuning these reactions to O₂ availability. However, it again establishes a baseline condition that may not accurately model the *in vivo* state. Even the permeability of cellular membranes to H₂O₂, NO, or O₂ may be influenced by O₂-mediated transcriptional regulation of AQPs, given their role in facilitating the diffusion of these molecules (Table 5).

The specific effects of O₂ on cellular ROS/RNS production described above point to the importance of culturing cells at physiologically relevant O₂ levels. In most cases, this requires lowering the incubator headspace O₂ levels. However, this

can cause pericellular and intracellular hypoxia, depending upon cell density and mitochondrial respiration rates. Therefore, it is further necessary to monitor pericellular O₂ levels and take steps to maintain them within a physiologically relevant range. We have found that standing O₂ gradients from the top of the media column to the pericellular region are present in undisturbed cell culture and that regular gentle mixing (e.g., via a rocker plate) can abolish these. Our simple solution to this problem will be presented in a future publication.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

References

- [1] E. O. Pettersen, L. H. Larsen, N. B. Ramsing, and P. Ebbesen, “Pericellular oxygen depletion during ordinary tissue culturing, measured with oxygen microsensors,” *Cell Proliferation*, vol. 38, no. 4, pp. 257–267, 2005.
- [2] R. H. Wenger, V. Kurtcuoglu, C. C. Scholz, H. H. Marti, and D. Hoogewijs, “Frequently asked questions in hypoxia research,” *Hypoxia*, vol. 3, pp. 35–43, 2015.
- [3] L. A. Maddalena, S. M. Selim, J. Fonseca, H. Messner, S. McGowan, and J. A. Stuart, “Hydrogen peroxide production is affected by oxygen levels in mammalian cell culture,” *Biochemical and Biophysical Research Communications*, vol. 493, no. 1, pp. 246–251, 2017.
- [4] M. Zscharnack, C. Poesel, J. Galle, and A. Bader, “Low oxygen expansion improves subsequent chondrogenesis of ovine bone-marrow-derived mesenchymal stem cells in collagen type I hydrogel,” *Cells Tissues Organs*, vol. 190, no. 2, pp. 81–93, 2009.
- [5] J. Milosevic, S. C. Schwarz, K. Krohn, M. Poppe, A. Storch, and J. Schwarz, “Low atmospheric oxygen avoids maturation, senescence and cell death of murine mesencephalic neural precursors,” *Journal of Neurochemistry*, vol. 92, no. 4, pp. 718–729, 2005.
- [6] V. Renault, L. E. Thornell, G. Butler-Browne, and V. Mouly, “Human skeletal muscle satellite cells: aging, oxidative stress and the mitotic clock,” *Experimental Gerontology*, vol. 37, no. 10–11, pp. 1229–1236, 2002.
- [7] M. Weiszenstein, N. Pavlikova, M. Elkalaf et al., “The effect of pericellular oxygen levels on proteomic profile and lipogenesis in 3T3-L1 differentiated preadipocytes cultured on gas-permeable cultureware,” *PLoS One*, vol. 11, no. 3, p. e0152382, 2016.
- [8] N. Haque, M. T. Rahman, N. H. Abu Kasim, and A. M. Alabsi, “Hypoxic culture conditions as a solution for mesenchymal stem cell based regenerative therapy,” *The Scientific World Journal*, vol. 2013, Article ID 632972, 12 pages, 2013.
- [9] R. I. Dmitriev, A. V. Zhdanov, G. Jasionek, and D. B. Papkovsky, “Assessment of cellular oxygen gradients with a panel of phosphorescent oxygen-sensitive probes,” *Analytical Chemistry*, vol. 84, no. 6, pp. 2930–2938, 2012.
- [10] H. S. Wong, P. A. Dighe, V. Mezera, P. A. Monternier, and M. D. Brand, “Production of superoxide and hydrogen peroxide from specific mitochondrial sites under different

- bioenergetic conditions,” *Journal of Biological Chemistry*, vol. 292, no. 41, pp. 16804–16809, 2017.
- [11] G. C. Brown and V. Borutaite, “There is no evidence that mitochondria are the main source of reactive oxygen species in mammalian cells,” *Mitochondrion*, vol. 12, no. 1, pp. 1–14, 2012.
- [12] J. A. Stuart, L. A. Maddalena, M. Merilovich, and E. L. Robb, “A midlife crisis for the mitochondrial free radical theory of aging,” *Longevity & Healthspan*, vol. 3, no. 1, p. 4, 2014.
- [13] E. T. Chouchani, V. R. Pell, E. Gaude et al., “Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS,” *Nature*, vol. 515, no. 7527, pp. 431–435, 2014.
- [14] A. Panday, M. K. Sahoo, D. Osorio, and S. Batra, “NADPH oxidases: an overview from structure to innate immunity-associated pathologies,” *Cellular & Molecular Immunology*, vol. 12, no. 1, pp. 5–23, 2015.
- [15] K. Bedard and K.-H. Krause, “The Nox family of ROS-generating NADPH oxidases: physiology and pathophysiology,” *Physiological Reviews*, vol. 87, no. 1, pp. 245–313, 2007.
- [16] K. Block, Y. Gorin, and H. E. Abboud, “Subcellular localization of Nox4 and regulation in diabetes,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 106, no. 34, pp. 14385–14390, 2009.
- [17] K. Shanmugasundaram, B. K. Nayak, W. E. Friedrichs, D. Kaushik, R. Rodriguez, and K. Block, “NOX4 functions as a mitochondrial energetic sensor coupling cancer metabolic reprogramming to drug resistance,” *Nature Communications*, vol. 8, no. 1, p. 997, 2017.
- [18] R. Kozziel, H. Pircher, M. Kratochwil et al., “Mitochondrial respiratory chain complex I is inactivated by NADPH oxidase Nox4,” *Biochemical Journal*, vol. 452, no. 2, pp. 231–239, 2013.
- [19] Y. Nisimoto, B. A. Diebold, D. Cosentino-Gomes, and J. D. Lambeth, “Nox4: a hydrogen peroxide-generating oxygen sensor,” *Biochemistry*, vol. 53, no. 31, pp. 5111–5120, 2014.
- [20] F. Rezende, O. Lowe, V. Helfinger et al., “Unchanged NADPH oxidase activity in Nox1-Nox2-Nox4 triple knockout mice: what do NADPH-stimulated chemiluminescence assays really detect?,” *Antioxidants & Redox Signaling*, vol. 24, no. 7, pp. 392–399, 2016.
- [21] F. Rezende, K.-K. Prior, O. Lowe et al., “Cytochrome P450 enzymes but not NADPH oxidases are the source of the NADPH-dependent lucigenin chemiluminescence in membrane assays,” *Free Radical Biology & Medicine*, vol. 102, pp. 57–66, 2017.
- [22] P. Navas, J. M. Villalba, and R. de Cabo, “The importance of plasma membrane coenzyme Q in aging and stress responses,” *Mitochondrion*, vol. 7, pp. S34–S40, 2007.
- [23] J. P. Gray, T. Eisen, G. W. Cline, P. J. S. Smith, and E. Heart, “Plasma membrane electron transport in pancreatic β -cells is mediated in part by NQO1,” *American Journal of Physiology-Endocrinology and Metabolism*, vol. 301, no. 1, pp. E113–E121, 2011.
- [24] P. M. Herst and M. V. Berridge, “Cell surface oxygen consumption: a major contributor to cellular oxygen consumption in glycolytic cancer cell lines,” *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, vol. 1767, no. 2, pp. 170–177, 2007.
- [25] C. Gómez-Díaz, J. M. Villalba, R. Pérez-Vicente, F. L. Crane, and P. Navas, “Ascorbate stabilization is stimulated in ρ^0 HL-60 cells by CoQ₁₀ increase at the plasma membrane,” *Biochemical and Biophysical Research Communications*, vol. 234, no. 1, pp. 79–81, 1997.
- [26] A. K. Samhan-Arias and C. Gutierrez-Merino, “Purified NADH-cytochrome b_5 reductase is a novel superoxide anion source inhibited by apocynin: sensitivity to nitric oxide and peroxyinitrite,” *Free Radical Biology & Medicine*, vol. 73, pp. 174–189, 2014.
- [27] D.-H. Hyun and G.-H. Lee, “Cytochrome b_5 reductase, a plasma membrane redox enzyme, protects neuronal cells against metabolic and oxidative stress through maintaining redox state and bioenergetics,” *Age*, vol. 37, no. 6, p. 122, 2015.
- [28] A. Martin-Montalvo, Y. Sun, A. Diaz-Ruiz et al., “Cytochrome b_5 reductase and the control of lipid metabolism and healthspan,” *npj Aging and Mechanisms of Disease*, vol. 2, no. 1, article 16006, 2016.
- [29] D.-H. Hyun, S. S. Emerson, D.-G. Jo, M. P. Mattson, and R. de Cabo, “Calorie restriction up-regulates the plasma membrane redox system in brain cells and suppresses oxidative stress during aging,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 103, no. 52, pp. 19908–19912, 2006.
- [30] D. James Morr e and D. M. Morr e, *Ecot-Nox Proteins: Growth, Cancer, and Aging*, Springer Press, New York, NY, USA, 2013.
- [31] G. P. Bienert and F. Chaumont, “Aquaporin-facilitated transmembrane diffusion of hydrogen peroxide,” *Biochimica et Biophysica Acta (BBA) - General Subjects*, vol. 1840, no. 5, pp. 1596–1604, 2014.
- [32] A. Almasalmeh, D. Krenc, B. Wu, and E. Beitz, “Structural determinants of the hydrogen peroxide permeability of aquaporins,” *The FEBS Journal*, vol. 281, no. 3, pp. 647–656, 2014.
- [33] I. Abreu-Rodriguez, R. Sanchez Silva, A. P. Martins et al., “Functional and transcriptional induction of aquaporin-1 gene by hypoxia; analysis of promoter and role of HIF-1 α ,” *PLoS One*, vol. 6, no. 12, article e28385, 2011.
- [34] J. D. Kawedia, F. Yang, M. A. Sartor, D. Gozal, M. Czyzyk-Krzaska, and A. G. Menon, “Hypoxia and hypoxia mimetics decrease aquaporin 5 (AQP5) expression through both hypoxia inducible factor-1 α and proteasome-mediated pathways,” *PLoS One*, vol. 8, no. 3, article e57541, 2013.
- [35] I. Medrano-Fernandez, S. Bestetti, M. Bertolotti et al., “Stress regulates aquaporin-8 permeability to impact cell growth and survival,” *Antioxidants & Redox Signaling*, vol. 24, no. 18, pp. 1031–1044, 2016.
- [36] Y. Wang and E. Tajkhorshid, “Nitric oxide conduction by the brain aquaporin AQP4,” *Proteins: Structure, Function, and Bioinformatics*, vol. 78, no. 3, pp. 661–670, 2010.
- [37] M. Herrera and J. L. Garvin, “Aquaporins as gas channels,” *Pflügers Archiv - European Journal of Physiology*, vol. 462, no. 4, pp. 623–630, 2011.
- [38] R. D. Fontijn, O. L. Volger, T. C. van der Pouw-Kraan et al., “Expression of nitric oxide-transporting aquaporin-1 is controlled by KLF2 and marks non-activated endothelium in vivo,” *PLoS One*, vol. 10, no. 12, article e0145777, 2015.
- [39] O. W. Griffith and D. J. Stuehr, “Nitric oxide synthases: properties and catalytic mechanism,” *Annual Review of Physiology*, vol. 57, no. 1, pp. 707–734, 1995.
- [40] M. A. Marletta, “Nitric oxide synthase structure and mechanism,” *Journal of Biological Chemistry*, vol. 268, no. 17, pp. 12231–12234, 1993.

- [41] W. K. Alderton, C. E. Cooper, and R. G. Knowles, "Nitric oxide synthases: structure, function and inhibition," *Biochemical Journal*, vol. 357, no. 3, pp. 593–615, 2001.
- [42] Y. T. Gao, S. P. Panda, L. J. Roman, P. Martásek, Y. Ishimura, and B. S. S. Masters, "Oxygen metabolism by neuronal nitric-oxide synthase," *Journal of Biological Chemistry*, vol. 282, no. 11, pp. 7921–7929, 2007.
- [43] Y. T. Gao, L. J. Roman, P. Martásek, S. P. Panda, Y. Ishimura, and B. S. S. Masters, "Oxygen metabolism by endothelial nitric-oxide synthase," *Journal of Biological Chemistry*, vol. 282, no. 39, pp. 28557–28565, 2007.
- [44] N. D. Roe and J. Ren, "Nitric oxide synthase uncoupling: a therapeutic target in cardiovascular diseases," *Vascular Pharmacology*, vol. 57, no. 5-6, pp. 168–172, 2012.
- [45] A. Rengasamy and R. A. Johns, "Determination of K_m for oxygen of nitric oxide synthase isoforms," *Journal of Pharmacology and Experimental Therapeutics*, vol. 276, no. 1, pp. 30–33, 1996.
- [46] N. Cantu-Medellin and E. E. Kelley, "Xanthine oxidoreductase-catalyzed reduction of nitrite to nitric oxide: Insights regarding where, when and how," *Nitric Oxide*, vol. 34, pp. 19–26, 2013.
- [47] I. M. Elayan, M. J. Axley, P. V. Prasad, S. T. Ahlers, and C. R. Aufer, "Effect of hyperbaric oxygen treatment on nitric oxide and oxygen free radicals in rat brain," *Journal of Neurophysiology*, vol. 83, no. 4, pp. 2022–2029, 2000.
- [48] C. C. McCormick, W. P. Li, and M. Calero, "Oxygen tension limits nitric oxide synthesis by activated macrophages," *Biochemical Journal*, vol. 350, no. 3, pp. 709–716, 2000.
- [49] D. D. Thomas, "Breathing new life into nitric oxide signaling: a brief overview of the interplay between oxygen and nitric oxide," *Redox Biology*, vol. 5, pp. 225–233, 2015.
- [50] J. R. Hickok, D. Vasudevan, K. Jablonski, and D. D. Thomas, "Oxygen dependence of nitric oxide-mediated signaling," *Redox Biology*, vol. 1, no. 1, pp. 203–209, 2013.
- [51] M. G. Battelli, L. Polito, M. Bortolotti, and A. Bolognesi, "Xanthine oxidoreductase-derived reactive species: physiological and pathological effects," *Oxidative Medicine and Cellular Longevity*, vol. 2016, Article ID 3527579, 8 pages, 2016.
- [52] N. Cantu-Medellin and E. E. Kelley, "Xanthine oxidoreductase-catalyzed reactive species generation: a process in critical need of reevaluation," *Redox Biology*, vol. 1, no. 1, pp. 353–358, 2013.
- [53] E. E. Kelley, N. K. H. Khoo, N. J. Hundley, U. Z. Malik, B. A. Freeman, and M. M. Tarpey, "Hydrogen peroxide is the major oxidant product of xanthine oxidase," *Free Radical Biology & Medicine*, vol. 48, no. 4, pp. 493–498, 2010.
- [54] E. E. Kelley, T. Hock, N. K. H. Khoo et al., "Moderate hypoxia induces xanthine oxidoreductase activity in arterial endothelial cells," *Free Radical Biology and Medicine*, vol. 40, no. 6, pp. 952–959, 2006.
- [55] N. Linder, E. Martelin, R. Lapatto, and K. O. Raivio, "Post-translational inactivation of human xanthine oxidoreductase by oxygen under standard cell culture conditions," *American Journal of Physiology-Cell Physiology*, vol. 285, no. 1, pp. C48–C55, 2003.
- [56] D. E. Edmondson, "Hydrogen peroxide produced by mitochondrial monoamine oxidase catalysis: biological implications," *Current Pharmaceutical Design*, vol. 20, no. 2, pp. 155–160, 2014.
- [57] K. F. Tipton, "90 years of monoamine oxidase: some progress and some confusion," *Journal of Neural Transmission*, pp. 1–33, 2018.
- [58] J. Munoz-Sanchez and M. E. Chanez-Cardenas, "A review on hemeoxygenase-2: focus on cellular protection and oxygen response," *Oxidative Medicine and Cellular Longevity*, vol. 2014, Article ID 604981, 16 pages, 2014.
- [59] H. L. Bonkovsky, J. F. Healey, and J. Pohl, "Purification and characterization of heme oxygenase from chick liver: comparison of the avian and mammalian enzymes," *European Journal of Biochemistry*, vol. 189, no. 1, pp. 155–166, 1990.
- [60] C. Schneider, D. A. Pratt, N. A. Porter, and A. R. Brash, "Control of oxygenation in lipoxygenase and cyclooxygenase catalysis," *Chemistry & Biology*, vol. 14, no. 5, pp. 473–488, 2007.
- [61] C. Sobolewski, C. Cerella, M. Dicato, L. Ghibelli, and M. Diederich, "The role of cyclooxygenase-2 in cell proliferation and cell death in human malignancies," *International Journal of Cell Biology*, vol. 2010, Article ID 215158, 21 pages, 2010.
- [62] Y. Liu and J. P. Roth, "A revised mechanism for human cyclooxygenase-2," *Journal of Biological Chemistry*, vol. 291, no. 2, pp. 948–958, 2016.
- [63] J. Qiao, L. Ma, J. Roth, Y. Li, and Y. Liu, "Kinetic basis for the activation of human cyclooxygenase-2 rather than cyclooxygenase-1 by nitric oxide," *Organic & Biomolecular Chemistry*, vol. 16, no. 5, pp. 765–770, 2018.
- [64] I. Ivanov, J. Saam, H. Kuhn, and H.-G. Holzhtutter, "Dual role of oxygen during lipoxygenase reactions," *The FEBS Journal*, vol. 272, no. 10, pp. 2523–2535, 2005.
- [65] I. Juránek, H. Suzuki, and S. Yamamoto, "Affinities of various mammalian arachidonate lipoxygenases and cyclooxygenases for molecular oxygen as substrate," *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*, vol. 1436, no. 3, pp. 509–518, 1999.
- [66] A. Carreau, B. E. Hafny-Rahbi, A. Matejuk, C. Grillon, and C. Kieda, "Why is the partial oxygen pressure of human tissues a crucial parameter? Small molecules and hypoxia," *Journal of Cellular and Molecular Medicine*, vol. 15, no. 6, pp. 1239–1253, 2011.
- [67] W. Wang, C. P. Winlove, and C. C. Michel, "Oxygen partial pressure in outer layers of skin of human finger nail folds," *The Journal of Physiology*, vol. 549, no. 3, pp. 855–863, 2003.
- [68] A. J. Brooks, J. Eastwood, I. J. Beckingham, and K. J. Girling, "Liver tissue partial pressure of oxygen and carbon dioxide during partial hepatectomy," *British Journal of Anaesthesia*, vol. 92, no. 5, pp. 735–737, 2004.
- [69] E. G. Mik, T. Johannes, C. J. Zuurbier et al., "In vivo mitochondrial oxygen tension measured by a delayed fluorescence lifetime technique," *Biophysical Journal*, vol. 95, no. 8, pp. 3977–3990, 2008.
- [70] R. S. Richardson, S. Duteil, C. Wary, D. W. Wray, J. Hoff, and P. G. Carlier, "Human skeletal muscle intracellular oxygenation: the impact of ambient oxygen availability," *The Journal of Physiology*, vol. 571, no. 2, pp. 415–424, 2006.
- [71] Y. Hirakawa, T. Yoshihara, M. Kamiya et al., "Quantitating intracellular oxygen tension in vivo by phosphorescence lifetime measurement," *Scientific Reports*, vol. 5, no. 1, article 17838, 2015.
- [72] D. J. Ceradini, A. R. Kulkarni, M. J. Callaghan et al., "Progenitor cell trafficking is regulated by hypoxic gradients through

- HIF-1 induction of SDF-1,” *Nature Medicine*, vol. 10, no. 8, pp. 858–864, 2004.
- [73] J. A. Spencer, F. Ferraro, E. Roussakis et al., “Direct measurement of local oxygen concentration in the bone marrow of live animals,” *Nature*, vol. 508, no. 7495, pp. 269–273, 2014.
- [74] A. V. Zhdanov, V. I. Ogurtsov, C. T. Taylor, and D. B. Papkovsky, “Monitoring of cell oxygenation and responses to metabolic stimulation by intracellular oxygen sensing technique,” *Integrative Biology*, vol. 2, no. 9, pp. 443–451, 2010.
- [75] H. Düssmann, S. Perez-Alvarez, U. Anilkumar, D. B. Papkovsky, and J. H. M. Prehn, “Single-cell time-lapse imaging of intracellular O₂ in response to metabolic inhibition and mitochondrial cytochrome-*c* release,” *Cell Death & Disease*, vol. 8, no. 6, article e2853, 2017.
- [76] R. I. Dmitriev and D. B. Papkovsky, “Optical probes and techniques for O₂ measurement in live cells and tissue,” *Cellular and Molecular Life Sciences*, vol. 69, no. 12, pp. 2025–2039, 2012.
- [77] G. T. Gabig, I. S. Bearman, and M. B. Babior, “Effects of oxygen tension and pH on the respiratory burst of human neutrophils,” *Blood*, vol. 53, pp. 1133–1139, 1979.
- [78] R. P. Brandes, N. Weissmann, and K. Schröder, “Nox family NADPH oxidases: molecular mechanisms of activation,” *Free Radical Biology & Medicine*, vol. 76, pp. 208–226, 2014.
- [79] S. Guo and X. Chen, “The human nox4: gene, structure, physiological function and pathological significance,” *Journal of Drug Targeting*, vol. 23, no. 10, pp. 888–896, 2015.
- [80] H. M. Abu-Soud, D. L. Rousseau, and D. J. Stuehr, “Nitric oxide binding to the heme of neuronal nitric-oxide synthase links its activity to changes in oxygen tension,” *Journal of Biological Chemistry*, vol. 271, no. 51, pp. 32515–32518, 1996.
- [81] H. M. Abu-Soud, J. Wang, D. L. Rousseau, J. M. Fukuto, L. J. Ignarro, and D. J. Stuehr, “Neuronal nitric oxide synthase self-inactivates by forming a ferrous-nitrosyl complex during aerobic catalysis,” *Journal of Biological Chemistry*, vol. 270, no. 39, pp. 22997–23006, 1995.
- [82] J. Santolini, A. L. Meade, and D. J. Stuehr, “Differences in three kinetic parameters underpin the unique catalytic profiles of nitric-oxide synthases I, II, and III,” *Journal of Biological Chemistry*, vol. 276, no. 52, pp. 48887–48898, 2001.
- [83] H. M. Abu-Soud, K. Ichimori, A. Presta, and D. J. Stuehr, “Electron transfer, oxygen binding, and nitric oxide feedback inhibition in endothelial nitric-oxide synthase,” *Journal of Biological Chemistry*, vol. 275, no. 23, pp. 17349–17357, 2000.
- [84] R. A. Dweik, D. Laskowski, H. M. Abu-Soud et al., “Nitric oxide synthesis in the lung. Regulation by oxygen through a kinetic mechanism,” *The Journal of Clinical Investigation*, vol. 101, no. 3, pp. 660–666, 1998.
- [85] M. A. Robinson, J. E. Baumgardner, V. P. Good, and C. M. Otto, “Physiological and hypoxic O₂ tensions rapidly regulate NO production by stimulated macrophages,” *American Journal of Physiology-Cell Physiology*, vol. 294, no. 4, pp. C1079–C1087, 2008.
- [86] M. D. Houslay and K. F. Tipton, “Rat liver mitochondrial monoamine oxidase. A change in the reaction mechanism on solubilization,” *Biochemical Journal*, vol. 145, no. 2, pp. 311–321, 1975.
- [87] M. Husain, D. E. Edmondson, and T. P. Singer, “Kinetic studies on the catalytic mechanism of liver monoamine oxidase,” *Biochemistry*, vol. 21, no. 3, pp. 595–600, 1982.
- [88] D. P. Jones, “Benzylamine metabolism at low O₂ concentrations: relative sensitivities of monoamine oxidase, aldehyde dehydrogenase and hippurate synthesis to hypoxia,” *Biochemical Pharmacology*, vol. 33, no. 3, pp. 413–417, 1984.
- [89] W. E. M. Lands, J. Sauter, and G. W. Stone, “Oxygen requirement for prostaglandin biosynthesis,” *Prostaglandins and Medicine*, vol. 1, no. 2, pp. 117–120, 1978.
- [90] A. Mukherjee, D. W. Brinkley, K.-M. Chang, and J. P. Roth, “Molecular oxygen dependent steps in fatty acid oxidation by cyclooxygenase-1,” *Biochemistry*, vol. 46, no. 13, pp. 3975–3989, 2007.
- [91] O. A. Ponomareva, N. A. Trushkin, I. S. Filimonov et al., “Prostaglandin H synthase kinetics in the two-phase aqueous-micellar system,” *Biochimica et Biophysica Acta (BBA) - Biomembranes*, vol. 1858, no. 9, pp. 2199–2207, 2016.
- [92] H. Tarhonskaya, A. P. Hardy, E. A. Howe et al., “Kinetic investigations of the role of factor inhibiting hypoxia-inducible factor (FIH) as an oxygen sensor,” *Journal of Biological Chemistry*, vol. 290, no. 32, pp. 19726–19742, 2015.
- [93] I. Diebold, A. Petry, J. Hess, and A. Gorlach, “The NADPH oxidase subunit NOX4 is a new target gene of the hypoxia-inducible factor-1,” *Molecular Biology of the Cell*, vol. 21, no. 12, pp. 2087–2096, 2010.
- [94] S. A. Saravanaperumal, J. H. Lee, S. A. Syed et al., “Hypoxia-inducible factor 1 alpha (HIF1A) stimulates neuronal nitric oxide synthase (NOS1) transcription by binding to multiple enhancers,” *Gastroenterology*, vol. 152, no. 5, pp. S565–S566, 2017.
- [95] F. Coulet, S. Nadaud, M. Agrapart, and F. Soubrier, “Identification of hypoxia-response element in the human endothelial nitric-oxide synthase gene promoter,” *Journal of Biological Chemistry*, vol. 278, no. 47, pp. 46230–46240, 2003.
- [96] L. Kalinowski, A. Janaszak-Jasiecka, A. Siekierzycka et al., “Posttranscriptional and transcriptional regulation of endothelial nitric-oxide synthase during hypoxia: the role of microRNAs,” *Cellular & Molecular Biology Letters*, vol. 21, no. 1, p. 16, 2016.
- [97] G. Melillo, T. Musso, A. Sica, L. S. Taylor, G. W. Cox, and L. Varesio, “A hypoxia-responsive element mediates a novel pathway of activation of the inducible nitric oxide synthase promoter,” *Journal of Experimental Medicine*, vol. 182, no. 6, pp. 1683–1693, 1995.
- [98] F. Jung, L. A. Palmer, N. Zhou, and R. A. Johns, “Hypoxic regulation of inducible nitric oxide synthase via hypoxia inducible factor-1 in cardiac myocytes,” *Circulation Research*, vol. 86, no. 3, pp. 319–325, 2000.
- [99] R. Hu, A. Dai, and S. Tan, “Hypoxia-inducible factor 1 alpha upregulates the expression of inducible nitric oxide synthase gene in pulmonary arteries of hypoxic rat,” *Chinese Medical Journal*, vol. 115, no. 12, pp. 1833–1837, 2002.
- [100] M. A. Sharpe and D. S. Baskin, “Monoamine oxidase B levels are highly expressed in human gliomas and are correlated with the expression of HIF-1 α and with transcription factors Sp1 and Sp3,” *Oncotarget*, vol. 7, no. 3, pp. 3379–3393, 2016.
- [101] P. J. Lee, B.-H. Jiang, B. Y. Chin et al., “Hypoxia-inducible factor-1 mediates transcriptional activation of the heme oxygenase-1 gene in response to hypoxia,” *Journal of Biological Chemistry*, vol. 272, no. 9, pp. 5375–5381, 1997.

- [102] C. S. Gonsalves and V. K. Kalra, "Hypoxia-mediated expression of 5-lipoxygenase-activating protein involves HIF-1 α and NF- κ B and microRNAs 135a and 199a-5p," *The Journal of Immunology*, vol. 184, no. 7, pp. 3878–3888, 2010.
- [103] L. Tie, N. Lu, X.-Y. Pan et al., "Hypoxia-induced up-regulation of aquaporin-1 protein in prostate cancer cells in a p38-dependent manner," *Cellular Physiology and Biochemistry*, vol. 29, no. 1-2, pp. 269–280, 2012.