Environmental proxies of antigen exposure explain variation in immune investment better than indices of pace-of-life

Nicholas P.C. Horrocks^{*1, 2}, Arne Hegemann¹, Stéphane Ostrowski³, Henry Ndithia^{1, 4}, Mohammed Shobrak⁵, Joseph B. Williams⁶, Kevin D. Matson¹ and B. Irene Tieleman¹

¹Animal Ecology Group, Centre for Evolutionary and Ecological Studies, University of Groningen, PO Box 11103, 9700 CC, Groningen, The Netherlands

²Behavioural Ecology Group, Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3EJ, United Kingdom

³Wildlife Conservation Society, 2300 Southern Boulevard, Bronx, NY 10460, USA

⁴Department of Ornithology, National Museums of Kenya, PO Box 40658, Nairobi, Kenya

⁵Biology Department, Science College, Taif University, PO Box 888, Taif 21974, Saudi Arabia

⁶Department of Evolution, Ecology and Organismal Biology, Ohio State University, Columbus, OH 43210, USA

*Author for correspondence (nh415@cam.ac.uk)

Author Contributions: NPCH and BIT conceived the study. All authors assisted with sample collection. NPCH and AH analysed the samples. NPCH analysed the data and wrote the manuscript; other authors provided editorial advice.

ABSTRACT

Investment in immune defences is predicted to co-vary with a variety of ecologically and evolutionarily relevant axes, with pace-of-life and environmental antigen exposure being two examples. These axes may themselves co-vary directly or inversely, and such relationships can lead to conflicting predictions regarding immune investment. If pace-of-life shapes immune investment then, following life history theory, slow-living, arid zone and tropical species should invest more in immunity than fast-living temperate species. Alternatively, if antigen exposure drives immune investment, then species in antigen-rich tropical and temperate environments are predicted to exhibit higher immune indices than species from antigen-poor arid locations. To test these contrasting predictions we investigated how variation in pace-of-life and antigen exposure influence immune investment in related lark species (*Alaudidae*) with differing life histories and predicted risks of exposure to environmental microbes and parasites. We used clutch size and total number of eggs laid year⁻¹ as indicators of pace-of-life, and aridity, and the climatic variables that influence aridity, as correlates of antigen abundance. We quantified immune investment by measuring four indices of innate immunity. Pace-of-life explained little of the variation in immune investment, and only one immune measure correlated significantly with pace-of-life, but not in the predicted direction. Conversely, aridity, our proxy for environmental antigen exposure, was predictive of immune investment, and larks in more mesic environments had higher immune indices than those living in arid, low-risk locations. Our study suggests that abiotic environmental variables with strong ties to environmental antigen exposure can be important correlates of immunological variation.

- 48   **Keywords:** Alaudidae, aridity, ecological immunology, lark, life history, pace-of-life
- 49   hypothesis.

INTRODUCTION

Explanations for variation in immune investment have often focused on the identification of evolutionary and ecological axes along which immune defences might co-vary. For example, ecological immunologists have exploited differences in pace-of-life to explain variation in immune defences. Limited resources and the costs associated with immunity suggest that immune investment must be counter-balanced against investment in other costly physiological processes such as growth and reproduction (Sheldon and Verhulst 1996; Ilmonen et al. 2000; Lochmiller and Deerenberg 2000; Norris and Evans 2000; Hegemann et al. 2013). Species at the fast end of the pace-of-life axis, with short lifespans and high reproductive rates, may allocate more of their limited resources to reproduction, and fewer to self maintenance activities such as immune defence. Conversely, 'slow-living' species that develop slowly, have low extrinsic mortality, and low reproductive rates, can prioritise self-maintenance activities and invest more heavily in immunity (Roff 1992; Stearns 1992; Ricklefs and Wikelski 2002).

Variation in exposure to environmental antigens represents another axis that might explain immune investment. Immune systems provide clear benefits in terms of protection against exogenous threats, including fitness-reducing micro- and macro-parasites. Immune investment might be greater when the risk of infection is higher (Tschirren and Richner 2006; Horrocks et al. 2011a), which could be associated with environment, time, and other ecological factors (Piersma 1997; Møller 1998; Guernier et al. 2004, Hegemann et al. 2012, 2013; Horrocks et al. 2012a, 2012b). For example, levels of environmental moisture shape endo- and ecto-parasitic communities, which show decreased prevalence, abundance and diversity in more arid environments (Little and Earlé 1995; Moyer et al. 2002; Valera et al. 2003; Guernier et al. 2004; Jex et al. 2007; Guerra et al. 2010; Froeschke et al. 2010; Pullan

and Brooker 2012). Combined with reduced moisture, the increased solar radiation and temperature extremes associated with arid environments also act to limit microbial assemblages (Tong and Lighthart 1997; Saranathan and Burt 2007; Burrows et al. 2009; Tang 2009; Bachar et al. 2010). If aridity is considered as a proxy for the level of antigenic exposure (Horrocks et al. 2014) then the requirement for immune investment should be greatest in cool, wet and humid environments where parasites and microbes are more likely to be encountered. This suggests a negative correlation between aridity and immune function.

Disentangling the relative contributions of pace-of-life and antigen exposure to immune variation is difficult because both axes may themselves co-vary (Horrocks et al. 2011a). Where pace-of-life and antigen exposure co-vary positively, predictions about immune investment coincide, even if the causal factor responsible for immunological variation is not clear. For example, relative to temperate birds, those in the tropics might invest more in immunity due to their slower pace-of-life (Martin II et al. 2006; Wiersma et al. 2007), because of increased exposure to environmental antigens such as parasites (Møller 1998; Guernier et al. 2004), or perhaps as a result of both factors. Where pace-of-life and antigen exposure co-vary negatively, conflicting predictions can arise. For example, the slow pace-of-life of desert-living birds (Tieleman et al. 2004) predicts strong investment in immune defences, similar to birds in the tropics, even though deserts and the tropics may pose contrasting risks in terms of exposure to antigens (Horrocks et al. 2011a). Investigating the drivers and correlates of immune variation in diverse environments requires careful consideration of study system characteristics. If the goal is to separate the contributions of pace-of-life and antigen exposure, then these two factors must be as un-confounded as possible.

We investigated how immune investment is influenced by pace-of-life and antigen exposure - expressed as environmental aridity - by studying related species of songbirds (larks; Alaudidae) that inhabit environments differing in aridity. Larks are ideally suited to this study because of their ecological similarities in different environments (del Hoyo et al. 2004), and because environmental moisture is already known to influence exposure to antigens in birds. Bacterial loads in nests, and infestation of nestlings by parasitic fly larvae correlate negatively with precipitation (Berger et al. 2003; Antoniazzi et al. 2011), while haematozoan infections and prevalence of lice and bacteria on feathers are reduced under more arid conditions (Little and Earlé 1995; Moyer et al. 2002; Valera et al. 2003; Saranathan and Burt 2007; Bush et al. 2009; Malenke et al. 2011; Horrocks et al. 2012b). Soil microbial abundance also correlates negatively with precipitation (Bachar et al. 2010; Drenovsky et al. 2010; Blankinship et al. 2011; Pasternak et al. 2013; Serna-Chavez et al. 2013) and soil microbes contaminate birds and their nests (Shawkey et al. 2005; Ruiz de Castañeda et al. 2011; Potter et al. 2013).

We measured lark species living in hot, hyper-arid deserts and cooler, wetter, mesic locations, as well as those from cold desert and tropical locations. Cold desert larks have clutch sizes typical of a fast pace-of-life yet live in an environment predicted to pose a low risk of antigen exposure. Tropical larks display life-history traits consistent with a slow pace-of-life yet live in potentially high antigen exposure settings (Tables 1 and 2). For the remaining lark species in our study pace-of-life increases with decreasing aridity, a finding that is unaffected by phylogeny (Tieleman et al. 2003; 2004). This means that the environments most associated with species exhibiting a slow pace-of-life (which may select *for* immune investment) are the same environments that present the lowest exposure to environmental antigens (which may select *against* immune investment). These contrasts

make larks particularly suited for teasing apart the roles of life history and environmental antigen exposure in shaping investment in immune defences.

We assessed immune investment by measuring circulating levels of four non-specific immune indices that any environmental antigen that has breached defensive barriers such as the skin or mucosa might encounter (Janeway et al. 2004). Haptoglobin and ovotransferrin are acute phase proteins with immunomodulatory properties that counter microbial challenges and limit microbial growth by directly sequestering iron (Xie et al. 2002; Arredouani et al. 2003). Natural antibodies opsonize invading microorganisms to facilitate phagocytosis and activate the complement system, which leads to cell lysis (Ochsenbein and Zinkernagel 2000). We used clutch size and number of eggs laid year⁻¹ as indicators of pace-of-life (Saether 1998; Ricklefs 2000), and aridity, precipitation and mean ambient temperature as proxies for environmental antigen exposure. We predicted that if immune investment is driven by pace-of-life, then slow-living, arid zone and tropical larks should invest relatively more in immune defences than fast-living species from temperate and cold-arid environments. If antigen exposure is more important for determining investment in immune defences, then we predicted that immune indices should be lowest in lark populations from arid locations, and be higher in temperate and tropical larks living in environments with greater abundance of microbes and macro-parasites.

MATERIALS AND METHODS

Study populations, sampling, and indicators of pace-of-life

We captured larks of 12 species in 11 climatically distinct locations during breeding and non-breeding periods from 2006 to 2009 (23 populations in total; Table 1). We collected <300µl

blood from the brachial vein of each bird and stored it on ice until processing by centrifugation to separate plasma and cellular fractions. The plasma was then frozen and stored at -20°C until it was used in immune assays. We gathered data on mean clutch size and number of eggs laid year⁻¹ (mean clutch size x mean number of clutches laid year⁻¹; Table 2) directly from our own study populations and from Tieleman et al. (2004), Cramp (1988), del Hoyo et al. (2004) and Hegemann et al. (2012, 2013). All birds were sampled under licence from the relevant authorities.

Climatic variables and aridity index

We obtained high-resolution (0.5 x 0.5 degree – approximately 55 x 55 km) gridded data on climatic variables for the period 1901-2009 from the Climatic Research Unit time series dataset (CRU TS 3.1; Harris et al. 2013). For each bird-sampling location we extracted mean annual values for precipitation (P, mm) and temperature (T, °C). We used these climatic variables to calculate de Martonne's aridity index A_M ($P / T + 10$; de Martonne 1926). A_M and climatic variables for each lark population are shown in Table 1. Low values of A_M indicate arid conditions, whereas higher values are associated with increasingly mesic environments.

Immune assays

We determined haptoglobin concentrations (mg ml⁻¹) using a functional assay that measures the haem-binding capacity of plasma (TP801; Tri-Delta Diagnostics, NJ, USA), following the 'manual method' instructions provided by the manufacturer and with incubation at 30°C for five minutes (Matson et al. 2012). We measured ovotransferrin concentrations (mg ml⁻¹) according to Horrocks et al. (2011b). Three of the 23 populations were not measured due to blood volume limitations (Table 2). We quantified natural

antibody-mediated agglutination titres and complement-mediated lysis titres against rabbit red blood cells (B-0009D, Harlan, UK), according to the assay of Matson et al. (2005).

Statistical analyses

We calculated mean values per population (i.e. per species per location) for each immune index and used these values to conduct regression analyses to investigate relationships between immune indices and life-history variables and aridity. Since our aridity index is a compound variable, we also ran separate models testing immune indices against each of the primary climatic variables precipitation and temperature. To account for potential non-independence among species due to common ancestry we used a phylogenetic generalised least squares (pgls) approach (Freckleton et al. 2002), simultaneously estimating maximum likelihood values of the parameter λ to test for phylogenetic signal in the model residuals (Pagel's lambda; Pagel 1999). A value of $\lambda = 0$ indicates no phylogenetic signal, whereas $\lambda = 1$ suggests that trait evolution is consistent with a strong effect of phylogeny. We based our phylogeny on the phylogenetic tree of larks (Alström et al. 2013). For the seven species with multiple sampled populations we added branches to the tree for each population to create polytomies, with the branch length for each population within a species set to zero. We found no evidence for phylogenetic signal in any of our datasets: λ never differed significantly from zero and was always significantly different from one. Setting all branch lengths to one did not change this finding. Therefore, we present all results and figures based on simple linear regression models (ordinary least squares, ols; Freckleton 2009; Revell 2010), with species and geographically distinct populations treated as independent points. Because sample size varied among species and populations (Table 1), we weighted regression models by the square root of the number of individuals sampled in each

population (Sokal and Rohlf 1995). Since some species or populations were only sampled during one period (breeding or non-breeding; Table 1) we ran analyses using restricted datasets containing values per period, as well as with the entire dataset of all values. The results of these analyses were qualitatively similar and so we only present results based on the entire dataset. To disentangle the roles of pace-of-life and environmental antigen exposure in shaping immune investment, these factors should not be correlated. This was the case in our dataset: pace-of-life indicators and aridity were not significantly correlated (clutch size vs. A_M : $P = 0.57$, $r = -0.12$; number of eggs laid year⁻¹ vs. A_M : $P = 0.16$, $r = 0.31$). All statistical analyses were conducted using R 2.15.2 (R Core Team 2012).

RESULTS

Relationships between immune indices and pace-of-life parameters were non-significant and generally weak (Table 3; Fig. 1). The notable exception to this was the relationship between number of eggs laid year⁻¹ and agglutination titres (Table 3; Fig. 1f).

Lark populations consistently exhibited lower immune defences in more arid locations (Fig. 2). These negative correlations with aridity were significant for haptoglobin concentrations and lysis titres, and approached significance for agglutination titres (Table 3; Fig. 2). Haptoglobin concentrations and lysis titres were also positively and significantly correlated with mean annual precipitation (Table 3; Fig. 2). Ovotransferrin concentrations showed no relationship with aridity or mean annual precipitation (Figs 2d-e). Lysis titres correlated significantly and positively with mean annual temperature: larks from warmer locations had higher lysis titres (Table 3; Fig. 2l). All other immune indices correlated

negatively with mean annual temperature but these relationships were not significant (Table 3; Fig. 2).

DISCUSSION

Contrary to the pace-of-life hypothesis, slow-living lark species did not invest more heavily in immune defences than their faster-living counterparts. Thus, our study adds to the list of publications that report no evidence for this hypothesis (Cutrera et al. 2010; Horrocks et al. 2012a; Versteegh et al. 2012, Palacios et al. 2013), and contradicts those studies that provide at least some support for an association between pace-of-life and immune investment (Tieleman et al. 2005; Martin et al. 2006; Lee et al. 2008; Sparkman and Palacios 2009; Previtali et al. 2012). Taxonomic and methodological differences between studies, including measurement of different immune components and the use of different proxies for pace-of-life, may explain some of this variation. However, even within a single immune index and taxonomic group, interpretation of immune investment in relation to pace-of-life is not straightforward. The sole significant relationship in our study that might suggest a link between pace-of-life and immune investment was between agglutination titres and number of eggs laid year⁻¹. However, this correlation was positive, and contrary to the pace-of-life hypothesis this suggests that lark species with a faster pace-of-life have higher agglutination ability. This finding contradicts earlier work showing greater agglutination ability in tropical species with a slower pace-of-life (Lee et al. 2008). It also conflicts with Versteegh et al. (2012), who found no relationship between agglutination titres and pace-of-life when studying stonechat subspecies in a common garden set-up. Natural antibody levels correlate positively with adaptive antibody responses (Parmentier et al. 2004) and these conflicting

patterns might arise because natural antibodies, although usually considered an innate immune defence, straddle the boundary between innate and adaptive immunity (Caroll and Prodeus 1998; Ochsenbein and Zinkernagel 2000). Studies that measure purely adaptive immune defences, such as induction of specific antibody responses, are needed to test this possibility. More generally, the equivocal nature of the evidence linking immune investment and pace-of-life suggests that further exploration of this topic is required.

In agreement with the antigen exposure hypothesis (Horrocks et al. 2011a), haptoglobin concentrations and lysis titres decreased with increasing aridity and were highest in populations of larks from less arid locations where exposure to abundant immunological challenges is expected to be higher (Little and Earlé 1995; Moyer et al. 2002; Valera et al. 2003; Guernier et al. 2004; Jex et al. 2007; Tang 2009; Bachar et al. 2010; Guerra et al. 2010; Froeschke et al. 2010). Agglutination titres showed a similar, but non-significant trend and only ovotransferrin concentrations showed no relationship with aridity. This might relate to the iron binding and transport function of ovotransferrin, which, aside from its anti-microbial and immunomodulatory properties, make ovotransferrin important for vascularisation and nerve and muscle growth in the developing embryo (Giansanti et al. 2012).

In addition to links with aridity, we found significant positive associations between haptoglobin concentrations and lysis titres and precipitation, and between lysis titres and temperature. Previous authors have linked the risk of antigen exposure to climatic factors (Guernier et al. 2004; Gage et al. 2008; Guerra et al. 2010) and other abiotic environmental variables (e.g. salinity; Figuerola 1999; Piersma 1997; Mendes et al. 2005). Our study extends this approach by showing that immune indices that are related to infection risk also correlate with abiotic environmental variation. This demonstrates that abiotic measures can

serve as useful proxies for antigenic exposure when direct biotic measurements are unavailable. Even so, it is important to recognise that other factors might influence immune investment, and may also co-vary with abiotic environmental variation. For example, physiological differences between birds along our aridity gradient, such as variation in metabolic rate (Tieleman et al. 2003) might influence immune responses, independent of any effect of antigen exposure. Temperature-constrained activity budgets (Tieleman and Williams 2002) and the reduced productivity associated with arid environments (Del Grosso et al. 2008; Serna-Chavez et al. 2013) could both restrict food availability in arid-living larks, leading to a resource trade-off that negatively impacts immune function. However, food availability might also be expected to influence clutch size and number of clutches year⁻¹ (Tieleman et al. 2004; Lepage and Lloyd 2004), yet neither pace-of-life indicator correlated significantly with aridity. Earlier work showed that lark species in hyper-arid regions encountered lower microbial abundances than more mesic, temperate-living larks, and had correspondingly lower immune indices (Horrocks et al. 2012a). This supports our argument that aridity is a good proxy for risk of antigen exposure and consequently immune investment. Future work that takes a similar approach and focuses on direct measurement of relevant antigens will shed additional light on the associations we have identified between aridity, risk of antigen exposure, and immune defence. Combined with experimental studies, this work is also necessary to determine whether patterns of correlation between antigen exposure and immune responses reflect evolved responses to an antigen-rich environment or are simply a reflection of current exposure to antigens.

Comparative studies, both within and among species and environments, represent a powerful approach for disentangling the roles of pace-of-life and antigen exposure in shaping immune defences and for understanding immune defence variation in more general

ecological and evolutionary terms. By employing a study system in which pace-of-life variation is uncoupled from variation in antigen exposure, we demonstrated that investment in innate immune defences might be related more to the likelihood of encountering an immunological challenge than to pace-of-life. Including relevant measures of environmental antigens and associated exposure risk in ecoimmunology studies will provide exciting opportunities for advancing our understanding of immunological variation.

ACKNOWLEDGEMENTS

We thank the following people and organisations for logistical support, and for permission to work with wild birds: HH Prince Bandar bin Saud, Secretary General of the Saudi Wildlife Commission, Mr Ahmad Al Bouq, Director of the National Wildlife Research Center and staff at Taif and Mahazat as-Sayd, Saudi Arabia; Staatsbosbeheer and volunteers on the Aekingerzand Lark Project, Netherlands; Muchai Muchane, National Museums Kenya, Friends of Kinangop Plateau, and Sarah Higgins, Kenya. Sample collection in Afghanistan was made possible by the generous support of the American people through the United States Agency for International Development (USAID) and its collaborative grantee the Wildlife Conservation Society (WCS). We thank Bob Ricklefs for commenting on an earlier version of the manuscript, and Dieter Luckas and Rose Thorogood for assistance with phylogenetic analyses. Two anonymous reviewers also gave useful comments on an earlier manuscript draft. Financial support came from the Schure-Beijerinck-Poppings Fonds (to NPCH and AH), BirdLife Netherlands (to BIT), NSF grant IBN 0212587 (to JBW), and VENI and VIDI grants from the Netherlands Organisation for Scientific Research (to KDM and BIT).

REFERENCES

- Alström P, Barnes KN, Olsson U, Barker FK, Bloomer P, Khan AA, Qureshi MA, Guillaumet A, Crochet P-A, Ryan PG (2013) Multilocus phylogeny of the avian family Alaudidae (larks) reveals complex morphological evolution, non-monophyletic genera and hidden species diversity. *Mol Phylogenet Evol* 69: 1043-1056 doi: 10.1016/j.ympev.2013.06.005
- Antoniazzi LR, Manzoli DE, Rohrmann D, Saravia MJ, Silvestri L, Beldomenico PM. (2011) Climate variability affects the impact of parasitic flies on Argentinean forest birds. *J Zool* 283: 126-134 doi: 10.1111/j.1469-7998.2010.00753.x
- Arredouani M, Matthijs P, Van Hoeyveld E, Kasran A, Baumann H, Ceuppens JL, Stevens E (2003) Haptoglobin directly affects T cells and suppresses T helper cell type 2 cytokine release. *Immunology* 108: 144–151 doi: 10.1046/j.1365-2567.2003.01569.x
- Bachar A, Al-Ashhab A, Soares M, Sklarz M, Angel R, Ungar E, Gillor O (2010) Soil microbial abundance and diversity along a low precipitation gradient. *Microb Ecol* 60: 453-461 doi: 10.1007/s00248-010-9727-1
- Berger S, Disko R, Gwinner H (2003) Bacteria in starling nests. *J Ornithol* 144: 317–322 doi: 10.1007/BF02465631
- Blankinship JC, Niklaus PA, Hungate BA (2011) A meta-analysis of responses of soil biota to global change. *Oecologia* 165: 553-565 doi 10.1007/s00442-011-1909-0
- Burrows SM, Elbert W, Lawrence MG, Poschl U (2009) Bacteria in the global atmosphere - Part 1: review and synthesis of literature data for different ecosystems. *Atmos Chem Phys* 9: 9263–9280 doi: 10.5194/acp-9-9263-2009
- Bush SE, Harbison CW, Slager DL, Townsend Peterson A, Price RD, Clayton DH. (2009) Geographic variation in the community structure of lice on Western Scrub-Jays. *J Parasitol* 95: 10-13

337 Carroll MC, Prodeus AP (1998) Linkages of innate and adaptive immunity. *Curr Opin Immunol*
 338 10: 36-40 doi: 10.1016/S0952-7915(98)80028-9
 339 Cramp S (1988) Tyrant Flycatchers to Thrushes. In: Cramp S (ed) *Handbook of the Birds of*
 340 *Europe, the Middle East and North Africa: The Birds of the Western Palearctic*. Oxford
 341 University Press, Oxford
 342 Cutrera AP, Zenuto RR, Luna F, Antenucci CD (2010) Mounting a specific immune response
 343 increases energy expenditure of the subterranean rodent *Ctenomys talarum* (tuco-tuco):
 344 implications for intraspecific and interspecific variation in immunological traits. *J Exp Biol*
 345 213: 715-724 doi: 10.1242/jeb.037887
 346 de Martonne E (1926) Une nouvelle fonction climatologique: L' indice d'aridité. *La*
 347 *Météorologie* 2: 449-458
 348 Del Grosso S, Parton W, Stohlgren T, Zheng D, Bachelet D, Prince S, Hibbard K, Olson R
 349 (2008) Global potential net primary production predicted from vegetation class,
 350 precipitation, and temperature. *Ecology* 89: 2117-2126 doi: 10.1890/07-0850.1
 351 del Hoyo J, Elliott A, Christie DA (2004) Cotingas to Pipits and Wagtails. *Handbook of the*
 352 *Birds of the World*. Lynx Edicions, Barcelona
 353 Drenovsky RE, Steenwerth KL, Jackson LE, Scow KM (2010) Land use and climatic factors
 354 structure regional patterns in soil microbial communities. *Global Ecol Biogeogr* 19: 27-39
 355 doi: 10.1111/j.1466-8238.2009.00486.x
 356 Figuerola J (1999) Effects of salinity on rates of infestation of waterbirds by haematozoa.
 357 *Ecography* 22: 681-685 doi: 10.1111/j.1600-0587.1999.tb00517.x
 358 Freckleton RP (2009) The seven deadly sins of comparative analysis. *J Evol Biol* 22: 1367-
 359 1375 doi: 10.1111/j.1420-9101.2009.01757.x

360 Freckleton RP, Harvey PH, Pagel M (2002) Phylogenetic analysis and comparative data: A
 361 test and review of evidence. *Am Nat* 160: 712-726 doi: 10.1086/343873

362 Froeschke G, Harf R, Sommer S, Matthee S (2010) Effects of precipitation on parasite burden
 363 along a natural climatic gradient in southern Africa – implications for possible shifts in
 364 infestation patterns due to global changes. *Oikos* 119: 1029-1039 doi: 10.1111/j.1600-
 365 0706.2009.18124.x

366 Gage KL, Burkot TR, Eisen RJ, Hayes EB (2008) Climate and vectorborne diseases. *Am J Prev*
 367 *Med* 35: 436-450 doi: 10.1016/j.amepre.2008.08.030

368 Giansanti F, Leboffe L, Pitari G, Ippoliti R, Antonini G (2012) Physiological roles of
 369 ovotransferrin. *BBA-Gen Subjects* 1820: 218-225 doi: 10.1016/j.bbagen.2011.08.004

370 Guernier V, Hochberg ME, Guegan JFO (2004) Ecology drives the worldwide distribution of
 371 human diseases. *PLoS Biol* 2: 740-746 doi: 10.1371/journal.pbio.0020141

372 Guerra CA, Howes RE, Patil AP, Gething PW, Van Boeckel TP, Temperley WH, Kabaria CW,
 373 Tatem AJ, Manh BH, Elyazar IRF, Baird JK, Snow RW, Hay SI (2010) The international limits
 374 and population at risk of *Plasmodium vivax* transmission in 2009. *PLoS Neglect Trop D* 4:
 375 e774 doi: 10.1371/journal.pntd.0000774

376 Harris I, Jones PD, Osborn TJ, Lister DH (2013) Updated high-resolution grids of monthly
 377 climatic observations. *Int J Climatol* 34: 623-642 doi:10.1002/joc.3711.

378 Hegemann A, Matson KD, Both C, Tieleman BI (2012) Immune function in a free-living bird
 379 varies over the annual cycle, but seasonal patterns differ between years. *Oecologia* 170:
 380 605-618 doi: 10.1007/s00442-012-2339-3

381 Hegemann A, Matson KD, Flinks H, Tieleman BI (2013) Offspring pay sooner, parents pay
 382 later: Experimental manipulation of body mass reveals trade-offs between immune
 383 function, reproduction and survival. *Front Zool* 10: 77 doi: 10.1186/1742-9994-10-77

384 Horrocks NPC, Matson KD, Tieleman BI (2011a) Pathogen pressure puts immune defense
 385 into perspective. *Integr Comp Biol* 51: 563-576 doi: 10.1093/icb/icr011
 386 Horrocks NPC, Tieleman BI, Matson KD (2011b) A simple assay for measurement of
 387 ovotransferrin – a marker of inflammation and infection in birds. *Methods Ecol Evol* 2:
 388 518-526 doi: 10.1111/j.2041-210X.2011.00096.x
 389 Horrocks NPC, Hegemann A, Matson KD, Hine K, Jaquier S, Shobrak M, Williams JB,
 390 Tinbergen JM, Tieleman BI (2012a) Immune indexes of larks from desert and temperate
 391 regions show weak associations with life history but stronger links to environmental
 392 variation in microbial abundance. *Physiol Biochem Zool* 85: 504-515 doi: 10.1086/666988
 393 Horrocks NPC, Matson KD, Shobrak M, Tinbergen JM, Tieleman BI. (2012b) Seasonal
 394 patterns in immune indices reflect microbial loads on birds but not microbes in the wider
 395 environment. *Ecosphere* 3: art 19 doi: 10.1890/ES11-00287.1
 396 Horrocks NPC, Hine K, Hegemann A, Ndithia HK, Shobrak M, Ostrowski S, Williams JB,
 397 Matson KD, Tieleman BI. (2014) Are antimicrobial defences in bird eggs related to
 398 climatic conditions associated with risk of trans-shell microbial infection? *Front Zool* 11:
 399 49 doi: 10.1186/1742-9994-11-49
 400 Ilmonen P, Taarna T, Hasselquist D (2000) Experimentally activated immune defence in
 401 female pied flycatchers results in reduced breeding success. *Proc R Soc Lond B* 267: 665–
 402 670 doi: 10.1098/rspb.2000.1053
 403 Janeway CA, Travers P, Walport M, Shlomchik J (2004) *Immunobiology: the immune system*
 404 *in health and disease*. Garland Publishing, New York
 405 Jex AR, Schneider MA, Rose HA, Cribb TH (2007) Local climate aridity influences the
 406 distribution of thelastomatoid nematodes of the Australian giant burrowing cockroach.
 407 *Parasitology* 134: 1401-1408 doi: 10.1017/S0031182007002727

408 Lee KA, Wikelski M, Robinson RD, Robinson TR, Klasing KC (2008) Constitutive immune
 409 defences correlate with life-history variables in tropical birds. *J Anim Ecol* 77: 356-363
 410 doi: 10.1111/j.1365-2656.2007.01347.x

411 Lepage D, Lloyd P (2004) Avian clutch size in relation to rainfall seasonality and stochasticity
 412 along an aridity gradient across South Africa. *Ostrich* 75: 259-268 doi:
 413 10.2989/00306520409485453

414 Little RM, Earlé RA (1995) Sandgrouse (Pterocleididae) and sociable weavers *Philetarius socius*
 415 lack avian haematozoa in semi-arid regions of South Africa. *J Arid Environ* 30: 367-370
 416 doi: 10.1016/S0140-1963(05)80011-4

417 Lochmiller RL, Deerenberg C (2000) Trade-offs in evolutionary immunology: just what is the
 418 cost of immunity? *Oikos* 88: 87–98 doi: 10.1034/j.1600-0706.2000.880110.x

419 Malenke JR, Newbold N, Clayton DH. (2011) Condition-specific competition governs the
 420 geographic distribution and diversity of ectoparasites. *Am Nat* 177: 522-534 doi:
 421 10.1086/658176

422 Martin II LB, Hasselquist D, Wikelski M (2006) Investment in immune defense is linked to
 423 pace of life in house sparrows. *Oecologia* 147: 565-575 doi: 10.1007/s00442-005-0314-y

424 Matson KD, Ricklefs RE, Klasing KC (2005) A hemolysis-hemagglutination assay for
 425 characterizing constitutive innate humoral immunity in wild and domestic birds. *Dev*
 426 *Comp Immunol* 29: 275-286 doi: 10.1016/j.dci.2004.07.006

427 Maston KD, Horrocks NPC, Versteegh MA, Tieleman BI (2012) Baseline haptoglobin
 428 concentrations are repeatable and predictive of certain aspects of a subsequent
 429 experimentally-induced inflammatory response. *Comp Biochem Phys A* 162: 7-15 doi:
 430 10.1016/j.cbpa.2012.01.010

431 Mendes L, Piersma T, Lecoq M, Spaans B, Ricklefs RE (2005) Disease-limited distributions?
 432 Contrasts in the prevalence of avian malaria in shorebird species using marine and
 433 freshwater habitats. *Oikos* 109: 396-404 doi: 10.1111/j.0030-1299.2005.13509.x
 434 Møller AP (1998) Evidence of larger impact of parasites on hosts in the tropics: investment
 435 in immune function within and outside the tropics. *Oikos* 82: 265-270 doi:
 436 10.2307/3546966
 437 Moyer BR, Drown DM, Clayton DH (2002) Low humidity reduces ectoparasite pressure:
 438 implications for host life history evolution. *Oikos* 97: 223-228 doi: 10.1034/j.1600-
 439 0706.2002.970208.x
 440 Norris K, Evans MR (2000) Ecological immunology: life history trade-offs and immune
 441 defense in birds. *Behav Ecol* 11: 19-26 doi: 10.1093/beheco/11.1.19
 442 Ochsenbein AF, Zinkernagel RM (2000) Natural antibodies and complement link innate and
 443 acquired immunity. *Immunol Today* 21: 624-630 doi: 10.1016/S0167-5699(00)01754-0
 444 Pagel M 1999 Inferring the historical patterns of biological evolution. *Nature* 401: 877-884
 445 doi: 10.1038/44766
 446 Palacios MG, Cunnick JE, Bronikowski AM (2013) Complex interplay of body condition, life
 447 history, and prevailing environment shapes immune defences of garter snakes in the
 448 wild. *Physiol Biochem Zool* 86: 547-558 doi: 10.1086/672371
 449 Parmentier HK, Lammers A, Hoekman JJ, Reilingh G V, Zaanen ITA, Savelkoul HFJ (2004)
 450 Different levels of natural antibodies in chickens divergently selected for specific
 451 antibody responses. *Dev Comp Immunol* 28: 39-49 doi: 10.1016/S0145-305X(03)00087-9
 452 Pasternak Z, Al-Ashhab A, Gatica J, Gafny R, Avraham S, Minz D, Gillor O, Jurkevitch E (2013)
 453 Spatial and temporal biogeography of soil microbial communities in arid and semiarid
 454 regions. *PLoS ONE* 8: e69705 doi: 10.1371/journal.pone.0069705

455 Piersma T (1997) Do global patterns of habitat use and migration strategies co-evolve with
 456 relative investments in immunocompetence due to spatial variation in parasite pressure?
 457 *Oikos* 80: 623-631 doi: 10.2307/3546640

458 Potter BA, Carlson BM, Adams AE, Voss MA (2013) An assessment of the microbial diversity
 459 present on the surface of naturally incubated House Wren eggs. *Open Ornithol J* 6: 32-39
 460 doi: 10.2174/1874453220130815001

461 Previtali MA, Ostfeld RS, Keesing F, Jolles AE, Hanselmann R, Martin LB (2012) Relationship
 462 between pace of life and immune responses in wild rodents. *Oikos* 121: 1483-1492 doi:
 463 10.1111/j.1600-0706.2012.020215.x

464 Pullan RL, Brooker SJ (2012) The global limits and population at risk of soil-transmitted
 465 helminths infections in 2010. *Parasit Vectors* 5: 81 doi: 10.1186/1756-3305-5-81

466 R Core Team (2013) R: A language and environment for statistical computing. R Foundation
 467 for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>

468 Revell LJ (2010) Phylogenetic signal and linear regression on species data. *Methods Ecol Evol*
 469 1: 319-329 doi: 10.1111/j.2041-210X.2010.00044.x

470 Ricklefs RE (2000) Density dependence, evolutionary optimization, and the diversification of
 471 avian life histories. *The Condor* 102: 9-22 doi: 10.1650/0010-
 472 5422(2000)102[0009:DDEOAT]2.0.CO;2

473 Ricklefs RE, Wikelski M (2002) The physiology-life history nexus. *Trends Ecol Evol* 17: 462-
 474 468 doi: 10.1016/S0169-5347(02)02578-8

475 Roff DA (1992) The evolution of life histories: theory and analysis. Chapman & Hall, New
 476 York

477 Ruiz-de-Castañeda R, Vela AI, Lobato E, Briones V, Moreno J (2011) Prevalence of potentially
 478 pathogenic culturable bacteria on eggshells and in cloacae of female Pied Flycatchers in a

479 temperate habitat in central Spain. *J Field Ornithol* 82: 215-224 doi: 10.1111/j.1557-
 480 9263.2011.00324.x
 481 Saether B-E (1988) Pattern of covariation between life-history traits of European birds.
 482 *Nature* 331: 616-617 doi: 10.1038/331616a0
 483 Saranathan V, Burt EH Jr (2007) Sunlight on feathers inhibits feather-degrading bacteria.
 484 *Wilson J Ornithol* 119: 239–245 doi: 10.1676/05-123.1
 485 Serna-Chavez HM, Fierer N, van Bodegom PM (2013) Global drivers and patterns of
 486 microbial abundance in soil. *Global Ecol Biogeogr* 22: 1162-1172 doi: 10.1111/geb.12070
 487 Shawkey MD, Mills KL, Dale C, Hill GE (2005) Microbial diversity of wild bird feathers
 488 revealed through culture-based and culture-independent techniques. *Microb Ecol* 50: 40-
 489 47 doi: 10.1007/s00248-004-0089-4
 490 Sheldon BC, Verhulst S (1996) Ecological immunology: costly parasite defences and trade-
 491 offs in evolutionary ecology. *Trends Ecol Evol* 11: 317-321 doi: 10.1016/0169-
 492 5347(96)10039-2
 493 Sokal RR, Rohlf FJ (1995) *Biometry: The principles and practice of statistics in biological*
 494 *research*. WH Freeman and Co, New York
 495 Sparkman AM, Palacios MG (2009) A test of life-history theories of immune defence in two
 496 ecotypes of the garter snake, *Thamnophis elegans*. *J Anim Ecol* 78: 1242-1248 doi:
 497 10.1111/j.1365-2656.2009.01587.x
 498 Stearns SC (1992) *The evolution of life histories*. Oxford University Press, Oxford
 499 Tang JW (2009) The effect of environmental parameters on the survival of airborne
 500 infectious agents. *J R Soc Interface* 6: S737-S746 doi: 10.1098/rsif.2009.0227.focus

501 Tieleman BI, Williams JB (2002) Effects of food supplementation on behavioural decisions of
 502 hoopoe-larks in the Arabian Desert: balancing water, energy and thermoregulation. *Anim*
 503 *Behav* 63: 519-529 doi: 10.1006/anbe.2001.1927
 504 Tieleman BI, Williams JB, Bloomer P (2003) Adaptation of metabolism and evaporative
 505 water loss along an aridity gradient. *Proc R Soc Lond B* 270: 207-214 doi:
 506 10.1098/rspb.2002.2205
 507 Tieleman BI, Williams JB, Visser GH (2004) Energy and water budgets of larks in a life history
 508 perspective: parental effort varies with aridity. *Ecology* 85: 1399-1410 doi: 10.1890/03-
 509 0170
 510 Tieleman BI, Williams JB, Ricklefs RE, Klasing KC (2005) Constitutive innate immunity is a
 511 component of the pace-of-life syndrome in tropical birds. *Proc R Soc Lond B* 272: 1715-
 512 1720 doi: 10.1098/rspb.2005.3155
 513 Tong Y, Lighthart B (1997) Solar radiation has a lethal effect on natural populations of
 514 culturable outdoor atmospheric bacteria. *Atmos Environ* 31: 897–900 doi:
 515 10.1016/S1352-2310(96)00235-X
 516 Tschirren B, Richner H (2006) Parasites shape the optimal investment in immunity. *Proc R*
 517 *Soc Lond B* 273: 1773-1777 doi: 10.1098/rspb.2006.3524
 518 Valera F, Carrillo CM, Barbosa A, Moreno E (2003) Low prevalence of haematozoa in
 519 Trumpeter finches *Bucanetes githagineus* from south-eastern Spain: additional support
 520 for a restricted distribution of blood parasites in arid lands. *J Arid Environ* 55: 209-213
 521 doi: 10.1016/S0140-1963(03)00041-7
 522 Versteegh MA, Schwabl I, Jaquier S, Tieleman BI (2012) Do immunological, endocrine and
 523 metabolic traits fall on a single Pace-of-Life axis? Covariation and constraints among
 524 physiological systems. *J Evol Biol* 25: 1864-1876 doi: 10.1111/j.1420-9101.2012.02574.x

525 Wiersma P, Muñoz-Garcia A, Walker A, Williams JB (2007) Tropical birds have a slow pace of
526 life. P Natl Acad Sci USA 104: 9340-9345 doi: 10.1073/pnas.0702212104
527 Xie H, Huff GR, Huff WE, Balog JM, Rath NC (2002) Effects of ovotransferrin on chicken
528 macrophages and heterophil-granulocytes. Dev Comp Immunol 26: 805-815 doi:
529 10.1016/S0145-305X(02)00028-9
530

531 **TABLES**

532 Table 1. Sample size (*n*), sampling period (breeding (B), non-breeding (NB), or sampled in both periods (both)), geographic origin and habitat
 533 description, and climatic variables for 12 species of lark. The climatic variables are the aridity index A_M ($P / T + 10$), and mean annual values for
 534 precipitation (P) and temperature (T). A lower value of A_M indicates a more arid environment.

#	species	<i>n</i>	sampling	latitude	longitude	country	habitat	A_M	P (mm)	T (°C)
a	hoopoe lark <i>Alaemon alaudipes</i>	61	both	22° 20' N	41° 44' E	Saudi Arabia	hot desert	2.29	81.19	25.38
b		4	B	19° 53' N	16° 18' W	Mauritania	hot desert	2.01	69.33	24.55
c	bar-tailed desert lark <i>Ammomanes cincturus</i>	56	both	22° 20' N	41° 44' E	Saudi Arabia	hot desert	2.29	81.19	25.38
d	black-crowned finchlark <i>Eremopterix nigriceps</i>	14	B	21° 15' N	40° 42' E	Saudi Arabia	hot desert	6.43	200.20	21.12
e		19	both	22° 20' N	41° 44' E	Saudi Arabia	hot desert	2.29	81.19	25.38
f	crested lark <i>Galerida cristata</i>	4	B	21° 15' N	40° 42' E	Saudi Arabia	hot desert	6.43	200.20	21.12
g		18	both	22° 20' N	41° 44' E	Saudi Arabia	hot desert	2.29	81.19	25.38
h		2	NB	34° 22' N	62° 11' E	Afghanistan	cold desert	8.66	226.44	16.14
i	Dunn's lark <i>Eremalauda dunni</i>	35	both	22° 20' N	41° 44' E	Saudi Arabia	hot desert	2.29	81.19	25.38
j	short-toed lark <i>Calandrella brachydactyla</i>	2	NB	22° 16' N	41° 45' E	Saudi Arabia	hot desert	2.29	81.19	25.38
k	bimaculated lark <i>Melanocorypha bimaculata</i>	6	NB	36° 54' N	66° 53' E	Afghanistan	cold desert	7.96	214.88	16.98
l		14	NB	34° 54' N	67° 11' E	Afghanistan	cold desert	26.92	389.80	4.48
m		7	NB	34° 22' N	62° 11' E	Afghanistan	cold desert	8.66	226.44	16.14
n		6	B	36° 43' N	67° 06' E	Afghanistan	cold desert	9.50	243.17	15.61

535

536 Table 1 continued.

#	species	<i>n</i>	sampling	latitude	longitude	country	habitat	<i>A_M</i>	<i>P</i> (mm)	<i>T</i> (°C)
o	calandra lark <i>Melanocorypha calandra</i>	3	NB	36° 54' N	66° 53' E	Afghanistan	cold desert	7.96	214.88	16.98
p		6	NB	34° 54' N	67° 11' E	Afghanistan	cold desert	26.92	389.80	4.48
q		11	NB	34° 22' N	62° 11' E	Afghanistan	cold desert	8.66	226.44	16.14
r	red-capped lark <i>Calandrella cinerea</i>	5	B	0° 51' S	36° 25' E	Kenya	tropical	19.64	593.98	20.25
s		8	B	0° 34' S	36° 29' E	Kenya	tropical	33.05	839.22	15.39
t	rufous-naped lark <i>Mirafra africana</i>	4	B	0° 48' S	36° 32' E	Kenya	tropical	19.64	593.98	20.25
u		2	B	0° 34' S	36° 29' E	Kenya	tropical	33.05	839.22	15.39
v	skylark <i>Alauda arvensis</i>	144	both	52° 56' N	6° 18' E	Netherlands	temperate	40.50	777.01	9.19
w	woodlark <i>Lullula arborea</i>	60	both	52° 56' N	6° 18' E	Netherlands	temperate	40.50	777.01	9.19

537

538

539 Table 2. Mean clutch size and number of clutches per year, concentrations of haptoglobin and ovotransferrin, and agglutination and lysis titres
540 for 23 populations of 12 lark species. Values for life-history variables are from this study and from the literature (data source column).

#	species	clutch size	clutches year ⁻¹	haptoglobin (mg ml ⁻¹)	ovotransferrin (mg ml ⁻¹)	agglutination (titre)	lysis (titre)	data source*
a	hoopoe lark	2.99	1	0.28	7.47	5.85	0.86	2
b		2.88	1	0.25	7.41	4.50	1.63	3
c	bar-tailed desert lark	3.24	1	0.29	9.08	6.13	0.38	2
d	black-crowned finchlark	2.57	1	0.49	5.43	7.03	1.52	3, 4
e		2.00	1	0.27	9.11	5.83	0.58	3, 4
f	crested lark	4.15	2	0.25	15.28	6.31	1.94	2
g		4.15	2	0.25	5.18	6.24	0.53	2
h		4.75	2	0.07	11.20	5.25	0.00	2, 3
i	Dunn's lark	2.88	1	0.49	9.76	6.65	1.63	2
j	short-toed lark	3.50	2	0.41	9.18	11.00	1.00	2
k	bimaculated lark	3.96	1.5	0.19	-	5.17	2.08	3
l		3.96	1.5	0.33	14.53	4.90	2.63	3
m		3.96	1.5	0.11	12.72	4.21	0.21	3
n		3.96	1.5	0.17	-	7.63	4.88	3

541

542

543 Table 2 continued.

#	species	clutch size	clutches year ⁻¹	haptoglobin (mg ml ⁻¹)	ovotransferrin (mg ml ⁻¹)	agglutination (titre)	lysis (titre)	data source*
o	calandra lark	4.20	2	0.08	-	7.25	3.58	2
p		4.20	2	0.06	6.01	6.46	1.75	2
q		4.20	2	0.07	9.78	5.90	1.25	2
r	red-capped lark	1.83	2	0.15	9.10	4.50	0.13	1
s		1.89	2	0.57	7.25	5.17	2.42	1
t	rufous-naped lark	2.11	1	0.74	9.08	6.31	3.69	1, 4
u		2.00	1	0.19	10.20	5.63	3.63	1, 4
v	skylark	3.56	3.5	0.48	-	7.82	2.26	1, 5
w	woodlark	4.02	2.5	0.46	9.41	7.20	2.21	1

544 * 1 Own data; 2 Tieleman et al. (2004); 3 Cramp (1988); 4 del Hoyo et al. (2004); 5 Hegemann et al. (2012, 2013).

545

546

Table 3. Results of linear models examining relationships between immune indices of 23 populations of 12 lark species in relation to pace-of-life proxies and to climatic proxies of environmental antigen exposure. *P* values < 0.05 are shown in bold.

response variable	explanatory variable	F	P
haptoglobin (mg ml ⁻¹)	mean clutch size	F _{1, 21} = 2.86	0.11
	total eggs year ⁻¹	0.21	0.66
	aridity index A _M	4.90	0.038
	mean annual precipitation (mm)	5.79	0.025
	mean annual temperature (°C)	0.31	0.59
ovotransferrin (mg ml ⁻¹)	mean clutch size	F _{1, 17} = 4.00	0.06
	total eggs year ⁻¹	1.74	0.21
	aridity index A _M	0.53	0.72
	mean annual precipitation (mm)	0.00	0.95
	mean annual temperature (°C)	2.79	0.11
agglutination (titre)	mean clutch size	F _{1, 21} = 1.00	0.33
	total eggs year ⁻¹	8.34	0.009
	aridity index A _M	3.14	0.09
	mean annual precipitation (mm)	2.31	0.14
	mean annual temperature (°C)	1.11	0.31
lysis (titre)	mean clutch size	F _{1, 21} = 0.23	0.64
	total eggs year ⁻¹	1.59	0.22
	aridity index A _M	6.58	0.018
	mean annual precipitation (mm)	7.03	0.015
	mean annual temperature (°C)	8.32	0.009

FIGURES

Figure 1. Haptoglobin (a-b) and ovotransferrin (c-d) concentrations and agglutination (e-f) and lysis (g-h) titres as a function of mean clutch size and number of eggs year⁻¹ in 23 populations of 12 lark species. Squares: hot desert larks; circles: cold desert larks; triangles: tropical larks; diamonds: temperate larks.

Figure 2. Haptoglobin (a-c) and ovotransferrin (d-f) concentrations and agglutination (g-i) and lysis (j-l) titres as a function of de Martonne's aridity index A_M , mean annual precipitation (mm) and mean annual temperature (°C) in 23 populations of 12 lark species measured along an environmental aridity gradient. A_M increases with decreasing aridity of the environment. Squares: hot desert larks; circles: cold desert larks; triangles: tropical larks; diamonds: temperate larks.