Ocean acidification does not impact shell growth or repair of the Antarctic brachiopod *Liothyrella uva* (Broderip, 1833)

Emma L. Cross^{a,b*}, Lloyd S. Peck^a, Elizabeth M. Harper^b

^a British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road, Cambridge, CB3 0ET, UK;

^b Department of Earth Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EQ, UK

* Corresponding author: Emma L Cross, British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road, Cambridge, CB3 0ET, UK. Email: emmoss@bas.ac.uk. Telephone: +44 (0)1223 333493.

Keywords: Climate change, calcification, CO₂, pH, carbonate saturation.

Abstract

Marine calcifiers are amongst the most vulnerable organisms to ocean acidification due to reduction in the availability of carbonate ions for skeletal/shell deposition. However, there are limited long-term studies on the possible impacts of increased $p\text{CO}_2$ on these taxa. A 7 month CO_2 perturbation experiment was performed on one of the most calcium carbonate dependent species, the Antarctic brachiopod *Liothyrella uva*, which inhabits the Southern Ocean where carbonate ion saturation levels are amongst the lowest on Earth. The effects of the predicted environmental conditions in 2050 and 2100 on the growth rate and ability to repair shell in *L.uva* were tested with four treatments; a low temperature control (0°C , pH7.98), a pH control (2°C , pH8.05), mid-century scenario (2°C , pH7.75) and end-century scenario (2°C , pH7.54). Environmental change impacts on shell repair are rarely studied, but here repair was not affected by either acidified conditions or temperature. Growth rate was also not impacted by low pH. Elevated temperature did, however, increase growth rates. The ability of *L.uva* to continue, and even increase shell production in warmer and acidified seawater suggests that this species can acclimate to these combined stressors and generate suitable conditions for shell growth at the site of calcification.

1. Introduction

Increasing CO_2 levels from anthropogenic activities over the past 250 years has altered our oceans through warming and also acidification (Caldeira and Wickett, 2003, 2005; IPCC, 2013; Orr et al., 2005). This latter process has received much

attention recently with the chemical implications now being fairly well described, although the biological and ecological consequences are less well described (Gattuso et al., 2013; Wittmann and Pörtner, 2013). However, there is a consensus that marine calcifying organisms are the most susceptible animal group to ocean acidification because the predicted reduction in the availability of carbonate ions will make it more difficult and more energetically expensive for shell production (Byrne, 2011; Byrne and Przeslawski, 2013; Doney et al., 2009; Watson et al., 2012). To date studies have reported varied responses of calcifying organisms to future predicted pH levels with an increasing number of studies indicating that some species are tolerant (Havenhand and Schlegel, 2009; Parker et al., 2012; Ries et al., 2009). However, it must be recognised that the majority of experiments have been conducted on relatively short time scales so the ability of organisms to acclimate or adapt is largely unknown (Byrne, 2011; Byrne and Przeslawski, 2013; Gattuso et al., 2013; Wittmann and Pörtner, 2013). Longer term studies are increasing though, which are providing insights into how organisms are coping with acidifying oceans (Form and Riebesell, 2012; Kelly et al., 2013; Pandolfi et al., 2011; Pespeni et al., 2013).

The fastest rates of change in carbonate chemistry are expected in the Southern Ocean (Caldeira and Wickett, 2005; McNeil and Matear, 2008). CO₂ is more soluble in cold water (Revelle and Fairbridge, 1957) resulting in naturally low carbonate ion saturation levels compared to temperate and tropical regions. Acid-base coefficients are also more sensitive in cold temperatures making this high latitude region a forerunner of biological ocean acidification impacts for other oceans (Fabry et al., 2009). Furthermore, the absence of shell-crushing predators, such as

crabs, lobsters and heavily jawed fish (Aronson et al., 2007) and the difficulty of extracting Ca^{2+} from seawater at low temperature (Aronson et al., 2007; Harper, 2000) has resulted in Antarctic species generally having thin, weakly calcified shells (Vermeij, 1978; Watson et al., 2012). This, added to the low physiological rates of Antarctic marine species (Peck et al., 2007), especially low metabolic rates (Peck and Conway, 2000), slow growth rates (Arntz et al., 1994), delayed reproduction (Meidlinger et al., 1998) and high longevity (Pearse et al., 1991), indicates that these organisms are likely to be amongst the most vulnerable species worldwide to acidifying oceans. Although there are several studies on the potential impacts of this aspect of climate change on the larval stage of Antarctic calcifying organisms (see review by Byrne (2011)), there are limited studies on adults (Cummings et al., 2011; McClintock et al., 2009), and the longest of these lasted 4 months.

Rhynchonelliform brachiopods are potentially the most calcium carbonate dependent group of marine animals because their calcareous skeleton and other support structures makes up over 90% of their dry mass (Peck, 1993, 2008). They have locally also been important organisms in shallow water communities, providing a habitat for a diverse range of epifauna including encrusting sponges and algae (Barnes and Peck, 1996), for the last 500 million years surviving several geological periods where the pH has fluctuated. Ocean pH has declined in the past 250 years at a rate of at least an order of magnitude faster than has occurred for millions of years (Doney and Schimel, 2007; Doney et al., 2009). Despite this, only two studies have addressed the potential climate change impacts on extant brachiopods (McClintock et al., 2009; Peck, 2008) and only the former investigated ocean acidification effects where significant dissolution in *Liothyrella uva* (our target taxon) valves was found

after only 14 days in pH 7.4 conditions. However, only empty valves were used so the biological response and ability of *L.uva* to compensate for the impacts of future pH conditions therefore remain to be investigated.

Liothyrella uva (Broderip, 1833) is a large (maximum recorded length is 55 mm), epifaunal, sessile, suspension-feeding terebratulide brachiopod with a circumpolar distribution (Peck et al., 2001). It is found down to 300 metres and is highly abundant in habitats protected from anchor ice and ice scour with reported densities up to 3000 individuals per m² (Foster, 1974; Peck et al., 2001). *L.uva* is typically found attached singly or in clumps to vertical and overhanging rocks around the South Orkney Islands, the Antarctic Peninsula and Peter I Island (Foster, 1974). Previous growth studies on *L.uva* have recorded slower growth rates than temperate rhychonelliform brachiopods and it can live for over 55 years (Peck and Brey, 1996). It also has a limited tolerance to elevated temperature, surviving up to 4.5°C (Peck et al., 2001).

The shell is essential to the existence of *L.uva*, providing protection from predators and preventing any encounters with harmful substances and the loss of body fluids (Harper et al., 2012). Any environmental insult negatively impacting the production, maintenance and/or repair of their shell could thus prove fatal. *L.uva* also becomes naturally damaged and their shells require repairing in the environment (Harper et al., 2009). Given this, and the current focus on ocean acidification, the aims of this study were to determine how shell growth rates and the frequency of shell repair following damage in *L.uva* were affected in a 7 month experiment using predicted mid and end century pH levels.

2. Materials and methods

2.1. Sampling collection

Specimens of *Liothyrella uva* (Broderip, 1833) were hand collected by SCUBA divers from Trolval Island, Ryder Bay, Antarctica (67° 35.44' S, 68° 12.44' W) at 15-25 m depth in May 2012. Animals remained in their conspecific clumps with only the pedicle of the central brachiopod attached to the cliff face being cut ensuring that the majority of specimens were not damaged during collection. Environmental conditions in Ryder Bay at 15-25m depth consist of seawater temperatures that range from -1.8 to +1.0°C, however, temperatures rarely exceed +0.5°C (Clarke et al., 2008), the pH range is 8.04-8.10 (McNeil and Mearns, 2008) and salinity is 33.0-34.0 (Clarke et al., 2008). Brachiopods were kept underwater during the short transportation from the sampling site to the marine laboratory in Rothera and in recirculating aquaria (0.0 ± 0.5°C) whilst being transported back to the UK. Specimens remained in an ambient recirculating seawater system in the UK in similar conditions for a further two weeks to habituate to aquarium conditions before the experiment began.

2.2. Experimental Design

This study was conducted in a recirculating CO₂ microcosm adapted from (Suckling et al., in press) at the British Antarctic Survey (BAS), UK. Four treatments were used where two functioned as lowered pH treatments (pH7.75 and pH7.54) based on the IPCC 'business-as-usual' scenario of the predicted reduction of 0.3-0.5 pH units from the present day average of pH8.05 in oceanic surface waters by 2100 (Table 1) (IPCC, 2013). The third was a pH control as the seawater remained at ambient pH (pH8.05). As a concurrent 2°C increase in temperature is expected to

occur alongside this forecasted decrease in pH by the end of the century (Mitchell et al., 1998), these three systems were maintained at 2°C throughout the experiment. The fourth remaining system was a temperature control which was held at the present-day average surface seawater temperature for Ryder Bay, 0°C (Clarke et al., 2008). The average pH of this treatment was pH7.98 which was slightly lower than, but close to, the pH of the pH control treatment probably as a result of the increased solubility of CO₂ and carbonates (CO₃²⁻) in seawater at the lower temperatures.

The pH of the lowered pH treatments was altered by intermittently bubbling CO₂ gas through a ceramic diffuser to maintain the pH at the predetermined pH levels via a solenoid valve connected to an Aqua Medic pH controlled computer and glass electrode (with plastic shaft) system. The pH control had a similar set up but without the pH control system. An Aqua Medic Ocean Runner power head 2000 circulated the seawater in the mixing tank to ensure a constant pH. Seawater was then gravity fed from each mixing tank at a rate of $0.65 \pm 0.03 \text{ L min}^{-1}$ into the experimental tank. Seawater temperature was manipulated by the use of temperature-controlled laboratories. Air temperature was maintained at -2.5°C in the laboratory with the pH control and both lowered pH treatments but the lifting pumps (Aqua Medic Ocean Runner 3500) and the mixing power heads (Aqua Medic Ocean Runner 2000) in each treatment's mixing tank caused the seawater temperature to raise to the desired ~2°C, with minimal variability (Table 1). The temperature control treatment was situated in the main BAS aquarium where the air temperature was set at -1.5°C and the absence of lifting and circulating pumps caused the seawater temperature to be maintained at ~0°C.

Seawater temperatures ($^{\circ}\text{C}$, Digital Testo 106) and pH_{NIST} (Aquamedic pH controlled computer and electrode system) were monitored and recorded daily. pH_{NIST} was also more accurately measured once a week with a temperature compensated HANNA bench top meter pH/ORP 115 v pH21-01). Salinity (Tropical Marine Centre V2 Handheld refractometer), TCO_2 (mmol L^{-1} ; Ciba Corning TCO_2 Analyzer 965, Olympic Analytical. UK) and nutrient content (silicate and phosphate; according to methods in Nickell et al. (2003)) of each treatment were also measured weekly. Twice a week, the Aqua Medic pH probes were calibrated with NIST certified pH buffers. Other carbonate system parameters, including the partial pressure of CO_2 (pCO_2) and the saturation values for calcite (Ω_{C}) and aragonite (Ω_{A}), were modelled from applying TCO_2 and pH_{NIST} data to the program CO2SYS (Lewis and Wallace, 1998) with refitted constants (Dickson and Millero, 1987; Mehrbach et al., 1973). Brachiopods in each treatment were fed weekly with microalgal concentrate of approximately $331 \times 10^4 \text{ cells L}^{-1}$ which is within the natural range of phytoplankton cell abundance along the west Antarctic Peninsula ($62 - 1150 \times 10^4 \text{ cells L}^{-1}$; (Garibotti et al., 2005; Garibotti et al., 2003). Water quality was maintained and alkalinity and other ions replenished through water changes twice a week and weekly siphoning of the aquaria to remove any debris.

2.3. Growth rates

One hundred and eleven specimens of *L.uva* across a wide size range (2.6-40.9 mm in length) and varying number of individuals in each conspecific clump (30 conspecific clumps with 2-15 individuals and 20 single specimens) were used in this experiment. Prior to the beginning of the experiment, shell lengths of each individual in each conspecific clump were measured to the nearest 0.1 mm using Vernier

calipers. Conspecific clumps were placed into one of the four treatments ensuring there was a similar size range of specimens and numbers of conspecific clumps and single individuals in each treatment. The experiment began in mid June 2012 and lasted for 7 months. At the mid-way and end of the experiment, the length of each specimen was measured again, recorded and the shell edge photographed. Growth rates were then calculated from the increase in length ($\mu\text{m day}^{-1}$).

2.4. Shell repair frequencies

Forty five specimens (8-12 individuals in each treatment) with a range of sizes (5.0-37.0 mm in length) were selected at random and damaged by creating a 1-2 mm deep notch at the valve edge using a metal file. Care was exercised to create notches of equivalent size and severity and not to break shells or cause other damage. After 3.5 and 7 months of the experiment, lengths were measured for each specimen and the damaged region of the shell edge photographed (Figure 1).

2.5. Statistical analyses

All data were analysed using Minitab (Statistical SoftwareTM Version 15). Growth rate data for each treatment were all shown to be significantly different from normal (Anderson-Darling Test; $P < 0.008$). These data were still not normally distributed after square root, logarithmic and double logarithmic transformations because of the presence of zeros in the dataset. Non-parametric Kruskal-Wallis tests were thus used to determine whether damage affected growth rate in each treatment and whether treatment affected growth. When a significant difference was found in the growth rate of undamaged individuals, a further Kruskal-Wallis Multiple Comparisons test was used to identify which treatments were statistically different from each other. Chi-

squared tests were used to determine if treatment affected the percentage of individuals that had completed shell repair, the percentage of individuals that had produced shell after fully repairing their notch and the percentage of individuals that did not grow throughout the experiment.

3. Results

Throughout this 7 month experiment, the saturation states of aragonite and calcite were just below the range usually reported for polar shallow coastal seawater (Table 1; Barry et al. (2010)). In both controls, aragonite was slightly undersaturated ($\Omega < 1$) whereas calcite was supersaturated ($\Omega > 1$). Saturation states with respect to aragonite and calcite in every other treatment were undersaturated (Table 1). The order of treatments with decreasing saturation state with respect to calcite was pH control > temperature control > pH7.75 > pH7.54. Despite these potentially challenging conditions, mortality across all treatments was low at 3.9% (total N=155) with no differences between treatments (temperature control = 4% (n=3), pH control = 8% (n=2), pH7.75 = 0% (n=0) and pH7.54 = 4% (n=1)). Also, the animals showed none of the previously reported signs of stress (slow snapping responses, remaining closed for extended periods, wide gaped when open; Peck (2001)) and responded rapidly to physical stimulation when disturbed, throughout the study. A few females in each treatment also managed to successfully complete brooding and release larvae with large numbers (up to 100) settling on shells of older brachiopods and continuing development to juveniles (E.C. unpubl. obs.). Survival of these newly settled juveniles between 3 and 7 months was also high (over 90%).

3.1. *Shell repair frequencies*

After 3 months, all damaged individuals had at least started shell repair with >50% of individuals fully repairing their notch across all treatments (Table 2). Following the completion of shell repair, >33% of specimens had also produced new shell at the anterior margin (Figure 1). After 7 months, >83% of individuals had completed shell repair in every treatment (Table 2) with no significant difference between treatments ($\chi^2 = 0.839$, $p = 0.840$). Punctae, shell perforations, which are part of the shell structure in this species were present in the repaired notches and the new shell growth. A high proportion (>64%) of specimens had also produced new shell at the growth margin in three of the four treatments, including in both acidified treatments. However, only a moderate proportion (38%) of specimens produced growth at the shell margin in the pH control treatment. Treatment appeared to have an effect on the proportion of individuals that had continued shell production after the completion of repair ($\chi^2 = 9.670$, $p = 0.022$), however, a further Chi-squared test on the same dataset but excluding the pH control indicated there was no significant difference ($\chi^2 = 0.815$, $p = 0.665$). All individuals that had completed shell repair and then made new shell growth after 3 months continued to produce new shell after 7 months (Figure 1).

3.2. *Growth rates*

In the undamaged individuals, the majority, but not all individuals, grew (Figure 2 and 3). In those that did grow, growth rates ranged from 2.4 to 33.7 $\mu\text{m day}^{-1}$ in all three treatments at the higher temperature (pH control, pH7.75 and pH7.54; Figure 2). However, growth rates in the lower temperature trial (temperature control treatment) were lower ranging from 1.2 to 13.2 $\mu\text{m day}^{-1}$ (Figure 3). The only clear ontogenetic

trend in every treatment was that individuals with initial lengths of >32.0 mm produced no growth. Growth rates of undamaged individuals in both the acidified treatments were not significantly different from the pH control (Kruskal-Wallis, $H = 3.96$, $p = 0.138$). A further Kruskal-Wallis test including the temperature control, showed a significant effect of treatment on the growth rate ($H = 13.06$, $p = 0.005$). A Kruskal-Wallis Multiple Comparisons test indicated that higher temperature increased the growth rate over the temperature control in two treatments (pH control: $Z = 2.20$, $p = 0.028$; pH7.75: $Z = 3.21$, $p = 0.001$), however, growth rate in the lowest pH treatment (pH7.54) was not significantly different from the temperature control treatment ($Z = 0.93$, $p = 0.352$).

In damaged individuals that produced new growth at the shell margin, growth rates ranged from 1.2 to 23.1 $\mu\text{m day}^{-1}$ across all treatments with no ontogenetic trend (Figure 2 and 3), and there was no effect of treatment on growth rate of damaged individuals (Kruskal-Wallis, $H = 1.24$, $P = 0.743$).

Growth rates were not significantly different in damaged and undamaged individuals in any treatment (Temperature control – $H = 0.24$, $P = 0.623$; pH control – $H = 2.82$, $P = 0.093$; pH7.75 – $H = 3.96$, $P = 0.046$; pH7.54 – $H = 0.06$, $P = 0.807$). Treatment also had no detectable effect on the proportion of damaged or undamaged individuals that did not grow throughout the experiment ($\chi^2 = 2.590$, $P = 0.459$).

4. Discussion

Clearly *Liothyrella uva* are able to tolerate predicted 2050 and 2100 pH levels as specimens showed no signs of prolonged stress and mortality was low. A similar

mortality rate was found in a study of growth rates of *Liothyrella uva* in their natural habitat ($2\% \text{ yr}^{-1}$; Peck et al. (1997)). The ability of the damaged individuals to repair their shells was not affected by acidified conditions or temperature and >90% of all injured specimens had completed shell repair and >63% had continued shell deposition after completion of repair after 7 months. This suggests that *Liothyrella uva* will be able to repair damage to their shells under the predicted pH and warming conditions in the next century. Similarly, shell repair rates in the gastropod *Subnirrella undulata* were also not affected by ocean acidification, however, the gastropod *Austrocochlea porcata* exhibited a decreased rate (Coleman et al., 2014). This study concluded that different species have varying tolerances and therefore the response is species-specific. Shell repair rates in living brachiopods have only been investigated in purposely damaged specimens of the temperate brachiopod, *Terebratulina retusa*, in ambient seawater where shell repair began after four weeks and caeca developed after eight weeks (Alexander et al., 1992). All notches in the Antarctic brachiopod in this study were made in the anterior margin where new shell is laid down, and so relatively easily repaired. The chemical environments differed between the treatments, but this did not affect the success rate of repairs in *Liothyrella uva*, possibly reflecting an increase in the rate of shell regeneration in the altered pH conditions as seen in some other species (e.g. Wood et al. (2008)). Energetic costs of repairing damaged shells, and whether the repair material has the same shell structure under altered pH conditions is under investigation.

Growth rates in both undamaged and damaged individuals were not affected in the pH control or lowered pH, indicating that *L. uva* should be able to continue shell production in the predicted future ocean acidification conditions even after the

disturbance of completing repair to shell damage. Other studies of shell growth in altered pH conditions, which have all been conducted on non-Antarctic molluscs, have demonstrated mixed responses. Early studies showed reduced growth in two congeneric bivalves, *Mytilus galloprovincialis* (Michaelidis et al., 2005) and *M. edulis* (Berge et al., 2006), after medium-term exposure of 90 and 44 days, respectively. However, acidified conditions as low as pH7.3 (Michaelidis et al., 2005) and pH6.7 (Berge et al., 2006) were used and no growth occurred in the latter. The negative effect in *M. edulis* started between pH7.4 and pH7.1 (Berge et al., 2006), a level which is below the range of pH used in the current work but also below the predicted end century acidified oceans. More recently, studies have predominantly used predicted mid and end century pH levels (>pH7.5). However, short or medium-term exposure (49-84 days, or less) is still generally the norm and this may not allow for acclimation and trans-generational effects. Decreased growth rates in some species of gastropods (*Urosalpinx cinerea*, *Littorina littorea*, *Nucella lamellosa*) are still seen in these more environmentally relevant experiments (Nienhuis et al., 2010; Ries et al., 2009). Although, lowered pH conditions have also had no impact on growth rates in some molluscs (*Arctica islandica* and *Mytilus edulis*) (Hiebenthal et al., 2012; Ries et al., 2009; Thomsen et al., 2010) and *Crepidula fornicata* has even demonstrated a positive effect to intermediate $p\text{CO}_2$ levels (605 and 903 ppm) (Ries et al., 2009). An extensive study on 18 benthic marine calcifiers concluded that the affect of ocean acidification on the calcification process is species-specific (Ries et al., 2009). The ability of *L. uva* to continue shell production in low pH and undersaturated with respect to aragonite and calcite, suggests that this species can generate suitable conditions at the site of calcification (Gazeau et al., 2013; Ries, 2011; Wittmann and Pörtner, 2013). It is believed that marine calcifiers can elevate pH in calcifying

compartments to facilitate calcium carbonate precipitation, however, the mechanisms are largely unknown, especially in the less-studied brachiopods. Potential methods are via either proton channelling (McConnaughey and Falk, 1991), Ca^{2+} -activated proton-translocating ATPase (Cohen and McConnaughey, 2003; McConnaughey and Falk, 1991; McConnaughey and Whelan, 1997), transcellular symporter and co-transporter proton-solute shuttling (McConnaughey and Whelan, 1997) or cellular extrusion of hydroxyl ions into the calcifying medium (Ries, 2011; Ries et al., 2009). Further research is needed to determine which mechanism is used in the study species, however, it is apparent that *Liothyrella uva* has a robust control on the calcification process similar to molluscs.

Increasing the temperature to the predicted warming conditions alongside ocean acidification positively affected the growth rate of *Liothyrella uva*. This suggests shell deposition could occur at a faster rate in 50-100 years as a consequence of elevated temperatures irrespective of the acidity of the oceans. The long-term upper temperature tolerance limit of the Antarctic brachiopod is 4.5°C (Peck, 1989) and higher shell growth rates have been shown with increasing temperatures in other species (Almada-Vilela et al., 1982; Lewis and Cerrato, 1997; Stor et al., 1982), therefore, warming to 2°C is unlikely to negatively impact shell production in *Liothyrella uva*. Several ocean acidification studies have involved combined effects with other environmental variables, particularly with temperature (Courtney et al., 2013; Findlay et al., 2008; Hiebenthal et al., 2012; Martin and Gattuso, 2009; Reynaud et al., 2003; Rodolfo-Metalpa et al., 2011). Organism responses to the interaction of these variables have been varied where temperature has both increased and decreased the effect size of ocean acidification, which has

been found in closely related molluscs, *M.edulis* and *M.galloprovincialis* (Hiebenthal et al., 2012; Rodolfo-Metalpa et al., 2011). Rodolfo-Metalpa et al. (2011) found a negative synergistic effect of low pH (pH_T 7.4) and elevated temperature (25°C) on the gross calcification rate of *M.galloprovincialis* whereas Hiebenthal et al. (2012) found that acidified conditions (pCO_2 1093.3 ± 123.6 μatm and 1654.5 ± 82.7 μatm) did not have an impact on shell growth of closely related *M.edulis*, however, elevated temperature did. Growth in *M.edulis* increased up to 20°C but then decreased at 25°C indicating that 25°C is above this species temperature threshold limit (Hiebenthal et al., 2012; Pörtner, 2008). Growth rates in the temperature control treatment here, which was used to compare the predicted scenarios to current Antarctic conditions, were in the same range (0-15 μm per day) as individuals of the same size (2.5-40.9 mm) in a study of *Liothyrella uva* growth rates in the wild (Peck et al., 1997). No growth occurred in individuals >32.0 mm in the current study. This is the approximate size that this species reaches sexual maturity (Meidlinger et al., 1998; Peck and Holmes, 1989), and our data possibly results from a shift in energy from somatic growth to reproduction.

Even though brachiopods arguably have the largest proportion of skeleton to tissue mass of any group and *Liothyrella uva* inhabits the Southern Ocean which has the naturally lowest carbonate ion saturation levels, our study shows that this species can survive, repair shell damage and deposit new shell after 7 months exposure to forecasted 2050 and 2100 pH conditions. However, there may be a cost to this apparent resilience under future climate change scenarios. The ophiuroid brittlestar, *Amphiura filiformis*, increased its metabolism and calcification to compensate for increased seawater acidity, which came as a substantial cost as muscle wastage

was reported to coincide with this increase (Wood et al., 2008). *L.uva* is a low energy species that uses low metabolic rates to survive the long period of low food availability in the Antarctic winter, although it can opportunistically exploit resuspended benthic microalgae when available (Peck et al., 2005). Increased metabolic costs from elevated temperature may, however, compromise abilities to survive periods of low food availability. The current study is a long-term CO₂ perturbation experiment measuring shell growth on a marine organism with a shell permanently exposed to seawater, therefore, providing insights into how these highly calcium carbonate dependent organisms can acclimate in future CO₂ scenarios. The main mechanisms that organisms can use to respond to environmental change are acclimation and genetic adaptation (Peck, 2011; Somero, 2010). Acclimation to altered temperatures can take 6 months in Antarctic marine invertebrates (Peck et al., 2014), and rates of acclimation to altered pH in negatively impacted species remain to be quantified. Comparable long term studies on other species from polar to tropical environments are essential to increase our knowledge of the capability of these integral organisms to succeed under changing environmental conditions.

Acknowledgements

The authors would like to thank the scientific dive team at the British Antarctic Survey Rothera Research Station for collecting specimens and also to Coleen Suckling, Simon Morley and Rebecca Smith for their help in the set up and maintenance of the ocean acidification experimental system. Emma Cross is supported by the NERC PhD Studentship (NE/T/A/2011).

References

- Alexander, R.R., James, M.A., Ansell, A.D., 1992. Survival and repair of surgical and natural shell damage in the articulate brachiopod *Terebratulina retusa* (Linnaeus). *Hist. Biol.* 6(3), 221-231.
- Almada-Vilela, P.C., Davenport, J., Gruffydd, L.D., 1982. The effects of temperature on the shell growth of young *Mytilus edulis* L. *J. Exp. Mar. Biol. Ecol.* 59, 275-288.
- Arntz, W.E., Brey, T., Gallardo, V.A., 1994. Antarctic zoobenthos. *Oceanogr. Mar. Biol. Annu. Rev.* 32, 251-303.
- Aronson, R.B., Thatje, S., Clarke, A., Peck, L.S., Blake, D.B., Wilga, C.D., Seibel, B.A., 2007. Climate change and invasibility of the Antarctic benthos. *Annu. Rev. Ecol., Evol. Syst.* 38, 129-154.
- Barnes, D.K.A., Peck, L.S., 1996. Epibiota and attachment substrata of deep-water brachiopods from Antarctica and New Zealand. *Phil. Trans. R. Soc. B.* 351, 677-687.
- Barry, J.P., Tyrell, T., Hansson, L., Plattner, G.K., Gattuso, J.P., 2010. Atmospheric CO₂ targets for ocean acidification perturbation experiments. In: Riebesell, U., Fabry, V.F., Hansson, L., Gattuso, J.P. (Eds.), *Guide to best practices for ocean acidification research and data reporting*. Publications Office of the European Union, pp. 53-66.
- Berge, J.A., Bjerkeng, B., Pettersen, O., Schaanning, M.T., Oxnevad, S., 2006. Effects of increased sea water concentrations of CO₂ on growth of the bivalve *Mytilus edulis* L. *Chemosphere* 62(4), 681-687.

- 422 Byrne, M., 2011. Impact of ocean warming and ocean acidification on marine
 423 invertebrate life history stages: vulnerabilities and potential for persistence in a
 424 changing ocean. *Oceanogr. Mar. Biol. Annu. Rev.* 49, 1-42.
- 425 Byrne, M., Przeslawski, R., 2013. Multistressor impacts of warming and acidification
 426 of the ocean on marine invertebrates' life histories. *Integ. Comp. Biol.* 53(4),
 427 582-596.
- 428 Caldeira, K., Wickett, M.E., 2003. Anthropogenic carbon and ocean pH. *Nature* 425,
 429 365.
- 430 Caldeira, K., Wickett, M.E., 2005. Ocean model predictions of chemistry changes
 431 from carbon dioxide emissions to the atmosphere and ocean. *J. Geophys.*
 432 *Res.* 110, C09S04.
- 433 Clarke, A., Meredith, M.P., Wallace, M.I., Brandon, M.A., Thomas, D.N., 2008.
 434 Seasonal and interannual variability in temperature, chlorophyll and
 435 macronutrients in northern Marguerite Bay, Antarctica. *Deep-Sea Research*
 436 *Part II: Topical Studies in Oceanography* 55(18-19), 1988-2006.
- 437 Cohen, A.L., McConnaughey, T.A., 2003. Geochemical perspectives on coral
 438 mineralization. In: Dove, P.M., Weiner, S., De Yoreo, J.J. (Eds.),
 439 *Biom mineralization. The Mineralogical Society of America*, pp. 151-187.
- 440 Coleman, D.W., Byrne, M., Davis, A.R., 2014. Molluscs on acid: gastropod shell
 441 repair and strength in acidifying oceans. *Mar. Ecol. Prog. Ser.* 509, 203-211.
- 442 Courtney, T., Westfield, I., Ries, J.B., 2013. CO₂-induced ocean acidification impairs
 443 calcification in the tropical urchin *Echinometra viridis*. *J. Exp. Mar. Biol. Ecol.*
 444 440, 169-175.
- 445 Cummings, V., Hewitt, J., Van Rooyen, A., Currie, K., Beard, S., Thrush, S., Norkko,
 446 J., Barr, N., Heath, P., Halliday, N.J., Sedcole, R., Gomez, A., McGraw, C.,

- 447 Metcalf, V., 2011. Ocean acidification at high latitudes: potential effects on
448 functioning of the Antarctic bivalve *Laternula elliptica*. PLoS ONE 6(1),
449 e16069.
- 450 Dickson, A.G., Millero, F.J., 1987. A comparison of the equilibrium-constants for the
451 dissociation of carbonic-acid in seawater media. Deep-Sea Res Pt I 34, 1733-
452 1743.
- 453 Doney, S.C., Schimel, D.S., 2007. Carbon and climate system coupling on
454 timescales from the Precambrian to the Anthropocene. Annual Review of
455 Environment and Resources 32, 31-66.
- 456 Doney, S.C., Fabry, V.J., Feely, R.A., Kleypas, J.A., 2009. Ocean acidification: the
457 other CO₂ problem. Ann. Rev. Mar. Sci. 1, 169-192.
- 458 Fabry, V.J., McClintock, J.B., Mathis, J.T., Grebmeier, J.M., 2009. Ocean acidification
459 at high latitudes: the bellwether. Oceanography 22(4), 160-171.
- 460 Findlay, H.S., Kendall, M.A., Spicer, J.I., Turley, C., Widdicombe, S., 2008. Novel
461 microcosm system for investigating the effects of elevated carbon dioxide and
462 temperature on intertidal organisms. Aquatic Biol. 3, 51-62.
- 463 Form, A.U., Riebesell, U., 2012. Acclimation to ocean acidification during long-term
464 CO₂ exposure in the cold-water coral *Lophelia pertusa*. Global Change Biol.
465 18(3), 843-853.
- 466 Foster, M.W., 1974. Recent Antarctic and subantarctic brachiopods. American
467 Geophysical Union, Washington D.C., USA, 189 pp.
- 468 Garibotti, I.A., Vernet, M., Ferrario, M.E., 2005. Annually recurrent phytoplanktonic
469 assemblages during summer in the seasonal ice zone west of the Antarctic
470 Peninsula (Southern Ocean). Deep-Sea Res Pt I 52(10), 1823-1841.

- 471 Garibotti, I.A., Vernet, M., Ferrario, M.E., Smith, R.C., Ross, R.M., Quentin, L.B.,
 472 2003. Phytoplankton spatial distribution patterns along the western Antarctic
 473 Peninsula (Southern Ocean). *Mar. Ecol. Prog. Ser.* 261, 21-39.
- 474 Gattuso, J.P., Mach, K.J., Morgan, G., 2013. Ocean acidification and its impacts: an
 475 expert survey. *Clim. Change* 117, 725-738.
- 476 Gazeau, F., Parker, L.M., Comeau, S., Gattuso, J.-P., O'Connor, W.A., Martin, S.,
 477 Pörtner, H.-O., Ross, P.M., 2013. Impacts of ocean acidification on marine
 478 shelled molluscs. *Mar. Biol.* 160(8), 2207-2245.
- 479 Harper, E.M., 2000. Are calcitic layers an effective adaptation against shell
 480 dissolution in the Bivalvia? *J. Zool. Lond.* 251(2), 179-186.
- 481 Harper, E.M., Peck, L.S., Hendry, K.R., 2009. Patterns of shell repair in articulate
 482 brachiopods indicate size constitutes a refuge from predation. *Mar. Biol.*
 483 156(10), 1993-2000.
- 484 Harper, E.M., Clark, M.S., Hoffman, J.I., Philipp, E.E., Peck, L.S., Morley, S.A., 2012.
 485 Iceberg scour and shell damage in the Antarctic bivalve *Laternula elliptica*.
 486 *PLoS ONE* 7(9), e46341.
- 487 Havenhand, J.N., Schlegel, P., 2009. Near-future levels of ocean acidification do not
 488 affect sperm motility and fertilisation kinetics in the oyster *Crassostrea gigas*.
 489 *Biogeosciences* 6, 3009-3015.
- 490 Hiebenthal, C., Philipp, E.E.R., Eisenhauer, A., Wahl, M., 2012. Effects of seawater
 491 $p\text{CO}_2$ and temperature on shell growth, shell stability, condition and cellular
 492 stress of Western Baltic Sea *Mytilus edulis* (L.) and *Arctica islandica* (L.). *Mar.*
 493 *Biol.* 160(8), 2073-2087.
- 494 IPCC, 2013. Climate Change 2013: The Physical Science Basis. In: Stocker, T.F.,
 495 Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia,

- 496 Y., Bex, V., Midgley, P.M. (Eds.), Working Group I Contribution to the Fifth
497 Assessment Report of the Intergovernmental Panel on Climate Change,
498 Cambridge, United Kingdom and New York, NY, USA, pp. 1552.
- 499 Kelly, M.W., Hofmann, G.E., Hoffmann, A., 2013. Adaptation and the physiology of
500 ocean acidification. *Funct. Ecol.* 27(4), 980-990.
- 501 Lewis, D.E., Cerrato, R.M., 1997. Growth uncoupling and the relationship between
502 shell growth and metabolism in the soft shell clam *Mya arenaria*. *Mar. Ecol.*
503 *Prog. Ser.* 158, 177-189.
- 504 Lewis, E., Wallace, D.W.R., 1998. Carbon Dioxide Information Analysis Center, Oak
505 Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee,
506 US.
- 507 Martin, S., Gattuso, J.-P., 2009. Response of Mediterranean coralline algae to ocean
508 acidification and elevated temperature. *Global Change Biol.* 15(8), 2089-2100.
- 509 McClintock, J.B., Angus, R.A., McDonald, M.R., Amsler, C.D., Catledge, S.A., Vohra,
510 Y.K., 2009. Rapid dissolution of shells of weakly calcified Antarctic benthic
511 macroorganisms indicates high vulnerability to ocean acidification. *Antarct.*
512 *Sci.* 21(05), 449.
- 513 McConnaughey, T.A., Falk, R.H., 1991. Calcium-proton exchange during algae
514 calcification. *Biol. Bull.* 180, 185-195.
- 515 McConnaughey, T.A., Whelan, J.F., 1997. Calcification generates protons for nutrient
516 and bicarbonate uptake. *Earth-Sci. Rev.* 42, 95-117.
- 517 McNeil, B.I., Matear, R.J., 2008. Southern Ocean acidification: a tipping point at 450-
518 ppm atmospheric CO₂. *PNAS* 105(48), 18860-18864.

- 519 Mehrbach, C., Culberson, C.H., Hawley, J.E., Pytkowicz, R.M., 1973. Measurement
520 of apparent dissociation constants of carbamic acid in seawater at
521 atmospheric pressure. *Limnol. Oceanogr.* 18, 897-907.
- 522 Meidinger, K., Tyler, P.A., Peck, L.S., 1998. Reproductive patterns in the Antarctic
523 brachiopod *Liothyrella uva*. *Mar. Biol.* 132, 153-162.
- 524 Michaelidis, B., Ouzounis, C., Paleras, A., Portner, H.O., 2005. Effects of long-term
525 moderate hypercapnia on acid-base balance and growth rate in marine
526 mussels *Mytilus galloprovincialis*. *Mar. Ecol. Prog. Ser.* 293, 109-118.
- 527 Mitchell, J.F.B., Johns, T.C., Senior, C.A., 1998. Transient response to increasing
528 greenhouse gases using models with and without flux adjustment, Hadley
529 Centre Technical Note 2. Available from Met Office, London Road Bracknell,
530 RG12 2SZ, UK.
- 531 Nickell, L.A., Black, K.D., Hughes, D.J., Overnell, J., Brand, T., Nickell, T.D., Breuer,
532 E., Martyn Harvey, S., 2003. Bioturbation, sediment fluxes and benthic
533 community structure around a salmon cage farm in Loch Creran, Scotland. *J.*
534 *Exp. Mar. Biol. Ecol.* 285-286, 221-233.
- 535 Nienhuis, S., Palmer, A.R., Harley, C.D., 2010. Elevated CO₂ affects shell dissolution
536 rate but not calcification rate in a marine snail. *Proc. R. Soc. B.* 277(1693),
537 2553-2558.
- 538 Orr, J.C., Fabry, V.J., Aumont, O., Bopp, L., Doney, S.C., Feely, R.A.,
539 Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R.M., Lindsay, K.,
540 Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R.G., Plattner,
541 G.K., Rodgers, K.B., Sabine, C.L., Sarmiento, J.L., Schlitzer, R., Slater, R.D.,
542 Totterdell, I.J., Weirig, M.F., Yamanaka, Y., Yool, A., 2005. Anthropogenic

- 543 ocean acidification over the twenty-first century and its impact on calcifying
544 organisms. *Nature* 437(7059), 681-686.
- 545 Pandolfi, J.M., Connolly, S.R., Marshall, D.J., Cohen, A.L., 2011. Projecting coral reef
546 futures under global warming and ocean acidification. *Science* 333(6041),
547 418-422.
- 548 Parker, L.M., Ross, P.M., O'Connor, W.A., Borysko, L., Raftos, D.A., Pörtner, H.,
549 2012. Adult exposure influences offspring response to ocean acidification in
550 oysters. *Global Change Biol.* 18, 82-92.
- 551 Pearse, J.S., McClintock, J.B., Bosch, I., 1991. Reproduction of Antarctic benthic
552 marine invertebrates: Tempos, modes and timing. *Am. Zool.* 31, 65-80.
- 553 Peck, L.S., 1989. Temperature and basal metabolism in two Antarctic marine
554 herbivores. *J. Exp. Mar. Biol. Ecol.* 127, 1-12.
- 555 Peck, L.S., 1993. The tissues of articulate brachiopods and their value to predators.
556 *Phil. Trans. R. Soc. B.* 339(1287), 17-32.
- 557 Peck, L.S., 2001. Physiology. In: Carlson, S., Sandy, M. (Eds.), *Brachiopods ancient*
558 *and modern: a tribute to G. Arthur Cooper. The Paleontology Society Papers,*
559 *New Haven, Connecticut: Yale University, pp. 89-104.*
- 560 Peck, L.S., 2008. Brachiopods and climate change. *Earth Env. Sci. T. R. So.* 98(3-4),
561 451-456.
- 562 Peck, L.S., 2011. Organisms and responses to environmental change. *Mar. Genom.*
563 4, 237-243.
- 564 Peck, L.S., Holmes, L.J., 1989. Seasonal and ontogenetic changes in tissue size in
565 the Antarctic brachiopod *Liothyrella uva* (Broderip, 1833). *J. Exp. Mar. Biol.*
566 *Ecol.* 134, 25-36.
- 567 Peck, L.S., Brey, T., 1996. Bomb signals in old Antarctic brachiopods. *Nature* 380.

- 568 Peck, L.S., Conway, L.Z., 2000. The myth of metabolic cold adaptation: oxygen
569 consumption in stenothermal Antarctic bivalves. Geol. Soc. London Spec.
570 Publ. 177, 441-450.
- 571 Peck, L.S., Brockington, S., Brey, T., 1997. Growth and metabolism in the Antarctic
572 brachiopod *Liothyrella uva*. Phil. Trans. R. Soc. B. 352, 851-858.
- 573 Peck, L.S., Meidlinger, K., Tyler, P.A., 2001. Developmental and settlement
574 characteristics of the Antarctic brachiopod *Liothyrella uva* (Broderip 1833). In:
575 Brunton, C.H.C., Cocks, L.R., Long, S.L. (Eds.), Brachiopods Past and
576 Present. Taylor and Francis, London and New York.
- 577 Peck, L.S., Barnes, D.K.A., Willmott, J., 2005. Responses to extreme seasonality in
578 food supply: diet plasticity in Antarctic brachiopods. Mar. Biol. 147(2), 453-
579 463.
- 580 Peck, L.S., Morley, S.A., Pörtner, H., Clark, M.S., 2007. Thermal limits of burrowing
581 capacity are linked to oxygen availability and size in the Antarctic clam
582 *Laternula elliptica*. Oecologia 154, 479-484.
- 583 Peck, L.S., Morley, S.A., Richard, J., Clark, M.S., 2014. Acclimation and thermal
584 tolerance in Antarctic marine ectotherms. J. Exp. Biol. 217, 16-22.
- 585 Pespeni, M.H., Sanford, E., Gaylord, B., Hill, T.M., Hosfelt, J.D., Jaris, H.K., LaVigne,
586 M., Lenz, E.A., Russell, A.D., Young, M.K., Palumbi, S.R., 2013. Evolutionary
587 change during experimental ocean acidification. PNAS 110(17), 6937-6942.
- 588 Pörtner, H., 2008. Ecosystem effects of ocean acidification in times of ocean
589 warming: a physiologist's view. Mar. Ecol. Prog. Ser. 373, 203-217.
- 590 Revelle, R.R., Fairbridge, R.W., 1957. Carbonates and carbon dioxide, Geological
591 Society of America Memoirs, pp. 239-296.

- Reynaud, S., Leclercq, N., Romaine-Lioud, S., Ferrier-Pages, C., Jaubert, J.,
 Gattuso, J.P., 2003. Interacting effects of CO₂ partial pressure and
 temperature on photosynthesis and calcification in a scleractinian coral. *Global
 Change Biol.* 9, 1660-1668.
- Ries, J.B., 2011. A physicochemical framework for interpreting the biological
 calcification response to CO₂-induced ocean acidification. *Geochim.
 Cosmochim. Acta* 75(14), 4053-4064.
- Ries, J.B., Cohen, A.L., McCorkle, D.C., 2009. Marine calcifiers exhibit mixed
 responses to CO₂-induced ocean acidification. *Geology* 37(12), 1131-1134.
- Rodolfo-Metalpa, R., Houlbreque, F., Tambutte, E., Boisson, F., Baggini, C., Patti,
 F.P., Jeffree, R., Fine, M., Foggo, A., Gattuso, J.P., Hall-Spencer, J.M., 2011.
 Coral and mollusc resistance to ocean acidification adversely affected by
 warming. *Nature Clim. Change* 1(6), 308-312.
- Somero, G.N., 2010. The physiology of climate change: how potentials for
 acclimatization and genetic adaptation will determine 'winners' and 'losers'. *J.
 Exp. Biol.* 213, 912-920.
- Stor, J.F., Costa, A.L., Prawel, D.A., 1982. Effects of temperature on calcium
 deposition in the hard-shelled clam, *Mercenaria mercenaria*. *J. Therm. Biol.* 7,
 57-61.
- Suckling, C.C., Clark, M.S., Richard, J., Morley, S.A., Thorne, M.A.S., Harper, E.M.,
 Peck, L.S., in press. Adult acclimation to combined temperature and pH
 stressors significantly enhances reproductive outcomes.
- Thomsen, J., Gutowska, M.A., Saphörster, J., Heinemann, A., Trübenbach, K.,
 Fietzke, J., Hiebenthal, C., Eisenhauer, A., Körtzinger, A., Wahl, M., Melzner,
 F., 2010. Calcifying invertebrates succeed in a naturally CO₂-rich coastal

617 habitat but are threatened by high levels of future acidification.
618 Biogeosciences 7(11), 3879-3891.

619 Vermeij, G.J., 1978. Biogeography and adaptation: patterns of marine life. Harvard
620 University Press, Cambridge, Massachusetts, USA.

621 Watson, S.-A., Peck, L.S., Tyler, P.A., Southgate, P.C., Tan, K.S., Day, R.W., Morley,
622 S.A., 2012. Marine invertebrate skeleton size varies with latitude, temperature
623 and carbonate saturation: implications for global change and ocean
624 acidification. Global Change Biol. 18(10), 3026-3038.

625 Wittmann, A.C., Pörtner, H., 2013. Sensitivities of extant animal taxa to ocean
626 acidification. Nature Clim. Change 3, 995-1001.

627 Wood, H.L., Spicer, J.I., Widdicombe, S., 2008. Ocean acidification may increase
628 calcification rates, but at a cost. Proc. R. Soc. B. 275(1644), 1767-1773.

629

Figure legends

Figure 1 – Examples of shell repair and further shell growth in damaged individuals after 3 months (top row) and 7 months (bottom row) in the following four treatments: pH control (a, e), temperature control (b, f), pH7.75 (c, g) and pH7.54 (d, h). The same individual is shown in each treatment at both growth periods where the arrow indicates the notch created at the start of the experiment. Scale bar = 100µm. (Colour on Web only)

Figure 2 – Growth rates of specimens that were left undamaged (●) and were damaged at the start of the experiment (○) after 7 months in the treatments kept at the higher temperature; (a) pH control ($\text{pH}8.05 \pm 0.03$, $T=1.7^{\circ}\text{C} \pm 0.3$), (b) pH7.75 ($\text{pH}7.75 \pm 0.03$, $T=1.9^{\circ}\text{C} \pm 0.4$) and (c) pH7.54 ($\text{pH}7.54 \pm 0.03$, $T=2.2^{\circ}\text{C} \pm 0.4$).

Figure 3 - Growth rates of specimens that were left undamaged (●) and were damaged at the start of the experiment (○) after 7 months in the low temperature control ($\text{pH}7.98 \pm 0.02$, $T=-0.3^{\circ}\text{C} \pm 0.1$).