



Treatment and outcomes of UK and German patients with relapsed intracranial germ cell tumors following uniform first-line therapy

Journal:	<i>International Journal of Cancer</i>
Manuscript ID	IJC-17-0134.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Murray, Matthew; Cambridge University Hospitals NHS Foundation Trust, Department of Paediatric Haematology and Oncology Bailey, Shivani; Cambridge University Hospitals NHS Foundation Trust, Department of Paediatric Haematology and Oncology Heinemann, Katja; University Children's Hospital, Department of Paediatric Haematology and Oncology Mann, Jillian; Birmingham Children's Hospital NHS Foundation Trust, Department of Paediatric Haematology and Oncology Göbel, Ulrich; German Paediatric Surveillance Unit (ESPED), Heinrich-Heine University, Medical Faculty Saran, Frank; Royal Marsden Hospital Sutton, Department of Oncology Hale, Juliet; Royal Victoria Infirmary, Department of Paediatric Haematology and Oncology Calaminus, Gabriele; University Children's Hospital, Department of Paediatric Haematology and Oncology Nicholson, James; Cambridge University Hospitals NHS Foundation Trust, Department of Paediatric Haematology and Oncology
Key Words:	germ cell tumor, high-dose chemotherapy, intracranial, non-germinoma, relapse, re-irradiation

SCHOLARONE™
Manuscripts

1
2
3 **Treatment and outcomes of UK and German patients with relapsed intracranial**
4 **germ cell tumors following uniform first-line therapy**
5
6

7 Matthew J. Murray^{1*}, Shivani Bailey^{1*}, Katja Heinemann², Jillian Mann³, Ulrich K
8 Göbel⁴, Frank Saran⁵, Juliet P. Hale⁶, Gabriele Calaminus², James C. Nicholson¹.
9
10

11
12
13
14 ¹ Department of Paediatric Haematology and Oncology, Cambridge University
15 Hospitals NHS Foundation Trust, Hills Road, Cambridge, CB2 0QQ, UK
16

17
18 ² Department of Paediatric Haematology and Oncology, University Children's
19 Hospital, Robert Koch Strasse, Bonn, Germany
20

21
22
23 ³ Department of Paediatric Haematology and Oncology, Birmingham Children's
24 Hospital, Steelhouse Lane, Birmingham, B4 6NH, UK
25

26
27 ⁴ German Paediatric Surveillance Unit (ESPED), Medical Faculty, Heinrich-Heine
28 University, Düsseldorf, Germany
29

30
31
32 ⁵ Department of Oncology, Royal Marsden Hospital, Downs Rd, Sutton, Surrey SM2
33 5PT, UK
34

35
36 ⁶ Department of Paediatric Haematology and Oncology, Royal Victoria Infirmary,
37 Queen Victoria Rd, Newcastle upon Tyne, Tyne and Wear, NE1 4LP, UK
38
39
40
41
42

43 **Corresponding author:** Dr Matthew Murray, Department of Paediatric Haematology
44 and Oncology, Cambridge University Hospital NHS Foundation Trust, Hills Road,
45 Cambridge, CB2 0QQ, UK. Email: mjm16@cam.ac.uk; Telephone: 0044 (0)1223
46 765066; Fax: 0044 (0)1223 586794
47
48
49
50

51
52 **Footnote:** * These authors contributed equally
53

54 **Funding:** None
55

56 **Conflict of Interest:** There are no conflicts to disclose
57
58
59
60

Novelty and impact statements

A retrospective assessment of the value of high-dose chemotherapy and autologous stem cell rescue (HDC+AuSCR) and re-irradiation in patients with relapsed intracranial germ cell tumors (GCTs). Those with germinoma had reasonable outcomes with either standard-dose chemotherapy with re-irradiation or HDC+AuSCR; those with relapsed non-germinomatous GCTs (NGGCTs) were only salvaged with HDC+AuSCR, but then only rarely. Counselling regarding poor outcomes for relapsed NGGCT patients is required prior to embarking on treatment.

Word count: 4,021; **Figures/Tables:** Four Figures, Two Tables

Running title: Relapsed intracranial germ cell tumors

Keywords: germ cell tumor; high-dose chemotherapy; intracranial; non-germinoma; relapse, re-irradiation.

Presented in part at: 15th International Symposium on Pediatric Neuro-oncology, Toronto, Canada, June 24-27, 2012 and 3rd International CNS Germ Cell Tumor Symposium, Cambridge, UK, April 17-20, 2013.

Abstract

We aimed to retrospectively assess treatments/outcomes, including the value of high-dose-chemotherapy and autologous-stem-cell-rescue (HDC+AuSCR) and re-irradiation, in a large, European patient-cohort with relapsed intracranial germ-cell-tumors (GCTs) receiving uniform first-line therapy, including radiotherapy as standard-of-care. Fifty-eight UK/German patients (48 male/10 female) with relapsed intracranial-GCTs [13 germinoma/45 non-germinomatous GCT(NGGCT)] treated 1996-2010 as per the SIOP-CNS-GCT-96 protocol were evaluated. For germinoma, six patients relapsed with germinoma and five with NGGCT (one palliative, one teratoma patient excluded). Five-year overall-survival (OS) for the whole-group (n=11) was 55%. Four of six germinoma relapses and two of five relapsing with NGGCT were salvaged; patients were salvaged with either standard-dose-chemotherapy (SDC) and re-irradiation or HDC+AuSCR with/without re-irradiation. Of 45 relapsed NGGCT patients, 13 were excluded (3 non-protocol adherence, 5 teratoma, 5 palliation). Five-year OS for the remaining 32 relapsed malignant NGGCT patients treated with curative intent was 9%(95%CI: 2-26%). By treatment received, 5-year OS for the 10 patients receiving SDC and 22 patients treated with intention for HDC+AuSCR was 0%(0-0%) and 14%(3-36%), respectively. The three relapsed NGGCT survivors had raised HCG markers alone; two received additional irradiation. Patients with relapsed germinoma had better 5-year OS than those with relapsed NGGCT (55%vs.9%; p=0.007). Patients with relapsed germinoma were salvaged both with SDC and re-irradiation or HDC+AuSCR with/without re-irradiation; which both represent valid treatment options. Outcomes for malignant relapse following initial diagnosis of NGGCT were exceptionally poor; the few

1
2
3 survivors received thiotepa-based HDC+AuSCR, which is a treatment option at first
4
5 malignant relapse for such patients, with further surgery/irradiation where feasible.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Introduction

Malignant intracranial germ cell tumors (GCTs) are a rare group of predominantly midline neoplasms¹, often classified into germinoma or non-germinomatous tumors (NGGCTs). The latter group is particularly heterogeneous and includes the malignant subtypes yolk sac tumor (YST) and choriocarcinoma (CHC) [which secrete the tumor markers alpha-fetoprotein (AFP) and human chorionic gonadotropin (HCG), respectively], as well as embryonal carcinoma (EC)². Most NGGCTs are mixed malignant GCTs, containing more than one subtype, which makes diagnostic work-up, classification and management more complex^{3,4}.

Germinomas comprise ~60% of all intracranial GCTs⁵, and are extremely sensitive to multimodal therapy⁶, including both chemotherapy and radiotherapy. Recent strategies have focused on maintaining excellent event-free (EFS) and overall (OS) survival rates, whilst reducing toxicity and late-effects^{5,7-9}. The European non-randomized trial, SIOP-CNS-GCT-96, compared craniospinal radiotherapy (CSI) alone (24 Gy plus 16 Gy tumor-boost), Option-A, with two courses of combination 'carboPEI' chemotherapy (carboplatin/etoposide/ifosfamide) and focal radiotherapy (40 Gy), Option-B, for localized intracranial germinomas⁵. Metastatic disease was treated with either CSI alone or combination chemotherapy (as for localized disease) and CSI. Outcomes were excellent, with five-year EFS of 94% and 88% for localized patients treated with Options-A and -B, respectively⁵. An excess of ventricular relapses for Option-B patients was addressed by the addition of whole ventricular irradiation for the current SIOP-CNS-GCT-II study⁵. In germinoma patients with metastatic disease, five-year EFS and OS was 98%⁵.

1
2
3 Patients with NGGCT on the SIOP-CNS-GCT-96 study received four courses of PEI
4 chemotherapy (cisplatin/etoposide/ifosfamide), followed by 54 Gy focal radiotherapy
5 for localized disease and 30 Gy CSI with 24 Gy tumor-boost for metastatic disease.
6
7 The rationale for treatment was based on studies that used a single chemotherapy
8 protocol to treat both extracranial and intracranial GCT, with the addition of radiation
9 for intracranial cases^{4,10}. Progression-free survival (PFS) was inferior to that observed
10 for germinoma, at 69% and 67% for localized and metastatic disease, respectively¹¹.
11
12 Recently, good outcomes for intracranial NGGCT patients were reported in a
13 Children's Oncology Group study¹².
14
15
16
17
18
19
20
21
22
23
24

25 As a consequence of the rarity of intracranial GCTs and the general success of their
26 treatment, relapses are rare. As a result, there is little available evidence to guide
27 optimal management¹³. A widely reported study of relapsed intracranial GCT
28 evaluated the efficacy of HDC+AuSCR in a small, highly heterogeneous group of
29 patients who had received variable first-line treatments¹⁴. Eight of the 21 patients
30 reported had not received prior radiotherapy, which would now be considered
31 standard-of-care². Furthermore, there was unavoidable selection bias in the patients
32 described, as those who experienced rapid tumor progression or did not have a good
33 response to re-induction chemotherapy did not proceed to HDC+AuSCR and were
34 therefore not reported¹⁴. As a result, the encouraging outcomes reported (78% and
35 33% survival for germinoma and NGGCT, respectively)¹⁴ are likely to be an
36 overestimate of the true salvage rates for patients who relapse following optimal first-
37 line treatment. A more recent small study of HDC+AuSCR for relapsed intracranial
38 GCT patients reported promising survival rates¹⁵, however, first-line treatments were
39 variable.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Here, we present data from the largest cohort of patients with relapsed intracranial GCTs described to date, treated with first-line treatment as per the European SIOP-CNS-GCT-96 trial protocol, which included radiotherapy as standard. The results described here will inform future approaches to the management of these patients, with the aim of optimizing survival.

For Peer Review

Patients and Methods

Methods. Both the UK and German groups, as major contributors to the SIOP-CNS-GCT-96 trial, treated sufficient numbers of patients according to the protocol to have a suitable patient cohort experiencing relapse to describe separately. In addition, UK and German groups made resources available to undertake this project. Accordingly, follow-up data on UK and German patients treated on the SIOP-CNS-GCT-96 trial, stored on a central database, was augmented in the UK by additional questionnaire-based data-collection, which captured additional non-trial patients who were treated according to the protocol. This anonymized service evaluation required no additional ethical approvals. Resources were not available to undertake additional questionnaire-based data collection to identify non-trial patients in Germany, or other countries who participated in the trial. Patients included in the study received treatment between 1996 and 2010. The SIOP-CNS-GCT-96 trial was open from 1996 and following closure in the UK in 2006, non-trial patients were treated on trial-based guidelines. Data were collected during 2011, with a further update to exclude further events in 2014. Data requested included full clinico-pathological details at primary diagnosis and relapse, confirmation of first-line treatment, relapse treatments received and outcome. The clinical characteristics of the subset of patients described here were similar to the trial group as a whole.

Overall patient cohort. A total of 58 relapsed central nervous system (CNS) GCT cases were identified from the UK (n=33) and Germany (n=25), comprising 48 male and 10 female patients, with an overall median age at diagnosis of 14.0 years (y) (range 1.4-30.0 y) and median follow-up from original diagnosis of 23 months (m)

1
2
3 (range 7-203 m). There were 13 cases with an original diagnosis of germinoma (10
4 UK/three German; nine male/four female) and 45 patients who relapsed following an
5 NGGCT diagnosis (23 UK/22 German; 39 male/six female). Of the 58 patients
6 identified in total, 41 (71%) were treated on trial (16 UK and 25 German patients).
7
8 The trial database only captured data up until the time of relapse, and therefore
9 subsequent information regarding treatments and outcome available on the database
10 was variable. Consequently, further anonymized information was sought from treating
11 centers. An additional 17 (29%) UK non-trial patients were identified by
12 questionnaire-based survey, hence the relative larger number of UK relapses. For
13 germinoma, the 10 UK patients comprised five trial and five non-trial patients (in
14 addition to the three German trial patients). For the 23 UK NGGCT cases, 11 patients
15 were treated on trial with a further 12 non-trial patients identified (in addition to the
16 22 German trial patients). From the trial database, overall outcomes for UK and
17 German patients receiving first-line treatment were similar to overall trial results for
18 patients from all participating countries (data not shown), and hence represented an
19 appropriate and non-biased cohort in which to study relapse. It should be noted that
20 the purpose of this work was not to assess the actual relapse rate for CNS GCTs and,
21 indeed, as both trial and non-trial patients were included, it was not possible to
22 provide an overall denominator defining the total number of patients treated upfront,
23 which are provided elsewhere ^{5,11}. See Figure 1 for a summary flow-diagram of
24 patients included in the study.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49
50
51 *Relapsed germinoma patients – initial diagnostic and staging work-up.* Clinico-
52 pathological data at diagnosis are summarized in Supplementary Table 1. The median
53 age at diagnosis of the 13 patients who relapsed following a germinoma diagnosis was
54
55
56
57
58
59
60

1
2
3 12.3 y (range 4.8-16.3 y). Primary disease site was pineal gland (n=5), suprasellar
4
5 region (n=4), bifocal (i.e. suprasellar and pineal regions, considered localized disease
6
7 if no evidence of disease elsewhere; n=3) and basal ganglia (n=1). All cases
8
9 underwent histological confirmation of germinoma diagnosis. One of 13 cases (8%)
10
11 was metastatic at diagnosis.
12
13

14
15
16 *Relapsed germinoma patients – first-line treatment received and response.* Data are
17
18 summarized in Supplementary Table 1. Seven of 13 patients (54%) received CSI as
19
20 per Option-A treatment on the SIOP-CNS-GCT-96 protocol, and 6/13 (46%) patients
21
22 received Option-B treatment with combined chemotherapy and focal radiotherapy. In
23
24 the absence of central radiological review to assess response in the SIOP-CNS-GCT-
25
26 96 trial, local radiological reports were utilized for both trial and non-trial patients in
27
28 this study. Eight of 13 patients (62%) achieved a complete response (CR) to first-line
29
30 therapy, 4/13 achieved a partial response (PR; 31%) and data were unavailable for
31
32 one patient (8%).
33
34
35
36
37

38
39 *Relapsed NGGCT patients – initial diagnostic and staging work-up.* Clinico-
40
41 pathological data are summarized in Supplementary Table 2. The median age at
42
43 diagnosis of the 45 patients who relapsed following an NGGCT diagnosis was 14.4 y
44
45 (range 1.4-30.0 y). Primary disease site was pineal (n=27), suprasellar (n=8), bifocal
46
47 (n=4), basal ganglia (n=3) and cerebral hemisphere (n=3). Forty-one of 45 cases
48
49 (91%) had markers above SIOP-CNS-GCT-96 defined thresholds (i.e. serum or CSF
50
51 AFP >25 ng/ml and/or HCG >50 IU/l) and the remaining four cases had negative
52
53 markers, so underwent a neurosurgical biopsy which histologically confirmed EC. Six
54
55
56
57
58
59
60

1
2
3 of 45 cases (13%) were metastatic at diagnosis, defined by imaging and/or CSF
4
5 cytology.
6
7

8
9
10 *Relapsed NGGCT patients – first-line treatment received and response.* Data are
11 summarized in Supplementary Table 2. Forty-two of 45 patients (93%) received
12 standard PEI chemotherapy, and three (7%) received non-protocol chemotherapy and
13 were excluded from subsequent relapse outcome analysis: one received ‘high-dose’
14 PEI due to presentation after trial closure with AFP ≥ 1000 ng/ml, as per the SIOP-
15 CNS-GCT-II trial treatment prior to its opening (physician choice), and two received
16 other chemotherapy (no indication given). Forty-three of 45 patients (96%) received
17 radiotherapy, one had progressive disease prior to receiving radiotherapy and the
18 other was considered too young (19m) at diagnosis. As per standard criteria¹⁶, 14 of
19 45 patients (31%) achieved CR at the end of first-line treatment, 20/45 (44%)
20 achieved partial remission (PR), 6/45 (13%) had stable disease (SD), 4/45 (9%) had
21 progressive disease (PD) and data were unavailable for one patient (2%). Five of the
22 26 patients (19%) with a PR/SD response underwent surgical resection for residual
23 disease.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 *Statistical analysis.* EFS was calculated from day of relapse to date of progression or
44 death, whichever was earlier, with censoring taken on the date of last clinical contact.
45 OS was calculated from date of relapse to date of death. Estimated EFS/OS rates were
46 calculated by Kaplan-Meier survival method. EFS/OS values were given with 95%
47 confidence intervals (CI) in parentheses, and differences in outcome between groups
48 assessed by log-rank test ($p \leq 0.05$ significant), where numbers were sufficient for such
49 statistical analyses/comparisons.
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Results

Relapsed patients – details of relapse. Data are summarized in Figure 1, Table 1, Table 2 and Supplementary Results.

Relapsed germinoma patients – relapse treatments and outcomes. Data are summarized in Figure 1 and Table 1. Two patients (both treated initially on Option-A) were excluded from further evaluation – one was a pure immature teratoma relapse and one declined active treatment. The majority of the 11 remaining patients who had a malignant relapse treated with curative intent following original germinoma diagnosis received re-induction chemotherapy with a platinum-based (carboplatin or cisplatin) strategy (Table 1), with variable subsequent treatments. This resulted in a 5-year EFS and OS of 5/11 (45%) and 6/11 (55%), respectively, with median EFS and OS of 20 m and 53 m for the whole group, respectively (Figure 2). Median OS for the six survivors was 84 m (range 53-194 m). We first compared OS by original first-line treatment, irrespective of subsequent relapse chemotherapy delivered. Of the five patients treated on Option-A, two survived (40%), with four of six surviving (67%) who initially received Option-B (Supplementary Table 1 and Table 1). Next, we compared the cohort by relapse chemotherapy/re-irradiation received. Three out of seven patients (43%) treated with standard-dose-chemotherapy (SDC) had complete responses and no further relapses/events; the figure was 2/4 (50%) who received HDC+AuSCR at first relapse. Two patients received HDC+AuSCR following further relapse after SDC, of whom one was salvaged, taking the OS of patients treated with SDC at initial relapse to 4/7 (57%) (Table 1). Six out of seven patients (86%) treated with SDC received additional radiotherapy (four focal, plus two patients who had CSI having received focal radiotherapy at original diagnosis). One SDC patient, who had

1
2
3 CSI at diagnosis, received no further radiotherapy at relapse. None of the four patients
4
5 who were treated with intent for HDC+AuSCR received additional radiotherapy at
6
7 relapse. Finally, we compared the group by presence or absence of positive tumor
8
9 markers at relapse. For marker-negative (i.e. germinoma) cases at relapse, 5-year EFS
10
11 and OS was 3/6 (50%) and 4/6 (67%), respectively. Three of these six patients were
12
13 treated with SDC (in addition to focal re-irradiation), of whom two (67%) were
14
15 salvaged – one with SDC alone and the other with additional HDC+AuSCR following
16
17 a second relapse. Three of the six patients were treated with HDC+AuSCR at first
18
19 relapse, with no further re-irradiation, of whom two (67%) were salvaged. Of the five
20
21 patients who relapsed and were marker-positive (i.e. classified as NGGCT at relapse),
22
23 two patients of four (50%) were successfully treated with SDC alone (Table 1). Of
24
25 these two survivors, one had a CNS relapse (treated with SDC and CSI) with a
26
27 maximum HCG level of only 55 IU/l (patient G12); the other had an exclusively
28
29 extracranial relapse due to tumor seeding through a ventriculo-peritoneal shunt, with
30
31 only germinoma confirmed on cytology of ascitic fluid and an HCG of 397 IU/l
32
33 (G13), described elsewhere¹⁷. The other germinoma patient who relapsed with
34
35 positive HCG (patient G2; maximum HCG value 734 IU/l), received SDC, but
36
37 subsequently died following further relapses after having received HDC+AuSCR.
38
39 One germinoma patient relapsing as NGGCT received HDC+AuSCR at first relapse,
40
41 who also subsequently died (Table 1).
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Relapsed NGGCT patients - relapse treatment and outcomes. Data are summarized in Figure 1 and Table 2. Of the 45 relapsed NGGCT patients, 13 were excluded, comprising non-protocol first-line chemotherapy (n=3), relapse with pure teratoma, treated with surgery alone (n=5), and a further five patients who received palliation

1
2
3 only (see Figure 1). As for relapsed germinoma, the majority of relapsed NGGCT
4
5 patients treated with curative intent received re-induction chemotherapy with a
6
7 platinum-based (carboplatin or cisplatin) strategy, with most of those patients then
8
9 proceeding to high-dose therapy (Table 1). Five-year EFS and OS for these 32
10
11 remaining malignant patients treated with curative intent was 9% (95% CI: 2-26%),
12
13 with a median EFS and OS of 8m and 12m, respectively (Figure 2). Median OS for
14
15 the three survivors was 105 m (range 88-127 m). Five-year OS for the 10 patients
16
17 who received SDC was 0% (0-0%), with median survival of 9 m; all died 3-43 m
18
19 following relapse (Figure 3). Of these 10 patients, four (40%) received CSI at first
20
21 relapse, two (20%) received focal re-irradiation and four (40%) received no further
22
23 radiotherapy. Five-year OS for the 22 patients treated with intention for
24
25 HDC+AuSCR was 14% (3-36%) with median survival of 12 m; only three patients
26
27 survived and the others all died 3-35 m following relapse (Figure 3). Of these 22
28
29 patients, five (23%) received focal re-irradiation at first relapse, eight (36%) received
30
31 CSI and nine (41%) received no further irradiation. Data analysis was performed on
32
33 the basis of physician intention to treat, but 6/22 patients (27%) did not receive
34
35 HDC+AuSCR because either they experienced further disease progression and died
36
37 before initiation of HDC, or because they were too clinically unwell to receive it. Of
38
39 the 16 patients who actually received HDC+AuSCR, 13 died (range 3-35 m), one is
40
41 alive with SD (at 88 m follow-up) and two patients are in CR (at 105 m and 127 m). It
42
43 is of note that the two CR patients both received additional radiation.
44
45
46
47
48
49
50
51

52 *Associations with survival in relapsed CNS GCTs.* Clinico-pathological factors were
53
54 interrogated for potential associations with clinical outcome. Firstly, patients who
55
56 relapsed following an original diagnosis of germinoma (n=11) had better 5-year EFS
57
58
59
60

1
2
3 and OS than those who relapsed following an original diagnosis of NGGCT (n=32)
4 (p=0.01 and p=0.007, respectively) (Figure 2). In the latter group, the only three
5 survivors were all noted to have raised serum/CSF HCG levels alone (defined as HCG
6 >50 IU/l at either diagnosis or relapse). Comparing the NGGCT relapse group treated
7 with curative intent (n=32), regardless of relapse treatment received (intention for
8 HDC+AuSCR or SDC), this 'raised HCG alone' group (n=7) had significantly better
9 5-year OS (3/7; 43%) than all other relapsed NGGCT patients (0/25; 0%), (p=0.003)
10 (Figure 4A). To avoid potential confounding by relapse treatment received, we next
11 repeated this HCG analysis for only the 22 relapsed NGGCT patients with intention to
12 treat with HDC+AuSCR. This also demonstrated a significantly better 5-year OS
13 (p=0.002) for the four patients with raised HCG alone (3/4 survivors; 75%) compared
14 with all other patients in the HDC+AuSCR cohort (0/18 survivors; 0%) (Figure 4B).
15 Within the 'raised HCG alone' group, there were, however, no significant differences
16 in the maximum levels of HCG between the patients who survived (n=3; levels 113,
17 4,664, 5,999 IU/L) and those who died (n=4; levels 608, 3,201, 3,559, 60,600 IU/L)
18 (p=0.49). For other associations with clinical factors, see Supplementary Results.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

Intracranial germ cell tumors (GCTs) are a rare, heterogeneous cancer group and first-line treatments are generally successful, so relapse is rare^{2,3,7}. Previous reports describing outcomes of patients with relapsed intracranial GCTs are therefore limited by either small sample size^{14,18-22}, selection bias and/or variable first-line treatments^{14,15,21,23}. For example, eight of 21 patients described in one study¹⁴ and six of 20 patients in another¹⁵ evaluating HDC+AuSCR for relapsed intracranial GCTs had not received first-line radiotherapy, which would now be considered standard-of-care for both intracranial germinoma and NGGCT patients^{2,4,22,24}. Consequently, treatments for relapsed disease remain varied with no consensus on optimal management². To address these limitations, we collected data from UK and German patients with relapsed intracranial GCTs, treated with standardized, first-line treatment as per the European SIOP-CNS-GCT-96 trial protocol. This has allowed us to describe the treatment received and subsequent outcome of the largest cohort of patients with relapsed intracranial GCTs to date. Appropriately, patients with non-uniform first-line treatment, those relapsing with teratoma only and those treated palliatively at relapse were excluded from further outcome analysis. The results will inform future approaches to the management of this disease, with the aim of maximizing survival.

Given the excellent EFS for patients with intracranial germinoma^{5,25,26}, relapses were a very rare occurrence indeed – indeed we only identified a total of 13 such cases across two large national groups (UK/Germany) contributing to the European SIOP-CNS-GCT-96 study. Consequently, this germinoma cohort represents a relatively

1
2
3 small and heterogeneous group, in terms of type of relapse (marker-negative vs.
4 marker-positive) and relapse treatments delivered (SDC vs. HDC+AuSCR; re-
5 irradiation vs. none). Full details are provided in Supplementary Discussion.
6
7
8
9
10 Outcomes for patients who relapsed following germinoma diagnosis were broadly
11 similar, whether treated with SDC or HDC+AuSCR. It should also be emphasised that
12
13
14 whilst no patient in this group receiving HDC+AuSCR had additional radiotherapy,
15
16 all such patients who received SDC had further radiotherapy (focal or CSI), except for
17
18 a single patient with extracranial relapse (patient G13). Re-irradiation is therefore
19
20 likely to make an important contribution to the similar outcomes observed between
21
22 the two chemotherapy groups. These data therefore support either treatment option for
23
24 these patients.
25
26

27
28
29 As expected, due to the inferior outcomes for patients with intracranial NGGCTs
30 compared with germinoma^{12,26}, relapses amongst this cohort were more common,
31
32 with 45 cases identified. Of the 32 cases with uniform first-line therapy and malignant
33
34 relapse treated with curative intent, outcomes were very poor. There were no
35
36 survivors in the patient group treated with SDC, and long-term OS was only 14% for
37
38 the group with physician intention-to-treat with HDC+AuSCR. Indeed, over a quarter
39
40 of this latter cohort did not actually receive HDC+AuSCR as they either progressed
41
42 during re-induction chemotherapy or were too unwell to receive it. The outcomes
43
44 described here are inferior to survival rates reported in previous studies: the 33% four-
45
46 year survival rate described in a cohort of 12 patients¹⁴ and 59% three-year survival in
47
48 a cohort of eleven patients¹⁵ with relapsed NGGCTs who received HDC+AuSCR
49
50 following variable first-line therapy. It should be noted that if only those patients who
51
52 actually went on to receive HDC+AuSCR with a thiotepa-based regimen are
53
54
55
56
57
58
59
60

1
2
3 considered, survival was 30% (three of 10 patients; Table 2) – similar to that
4
5 previously described¹⁴. However, the overall inferior outcomes described here are
6
7 more likely to be representative of this patient cohort as a whole. This is due to our
8
9 study ensuring that outcome analyses were only performed on patients receiving
10
11 uniform first-line treatment, and inclusion of cases from the time-of-relapse on a
12
13 curative intention-to-treat basis, and not just those who responded well enough to
14
15 proceed to HDC+AuSCR. Accordingly, patients need to be counselled regarding
16
17 these poor outcomes at the time of relapse.
18
19

20
21
22
23 Of note, the only three survivors within the NGGCT relapse group had raised HCG
24
25 alone, either at original diagnosis or at relapse. All other patients, i.e. those with
26
27 raised AFP, combined raised AFP and HCG, or those who were marker-negative but
28
29 had EC confirmed on biopsy, all died following relapse, regardless of salvage therapy
30
31 delivered. Furthermore, at relapse, those patients with raised HCG alone had
32
33 improved OS, both for the group as a whole, and when just those patients with
34
35 intention-to-treat with HDC+AuSCR were considered, where three of four patients
36
37 with raised HCG were long-term survivors. Survival did not appear to be related to
38
39 the actual HCG level however, because one of the three surviving patients had a
40
41 maximum HCG level of 113 IU/L and the other two had levels >4,500 IU/L.
42
43 Although the first patient had a level again potentially consistent with HCG
44
45 production from syncytiotrophoblast within a pure germinoma, the other two patients
46
47 had levels that all national groups would consider to represent the presence of CHC
48
49 within the tumor. These observations do highlight, however, the importance of
50
51 attempting to agree an internationally agreed cut-off/threshold for tumor-markers, in
52
53 order to better understand the correlation between marker levels, histology and
54
55
56
57
58
59
60

1
2
3 clinical outcome^{2,27}. Such agreement will facilitate trials to improve survival rates for
4
5 patients with NGGCT whilst avoiding unnecessary treatment for those with
6
7 germinoma. HDC+AuSCR is therefore a treatment option for patients diagnosed
8
9 initially with NGGCT and experiencing a malignant relapse, if curative intent is the
10
11 aim¹³. Additional surgery and/or radiotherapy should also be delivered where feasible
12
13 – it should be noted that of the three survivors, the two who had CR both received re-
14
15 irradiation. For commentary on EC subtype relapses²⁸, see Supplementary Discussion.
16
17
18
19

20
21 Recent advances in our biological understanding^{29–33} are likely to assist clinical
22
23 management decisions in the future, through improved risk stratification systems and
24
25 the identification of molecular targets that will help determine the selection of
26
27 appropriate novel therapeutic agents, particularly for the NGGCT patient population
28
29 that have inferior outcomes. Recent use of exome sequencing to identify common
30
31 genomic mutations is one such example^{29,30}. Collection of serum/plasma and CSF
32
33 may also facilitate non-invasive diagnosis using microRNA expression levels^{31,32} and
34
35 risk stratification through identification of tumor mutations via CSF tumor DNA
36
37 (ctDNA) analysis³³. Embedding such non-invasive biological studies in future clinical
38
39 trials is a particularly attractive strategy for trial groups where NGGCT tissue
40
41 specimens are a scarce resource due to reliance on typical radiology and tumor
42
43 markers for diagnosis.
44
45
46
47
48

49
50 The main limitation of this study, as in previous reports¹⁴, is its retrospective nature.
51
52 However, as relapse is a rare outcome of this infrequent malignancy, a prospective
53
54 randomised trial of relapsed CNS GCT patients is unlikely to be feasible. In future, it
55
56 will be important to ensure that prospective data collection is embedded routinely
57
58
59
60

1
2
3 within treatment guidelines, to facilitate the identification of optimal re-induction
4
5 strategies and further assess the role of HDC in this setting.
6
7

8
9
10 In conclusion, we present the largest cohort of patients with relapsed intracranial
11
12 GCTs reported to date. This study benefits from the uniform manner in which patients
13
14 received first-line treatment, including radiotherapy as standard-of-care. Patients with
15
16 an original diagnosis of intracranial germinoma experiencing relapse were
17
18 salvageable with SDC and re-irradiation and HDC+AuSCR with and without re-
19
20 irradiation. Patients relapsing following an original diagnosis of intracranial NGGCT
21
22 fared very poorly; only patients with raised HCG levels alone survived in this series,
23
24 and only then with HDC+AuSCR. Thiotepa-based HDC+AuSCR, with additional
25
26 surgery and radiotherapy where feasible, is a treatment option at first malignant
27
28 relapse for such patients, albeit with counselling regarding poor outcomes.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Calaminus G, Bamberg M, Baranzelli MC, et al. Intracranial germ cell tumors: a comprehensive update of the European data. *Neuropediatrics*. 1994;25(1):26-32.
2. Murray MJ, Bartels U, Nishikawa R, Fangusaro J, Matsutani M, Nicholson JC. Consensus on the management of intracranial germ-cell tumours. *Lancet Oncol*. 2015;16(9):e470-477.
3. Matsutani M, Sano K, Takakura K, et al. Primary intracranial germ cell tumors: a clinical analysis of 153 histologically verified cases. *J Neurosurg*. 1997;86:446-455.
4. Calaminus G, Andreussi L, Garré ML, Kortmann RD, Schober R, Göbel U. Secreting germ cell tumors of the central nervous system (CNS). First results of the cooperative German/Italian pilot study (CNS sGCT). *Klin Padiatr*. 1997;209(4):222-227.
5. Calaminus G, Kortmann R, Worch J, et al. SIOP CNS GCT 96: final report of outcome of a prospective, multinational nonrandomized trial for children and adults with intracranial germinoma, comparing craniospinal irradiation alone with chemotherapy followed by focal primary site irradiation for pat. *Neuro Oncol*. 2013;15(6):788-796.
6. Sano K, Matsutani M. Microsurgery of teratoma and germinoma involving the diencephalon and the brain stem. *Neurosurg Rev*. 1983;6(2):51-55.
7. Bamberg M, Kortmann RD, Calaminus G, et al. Radiation therapy for intracranial germinoma: Results of the German cooperative prospective trials MAKEI 83/86/89. *J Clin Oncol*. 1999;17(8):2585-2592.

- 1
2
3 8. Alapetite C, Brisse H, Patte C, et al. Pattern of relapse and outcome of non-
4 metastatic germinoma patients treated with chemotherapy and limited field
5 radiation: the SFOP experience. *Neuro Oncol.* 2010;12(12):1318-1325.
6
7
- 8
9
10 9. Cheng S, Kilday J-P, Laperriere N, et al. Outcomes of children with central
11 nervous system germinoma treated with multi-agent chemotherapy followed by
12 reduced radiation. *J Neurooncol.* 2016;127(1):173-180.
13
14
- 15
16
17 10. Gobel U, Bamberg M, Calaminus G, et al. Improved prognosis of intracranial
18 germ cell tumors by intensified therapy: results of the MAKEI 89 therapy
19 protocol. *Klin Padiatr.* 1993;205(4):217-224.
20
21
- 22
23
24 11. Calaminus G, Frappaz D, Kortmann RD, et al. Risk adapted irradiation is
25 feasible in intracranial non-germinomatous germ cell tumours (NGGCT): final
26 results of SIOP CNS GCT 96. *Neuro Oncol.* 2012;14((suppl 1)):i49:55.
27
28
- 29
30
31 12. Goldman S, Bouffet E, Fisher PG, et al. Phase II Trial Assessing the Ability of
32 Neoadjuvant Chemotherapy With or Without Second-Look Surgery to
33 Eliminate Measurable Disease for Nongerminomatous Germ Cell Tumors: A
34 Children's Oncology Group Study. *J Clin Oncol.* 2015;33(22):2464-2471.
35
36
- 37
38
39 13. Bouffet E. The Role of Myeloablative Chemotherapy with Autologous
40 Hematopoietic Cell Rescue in Central Nervous System Germ Cell Tumors.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
14. Modak S, Gardner S, Dunkel IJ, et al. Thiotepa-Based High-Dose
Chemotherapy With Autologous Stem-Cell Rescue in Patients With Recurrent
or Progressive CNS Germ Cell Tumors. *J Clin Oncol.* 2004;22(10):1934-1943.
15. Baek HJ, Park HJ, Sung KW, et al. Myeloablative chemotherapy and
autologous stem cell transplantation in patients with relapsed or progressed

- 1
2
3 central nervous system germ cell tumors: Results of Korean Society of
4 Pediatric Neuro-Oncology (KSPNO) S-053 study. *J Neurooncol.*
5 2013;114(3):329-338.
6
7
8
9
10 16. Eisenhauer E a. A, Therasse P, Bogaerts J, et al. New response evaluation
11 criteria in solid tumours: Revised RECIST guideline (version 1.1). *Eur J*
12 *Cancer.* 2009;45(2):228-247.
13
14
15
16
17 17. Murray MJ, Metayer LE, Mallucci CL, et al. Intra-abdominal metastasis of an
18 intracranial germinoma via ventriculo-peritoneal shunt in a 13-year-old female.
19 *Br J Neurosurg.* 2011;25(6):747-749.
20
21
22
23
24 18. Nichols CR, Andersen J, Lazarus HM, et al. High-dose carboplatin and
25 etoposide with autologous bone marrow transplantation in refractory germ cell
26 cancer: an Eastern Cooperative Oncology Group protocol. *J Clin Oncol.*
27 1992;10(4):558-563.
28
29
30
31
32
33
34 19. Nichols CR, Tricot G, Williams SD, et al. Dose-intensive chemotherapy in
35 refractory germ cell cancer--a phase I/II trial of high-dose carboplatin and
36 etoposide with autologous bone marrow transplantation. *J Clin Oncol.*
37 1989;7(7):932-939.
38
39
40
41
42
43
44 20. Wolff SN, Johnson DH, Hainsworth JD, Greco FA. High-dose VP-16-213
45 monotherapy for refractory germinal malignancies: A phase II study. *J Clin*
46 *Oncol.* 1984;2(4):271-274.
47
48
49
50
51 21. Gururangan S, Dunkel IJ, Goldman S, et al. Myeloablative chemotherapy with
52 autologous bone marrow rescue in young children with recurrent malignant
53 brain tumors. *J Clin Oncol.* 1998;16(7):2486-2493.
54
55
56
57
58 22. Baranzelli MC, Patte C, Bouffet E, et al. An attempt to treat pediatric
59
60

- 1
2
3 intracranial alphaFP and betaHCG secreting germ cell tumors with
4 chemotherapy alone. SFOP experience with 18 cases. Société Française
5 d'Oncologie Pédiatrique. *J Neurooncol.* 1998;37(3):229-239.
6
7
8
9
10 23. Beyer J, Kramar A, Mandanas R, et al. High-dose chemotherapy as salvage
11 treatment in germ cell tumors: A multivariate analysis of prognostic variables.
12 *J Clin Oncol.* 1996;14(10):2638-2645.
13
14
15
16
17 24. Balmaceda C, Heller G, Rosenblum M, et al. Chemotherapy without
18 irradiation--a novel approach for newly diagnosed CNS germ cell tumors:
19 results of an international cooperative trial. The First International Central
20 Nervous System Germ Cell Tumor Study. *J Clin Oncol.* 1996;14(11):2908-
21 2915.
22
23
24
25
26
27
28
29 25. Bouffet E, Baranzelli MC, Patte C, et al. Combined treatment modality for
30 intracranial germinomas: results of a multicentre SFOP experience. Société
31 Française d'Oncologie Pédiatrique. *Br J Cancer.* 1999;79(7-8):1199-1204.
32
33
34
35
36 26. Matsutani M. Combined chemotherapy and radiation therapy for CNS germ
37 cell tumors--the Japanese experience. *J Neurooncol.* 2001;54(3):311-316.
38
39
40
41 27. Murray MJ, Horan G, Lewis S, Nicholson JC. Highlights from the Third
42 International Central Nervous System Germ Cell Tumour symposium: laying
43 the foundations for future consensus. *Ecancermedicalscience.* 2013;7:333.
44
45
46
47
48 28. da Silva NS, Cappellano AM, Diez B, et al. Primary chemotherapy for
49 intracranial germ cell tumors: results of the third international CNS germ cell
50 tumor study. *Pediatr Blood Cancer.* 2010;54(3):377-383.
51
52
53
54
55
56 29. Wang L, Yamaguchi S, Burstein MD, et al. Novel somatic and germline
57 mutations in intracranial germ cell tumours. *Nature.* 2014;511(7508):241-245.
58
59
60

- 1
2
3 30. Fukushima S, Otsuka A, Suzuki T, et al. Mutually exclusive mutations of KIT
4 and RAS are associated with KIT mRNA expression and chromosomal
5 instability in primary intracranial pure germinomas. *Acta Neuropathol.*
6 2014;127(6):911-925.
7
8
9
10
11
12 31. Murray MJ, Bell E, Raby KL, et al. A pipeline to quantify serum and
13 cerebrospinal fluid microRNAs for diagnosis and detection of relapse in
14 paediatric malignant germ cell tumours. *Br J Cancer.* 2016;114:151-162.
15
16
17
18
19 32. Terashima K, Shen J, Luan J, et al. microRNA 371-373 and 302a in
20 cerebrospinal fluid are potential tumor-derived biomarkers for intracranial
21 germ cell tumors. *Neuro Oncol.* 2013;15:iii152.
22
23
24
25
26
27 33. De Mattos-Arruda L, Mayor R, Ng CKY, et al. Cerebrospinal fluid-derived
28 circulating tumour DNA better represents the genomic alterations of brain
29 tumours than plasma. *Nat Commun.* 2015;6:8839.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Legends to Figures and Tables

Figure 1. Summary flow-diagram listing the 58 patients described in the study and their basic clinical characteristics.

Figure 2. Kaplan-Meier curves comparing five-year A) event-free survival and B) overall survival for patients treated with curative intent: intracranial malignant relapse of germinoma (solid line) versus non-germinomatous germ-cell-tumor (NGGCT; broken line).

Figure 3. Kaplan-Meier curves comparing five-year overall survival for patients treated with curative intent at first malignant relapse of intracranial non-germinomatous germ-cell-tumor (NGGCT): intention for high-dose chemotherapy (solid line) versus those receiving standard-dose chemotherapy (broken line). N/S = non-significant.

Figure 4. Kaplan-Meier curves comparing five-year overall survival for patients treated with curative intent at first malignant relapse of intracranial non-germinomatous germ-cell-tumor (NGGCT) with: elevated levels of HCG alone (solid line) versus all other patients in the cohort* (broken line). Graphs show data for A) all 32 patients regardless of malignant relapse treatment schedule and B) the 22 patients treated with intention for high-dose chemotherapy only. *i.e., all those patients with elevated levels of AFP, combined elevations of AFP/HCG and/or with a tissue diagnosis of EC either in the serum or CSF, at diagnosis and/or relapse.

1
2
3
4
5
6 **Table 1.** Details of the intracranial germinoma patients at relapse, description of
7
8 relapse treatments received and outcomes.
9

10
11
12
13 **Table 2.** Details of the intracranial non-germinomatous germ-cell-tumor (NGGCT)
14
15 patients at relapse, description of relapse treatments received and outcomes.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Figure 1. Summary flow-diagram listing the 58 patients described in the study and their basic clinical characteristics.




Figure 2. Kaplan-Meier curves comparing five-year A) event-free survival and B) overall survival for patients treated with curative intent: intracranial malignant relapse of germinoma (solid line) versus non-germinomatous germ cell tumor (NGGCT; broken line).





Figure 3. Kaplan-Meier curves comparing five-year overall survival for patients treated with curative intent at first malignant relapse of intracranial non-germinomatous germ cell tumor (NGGCT): intention for high-dose chemotherapy (solid line) versus those receiving standard-dose chemotherapy (broken line). N/S = non-significant.




Figure 4. Kaplan-Meier curves comparing five-year overall survival for patients treated with curative intent at first malignant relapse of intracranial non-germinomatous germ cell tumor (NGGCT) with: elevated levels of HCG alone (solid line) versus all other patients in the cohort (broken line). Graphs show data for A) all 32 patients regardless of malignant relapse treatment schedule and B) the 22 patients treated with intention for high-dose chemotherapy only.





Table 1. Details of the intracranial germinoma patients at relapse, description of relapse treatments received and outcomes.

Patient number	Site(s) of disease at relapse	Elevated tumor markers identified at relapse?	Diagnosis at relapse	Surgery for relapse?	HDC or SDC?	Chemotherapy received	Radiotherapy received	Response	Morbidity described?	Further relapses?	Outcome	Time (from relapse) of death or latest follow up (months)	Included in further analysis?
G1	Ventricular	No	Presumed germinoma	No	SDC	Re-induction (carboPEI)	Focal	CR	None	None	DOC (sepsis, brainstem impairment)	16	Yes
G4	Distant intracranial	No	Immature teratoma (histology)	Total resection	SDC	Re-induction (carboPEI)	No	CR	Metabolic imbalance	3 (treatment included HDC)	DOD	10	No (immature teratoma)
G5	Primary	No	Germinoma (histology)	Total resection	HDC	Re-induction (PEI); HDC (carbo/etop)	No	CR	None	None	AND	194	Yes
G7	Primary	No	Presumed germinoma	No	HDC	Re-induction (carboPEI); HDC (thiotepa/etop)	None	CR	Infection	None	DOC (pneumonitis)	5	Yes
G8	Ventricular	No	Presumed germinoma	No	SDC	Re-induction (PEI)	Focal	CR	None	1 (treatment included HDC)	AND	92	Yes
G9	Distant intracranial	No	Presumed germinoma	No	N/A	None	None	N/A	N/A	None	DOD	0	No (palliative)
G10	Extracranial (skeletal)	No	Germinoma (histology)	Biopsy	HDC	Re-induction (PEI); HDC (thiotepa/etop)	None	CR	Panhypopituitarism	None	AND	53	Yes
G11	Distant intracranial	No	Germinoma (histology)	Partial resection	SDC	Re-induction (other)	Focal	CR	Panhypopituitarism	None	AND	148	Yes
G2	Leptomeningeal	HCG	CHC (markers)	No	SDC	Re-induction (other)	CSI	CR	Electrolyte imbalance, seizures	2 (treatment included HDC)	DOD	47	Yes
G3	Primary	AFP	YST (markers)	No	HDC	Re-induction (PEI); HDC (not specified)	No	PD	Febrile neutropenia	None	DOD	8	Yes
G6	Primary	AFP	Germinoma (histology); YST (markers)	Total resection	SDC	Re-induction (PEI)	Focal	PD	None	1 (treatment not specified)	DOD	8	Yes
G12	Ventricular and leptomeningeal	HCG	CHC (markers)	No	SDC	Re-induction (PEI)	CSI	CR	DI	None	AND	62	Yes
G13	Extracranial (abdominal via VPS)	HCG	Germinoma (cytology of ascites) but CHC (markers)	Total resection of residual after chemotherapy	SDC	Re-induction (JEB)	No	CR	None	None	AND	75	Yes

Abbreviations:

SDC = standard-dose chemotherapy, HDC = high-dose chemotherapy

CHC = choriocarcinoma

YST = yolk sac tumor

MGCT = malignant germ cell tumor

CarboPEI = carboplatin, etoposide, ifosfamide

PEI = cisplatin, etoposide, ifosfamide

carbo/etop = carboplatin, etoposide

thiotepa/etop = thiotepa, etoposide

JEB = carboplatin, etoposide, bleomycin

CSI = craniospinal irradiation

CR = complete response, PR = partial response, PD = progressive disease

AFP = alpha-fetoprotein; defined as elevated in serum or cerebrospinal fluid when levels >25 ng/ml

HCG = human chorionic gonadotrophin, defined as elevated in serum or cerebrospinal fluid when levels >50 IU/l

DOD = died of disease, DOC = died of complication, AND = alive with no disease

N/A = not applicable

DI = diabetes insipidus

VPS = ventriculo-peritoneal shunt

Table 2: Details of the intracranial non-germinomatous germ-cell-tumor (NGGCT) patients at relapse, description of relapse treatments received and outcomes.

Patient Number	Site(s) of disease at relapse	Elevated AFP serum or CSF levels identified at relapse?	Elevated HCG serum or CSF levels identified at relapse?	Diagnosis and additional histological confirmation	Surgery for relapse?	Intention to treat with HDC + AuSCR	Received HDC?	Chemotherapy received	Subsequent radiotherapy	Response	Morbidity	Number of further relapses and treatment received	Outcome	Time of death from 1 st relapse (months)	Included in analysis?
7 8 9 10 11 12	NG1 Distant intracranial	+	-	YST (markers)	No	Yes	No	Re-induction (carboPEI); HDC not received – disease progression	Focal	PD	Neurological (seizures)	N/A	DOD	13	Yes
13 14 15 16 17 18 19	NG2 Primary	+	-	YST (markers)	Subtotal resection	Yes	Yes	Re-induction (carboPEI); HDC (thiotepa-based)	Focal	PR	Toxicity from chemotherapy (not specified)	None	DOD	7	Yes
20 21 22 23 24 25	NG3 Distant intracranial	-	++	CHC (markers)	No	Yes	Yes	Re-induction (carboPEI); HDC (etop)	None	CR	None	1 (HDC)	DOD	18	Yes
26 27 28 29 30 31 32	NG4 Not specified	-	-	Teratoma (histology)	Subtotal resection	N/A	N/A	None	None	PR	None	1 (surgery)	DOD	30	No (teratoma)
33 34 35 36 37 38 39	NG5 Primary, ventricular, spinal	+	-	Germinoma (histology), YST (markers)	Subtotal resection	Yes	Yes	Not specified	None	PD	None	None	DOD	6	Yes
40 41 42 43 44	NG6 Primary	-	-	Teratoma (histology)	Total resection	N/A	N/A	None	None	CR	None	None	AND	65	No (teratoma)
45 46 47 48 49	NG7 Primary	-	-	Presumed NGGCT	No	N/A	N/A	None	None	N/A	N/A	N/A	DOD	2	No (palliative)
50 51 52 53 54 55 56	NG8 Primary	-	-	Presumed NGGCT	No	Yes	No	Re-induction (not specified); HDC not received – disease progression	None	PD	Infection	N/A	DOD	3	Yes
57 58 59 60 61 62 63	NG9 Primary	+	-	YST (markers)	No	No	N/A	Re-induction (carbo/etop)	Focal	PD	Infection	None	DOD	19	Yes
64 65 66 67 68 69 70	NG10 Primary, distant intracranial, spinal	+++	-	YST (markers)	Total resection	Yes	Yes	Re-induction (carbo/etop); HDC (PEI)	CSI	PD	None	1 (not specified)	DOD	6	Yes
71 72 73 74 75 76 77	NG11 Primary, spinal	++	-	YST (markers)	No	Yes	Yes	Re-induction (carbo/etop); HDC (thiotepa-based)	CSI	PR	Infection	2 (SDC)	DOD	35	Yes
78 79 80 81 82 83 84	NG12 Distant intracranial	-	+	CHC (markers)	No	No	N/A	Re-induction (carbo/etop)	CSI	CR	Neurological (Guillain-Barré syndrome)	1 (palliative)	DOD	12	Yes
85 86 87 88 89 90 91	NG13 Primary, spinal	-	-	Presumed NGGCT	No	Yes	No	Re-induction (PEI); HDC not received – disease progression	None	PD	Not specified	N/A	DOD	3	Yes
92 93 94 95 96 97 98	NG14 Primary	+	-	YST (markers)	No	Yes	Yes	Re-induction (carbo/etop); HDC (etop)	None	CR	Infection	1 (SDC)	DOD	14	Yes
99 100 101 102 103 104 105	NG15 Distant intracranial, spinal	-	++	CHC (markers)	No	Yes	Yes	Re-induction (carboPEI); HDC (thiotepa-based)	None	SD	None	None	ASD	88	Yes
106 107 108 109 110 111 112	NG16 Primary, spinal	-	-	Presumed NGGCT	Metastases resected	Yes	Yes	Re-induction and HDC (HD-PEI)	CSI	CR	Not specified	1 (palliative)	DOD	11	Yes

1 2	NG17	Primary	+	-	YST (markers)	No	No	N/A	Ventricular etoposide	CSI	SD	None	1 (palliative)	DOD	8	Yes
3 4 5	NG18	Primary, spinal	-	-	Germinoma (histology)	Subtotal resection	No	N/A	Re-induction (carbo/etop)	None	PR	Infection, respiratory and renal failure	None	DOD	10	Yes
6 7 8	NG19	Distant intracranial	Yes (levels not specified)	-	YST (markers)	No	No	N/A	Re-induction (carbo/etop)	CSI	PD	Not specified	1 (SDC)	DOD	9	No (non-protocol first-line treatment)
9 10	NG20	Primary	-	-	Teratoma (histology)	Total resection	No	N/A	None	None	CR	None	None	AND	79	No (teratoma)
11 12	NG21	Distant intracranial	-	-	Mixed MGCT (histology)	Total resection	No	N/A	None	None	PD	None	5 (SDC)	DOD	35	Yes
13 14 15	NG22	Spinal	-	++	CHC (markers), EC on initial biopsy	No	No	N/A	Re-induction (etop)	None	PD	Not specified	None	DOD	3	Yes
16 17 18	NG23	Spinal	+++	-	YST (markers)	Biopsy	Yes	Yes	Re-induction (carbo/cyclophos); HDC (thiotepa-based)	Focal	PR	None	None	DOD	11	Yes
19 20	NG24	Primary, spinal	+	-	YST (markers)	No	N/A	N/A	Palliative (oral etoposide)	Focal (to spine)	PD	Infection, seizures	None	DOD	7	No (palliative)
21	NG25	Ventricular	-	-	Presumed germinoma	No	N/A	N/A	None	None	N/A	N/A	None	DOD	0	No (palliative)
22 23	NG26	Distant intracranial	++	-	YST (markers)	Total resection	No	N/A	Re-induction (JEB)	Focal	CR	Memory loss	1 (not specified)	DOD	43	Yes
24 25	NG27	Primary	-	++	CHC (markers)	No	Yes	Yes	Re-induction (carbo/etop); HDC (thiotepa-based)	None	CR	DI	1 (SDC)	DOD	34	Yes
26 27	NG28	Ventricular	-	+++	CHC (markers)	No	Yes	Yes	Re-induction (carbo/etop); HDC (thiotepa-based)	Focal	CR	Extreme lethargy	None	AND	127	Yes
28 29 30 31 32	NG29	Ventricular, spinal	-	+	CHC (markers)	No	Yes	Yes	Re-induction (paclitaxel/etop/cisplatin); HDC (etop/carbo/cyclophos)	None	CR	Hearing loss, mucositis	1 (palliative)	DOD	17	Yes
33 34	NG30	Primary	+	-	YST (markers)	No	Yes	Yes	Re-induction (carbo/etop); HDC (thiotepa-based)	None	PD	Bone marrow suppression	1 (HDC)	DOD	11	Yes
35 36 37	NG31	Primary, spinal	+	-	YST (markers), EC on initial biopsy	No	N/A	N/A	None	CSI	PD	None	None	DOD	1	No (palliative)
38 39 40	NG32	Spinal	+	-	YST (markers)	No	Yes	No	Re-induction (cisplatin/etop/vinc); HDC not received – disease progression	CSI	PD	None	None	DOD	12	Yes
41 42 43	NG33	Distant intracranial	++	-	YST (markers), EC on initial biopsy	No	No	N/A	Re-induction (carbo/etop)	None	PD	Toxicity from chemotherapy (not specified)	None	DOD	6	Yes
44	NG34	Distant	-	-	Presumed	No	Yes	No	Re-induction	CSI	PR	Slow count	None	DOD	9	Yes

1				NGGCT, EC on initial biopsy				(carbo/etop); HDC not received – toxicity of treatment			recovery					
2	intracranial, spinal															
3																
4	NG35	Ventricular	-	-	Presumed germinoma (non-diagnostic biopsy)	Biopsy	Yes	Yes	Re-induction (carbo/etop); HDC (thiotepa-based)	CSI	CR	Mild LD, panhypopituitarism	None	AND	105	Yes
5																
6																
7																
8	NG36	Primary	+++	-	YST (markers)	No	Yes	Yes	Re-induction (carbo/etop); HDC (carbo/etop)	Not specified	PD	None	None	DOD	9	No (non-protocol first-line treatment)
9																
10																
11	NG37	Spinal	-	-	CHC (cytology)	No	No	N/A	Re-induction (carbo/etop); then vinblastine/etop	CSI	PD	Tumour caused cord compression	None	DOD	4	Yes
12																
13	NG38	Spinal	++	-	YST (markers)	No	Yes	Yes	Re-induction (carbo/etop/cyclophos); HDC (thiotepa-based)	CSI	PD	Minimal	1 (HDC)	DOD	12	Yes
14																
15																
16	NG39	Primary	-	-	Teratoma (histology)	Partial resection	No	N/A	None	None	PR	Seizures, hemiplegia, LD, panhypopituitarism	None	ASD	120	No (teratoma)
17																
18																
19																
20	NG40	Primary	+	-	YST (markers)	No	Yes	Yes	Re-induction (carbo/etop/cyclophos); HDC (thiotepa-based)	Focal	PR	Intensive care after HDC	1 (palliative)	DOD	21	Yes
21																
22																
23	NG41	Spinal	++	-	YST (markers)	Not specified	No	N/A	Re-induction (not specified)	Not specified	Not specified	Leucopenia	None	DOD	2	No (non-protocol first-line treatment)
24																
25																
26	NG42	Leptomeningeal	-	+	CHC (markers)	No	N/A	N/A	None – palliative treatment only	None	N/A	N/A	None	DOD	1	No (palliative)
27																
28	NG43	Spinal	+	-	YST (markers)	Total resection	Yes	No	Re-induction (JEB); HDC not received – clinically too unwell	CSI	CR	Candida endocarditis, DI, shunt problems	1 (not specified)	DOD	0	Yes
29																
30	NG44	Primary, distant intracranial	-	-	Teratoma (histology)	Total resection	N/A	N/A	None	None	PD	None	None	DOD	3	No (teratoma)
31																
32																
33	NG45	Leptomeningeal	-	-	Presumed NGGCT	None	No	N/A	None	CSI	PD	None	None	DOD	4	Yes

Abbreviations:

1 NGGCT = non-germinomatous germ cell tumor
2 YST = yolk sac tumor
3 CHC = choriocarcinoma

4 SDC = standard-dose chemotherapy, HDC = high-dose chemotherapy

5
6 CSF = cerebrospinal fluid
7 AFP = alpha-fetoprotein; defined as elevated in serum or cerebrospinal fluid when levels >25 ng/ml
8 HCG = human chorionic gonadotrophin, defined as elevated in serum or cerebrospinal fluid when levels >50 IU/l

9 Marker key:

10 AFP:

11 - refers to levels ≤ 25 ng/ml
12 + refers to levels >25 but <200 ng/ml
13 ++ refers to levels ≥ 200 but <1000 ng/ml
14 +++ refers to levels ≥ 1000 ng/ml

15 HCG:

16 - refers to levels ≤ 50 IU/l
17 + refers to levels >50 but <200 IU/l
18 ++ refers to levels ≥ 200 but <1000 IU/l
19 +++ refers to levels ≥ 1000 IU/l

20 PEI = cisplatin, etoposide, ifosfamide
21 carboPEI = carboplatin, etoposide, ifosfamide
22 PEI = cisplatin, etoposide, ifosfamide
23 carbo/etop = carboplatin, etoposide
24 carbo/etop/cyclophos = carboplatin, etoposide, cyclophosphamide
25 thiotepa/etop = thiotepa, etoposide
26 cisplatin/etop/vinc = cisplatin, etoposide, vincristine
27 paclitaxel, etop/carbo/cyclophos = paclitaxel, etoposide, carboplatin, cyclophosphamide
28 JEB = carboplatin, etoposide, bleomycin

29 CSI = craniospinal irradiation
30 RT = radiotherapy

31 CR = complete response, PR = partial response, SD = stable disease, PD = progressive disease
32 DOD = died of disease, DOC = died of complication, ASD = alive with stable disease, AND = alive with no disease

33 LD = learning difficulties
34 DI = diabetes insipidus

35 N/A = not applicable
36
37
38
39
40
41
42
43
44
45
46
47
48
49