STRONGLY INTERACTING BLOW UP BUBBLES FOR THE MASS CRITICAL NLS

YVAN MARTEL AND PIERRE RAPHAËL

ABSTRACT. We consider the mass critical two dimensional nonlinear Schrödinger equation (NLS)

$$i\partial_t u + \Delta u + |u|^2 u = 0, \quad t \in \mathbb{R}, \quad x \in \mathbb{R}^2.$$

Let Q denote the positive ground state solitary wave satisfying $\Delta Q - Q + Q^3 = 0$. We construct a new class of multi-solitary wave solutions: given any integer $K \geq 2$, there exists a global (for $t > 0$) solution $u(t)$ of (NLS) that decomposes asymptotically into a sum of solitary waves centered at the vertices of a K -sided regular polygon and concentrating at a logarithmic rate as $t \rightarrow +\infty$ so that the solution blows up in infinite time with the rate

$$\|\nabla u(t)\|_{L^2} \sim |\log t| \quad \text{as } t \rightarrow +\infty.$$

This special behavior is due to strong interactions between the waves, in contrast with previous works on multi-solitary waves of (NLS) where interactions do not affect the blow up rate. Using the pseudo-conformal symmetry of the (NLS) flow, this yields the first example of solution $v(t)$ of (NLS) blowing up in finite time with a rate strictly above the pseudo-conformal one, namely,

$$\|\nabla v(t)\|_{L^2} \sim \left| \frac{\log |t|}{t} \right| \quad \text{as } t \uparrow 0.$$

Such solution concentrates K bubbles at a point $x_0 \in \mathbb{R}^2$, i.e. $|v(t)|^2 \rightharpoonup K\|Q\|_{L^2}^2 \delta_{x_0}$ as $t \uparrow 0$.

1. INTRODUCTION

1.1. General setting. We consider in this paper the mass critical two dimensional non linear Schrödinger equation (NLS)

$$i\partial_t u + \Delta u + |u|^2 u = 0, \quad (t, x) \in \mathbb{R} \times \mathbb{R}^2. \quad (1.1)$$

It is well-known (see e.g. [?] and the references therein) that for any $u_0 \in H^1(\mathbb{R}^2)$, there exists a unique maximal solution $u \in \mathcal{C}((-T_*, T^*), H^1(\mathbb{R}^2))$ of (??) with $u(0) = u_0$. Moreover, the following blow up criterion holds

$$T^* < +\infty \quad \text{implies} \quad \lim_{t \uparrow T^*} \|\nabla u(t)\|_{L^2} = +\infty. \quad (1.2)$$

The mass (i.e. the L^2 norm) and the energy E of the solution are conserved by the flow, where

$$E(u) = \frac{1}{2} \int_{\mathbb{R}^2} |\nabla u|^2 - \frac{1}{4} \int_{\mathbb{R}^2} |u|^4$$

From a variational argument, the unique (up to symmetry) ground state solution to

$$\Delta Q - Q + Q^3 = 0, \quad Q \in H^1(\mathbb{R}^2), \quad Q > 0, \quad Q \text{ is radially symmetric}$$

attains the best constant C in the following Gagliardo–Nirenberg inequality

$$\forall u \in H^1(\mathbb{R}^2), \quad \|u\|_{L^4}^4 \leq C \|u\|_{L^2}^2 \|\nabla u\|_{L^2}^2 \quad (1.3)$$

(see [?, ?, ?]). As a consequence, one has

$$\forall u \in H^1(\mathbb{R}^2), \quad E(u) \geq \frac{1}{2} \|\nabla u\|_{L^2}^2 \left(1 - \frac{\|u\|_{L^2}^2}{\|Q\|_{L^2}^2} \right). \quad (1.4)$$

Together with the conservation of mass and energy and the blow up criterion (??), this implies the global existence of any solution with initial data $\|u_0\|_2 < \|Q\|_2$. Actually it is now known that in this case, the solution scatters i.e. behaves asymptotically in large time as a solution of the linear equation, see [?, ?] and references therein.

We also know that $\|u\|_{L^2} = \|Q\|_{L^2}$ corresponds to the mass threshold for global existence since the pseudo-conformal symmetry of the (NLS) equation

$$v(t, x) = \frac{1}{|t|} u \left(\frac{1}{|t|}, \frac{x}{|t|} \right) e^{-i \frac{|x|^2}{4|t|}} \quad (1.5)$$

applied to the solitary wave solution $u(t, x) = e^{it} Q(x)$ yields the existence of an explicit single bubble blow up solution $S(t)$ with minimal mass

$$S(t, x) = \frac{1}{|t|} Q \left(\frac{x}{|t|} \right) e^{-i \frac{|x|^2}{4|t|}} e^{\frac{i}{|t|}}, \quad \|S(t)\|_{L^2} = \|Q\|_{L^2}, \quad \|\nabla S(t)\|_{L^2} \underset{t \rightarrow 0^-}{\sim} \frac{1}{|t|}. \quad (1.6)$$

We refer to [?] for more properties of the pseudo-conformal transform. From [?], minimal mass blow up solutions are *classified* in $H^1(\mathbb{R}^2)$:

$$\|u(t)\|_{L^2} = \|Q\|_{L^2} \quad \text{and} \quad T^* < +\infty \quad \text{imply} \quad u \equiv S \text{ up to the symmetries of the flow.}$$

Recall also the following well-known general sufficient criterion for finite time blow up: for initial data $u_0 \in \Sigma = H^1 \cap L^2(|x|^2 dx)$, the virial identity

$$\frac{d^2}{dt^2} \int_{\mathbb{R}^2} |x|^2 |u|^2 = 16E(u_0) \quad (1.7)$$

implies blow up in finite time provided $E(u_0) < 0$ (by (??), this implies necessarily $\|u_0\|_{L^2} > \|Q\|_{L^2}$).

1.2. Single bubble blow up dynamics. We focus now on the case of mass slightly above the threshold, i.e.

$$\|Q\|_{L^2} < \|u_0\|_{L^2} < \|Q\|_{L^2} + \alpha_0, \quad 0 < \alpha_0 \ll 1. \quad (1.8)$$

We first recall in this context that a large class of finite time blow up solutions was constructed in [?] (see also [?, ?]) as weak perturbation of the minimal mass solution $S(t)$. In particular, these solutions blow up with the pseudo-conformal blow up rate

$$\|\nabla u(t)\|_{L^2} \underset{t \sim T^*}{\sim} \frac{1}{T^* - t}. \quad (1.9)$$

Second, recall that the series of works [?, ?, ?, ?, ?, ?] provides a thorough study of the *stable* blow up dynamics under condition (??), corresponding to the so called *log-log* blow up regime

$$\|\nabla u(t)\|_{L^2} \underset{t \sim T^*}{\sim} c^* \sqrt{\frac{\log |\log(T^* - t)|}{T^* - t}}. \quad (1.10)$$

Third, it is proved in [?] (see also [?]) that solutions constructed in [?] are unstable and correspond in some sense to a threshold between the above log-log blow up and scattering.

Finally, recall that under (??), a universal gap on the blow up speed was proved in [?]: given a finite time blow up solution satisfying (??), either it blows up in the log–log regime (??), or it blows up faster than the pseudo–conformal rate

$$\|\nabla u(t)\|_{L^2} \gtrsim \frac{1}{T^* - t}.$$

(See also [?, ?].) However, the existence of solutions blowing up strictly faster than the conformal speed is a long lasting open problem, which is equivalent, by the pseudo–conformal symmetry (??), to the existence of global solutions blowing up in infinite time.

1.3. Multi bubbles blow up dynamics. For larger L^2 mass, it is conjectured (see e.g. [?]) that any finite time blow up solution concentrates at the blow up time universal quanta of mass $m_j > 0$ at a finite number of points $x_k \in \mathbb{R}^2$, i.e.

$$|u(t)|^2 \rightharpoonup \sum_{k=1}^K m_k \delta_{x_k} + |u^*|^2 \quad \text{as } t \uparrow T^*,$$

where $u^* \in L^2$ is a (possibly zero) residual. The first example of multiple point blow up solution is given in [?]: let $K \geq 1$ and let $(x_k)_{1 \leq k \leq K}$ be K arbitrary distinct points of $\mathbb{R}^2$, there exist a finite time blow up solution $u(t)$ of (??) with

$$\left\| u(t) - \sum_{k=1}^K S(t, \cdot - x_k) \right\|_{H^1} \rightarrow 0, \quad |u(t)|^2 \rightharpoonup \|Q\|_{L^2}^2 \sum_{k=1}^K \delta_{x_k} \quad \text{as } t \uparrow 0.$$

In particular, $u(t)$ blows up with the pseudo–conformal rate

$$\|\nabla u(t)\|_{L^2} \sim \frac{1}{|t|} \quad \text{as } t \uparrow 0.$$

Other general constructions of multi bubble blow up are provided by [?, ?] in the context of the log–log regime. Observe that these works deal with *weak* interactions in the sense that the blow up dynamics of each bubble is not perturbed at the main order by the presence of the other (distant) bubbles.

1.4. Main results. In this paper we construct the first example of infinite time blow up solution of (NLS), related to the *strong* interactions of an arbitrary number $K \geq 2$ of bubbles. As a consequence, using the pseudo–conformal transform, we also obtain the first example of solution blowing up in finite time strictly faster than the conformal blow up rate. Such a solution concentrates the K bubbles at one point at the blow up time.

Theorem 1 (Infinite time blow up). *Let $K \geq 2$ be an integer. There exists a solution $u \in \mathcal{C}([0, +\infty), \Sigma)$ of (??) which decomposes asymptotically into a sum of K solitary waves*

$$\left\| u(t) - e^{i\gamma(t)} \sum_{k=1}^K \frac{1}{\lambda(t)} Q\left(\frac{\cdot - x_k(t)}{\lambda(t)}\right) \right\|_{H^1} \rightarrow 0, \quad \lambda(t) = \frac{1 + o(1)}{\log t} \quad \text{as } t \rightarrow +\infty, \quad (1.11)$$

where the translation parameters $x_k(t)$ converge as $t \rightarrow +\infty$ to the vertices of a K -sided regular polygon, and where $\gamma(t)$ is some phase parameter. In particular,

$$\|\nabla u(t)\|_{L^2} = K^{\frac{1}{2}} \|\nabla Q\|_{L^2} (1 + o(1)) \log t \quad \text{as } t \rightarrow +\infty. \quad (1.12)$$

Corollary 2 (Finite time collision). *Let $u(t) \in C([0, +\infty), \Sigma)$ be given by Theorem ?? and let $v \in \mathcal{C}((-\infty, 0), \Sigma)$ be the pseudo conformal transform of $u(t)$ defined by (??). Then $v(t)$ blows up at $T^* = 0$ with*

$$\|\nabla v(t)\|_{L^2} = K^{\frac{1}{2}} \|\nabla Q\|_{L^2} (1 + o(1)) \left| \frac{\log |t|}{t} \right|, \quad |v|^2 \rightharpoonup K \|Q\|_{L^2}^2 \delta_0 \quad \text{as } t \uparrow 0. \quad (1.13)$$

Comments on the main results.

1. *Dynamics with multiple nonlinear objects.* Multiple bubble solutions with *weak interactions* and asymptotically free Galilean motion have been constructed in various settings, both in stable and unstable contexts, see in particular [?, ?, ?, ?, ?, ?, ?, ?]. As a typical example of weakly interacting dynamics, for the nonlinear Schrödinger equations

$$i\partial_t u + \Delta u + |u|^{p-1}u = 0, \quad x \in \mathbb{R}^d, \quad 1 < p < 1 + \frac{4}{d-2}, \quad (1.14)$$

there exist multi solitary wave solutions satisfying for large t ,

$$\left\| u(t) - \sum_{k=1}^K e^{-i(\frac{1}{2}\nu_k \cdot x - \frac{1}{4}|\nu_k|^2 t + \omega_k t + \gamma_k)} \omega_k^{\frac{1}{p-1}} Q\left(\omega_k^{\frac{1}{2}}(\cdot - \nu_k t)\right) \right\|_{H^1} \lesssim e^{-\gamma t}, \quad \gamma > 0, \quad (1.15)$$

for any given set of parameters $\{\nu_k, \omega_k, \gamma_k\}$ with the decoupling condition $\nu_k \neq \nu_{k'}$ if $k \neq k'$ (see [?, ?]).

In [?], two different regimes with *strong interactions* related to the two body problem of gravitation are exhibited for the Hartree model (hyperbolic and parabolic asymptotic motions). We also refer to [?, ?] for works related to sharp interaction problems in the setting of the subcritical (gKdV) equation. We thus see the present work as the first intrusion into the study of strongly interacting non radial multi solitary wave motions for (NLS). Note that the solution given by Theorem ?? is a minimal threshold dynamics and its behavior is unstable by perturbation of the data. An important direction of further investigation is the derivation of *stable* strongly interacting multiple bubbles blow up dynamics.

We observe from the proof of Corollary ?? that the K bubbles of the solution collide at the same point at the blow up time providing the first example of collision at blow up for (NLS). Note that the geometry of the trajectories of the blow up points (straight lines from the origin to the egde of the K -sided regular polygon) is an essential feature of these solutions. A related one dimensional mechanism is involved in the derivation of degenerate blow up curves in the context of “type I” blow up for the wave equation, see [?]. For the nonlinear heat equation in one dimension, solutions for which two points of maximum collide at blow up are constructed in [?]. There are also analogies of the present work with the construction of stationary solutions with mass concentrated along specific nonlinear grids, see [?]. In the context of two dimensional incompressible fluid mechanics, special solutions to the vortex point system are studied as a simplified model for dynamics of interacting and possibly colliding vortex, see for example [?] for an overview of these problems.

2. *Construction of minimal mass solutions.* The proof of Theorem ?? follows the now standard strategy of constructing *minimal* dynamics by approximate solutions and compactness, initiated in [?] and extended in various ways and contexts by [?, ?, ?, ?]. The proof of Theorem ?? combines in a blow up context the approach developed for multibubble flows in [?, ?] and a specific strategy to construct minimal blow up solutions for (NLS) type equations introduced in [?, ?]. An key ingredient of the proof is the precise tuning of the interactions between the waves. Note that we restrict ourselves to space dimension 2 for simplicity, but

similar results hold for the mass critical (NLS) equation in any space dimension with same proof. For the mass subcritical and supercritical nonlinear Schrödinger equations (??), we expect using similar approach the existence of bounded strongly interacting multi solitary waves, with logarithmic relative distances, i.e. non free Galilean motion. Interestingly enough, the existence of such solutions is *ruled out* in the mass critical case by the virial law (??). Indeed, the scaling instability direction of the critical case is excited by the interactions which leads to the infinite time concentration displayed in Theorem ?? . Note that the construction of of Theorem ?? is performed near $t = +\infty$ (by translation invariance, it is then obvious to obtain a solution on the time interval $[0, +\infty)$). An interesting question is to understand the behaviour of such solutions for $t \leq 0$.

3. *Zero energy global solutions.* From the proof of Theorem ??, the solution u has zero energy. In [?], it is proved that any zero energy solution satisfying (??) blows up in finite time with the log–log regime. Thus, in the neighborhood of Q , $e^{it}Q$ is the only global zero energy solution. For the critical (gKdV) equation, a similar result holds, though in a stronger topology (see [?]). Note that the existence of global in time zero energy solutions is strongly related to Liouville type theorems and to blow up profile, see [?, ?]. For (NLS), the only known examples of global in positive time zero energy solutions so far were the time periodic solutions $e^{it}P$ where P is any solution to the stationary equation $\Delta P - P + P^3 = 0$. Therefore, the existence of such a non trivial global (for positive time) zero energy solution $u(t)$ is surprising. For other works related to minimal mass solutions and their key role in the dynamics of the flow, we refer to [?, ?, ?, ?, ?, ?, ?].

4. *Blow up speed for (NLS).* The question of determining all possible blow up rates for solutions of nonlinear dispersive equations is in general intricate. For the (NLS) equation (??) in the mass supercritical–energy subcritical range, a universal sharp upper bound on the blow up rate has been derived in [?] for radial data, but no such bound exists for the mass critical problem. For (NLS) with a double power non linearity of the form $|u|^{p-1}u + |u|^2u$ where $1 < p < 3$, the minimal mass solution has a surprising blow up rate different from the conformal rate, see [?]. For the mass critical (gKdV) equation, solutions arbitrarily close to the solitary wave with arbitrarily fast blow up speed have been constructed in [?]. Recall that constructions of blow up solutions with various blow up rate are also available in the energy critical and super–critical context, see [?, ?, ?, ?, ?, ?]. However, such general constructions seem by now out of reach for the mass critical (NLS) problem. In this context, the derivation of the anomalous blow up speed (??), in spite of its rigidity, is an interesting new fact. We will see in the proof how such a blow up rate is related to strong coupling between the solitary waves.

1.5. **Notation.** Let $\Sigma = H^1 \cap L^2(|x|^2 dx)$. The L^2 scalar product of two complex valued functions $f, g \in L^2(\mathbb{R}^2)$ is denoted by

$$\langle f, g \rangle = \operatorname{Re} \left(\int_{\mathbb{R}^2} f(x) \bar{g}(x) dx \right).$$

In this paper, K is an integer with $K \geq 2$. For brevity, $\sum_k$ denotes $\sum_{k=1}^K$. For $k = 1, \dots, K$, e_k denotes the unit vector of $\mathbb{R}^2$ corresponding to the complex number $e^{i \frac{2\pi(k-1)}{K}}$. We define the constant $\kappa = \kappa(K)$ by

$$\kappa = \left| 1 - e^{i \frac{2\pi}{K}} \right| = (2 - 2 \cos(2\pi/K))^{1/2} > 0. \quad (1.16)$$

Recall that we denote by $Q(x) := Q(|x|)$ the unique radial positive ground state of (??):

$$Q'' + \frac{Q'}{r} - Q + Q^3 = 0, \quad Q'(0) = 0, \quad \lim_{r \rightarrow +\infty} Q(r) = 0. \quad (1.17)$$

It is well-known and easily checked by ODE arguments that for some constant $c_Q > 0$,

$$\text{for all } r > 1, \quad \left| Q(r) - c_Q r^{-\frac{1}{2}} e^{-r} \right| + \left| Q'(r) + c_Q r^{-\frac{1}{2}} e^{-r} \right| \lesssim r^{-\frac{3}{2}} e^{-r}. \quad (1.18)$$

We set

$$I_Q = \int Q^3(x) e^{x_1} dx, \quad x = (x_1, x_2). \quad (1.19)$$

We denote by $\mathcal{Y}$ the set of smooth functions f such that

$$\text{for all } p \in \mathbb{N}, \text{ there exists } q \in \mathbb{N}, \text{ s.t. for all } x \in \mathbb{R}^2 \quad |f^{(p)}(x)| \lesssim |x|^q e^{-|x|}. \quad (1.20)$$

Let Λ be the generator of L^2 -scaling in two dimensions:

$$\Lambda f = f + x \cdot \nabla f.$$

The linearization of (??) around Q involves the following Schrödinger operators:

$$L_+ := -\Delta + 1 - 3Q^2, \quad L_- := -\Delta + 1 - Q^2.$$

Denote by $\rho \in \mathcal{Y}$ the unique radial solution H^1 to

$$L_+ \rho = \frac{|x|^2}{4} Q \quad (1.21)$$

which satisfies on $\mathbb{R}^2$

$$|\rho(x)| + |\nabla \rho(x)| \lesssim (1 + |x|^3) Q(x). \quad (1.22)$$

We recall the generalized null space relations (see [?])

$$\begin{aligned} L_- Q &= 0, \quad L_+(\Lambda Q) = -2Q, \quad L_- (|x|^2 Q) = -4\Lambda Q, \quad L_+ \rho = \frac{|x|^2}{4} Q, \\ L_+(\nabla Q) &= 0, \quad L_-(xQ) = -2\nabla Q, \end{aligned} \quad (1.23)$$

and the classical (see e.g. [?, ?, ?, ?, ?]) coercivity property: there exists $\mu > 0$ such that for all $\eta \in H^1$,

$$\begin{aligned} \langle L_+ \operatorname{Re} \eta, \operatorname{Re} \eta \rangle + \langle L_- \operatorname{Im} \eta, \operatorname{Im} \eta \rangle &\geq \mu \|\eta\|_{H^1}^2 \\ &- \frac{1}{\mu} [\langle \eta, Q \rangle^2 + \langle \eta, |x|^2 Q \rangle^2 + \langle \eta, xQ \rangle^2 + \langle \eta, i\rho \rangle^2 + \langle \eta, i\nabla Q \rangle^2]. \end{aligned} \quad (1.24)$$

1.6. Outline of the paper. The main goal of Sect. ?? is to construct a symmetric K -bubble approximate solution to (NLS) and to extract the formal evolution system of the geometrical parameters of the bubbles. The key observation is that this system contains forcing terms due to the nonlinear interactions of the waves, and has a special solution corresponding at the main order to the regime of Theorem ?? (see Sect. ??). In Sect. ??, we prove uniform estimates on particular *backwards solutions* of (NLS) related to the special regime of Theorem ??. We proceed in two main steps. First, we control the residue term by energy arguments in the context of multi-bubbles. Second, a careful adjustment of the *final data* yields a uniform control of the geometrical parameters. In Sect. ??, we finish the proof of Theorem ?? by compactness arguments on a suitable sequence of backwards solutions of (NLS) satisfying the uniform estimates of Sect. ??.

Acknowledgements. Y.M. was partly supported by ERC 291214 BLOWDISOL. P.R. was supported by the ERC-2014-CoG 646650 SingWave and the Institut Universitaire de France. This work was done while the authors were members of the MSRI (Fall 2015). They warmly thank MSRI for its hospitality.

2. APPROXIMATE SOLUTION

In this section, we first construct a symmetric K -bubble approximate solution to (NLS) and extract the evolution system of the geometrical parameters of the bubbles. This system contains forcing terms due to the nonlinear interactions of the waves. Second, we write explicitly a special formal solution of this system that will serve as a guideline for the construction of the special solution $u(t)$ of Theorem ???. Third, we state a standard modulation lemma around the approximate solution. Recall that the integer $K \geq 2$ is fixed.

2.1. Approximate solution and nonlinear forcing. Consider a time dependent C^1 function $\vec{p}$ of the form

$$\vec{p} = (\lambda, z, \gamma, \beta, b) \in (0, +\infty)^2 \times \mathbb{R}^3,$$

with $|b| + |\beta| \ll 1$ and $z \gg 1$. We renormalize the flow by considering

$$u(t, x) = \frac{e^{i\gamma(s)}}{\lambda(s)} v(s, y), \quad dt = \lambda^2(s) ds, \quad y = \frac{x}{\lambda(s)}, \quad (2.1)$$

so that

$$i\partial_t u + \Delta u + |u|^2 u = \frac{e^{i\gamma}}{\lambda^3} \left[i\dot{v} + \Delta v - v + |v|^2 v - i\frac{\dot{\lambda}}{\lambda} \Lambda v + (1 - \dot{\gamma})v \right] \quad (2.2)$$

($\dot{v}$ denotes derivation with respect to s). We introduce the following $\vec{p}$ -modulated ground state solitary waves, for any $k \in \{1, \dots, K\}$,

$$P_k(s, y) = e^{i\Gamma_k(s, y - z_k(s))} Q_{a(z(s))}(y - z_k(s)), \quad (2.3)$$

for

$$\beta_k = \beta e_k, \quad z_k = z e_k, \quad \Gamma_k(s, y) = \beta_k \cdot y - \frac{b}{4} |y|^2, \quad (2.4)$$

and where we have fixed

$$Q_a = Q + a\rho, \quad a(z) = -c_a z^{\frac{1}{2}} e^{-\kappa z}, \quad c_a = \begin{cases} \frac{\kappa^{\frac{1}{2}} c_Q I_Q}{4\langle \rho, Q \rangle} > 0 & \text{for } K = 2 \\ \frac{\kappa^{\frac{1}{2}} c_Q I_Q}{2\langle \rho, Q \rangle} > 0 & \text{for } K \geq 3 \end{cases} \quad (2.5)$$

Note that the introduction of such modulated Q_a corresponds to the intrinsic instability of the pseudo-conformal blow up regime ($a = 0$ leads to $b(s) = s^{-1}$). Similar exact Q_a (at any order of a) were introduced in [?]. The explicit above choice of $a(z)$ corresponds to direct integration of the nonlinear interactions at the main order, as explained in Sect. ??. We also refer to (??) in the proof of Lemma ?? where this choice of $a(z)$ leads to an almost conservation of the mass for the approximate solution $\mathbf{P}$ defined below. Note that the different formula for c_a depending on the value of K corresponds to the fact that for $K \geq 3$, each given soliton has exactly two closest neighbor solitons.

Let

$$\mathbf{P}(s, y) = \mathbf{P}(y; (z(s), b(s), \beta(s))) = \sum_k P_k(s, y). \quad (2.6)$$

Then, $\mathbf{P}$ is an approximate solution of the rescaled equation in the following sense.

Lemma 3 (Leading order approximate flow). *Let the vectors of modulation equations be*

$$\vec{m}_k^a = \begin{pmatrix} b + \frac{\dot{\lambda}}{\lambda} \\ \dot{z}_k - 2\beta_k + \frac{\dot{\lambda}}{\lambda} z_k \\ \dot{\gamma} - 1 + |\beta_k|^2 - \frac{\dot{\lambda}}{\lambda}(\beta_k \cdot z_k) - (\beta_k \cdot \dot{z}_k) \\ \dot{\beta}_k - \frac{\dot{\lambda}}{\lambda}\beta_k + \frac{b}{2}(\dot{z}_k - 2\beta_k + \frac{\dot{\lambda}}{\lambda} z_k) \\ \dot{b} + b^2 - 2b(b + \frac{\dot{\lambda}}{\lambda}) - a \end{pmatrix}, \quad \vec{M}V = \begin{pmatrix} -i\Lambda V \\ -i\nabla V \\ -V \\ -yV \\ \frac{|y|^2}{4}V \end{pmatrix}. \quad (2.7)$$

Then the error to the renormalized flow (??) at $\mathbf{P}$,

$$\mathcal{E}_{\mathbf{P}} = i\dot{\mathbf{P}} + \Delta\mathbf{P} - \mathbf{P} + |\mathbf{P}|^2\mathbf{P} - i\frac{\dot{\lambda}}{\lambda}\Lambda\mathbf{P} + (1 - \dot{\gamma})\mathbf{P} \quad (2.8)$$

decomposes as

$$\mathcal{E}_{\mathbf{P}} = \sum_k [e^{i\Gamma_k}\Psi_k](y - z_k), \quad \Psi_k = \vec{m}_k^a \cdot \vec{M}Q_a + i\dot{z}a'(z)\rho + G_k + \Psi_{Q_a}, \quad (2.9)$$

where

$$\|G_k\|_{L^\infty} \lesssim z^{-\frac{1}{2}}e^{-\kappa z}, \quad \|\Psi_{Q_a}\|_{L^\infty} \lesssim |a|^2, \quad (2.10)$$

and

$$\left| \langle G_k, iQ_a \rangle + \kappa c_a \langle \rho, Q \rangle b z^{\frac{3}{2}} e^{-\kappa z} \right| \lesssim (|\beta|^2 z^2 + |b|^2 z^4 + |\beta|z + |b|z) z^{-\frac{1}{2}} e^{-\kappa z} + z^3 e^{-2\kappa z}. \quad (2.11)$$

Proof of Lemma ??. **step 1.** Equation for P_k . Let

$$\mathcal{E}_{P_k} = i\dot{P}_k + \Delta P_k - P_k + |P_k|^2 P_k - i\frac{\dot{\lambda}}{\lambda}\Lambda P_k + (1 - \dot{\gamma})P_k.$$

Let $y_{z_k} = y - z_k$. By direct computations

$$\begin{aligned} i\dot{P}_k = & \left[e^{i\Gamma_k} \left(i\dot{z}a'(z)\rho - (\dot{\beta}_k \cdot y_{z_k})Q_a + (\dot{z}_k \cdot \beta_k)Q_a \right. \right. \\ & \left. \left. + \frac{\dot{b}}{4}|y_{z_k}|^2 Q_a - \frac{b}{2}(\dot{z}_k \cdot y_{z_k})Q_a - i(\dot{z}_k \cdot \nabla Q_a) \right) \right] (y_{z_k}), \end{aligned}$$

$$\begin{aligned} \Delta P_k = & \left[e^{i\Gamma_k} \left(\Delta Q_a - |\beta_k|^2 Q_a - \frac{b^2}{4}|y_{z_k}|^2 Q_a - ibQ_a \right. \right. \\ & \left. \left. + b(\beta_k \cdot y_{z_k})Q_a + 2i(\beta_k \cdot \nabla Q_a) - ib(y_{z_k} \cdot \nabla Q_a) \right) \right] (y_{z_k}), \end{aligned}$$

$$\begin{aligned} \Lambda P_k = & \left[e^{i\Gamma_k} \left(\Lambda Q_a + i(\beta_k \cdot y_{z_k})Q_a - i\frac{b}{2}|y_{z_k}|^2 Q_a + (y_{z_k} \cdot \nabla Q_a) \right. \right. \\ & \left. \left. + i(z_k \cdot \beta_k)Q_a - i\frac{b}{2}(z_k \cdot y_{z_k})Q_a + (z_k \cdot \nabla Q_a) \right) \right] (y_{z_k}). \end{aligned}$$

Thus,

$$\begin{aligned} \mathcal{E}_{P_k} = & \left[e^{i\Gamma_k} \left(-i(b + \frac{\dot{\lambda}}{\lambda})\Lambda Q_a - i(\dot{z}_k - 2\beta_k + \frac{\dot{\lambda}}{\lambda}z_k) \cdot \nabla Q_a \right. \right. \\ & - (\dot{\gamma} - 1 + |\beta_k|^2 - \frac{\dot{\lambda}}{\lambda}(\beta_k \cdot z_k) - (\beta_k \cdot \dot{z}_k))Q_a - (\dot{\beta}_k - \frac{\dot{\lambda}}{\lambda}\beta_k + \frac{b}{2}(\dot{z}_k - 2\beta_k + \frac{\dot{\lambda}}{\lambda}z_k)) \cdot y_{z_k} Q_a \\ & \left. \left. + \frac{1}{4}(\dot{b} + b^2 - 2b(b + \frac{\dot{\lambda}}{\lambda}))|y_{z_k}|^2 Q_a + i\dot{z}a'(z)\rho + \Delta Q_a - Q_a + |Q_a|^2 Q_a \right) \right] (y_{z_k}) \end{aligned}$$

By $\Delta Q - Q + Q^3 = 0$ and the definition of ρ , $L_+\rho = -\Delta\rho + \rho - 3Q^2\rho = \frac{1}{4}|y|^2Q$ (see (??)), we have

$$\Delta Q_a - Q_a + |Q_a|^2 Q_a = -\frac{a}{4}|y|^2 Q_a + \Psi_{Q_a},$$

where

$$\Psi_{Q_a} = |Q_a|^2 Q_a - Q^3 - 3aQ^2\rho + \frac{a^2}{4}|y|^2\rho. \quad (2.12)$$

We have thus obtained the P_k equation

$$\mathcal{E}_{P_k} = \left[e^{i\Gamma_k} \left(\vec{m}_k^a \cdot \vec{M} Q_a + i\dot{z}a'(z)\rho + \Psi_{Q_a} \right) \right] (y - z_k), \quad (2.13)$$

where $\vec{m}_k^a$ and $\vec{M}$ are defined in (??).

step 2. Equation for $\mathbf{P}$. From step 1 and the definition of $\mathcal{E}_{\mathbf{P}}$ in (??), it follows that

$$\mathcal{E}_{\mathbf{P}} = \sum_k \mathcal{E}_{P_k} + |\mathbf{P}|^2 \mathbf{P} - \sum_k |P_k|^2 P_k.$$

Observe that

$$|\mathbf{P}|^2 \mathbf{P} - \sum_k |P_k|^2 P_k = \sum_{j,k,l} P_k P_j \bar{P}_l - \sum_k |P_k|^2 P_k = \sum_k F_k,$$

with

$$F_k = 2|P_k|^2 \sum_{j \neq k} P_j + P_k^2 \sum_{j \neq k} \bar{P}_j + \bar{P}_k \sum_{j \neq k, l \neq k, j \neq l} P_j P_l = [e^{i\Gamma_k} G_k](y - z_k),$$

where we have set

$$G_k = 2G_k^{(I)} + \bar{G}_k^{(I)} + G_k^{(II)}, \quad (2.14)$$

and

$$\begin{aligned} G_k^{(I)}(y) &= [e^{-i\Gamma_k} Q_a^2](y) \sum_{j \neq k} [e^{i\Gamma_j} Q_a](y - (z_j - z_k)), \\ G_k^{(II)}(y) &= [e^{-2i\Gamma_k} Q_a](y) \sum_{j \neq k, l \neq k, j \neq l} ([e^{i\Gamma_j} Q_a](y - (z_j - z_k)) \cdot [e^{i\Gamma_l} Q_a](y - (z_l - z_k))). \end{aligned}$$

Therefore,

$$\mathcal{E}_{\mathbf{P}} = \sum_k [e^{i\Gamma_k} \Psi_k](y - z_k) \quad \text{where} \quad \Psi_k = \vec{m}_k^a \cdot \vec{M} Q_a + i\dot{z}a'(z)\rho + G_k + \Psi_{Q_a}. \quad (2.15)$$

step 3 Nonlinear interaction estimates. In order to estimate the various terms in (??), we will use the following interaction estimates: let $\omega, \tilde{\omega} \in \mathbb{R}^2$, $|\omega| \gg 1$, $|\tilde{\omega}| \gg 1$, let $q \geq 0$, then:

$$\int (1 + |y|^q) Q^3(y) Q(y - \omega) dy \lesssim |\omega|^{-\frac{1}{2}} e^{-|\omega|}. \quad (2.16)$$

$$\int (1 + |y|^q) Q^2(y) Q(y - \omega) Q(y - \tilde{\omega}) dy \lesssim e^{-\frac{3}{2}|\omega|} + e^{-\frac{3}{2}|\tilde{\omega}|}. \quad (2.17)$$

$$\left| \int Q^3(y) Q(y - \omega) dy - c_Q I_Q |\omega|^{-\frac{1}{2}} e^{-|\omega|} \right| \lesssim |\omega|^{-\frac{3}{2}} e^{-|\omega|}, \quad (2.18)$$

with I_Q given by (??).

Proof of (??). From (??), observe that

$$Q(y) Q(y - \omega) \lesssim (1 + |y|)^{-\frac{1}{2}} (1 + |y - \omega|)^{-\frac{1}{2}} e^{-|y|} e^{-|\omega| + |y|} \lesssim |\omega|^{-\frac{1}{2}} e^{-|\omega|}. \quad (2.19)$$

Thus,

$$\int (1 + |y|^q) Q^3(y) Q(y - \omega) dy \lesssim |\omega|^{-\frac{1}{2}} e^{-|\omega|} \int (1 + |y|^q) Q^2(y) dy \lesssim |\omega|^{-\frac{1}{2}} e^{-|\omega|}.$$

Proof of (??). From (??),

$$\begin{aligned} \int (1 + |y|^q) Q^2(y) Q(y - \omega) Q(y - \tilde{\omega}) dy &\lesssim \int (1 + |y|^q) Q^2(y) Q^{\frac{3}{4}}(y - \omega) Q^{\frac{3}{4}}(y - \tilde{\omega}) dy \\ &\lesssim e^{-\frac{3}{4}|\omega|} e^{-\frac{3}{4}|\tilde{\omega}|} \int (1 + |y|^q) Q^{\frac{1}{2}}(y) dy \lesssim e^{-\frac{3}{4}|\omega|} e^{-\frac{3}{4}|\tilde{\omega}|}. \end{aligned}$$

Proof of (??). First, using (??),

$$\int_{|y| > \frac{3}{4}|\omega|} Q^3(y) Q(y - \omega) dy \lesssim e^{-\frac{9}{4}|\omega|} \int Q(y - \omega) dy \lesssim e^{-\frac{9}{4}|\omega|}.$$

Second, for $|y| < \frac{3}{4}|\omega|$, we use (??) to write

$$\left| Q(y - \omega) - c_Q |y - \omega|^{-\frac{1}{2}} e^{-|y - \omega|} \right| \lesssim |y - \omega|^{-\frac{3}{2}} e^{-|y - \omega|} \lesssim |\omega|^{-\frac{3}{2}} e^{-|\omega| + |y|}.$$

In particular,

$$\left| \int_{|y| < \frac{3}{4}|\omega|} Q^3(y) Q(y - \omega) dy - c_Q \int_{|y| < \frac{3}{4}|\omega|} Q^3(y) |y - \omega|^{-\frac{1}{2}} e^{-|y - \omega|} dy \right| \lesssim |\omega|^{-\frac{3}{2}} e^{-|\omega|}.$$

Still for $|y| < \frac{3}{4}|\omega|$, the expansion

$$|y - \omega|^2 = |\omega|^2 - 2y \cdot \omega + |y|^2$$

implies

$$\left| |y - \omega|^{-\frac{1}{2}} - |\omega|^{-\frac{1}{2}} \right| \lesssim |\omega|^{-\frac{3}{2}} |y|$$

and

$$\left| |y - \omega| - |\omega| + y \cdot \frac{\omega}{|\omega|} \right| \lesssim |\omega|^{-1} |y|^2.$$

Thus,

$$\left| e^{-|y - \omega|} - e^{-|\omega| + y \cdot \frac{\omega}{|\omega|}} \right| \lesssim |\omega|^{-1} |y|^2 \left(e^{-|y - \omega|} + e^{-|\omega| + y \cdot \frac{\omega}{|\omega|}} \right) \lesssim |\omega|^{-1} |y|^2 e^{-|\omega|} e^{|y|}.$$

Therefore,

$$\left| |y - \omega|^{-\frac{1}{2}} e^{-|y - \omega|} - |\omega|^{-\frac{1}{2}} e^{-|\omega| + y \cdot \frac{\omega}{|\omega|}} \right| \lesssim |\omega|^{-\frac{3}{2}} (1 + |y|^2) e^{-|\omega|} e^{|y|},$$

and so

$$\begin{aligned} & \left| \int_{|y| < \frac{3}{4}|\omega|} Q^3(y) |y - \omega|^{-\frac{1}{2}} e^{-|y-\omega|} dy - |\omega|^{-\frac{1}{2}} e^{-|\omega|} \int_{|y| < \frac{3}{4}|\omega|} Q^3(y) e^{y \cdot \frac{\omega}{|\omega|}} dy \right| \\ & \lesssim |\omega|^{-\frac{3}{2}} e^{-|\omega|} \int (1 + |y|^2) e^{-2|y|} dy \lesssim |\omega|^{-\frac{3}{2}} e^{-|\omega|}. \end{aligned}$$

Also, we see that

$$|\omega|^{-\frac{1}{2}} e^{-|\omega|} \int_{|y| > \frac{3}{4}|\omega|} Q^3(y) e^{y \cdot \frac{\omega}{|\omega|}} dy \lesssim |\omega|^{-\frac{1}{2}} e^{-|\omega|} \int_{|y| > \frac{3}{4}|\omega|} e^{-2|y|} dy \lesssim e^{-2|\omega|}.$$

Since for all $\omega \neq 0$ (see (??)),

$$I_Q = \int Q^3(y) e^{y \cdot \frac{\omega}{|\omega|}} dy,$$

we have proved (??).

step 4. Estimates of G_k and Ψ_{Q_a} . We are now in position to prove (??) and (??). The estimate on Ψ_{Q_a} in (??) follows directly from its definition (??). To estimate G_k as in (??), we first note that from (??), (??) and the definition of $a(z)$ in (??), we have, for some $q > 0$,

$$|Q_a| \lesssim |y|^{-\frac{1}{2}} e^{-|y|} + |a(z)| |y|^q e^{-|y|} \lesssim (1 + |y|)^{-\frac{1}{2}} e^{-|y|} + z^{\frac{1}{2}} e^{-\kappa z} (1 + |y|)^q e^{-|y|}.$$

Moreover, for $j \neq k$, from the definition of κ in (??),

$$|z_j - z_k| = z |e_k - e_j| \geq \kappa z.$$

From this, it follows easily that for $j \neq k$,

$$|Q_a(y)| |Q_a(y - (z_k - z_j))| \lesssim z^{-\frac{1}{2}} e^{-\kappa z},$$

which in light of the explicit formula (??) yields the control of G_k in (??).

We now turn to the proof of (??) which requires a more careful analysis of the interaction terms. We first compute the main order of the contribution of $G_1^{(1)}$ to $\langle G_1, iQ \rangle$. For $j = 2, \dots, K$,

$$\begin{aligned} & \langle [e^{-i\Gamma_1} Q_a^2](y) [e^{i\Gamma_j} Q_a](y - (z_j - z_1)), iQ_a \rangle \\ & = \int Q_a^3(y) Q_a(y - z(e_j - e_1)) \sin(\Gamma_j(y - z(e_j - e_1)) - \Gamma_1(y)) dy. \end{aligned}$$

First, by the decay of ρ (see (??)), (??) and the definition of $a(z)$ in (??), we have

$$\int |Q_a^3(y) Q_a(y - z(e_j - e_1)) - Q^3(y) Q(y - z(e_j - e_1))| dy \lesssim |a| z^{\frac{5}{2}} e^{-\kappa z} \lesssim z^3 e^{-2\kappa z}.$$

Next, note that, since $\Gamma_j = \beta_j \cdot y - \frac{b}{4}|y|^2$,

$$\begin{aligned} & |\sin(\Gamma_j(y - z(e_j - e_1)) - \Gamma_1(y)) - (\Gamma_j(y - z(e_j - e_1)) - \Gamma_1(y))| \\ & \lesssim |\Gamma_j(y - z(e_j - e_1))|^2 + |\Gamma_1(y)|^2 \lesssim |\beta|^2(|y|^2 + z^2) + |b|^2(|y|^4 + z^4), \end{aligned}$$

and

$$\left| (\Gamma_j(y - z(e_j - e_1)) - \Gamma_1(y)) + \frac{b}{4}|z(e_j - e_1)|^2 \right| \lesssim |\beta|(|y| + z) + |b|(|y|^2 + |y|z).$$

Thus, using (??),

$$\begin{aligned} & \int Q^3(y)Q(y - z(e_j - e_1)) \left| \sin(\Gamma_j(y - z(e_j - e_1)) - \Gamma_1(y)) + \frac{b}{4}|z(e_j - e_1)|^2 \right| dy \\ & \lesssim (|\beta|^2 z^2 + |b|^2 z^4 + |\beta|z + |b|z)z^{-\frac{1}{2}}e^{-\kappa z}. \end{aligned}$$

Therefore, we have proved

$$\begin{aligned} & \left| \langle [e^{i\Gamma_1} Q_a^2](y) [e^{i\Gamma_j} Q_a](y - z(e_j - e_1)), iQ_a \rangle + \frac{b}{4}|z(e_j - e_1)|^2 \int Q^3(y)Q(y - z(e_j - e_1)) \right| \\ & \lesssim (|\beta|^2 z^2 + |b|^2 z^4 + |\beta|z + |b|z)z^{-\frac{1}{2}}e^{-\kappa z} + z^3 e^{-2\kappa z}. \end{aligned}$$

For $j = 2$ and $j = K$, we have $|z(e_j - e_1)| = \kappa z$, and so using (??),

$$\begin{aligned} & \left| \langle [e^{i\Gamma_1} Q_a^2](y) [e^{i\Gamma_j} Q_a](y - z(e_j - e_1)), iQ_a \rangle + \frac{b}{4}c_Q I_Q \kappa^{\frac{3}{2}} z^{\frac{3}{2}} e^{-\kappa z} \right| \\ & \lesssim (|\beta|^2 z^2 + |b|^2 z^4 + |\beta|z + |b|z)z^{-\frac{1}{2}}e^{-\kappa z} + z^3 e^{-2\kappa z}. \end{aligned}$$

For $K \geq 4$ and $j = 3, \dots, K-1$, we have $|e_j - e_1| > \kappa'$, for some $\kappa' > \kappa$. Thus the following bound follows from similar computations

$$\begin{aligned} & |\langle [e^{i\Gamma_1} Q_a^2](y) [e^{i\Gamma_j} Q_a](y - z(e_j - e_1)), iQ_a \rangle| \\ & \lesssim (|\beta|^2 z^2 + |b|^2 z^4 + |\beta|z + |b|z^2)z^{-\frac{1}{2}}e^{-\kappa' z} + z^3 e^{-2\kappa z}. \end{aligned}$$

Note that $\langle 2G_1^{(I)} + \overline{G}_1^{(I)}, iQ_a \rangle = \langle G_1^{(I)}, iQ_a \rangle$. We finally bound the contribution of $G_1^{(II)}$. For $j \neq 1$, $l \neq 1$ and $l \neq j$,

$$\begin{aligned} & \langle [e^{-2i\Gamma_k} Q_a](y) [e^{i\Gamma_j} Q_a](y - (z_j - z_k)) [e^{i\Gamma_l} Q_a](y - (z_l - z_k)), iQ_a \rangle \\ & = \int Q_a^2(y)Q_a(y - (z_j - z_k))Q_a(y - (z_l - z_k)) \\ & \quad \times \sin(\Gamma_j(y - (z_j - z_k)) + \Gamma_l(y - (z_l - z_k)) - 2\Gamma_k(y)) dy. \end{aligned}$$

By (??), the bound on $|a|$ and $|\Gamma_j| \lesssim |\beta|(|y| + z) + |b|(|y|^2 + z^2)$, this term is bounded by $(|\beta|z + |b|z^2)e^{-\frac{3}{2}\kappa z}$.

Gathering these estimates, using the definition of the constant c_a in (??) which takes into account the two different cases $K = 2$ and $K \geq 2$ (for $K = 2$, the soliton P_1 has nonlinear interaction with only one other soliton, while for $K \geq 3$, it has exactly two closest neighboring solitons, P_2 and P_K), we obtain finally

$$|\langle G_k, iQ_a \rangle + \kappa c_a \langle \rho, Q \rangle b z^{\frac{3}{2}} e^{-\kappa z}| \lesssim (|\beta|^2 z^2 + |b|^2 z^4 + |\beta|z + |b|z)z^{-\frac{1}{2}}e^{-\kappa z} + z^3 e^{-2\kappa z},$$

which completes the proof of (??). $\square$

2.2. Formal resolution of the modulation system with forcing. From Lemma ??, we derive a simplified modulation system with forcing term and we determine one of its approximate solution that is relevant for the regime of Theorem ?. Moreover, we justify the special choice of function $a(z)$ in (??). Formally, i.e. assuming that $\mathbf{P}$ is a solution of (??) up to error terms of lower order than the ones in (??) (making this rigorous will be the object of the bootstrap estimates in Sect. ??), we have the following bounds ($\vec{m}_k^a$ is defined in (??))

$$|\vec{m}_1^a| \lesssim z^{-\frac{1}{2}}e^{-\kappa z}. \quad (2.20)$$

Indeed, (??) is obtained from (??)–(??) by projecting $\mathcal{E}_{\mathbf{P}}$ onto directions related to the generalized null space (??) (see Lemma ?? for rigorous computations). To simplify the discussion, we drop the equation of γ , which is not coupled with any other equation and has no influence on the regime. Next, we see that using the first line of $\vec{m}_1^a$, i.e. $|b + \frac{\dot{\lambda}}{\lambda}| \lesssim z^{-\frac{1}{2}}e^{-\kappa z}$, we can replace $\frac{\dot{\lambda}}{\lambda}$ by $-b$ in all the other estimates. Similarly, we insert the estimate on $\dot{z}$ from the second line into the estimate for $\dot{\beta}$. We obtain the following simplified system

$$|b + \frac{\dot{\lambda}}{\lambda}| + |\dot{z} - 2\beta - bz| + |\dot{\beta} + b\beta| + |\dot{b} + b^2 - a| \lesssim z^{-\frac{1}{2}}e^{-\kappa z}. \quad (2.21)$$

It is easy to check the following estimates

Lemma 4. *Let $(z_{\text{app}}, \lambda_{\text{app}}(s), \beta_{\text{app}}, b_{\text{app}}(s))$ be such that*

$$\begin{aligned} \lambda_{\text{app}}(s) &= \log^{-1}(s), \quad z_{\text{app}}^{-\frac{3}{2}}(s)e^{\kappa z_{\text{app}}(s)} = \frac{\kappa c_a}{2}s^2, \\ |\beta_{\text{app}}(s)| &\lesssim s^{-1}\log^{-\frac{3}{2}}(s), \quad b_{\text{app}}(s) = s^{-1}\log^{-1}(s). \end{aligned} \quad (2.22)$$

Then,

$$\begin{aligned} z_{\text{app}}(s) &\sim \frac{2}{\kappa}\log(s), \quad |b_{\text{app}} + \frac{\dot{\lambda}_{\text{app}}}{\lambda_{\text{app}}}| = 0, \quad |\dot{z}_{\text{app}} - 2\beta_{\text{app}} - b_{\text{app}}z_{\text{app}}| \lesssim s^{-1}\log^{-\frac{1}{2}}(s), \\ |a(z_{\text{app}}) + s^{-2}\log^{-1}(s)| &\lesssim s^{-2}\log^{-\frac{3}{2}}(s), \quad |\dot{b}_{\text{app}} + b_{\text{app}}^2 - a(z_{\text{app}})| \lesssim s^{-2}\log^{-\frac{3}{2}}(s). \end{aligned} \quad (2.23)$$

The above estimates mean that (??) is a reasonable guess for the first order asymptotics as $s \rightarrow +\infty$ of some particular solutions of (??) (we refer to Sect. ?? for a rigorous integration of (??)). Note that we do not actually determine the main order of $\beta(s)$; to do this, more interaction computations would be necessary. However, since $|\dot{\beta} + b\beta| \lesssim z^{-\frac{1}{2}}e^{-\kappa z}$, formally, we obtain $|\dot{\beta}| \lesssim s^{-2}\log^{-2}(s)$, which justifies a bootstrap on $\beta(s)$ of the form $|\beta(s)| \ll s^{-1}\log^{-2}(s)$. Note also that there exist solutions of (??) with different asymptotics, corresponding to (NLS) solutions like $v(t)$ of Corollary ??.

To complete this formal discussion, we justify the choice of $a(z)$ in (??) in the regime given by (??). Indeed, projecting Ψ_1 onto the direction iQ_a , from (??), we obtain at the leading order

$$|\dot{z}a'(z) - \kappa c_a b z^{\frac{3}{2}}e^{-\kappa z}| \lesssim (|\beta|^2 z^2 + |b|^2 z^4 + |\beta|z + |b|z)z^{-\frac{1}{2}}e^{-\kappa z} + z^3e^{-2\kappa z}. \quad (2.24)$$

In the regime suggested by (??), since $|\beta| \ll |b|z$, we have $bz \sim \dot{z}$ and thus, simplifying $\dot{z}$, we obtain

$$|a'(z) - \kappa c_a z^{\frac{1}{2}}e^{-\kappa z}| \lesssim z^{-\frac{1}{2}}e^{-\kappa z},$$

which justifies the definition (??) by integrating in z .

2.3. Modulation of the approximate solution. We state a standard modulation result around $\mathbf{P}$. We restrict ourselves to the case of solutions invariant by the rotation preserving $\mathbf{P}$. Denote by τ_K the rotation of center 0 and angle $\frac{2\pi}{K}$ in $\mathbb{R}^2$. Since Q and ρ are radial, by definition of P_k and β_k , z_k in (??) and (??), we have for $k \in \{1, \dots, K-1\}$, $P_k(y) = P_{k+1}(\tau_K y)$ and $P_K(y) = P_1(\tau_K y)$. In particular, it follows that $\mathbf{P}(\tau_K y) = \mathbf{P}(y)$, i.e. $\mathbf{P}$ is invariant by the rotation τ_K . Note also that equation (??) is invariant by rotation. In particular, if a solution of (??) is invariant by the rotation τ_K at some time, then it is invariant by rotation at any time. In this context, the following modulation result relies on a standard argument based on the Implicit Function Theorem (see e.g. Lemma 2 in [?]) and we omit its proof.

Lemma 5 (Modulation around $\mathbf{P}$). *Let I be some time interval. Let $u \in C(I, H^1(\mathbb{R}^2))$ be a solution of (??) invariant by the rotation τ_K and such that*

$$\sup_{t \in I} \left\| e^{-i\tilde{\gamma}(t)} \tilde{\lambda}(t) u(t, \tilde{\lambda}(t) \cdot) - \sum_k Q(\cdot - e_k \tilde{z}(t)) \right\|_{H^1} < \delta \quad (2.25)$$

for some $\tilde{\lambda}(t) > 0$, $\tilde{\gamma}(t) \in \mathbb{R}^2$, $\tilde{z}(t) > \delta^{-1}$, where $\delta > 0$ is small enough. Then, there exist a C^1 function

$$\vec{p} = (\lambda, z, \gamma, \beta, b) : I \rightarrow (0, \infty)^2 \times \mathbb{R}^3,$$

such that, for $\mathbf{P}(t, y) = \mathbf{P}(y; z(t), b(t), \beta(t))$ as defined in (??), the solution $u(t)$ decomposes on I as

$$u(t, x) = \frac{e^{i\gamma(t)}}{\lambda(t)} (\mathbf{P} + \varepsilon)(t, y), \quad y = \frac{x}{\lambda(t)}, \quad (2.26)$$

where for all $t \in I$,

$$|b(t)| + |\beta(t)| + \|\varepsilon(t)\|_{H^1} + |z(t)|^{-1} \lesssim \delta, \quad (2.27)$$

and, setting $\varepsilon(t, y) = [e^{i\Gamma_1} \eta_1](t, y - z_1)$,

$$\langle \eta_1(t), |y|^2 Q \rangle = \langle \eta_1(t), y Q \rangle = \langle \eta_1(t), i\rho \rangle = \langle \eta_1(t), i\nabla Q \rangle + \langle \eta_1(t), i\Lambda Q \rangle = 0. \quad (2.28)$$

Moreover, ε is also invariant by the rotation τ_K .

Note that the choice of the special orthogonality conditions (??) is related to the generalized null space of the linearized equation around Q , (??) and to the coercivity property (??). See the proof of Lemma ?? for a technical justification of these choices (see also [?]).

3. BACKWARDS UNIFORM ESTIMATES

In this section, we prove uniform estimates on particular backwards solutions. The key point is to carefully adjust their final data to obtain uniform estimates corresponding to the special regime of Theorem ?? and Lemma ??.

Let $(\lambda^{in}, z^{in}, b^{in}) \in (0, +\infty)^2 \times \mathbb{R}$ to be chosen with $\lambda^{in} \ll 1$, $z^{in} \gg 1$, $|b^{in}| \ll 1$. Let $u(t)$ for $t \leq 0$ be the solution of (??) with data (see (??))

$$u(0, x) = \frac{1}{\lambda^{in}} \mathbf{P}^{in} \left(\frac{x}{\lambda^{in}} \right) \quad \text{where} \quad \mathbf{P}^{in}(y) = \mathbf{P}(y; (z^{in}, b^{in}, 0)) \quad (3.1)$$

(we arbitrarily fix $\gamma^{in} = \beta^{in} = 0$). Note that $u(0)$ satisfies (??) and, by continuity of the solution of (??) in H^1 , it exists and satisfies (??) on some maximal time interval $(t^{mod}, 0]$, where $t^{mod} \in [-\infty, 0)$. Note also that by invariance by rotation of equation (??), $u(t)$ is invariant by the rotation τ_K . On $(t^{mod}, 0]$, we consider $(\vec{p}, \varepsilon)$ the decomposition of u defined from Lemma ?. For $s^{in} \gg 1$, we normalize the rescaled time s as follows, for $t \in (t^{mod}, 0]$,

$$s = s(t) = s^{in} - \int_t^0 \frac{d\tau}{\lambda^2(\tau)}. \quad (3.2)$$

Observe from (??) that

$$\begin{aligned} \lambda(s^{in}) &= \lambda^{in}, & b(s^{in}) &= b^{in}, & z(s^{in}) &= z^{in}, \\ \gamma(s^{in}) &= 0, & \beta(s^{in}) &= 0, & \varepsilon(s^{in}) &\equiv 0. \end{aligned} \quad (3.3)$$

Proposition 6 (Uniform backwards estimates). *There exists $s_0 > 10$ such that for all $s^{in} > s_0$, there exists a choice of parameters $(\lambda^{in}, z^{in}, b^{in})$ with*

$$\begin{aligned} \left| \left(\frac{2}{\kappa c_a} \right)^{\frac{1}{2}} (z^{in})^{-\frac{3}{4}} e^{\frac{\kappa}{2} z^{in}} - s^{in} \right| &< s^{in} \log^{-\frac{1}{2}}(s^{in}), \\ \lambda^{in} &= \log^{-1}(s^{in}), \quad b^{in} = \left(\frac{2c_a}{\kappa} \right)^{\frac{1}{2}} (z^{in})^{-\frac{1}{4}} e^{-\frac{\kappa}{2} z^{in}}, \end{aligned} \quad (3.4)$$

such that the solution u of (??) corresponding to (??) exists and satisfies (??) on the rescaled interval of time $[s_0, s^{in}]$, the rescaled time s being defined in (??). Moreover, the decomposition of u given by Lemma ?? on $[s_0, s^{in}]$

$$u(s, x) = \frac{e^{i\gamma(s)}}{\lambda(s)} (\mathbf{P} + \varepsilon)(s, y), \quad y = \frac{x}{\lambda(s)},$$

satisfies the following uniform estimates, for all $s \in [s_0, s^{in}]$,

$$\begin{aligned} \left| z(s) - \frac{2}{\kappa} \log(s) \right| &\lesssim \log(\log(s)), \quad |\lambda(s) - \log^{-1}(s)| \lesssim \log^{-\frac{3}{2}}(s), \\ |b(s) - s^{-1} \log^{-1}(s)| + |\beta(s)| + \|\varepsilon(s)\|_{H^1} &\lesssim s^{-1} \log^{-\frac{3}{2}}(s), \quad |a(s)| \lesssim s^{-2} \log^{-1}(s). \end{aligned} \quad (3.5)$$

The key point in Proposition ?? is that s_0 and the constants in (??) are independent of s^{in} as $s^{in} \rightarrow +\infty$. Observe that estimates (??) match the discussion of Sect. ??.

The rest of this section is devoted to the proof of Proposition ?. The proof relies on a bootstrap argument, integration of the differential system of geometrical parameters and energy estimates. We estimate ε by standard energy arguments in the framework of multi-bubble solutions. The particular regime of the geometrical parameters does not create any further difficulty. On the contrary, the special behavior $b(s) \sim s^{-1} \log^{-1}(s)$ simplifies this part of the proof (see step 2 of the proof of Proposition ??). We control the geometrical parameters of the bubbles in the bootstrap regime adjusting the final data $(\lambda^{in}, z^{in}, b^{in})$.

3.1. Bootstrap bounds. The proof of Proposition ?? follows from bootstrapping the following estimates, chosen in view of the formal computations in Sect. ??,

$$\begin{aligned} \left| \left(\frac{2}{\kappa c_a} \right)^{\frac{1}{2}} z^{-\frac{3}{4}} e^{\frac{\kappa}{2} z} - s \right| &\leq s \log^{-\frac{1}{2}}(s), \\ \frac{1}{2} s^{-1} \log^{-1}(s) &\leq b(s) \leq 2s^{-1} \log^{-1}(s), \\ |\beta(s)| &\leq s^{-1} \log^{-\frac{3}{2}}(s), \quad \|\varepsilon(s)\|_{H^1} \leq s^{-1} \log^{-\frac{3}{2}}(s). \end{aligned} \quad (3.6)$$

Note that the estimate on z in (??) immediately implies that, for s large

$$e^{-\kappa z} \lesssim s^{-2} \log^{-\frac{3}{2}}(s), \quad \left| z(s) - \frac{2}{\kappa} \log(s) \right| \lesssim \log(\log(s)), \quad |a(s)| \lesssim s^{-2} \log^{-1}(s). \quad (3.7)$$

For $s_0 > 10$ to be chosen large enough (independent of s^{in}), and all $s^{in} \gg s_0$, we define

$$s^* = \inf\{\tau \in [s_0, s^{in}]; \text{ (??) holds on } [\tau, s^{in}]\}. \quad (3.8)$$

3.2. Control of the modulation equations. We claim the following bounds on the modulation system $\vec{m}_1^a$ and on the error $\mathcal{E}_{\mathbf{P}}$ given by (??), (??)–(??) in the bootstrap regime (??).

Lemma 7 (Pointwise control of the modulation equations and the error). *The following estimates hold on $[s^*, s^{in}]$.*

$$|\vec{m}_1^a(s)| \lesssim s^{-2} \log^{-2}(s). \quad (3.9)$$

$$|\langle \eta_1(s), Q \rangle| \lesssim s^{-2} \log^{-2}(s), \quad (3.10)$$

$$|\dot{z} - bz| \lesssim s^{-1} \log^{-1}(s), \quad |\dot{\beta}| + |\dot{b} - a| \lesssim s^{-2} \log^{-2}(s). \quad (3.11)$$

Moreover, for all $s \in [s^*, s^{in}]$, for all $y \in \mathbb{R}^2$,

$$|\mathcal{E}_{\mathbf{P}}(s, y)| + |\nabla \mathcal{E}_{\mathbf{P}}(s, y)| \lesssim s^{-2} \log^{-2}(s) \sum_k Q^{1/2}(y - z_k(s)). \quad (3.12)$$

Proof of Lemma ??. The proofs of the first two estimates are to be combined. Since $\varepsilon(s^{in}) \equiv 0$, we may define

$$s^{**} = \inf\{s \in [s^*, s^{in}]; |\langle \eta_1(\tau), Q \rangle| \leq C^{**} \tau^{-2} \log^{-2}(\tau) \text{ holds on } [s, s^{in}]\},$$

for some constant $C^{**} > 0$ to be chosen large enough. We work on the interval $[s^{**}, s^{in}]$.

step 1 Equation of ε and change of variable. Let $v = \mathbf{P} + \varepsilon$ in (??). It follows from (??), (??) that

$$i\dot{\varepsilon} + \Delta\varepsilon - \varepsilon + (|\mathbf{P} + \varepsilon|^2(\mathbf{P} + \varepsilon) - |\mathbf{P}|^2\mathbf{P}) - i\frac{\dot{\lambda}}{\lambda}\Lambda\varepsilon + (1 - \dot{\gamma})\varepsilon + \mathcal{E}_{\mathbf{P}} = 0. \quad (3.13)$$

By rotation symmetry (see Lemma ??) it is enough to understand the orthogonality for η_1 . Thus, we change the space variable to match the one of the bubble P_1 . Recall that we have defined $\varepsilon(s, y) = [e^{i\Gamma_1}\eta_1](s, y - z_1)$. Denote

$$\mathbf{P}(s, y) = [e^{i\Gamma_1}\mathbf{P}_1](s, y - z_1), \quad \mathcal{E}_{\mathbf{P}}(s, y) = [e^{i\Gamma_1}\mathcal{E}_{\mathbf{P}_1}](s, y - z_1).$$

We rewrite the equation of ε into the following equation for η_1 (see also step 1 of the proof of Lemma ??)

$$i\dot{\eta}_1 + \Delta\eta_1 - \eta_1 + (|\mathbf{P}_1 + \eta_1|^2(\mathbf{P}_1 + \eta_1) - |\mathbf{P}_1|^2\mathbf{P}_1) + \vec{m}_1^0 \cdot \vec{M}\eta_1 + \mathcal{E}_{\mathbf{P}_1} = 0, \quad (3.14)$$

Recall also that η_1 satisfies the orthogonality conditions (??).

step 2 General null space like computation. Let $A(y)$ and $B(y)$ be two real-valued functions in $\mathcal{Y}$. We claim the following estimate on $[s^{**}, s^{in}]$

$$\left| \frac{d}{ds} \langle \eta_1, A + iB \rangle - \left[\langle \eta_1, iL_-A - L_+B \rangle - \langle \vec{m}_1^a \cdot \vec{M}Q, iA - B \rangle \right] \right| \lesssim s^{-2} \log^{-2}(s) + s^{-1} |\vec{m}_1^a|. \quad (3.15)$$

We compute from (??),

$$\begin{aligned} \frac{d}{ds} \langle \eta_1, A + iB \rangle &= \langle \dot{\eta}_1, A + iB \rangle = \langle i\dot{\eta}_1, iA - B \rangle \\ &= \langle -\Delta\eta_1 + \eta_1 - (2Q^2\eta_1 + Q\bar{\eta}_1), iA - B \rangle \\ &= \langle |\mathbf{P}_1 + \eta_1|^2(\mathbf{P}_1 + \eta_1) - |\mathbf{P}_1|^2\mathbf{P}_1 - 2Q^2\eta_1 - Q^2\bar{\eta}_1, iA - B \rangle \\ &= \langle \vec{m}_1^0 \cdot \vec{M}\eta_1, iA - B \rangle - \langle \mathcal{E}_{\mathbf{P}_1}, iA - B \rangle. \end{aligned}$$

First, since A and B are real-valued, we have

$$\langle -\Delta\eta_1 + \eta_1 - (2Q^2\eta_1 + Q\bar{\eta}_1), iA - B \rangle = \langle \eta_1, iL_-A - L_+B \rangle.$$

Second, note that

$$\begin{aligned} & |\mathbf{P}_1 + \eta_1|^2(\mathbf{P}_1 + \eta_1) - |\mathbf{P}_1|^2\mathbf{P}_1 - 2Q^2\eta_1 - Q^2\bar{\eta}_1 \\ &= 2(|\mathbf{P}_1|^2 - Q^2)\eta_1 + (\mathbf{P}_1^2 - Q^2)\bar{\eta}_1 + 2\mathbf{P}_1|\eta_1|^2 + \bar{\mathbf{P}}_1\eta_1^2 + 2\mathbf{P}_1|\eta_1|^2 + |\eta_1|^2\eta_1, \end{aligned}$$

and recall the expression of $\mathbf{P}_1$

$$\mathbf{P}_1 = Q + a\rho + \sum_{k=2}^K e^{i(\Gamma_k(y-(z_k-z_1))-\Gamma_1(y))} Q_a(y - (z_k - z_1)).$$

Therefore, using $A, B \in \mathcal{Y}$, (??)–(??) and $|z_k - z_j| \geq \kappa z$, for $k \neq j$, we have, for some $q > 0$,

$$\begin{aligned} & |\langle (|\mathbf{P}_1|^2 - Q^2)\eta_1, iA - B \rangle| + |\langle (\mathbf{P}_1^2 - Q^2)\bar{\eta}_1, iA - B \rangle| \\ & \lesssim (|a| + z^q e^{-\kappa z}) \|\eta_1\|_{L^2} \lesssim s^{-3} \log^q(s). \end{aligned}$$

Next,

$$\begin{aligned} & |\langle 2\mathbf{P}_1|\eta_1|^2 + \bar{\mathbf{P}}_1\eta_1^2 + 2\mathbf{P}_1|\eta_1|^2, iA - B \rangle| \lesssim \|\varepsilon\|_{L^2}^2 \lesssim s^{-2} \log^{-3}(s), \\ & |\langle |\eta_1|^2\eta_1, iA - B \rangle| \lesssim \langle |\varepsilon|^3, |A| + |B| \rangle \lesssim \|\varepsilon\|_{H^1}^3 \lesssim s^{-3} \log^{-\frac{9}{2}}(s). \end{aligned}$$

Third, we have, using (??)–(??), integration by parts,

$$\begin{aligned} & |\langle \vec{m}_1^0 \cdot \vec{M}\eta_1, iA - B \rangle| \lesssim \left| \langle \vec{m}_1^a \cdot \vec{M}\eta_1, iA - B \rangle \right| + \left| \langle (\vec{m}_1^a - \vec{m}_1^0) \cdot \vec{M}\eta_1, iA - B \rangle \right| \\ & \lesssim s^{-1} \log^{-\frac{3}{2}}(s) |\vec{m}_1^a| + s^{-3} \log^{-\frac{5}{2}}(s). \end{aligned}$$

Finally, we claim the following estimate, which is enough to complete the proof of (??).

$$\left| \langle \mathcal{E}_{\mathbf{P}_1}, iA - B \rangle - \langle \vec{m}_1^a \cdot \vec{M}Q, iA - B \rangle \right| \lesssim s^{-2} \log^{-2}(s) + s^{-1} |\vec{m}_1^a|. \quad (3.16)$$

Indeed, recall the expression of $\mathcal{E}_{\mathbf{P}_1}$ (from (??)–(??))

$$\begin{aligned} \mathcal{E}_{\mathbf{P}_1} &= \Psi_1 + \sum_{k=2}^K e^{i(\Gamma_k(y-(z_k-z_1))-\Gamma_1(y))} \Psi_k(y - (z_k - z_1)). \\ &= \vec{m}_1^a \cdot \vec{M}Q_a + i\dot{z}a'(z)\rho + G_1 + \Psi_{Q_a} + \sum_{k=2}^K e^{i(\Gamma_k(y-(z_k-z_1))-\Gamma_1(y))} \Psi_k(y - (z_k - z_1)). \end{aligned}$$

First, since $Q_a = Q + a\rho$, by (??)–(??),

$$\left| \langle \vec{m}_1^a \cdot \vec{M}(Q_a - Q), iA - B \rangle \right| \lesssim |a| |\vec{m}_1^a| \lesssim s^{-2} \log^{-1}(s) |\vec{m}_1^a|.$$

Second, from (??)–(??),

$$\begin{aligned} & |\langle \dot{z}a'(z)\rho, iA - B \rangle| \lesssim |a'(z)| |\dot{z}| \lesssim s^{-2} \log^{-1}(s) (|\vec{m}_1^a| + |\beta| + \left| \frac{\dot{\lambda}}{\lambda} \right| z) \\ & \lesssim s^{-2} \log^{-1}(s) ((z+1) |\vec{m}_1^a| + |\beta| + |b|z) \\ & \lesssim s^{-2} |\vec{m}_1^a| + s^{-3} \log^{-1}(s). \end{aligned} \quad (3.17)$$

Third, from (??) and (??)–(??),

$$|\langle G_1, iA - B \rangle| \lesssim \|G_1\|_{L^\infty} \lesssim z^{-\frac{1}{2}} e^{-\kappa z} \lesssim s^{-2} \log^{-2}(s).$$

Fourth, from (??) and (??)–(??),

$$|\langle \Psi_{Q_a}, iA - B \rangle| \lesssim \|\Psi_{Q_a}\|_{L^\infty} \lesssim |a|^2 \lesssim s^{-4} \log^{-2}(s).$$

Last, since $A, B \in \mathcal{Y}$, for $k \geq 2$, we have

$$|\langle e^{i(\Gamma_k(y-(z_k-z_1))-\Gamma_1(y))}(\vec{m}_k^a \cdot \vec{M}Q_a(\cdot - (z_k - z_1))), iA - B \rangle| \lesssim s^{-1}|\vec{m}_k^a|,$$

and, proceeding as before for the other terms in Ψ_k , we obtain

$$|\langle e^{i(\Gamma_k(y-(z_k-z_1))-\Gamma_1(y))}\Psi_k(y - (z_k - z_1)), iA - B \rangle| \lesssim s^{-1}|\vec{m}_k^a| + s^{-2}\log^{-2}(s).$$

The proof of (??) is complete.

step 3 Modulation equations. We now use (??) and (??) to control the modulation vector $\vec{m}_1^a$. Using (??), we draw one by one the consequences of the orthogonality relations (??). Note that the special orthogonality conditions (??), related to cancellations (see (??)) are crucial in these computations.

$\langle \eta_1, |y|^2 Q \rangle = 0$. Let $A = |y|^2 Q$ and $B = 0$. Since $L_-(|y|^2 Q) = -4\Lambda Q$, $\langle \eta_1, i\Lambda Q \rangle = 0$ and $\langle \vec{m}_1^a \cdot \vec{M}Q, i|y|^2 Q \rangle = -(b + \frac{\dot{\lambda}}{\lambda})\langle \Lambda Q, |y|^2 Q \rangle = c_1(b + \frac{\dot{\lambda}}{\lambda})$, where $c_1 \neq 0$, we obtain

$$\left| b + \frac{\dot{\lambda}}{\lambda} \right| \lesssim s^{-2}\log^{-2}(s) + s^{-1}|\vec{m}_1^a|. \quad (3.18)$$

$\langle \eta_1, yQ \rangle = 0$. Let $A = yQ$ and $B = 0$. Since $L_-(yQ) = -2\nabla Q$, $\langle \eta_1, i\nabla Q \rangle = 0$ and $\langle \vec{m}_1^a \cdot \vec{M}Q, iyQ \rangle = -(\dot{z} - 2\beta + \frac{\dot{\lambda}}{\lambda}z)\langle \nabla Q, yQ \rangle = c_2(\dot{z} - 2\beta + \frac{\dot{\lambda}}{\lambda}z)$, where $c_2 \neq 0$, we obtain

$$\left| \dot{z} - 2\beta + \frac{\dot{\lambda}}{\lambda}z \right| \lesssim s^{-2}\log^{-2}(s) + s^{-1}|\vec{m}_1^a|. \quad (3.19)$$

$\langle \eta_1, i\rho \rangle = 0$. Let $A = 0$ and $B = \rho$. Since $L_+\rho = |y|^2 Q$, $\langle \eta_1, |y|^2 Q \rangle = 0$ and

$$\langle \vec{m}_1^a \cdot \vec{M}Q, |y|^2 Q \rangle = c_3(\dot{\gamma} - 1 + \beta^2 - \frac{\dot{\lambda}}{\lambda}\beta z - \beta\dot{z}) + c_4(\dot{b} + b^2 - 2b(b + \frac{\dot{\lambda}}{\lambda}) - a),$$

where $c_3, c_4 \neq 0$, we obtain, for some c ,

$$\left| (\dot{\gamma} - 1 + \beta^2 - \frac{\dot{\lambda}}{\lambda}\beta z - \beta\dot{z}) + c(\dot{b} + b^2 - 2b(b + \frac{\dot{\lambda}}{\lambda}) - a) \right| \lesssim s^{-2}\log^{-2}(s) + s^{-1}|\vec{m}_1^a|. \quad (3.20)$$

$\langle \eta_1, i\nabla Q \rangle = 0$. Let $A = 0$ and $B = \nabla Q$. Since $L_+\nabla Q = 0$, and $\langle \vec{m}_1^a \cdot \vec{M}Q, \nabla Q \rangle = c_5(\dot{\beta} - \frac{\dot{\lambda}}{\lambda}\beta + \frac{b}{2}(\dot{z} - 2\beta + \frac{\dot{\lambda}}{\lambda}z))$, where $c_5 \neq 0$, we obtain

$$\left| \dot{\beta} - \frac{\dot{\lambda}}{\lambda}\beta + \frac{b}{2}(\dot{z} - 2\beta + \frac{\dot{\lambda}}{\lambda}z) \right| \lesssim s^{-2}\log^{-2}(s) + s^{-1}|\vec{m}_1^a|. \quad (3.21)$$

$\langle \eta_1, i\Lambda Q \rangle = 0$. Let $A = 0$ and $B = \Lambda Q$. Note that $L_+(\Lambda Q) = -2Q$, and by the definition of s^{**} , $|\langle \eta_1, Q \rangle| \lesssim C^{**}s^{-2}\log^{-2}(s)$. Moreover, $\langle \vec{m}_1^a \cdot \vec{M}Q, \nabla Q \rangle = c_6(\dot{b} + b^2 - 2b(b + \frac{\dot{\lambda}}{\lambda}) - a)$, where $c_6 \neq 0$, so that we obtain

$$\left| \dot{b} + b^2 - 2b(b + \frac{\dot{\lambda}}{\lambda}) - a \right| \lesssim C^{**}s^{-2}\log^{-2}(s) + s^{-1}|\vec{m}_1^a|. \quad (3.22)$$

Combining (??)–(??), we have proved, for all $s \in [s^{**}, s^{in}]$,

$$|\vec{m}_1^a(s)| \lesssim C^{**}s^{-2}\log^{-2}(s). \quad (3.23)$$

step 4 Minimal mass property of the approximate solution. The proof of the degeneracy estimate (??) relies on the following minimal mass property for the ansatz $\mathbf{P}$ under the bootstrap assumptions (??):

$$\|\mathbf{P}(s)\|_{L^2} - \|\mathbf{P}^{in}\|_{L^2} \lesssim s^{-2} \log^{-2}(s). \quad (3.24)$$

Note that the implicit constant on the right-hand side does not depend on C^{**} . By the definition (??) of $\mathcal{E}_{\mathbf{P}}$, we have

$$\frac{1}{2} \frac{d}{ds} \|\mathbf{P}\|_{L^2}^2 = \langle i\dot{\mathbf{P}}, i\mathbf{P} \rangle = \langle \mathcal{E}_{\mathbf{P}}, i\mathbf{P} \rangle.$$

In view of the formula for $\mathcal{E}_{\mathbf{P}}$ (??), and the definition of $\mathbf{P} = \sum_j P_j$, (??) follows by integration of the following estimate: for all $j, k \in \{1, \dots, K\}$,

$$|\langle [e^{i\Gamma_k} \Psi_k](y - z_k), i[e^{i\Gamma_j} Q_a](y - z_j) \rangle| \lesssim s^{-3} \log^{-2}(s). \quad (3.25)$$

Proof of (??). We start by proving (??) in the case $j = k = 1$. From (??):

$$\langle [e^{i\Gamma_k} \Psi_k](y - z_k), i[e^{i\Gamma_k} Q_a](y - z_k) \rangle = \langle \Psi_1, iQ_a \rangle = \langle \vec{m}_1^a \cdot \vec{M}Q_a + i\dot{z}a'(z)\rho + G_1 + \Psi_{Q_a}, iQ_a \rangle.$$

Note that $\langle \vec{M}Q, iQ \rangle = 0$. Thus, by (??), (??)–(??),

$$|\langle \vec{m}_1^a \cdot \vec{M}Q_a, iQ_a \rangle| \lesssim |a| |\vec{m}_1^a| \lesssim C^{**} s^{-4} \log^{-3}(s) \lesssim s^{-3} \log^{-3}(s).$$

Next, we claim the following estimate, which justifies the special choice of $a(z)$ done in (??) (see also Sect. ??)

$$|\langle i\dot{z}a'(z)\rho + G_1, iQ_a \rangle| \lesssim s^{-3} \log^{-2}(s). \quad (3.26)$$

Indeed, first by (??)–(??) and (??),

$$|\langle G_1, iQ_a \rangle + \kappa c_a \langle \rho, Q \rangle b z^{\frac{3}{2}} e^{-\kappa z}| \lesssim s^{-3} \log^{-2}(s). \quad (3.27)$$

Second, we note that by (??) and (??)–(??),

$$|\dot{z} - bz| \lesssim s^{-1} \log^{-1}(s), \quad (3.28)$$

and that by the definition of $a(z)$ in (??),

$$|a'(z) - c_a \kappa z^{\frac{1}{2}} e^{-\kappa z}| \lesssim z^{-\frac{1}{2}} e^{-\kappa z} \lesssim s^{-2} \log^{-2}(s). \quad (3.29)$$

Gathering (??)–(??), we obtain (??). Finally, since Q_a and Ψ_{Q_a} given by (??) are real-valued, we have the cancellation

$$\langle \Psi_{Q_a}, iQ_a \rangle = 0.$$

The collection of above estimates concludes the proof of (??) for $j = k = 1$.

We now prove (??) in the case $k = 1$ and $j \in \{2, \dots, K\}$. Note that

$$\begin{aligned} & |\langle [e^{i\Gamma_k} \Psi_k](y - z_k), i[e^{i\Gamma_j} Q_a](y - z_j) \rangle| \\ &= |\langle \vec{m}_1^a \cdot \vec{M}Q_a + i\dot{z}a'(z)\rho + G_1 + \Psi_{Q_a}, ie^{i(\Gamma_j(y-(z_j-z_k))-\Gamma_k(y))} Q_a(y-(z_j-z_k)) \rangle| \end{aligned}$$

First, by (??), for some $q > 0$,

$$\begin{aligned} & |\langle \vec{m}_1^a \cdot \vec{M}Q_a, ie^{i(\Gamma_j(y-(z_j-z_k))-\Gamma_k(y))} Q_a(y-(z_j-z_k)) \rangle| \\ & \lesssim |\vec{m}_1^a| z^q e^{-\kappa z} \lesssim C^{**} s^{-4} \log^q(s) \lesssim s^{-3} \log^{-2}(s). \end{aligned}$$

Second, using similar arguments, for some $q > 0$,

$$|\langle i\dot{z}a'(z)\rho + G_1 + \Psi_{Q_a}, ie^{i(\Gamma_j(y-(z_j-z_k))-\Gamma_k(y))} Q_a(y-(z_j-z_k)) \rangle| \lesssim s^{-4} \log^q(s).$$

The collection of above estimates concludes the proof of (??).

step 5 Proof of (??). The conservation of mass for the solution u and (??) imply:

$$\|u(s)\|_{L^2} = \|u(s^{in})\|_{L^2} = \|\mathbf{P}^{in}\|_{L^2}.$$

By (??),

$$\langle \varepsilon(s), \mathbf{P} \rangle = \frac{1}{2} (\|u(s)\|_{L^2}^2 - \|\mathbf{P}(s)\|_{L^2}^2 - \|\varepsilon(s)\|_{L^2}^2).$$

Therefore, using (??)–(??) and (??), we obtain

$$|\langle \varepsilon(s), \mathbf{P} \rangle| \lesssim s^{-2} \log^{-2}(s).$$

Now, we use the symmetry $\langle \varepsilon, P_k \rangle = \langle \varepsilon, P_j \rangle = K^{-1} \langle \varepsilon(s), \mathbf{P} \rangle$ for all $j, k \in \{1, \dots, K\}$. Moreover, by (??)–(??),

$$\langle \varepsilon(s), P_1 \rangle = \langle \eta_1, Q_a \rangle = \langle \eta_1, Q \rangle + O(|a| \|\varepsilon\|_{L^2}) = \langle \eta_1, Q \rangle + O(s^{-3} \log^{-\frac{5}{2}}(s)).$$

Gathering this information, we obtain $|\langle \eta_1, Q \rangle| \lesssim s^{-2} \log^{-2}(s)$, i.e. estimate (??). In particular, choosing C^{**} large enough, we have $s^{**} = s^*$.

step 6 Conclusion. The estimate (??) is a direct consequence of (??) and (??)–(??). We now turn to the proof of (??). Using (??), (??)–(??) and (??),

$$|\dot{z}a'(z)\rho| \lesssim Q^{\frac{1}{2}}(|b|z + s^{-1} \log^{-1}(s))s^{-2} \log^{-1}(s) \lesssim Q^{\frac{1}{2}}s^{-3} \log^{-1}(s).$$

By (??),

$$|\vec{m}_1^a \cdot \vec{M}Q_a| \lesssim Q^{\frac{1}{2}}s^{-2} \log^{-2}(s).$$

Next, by the definition of G_k in (??), the decay $|\rho| \lesssim Q^{\frac{7}{8}}$ (see (??)) and $|e_k - e_1| \geq \kappa$ for $k \neq 1$,

$$\begin{aligned} |G_1| &\lesssim Q^{\frac{1}{2}} \sum_{k=2}^K \left(Q^{\frac{5}{4}}(y)Q(y - z(e_k - e_1)) + |a|Q^{\frac{5}{4}}(y)Q^{\frac{7}{8}}(y - z(e_k - e_1)) \right) \\ &\lesssim Q^{\frac{1}{2}}(z^{-\frac{1}{2}}e^{-\kappa z} + s^{-2}e^{-\frac{\kappa}{2}z}) \lesssim Q^{\frac{1}{2}}s^{-2} \log^{-2}(s). \end{aligned}$$

Finally, by the definition of Ψ_{Q_a} in (??),

$$|\Psi_{Q_a}| \lesssim Q^{\frac{1}{2}}|a|^2 \lesssim Q^{\frac{1}{2}}s^{-4} \log^{-2}(s).$$

The same estimates hold for $\nabla \mathcal{E}$, which finishes the proof of (??). $\square$

3.3. Energy functional. Consider the nonlinear energy functional for ε

$$\mathbf{H}(s, \varepsilon) = \frac{1}{2} \int \left(|\nabla \varepsilon|^2 + |\varepsilon|^2 - \frac{1}{2} (|\mathbf{P} + \varepsilon|^4 - |\mathbf{P}|^4 - 4|\mathbf{P}|^2 \operatorname{Re}(\varepsilon \bar{\mathbf{P}})) \right).$$

Pick a smooth function $\chi : [0, +\infty) \rightarrow [0, \infty)$, non increasing, with $\chi \equiv 1$ on $[0, \frac{1}{10}]$, $\chi \equiv 0$ on $[\frac{1}{8}, +\infty)$. We define the localized momentum:

$$\mathbf{J} = \sum_k J_k, \quad J_k(s, \varepsilon) = b \operatorname{Im} \int (z_k \cdot \nabla \varepsilon) \bar{\varepsilon} \chi_k, \quad \chi_k(s, y) = \chi(\log^{-1}(s)|y - z_k(s)|).$$

Finally, set

$$\mathbf{F}(s, \varepsilon) = \mathbf{H}(s, \varepsilon) - \mathbf{J}(s, \varepsilon).$$

The functional F is coercive in ε at the main order and it is an almost conserved quantity for the problem.

Proposition 8 (Coercivity and time control of the energy functional). *For all $s \in [s^*, s^{in}]$,*

$$\mathbf{F}(s, \varepsilon(s)) \gtrsim \|\varepsilon(s)\|_{H^1}^2 + O(s^{-4} \log^{-4}(s)), \quad (3.30)$$

and

$$\left| \frac{d}{ds} [\mathbf{F}(s, \varepsilon(s))] \right| \lesssim s^{-2} \log^{-2}(s) \|\varepsilon(s)\|_{H^1} + s^{-1} \log^{-1}(s) \|\varepsilon(s)\|_{H^1}^2. \quad (3.31)$$

Proof of Proposition ??. **step 1** Coercivity. The proof of the coercivity (??) is a standard consequence of the coercivity property (??) around one solitary wave with the orthogonality properties (??), (??), and an elementary localization argument. Hence we briefly sketch the argument. First, using the coercivity property (??) and the orthogonality properties (??), (??) and localization arguments, we have

$$\mathbf{H}(s, \varepsilon) \gtrsim \|\varepsilon\|_{H^1}^2 + O(s^{-4} \log^{-4}(s)). \quad (3.32)$$

Note that the error term in $O(s^{-4} \log^{-4}(s))$ is due to the bound (??). We refer to the proof of Lemma 4.1 in Appendix B of [?] for a similar proof. Second, we note that by (??)–(??), $|\mathbf{J}(s, \varepsilon)| \lesssim |b|z \|\varepsilon\|_{H^1}^2 \lesssim s^{-1} \|\varepsilon\|_{H^1}^2$, and (??) follows.

step 2 Variation of the energy. We estimate the time variation of the functional $\mathbf{H}$ and claim: for all $s \in [s^*, s^{in}]$,

$$\left| \frac{d}{ds} [\mathbf{H}(s, \varepsilon(s))] - \sum_k \langle \dot{z}_k \cdot \nabla P_k, 2|\varepsilon|^2 P_k + \varepsilon^2 \bar{P}_k \rangle \right| \lesssim s^{-2} \log^{-2}(s) \|\varepsilon(s)\|_{H^1} + s^{-1} \log^{-1}(s) \|\varepsilon(s)\|_{H^1}^2. \quad (3.33)$$

The time derivative of $s \mapsto H(s, \varepsilon(s))$ splits into two parts

$$\frac{d}{ds} [\mathbf{H}(s, \varepsilon(s))] = D_s \mathbf{H}(s, \varepsilon(s)) + \langle D_\varepsilon \mathbf{H}(s, \varepsilon(s)), \dot{\varepsilon}_s \rangle,$$

where D_s denotes differentiation of $\mathbf{H}$ with respect to s and D_ε denotes differentiation of $\mathbf{H}$ with respect to ε . First compute:

$$D_s \mathbf{H} = -\langle \dot{\mathbf{P}}, |\mathbf{P} + \varepsilon|^2 (\mathbf{P} + \varepsilon) - |\mathbf{P}^2| \mathbf{P} - (2\varepsilon |\mathbf{P}|^2 + \bar{\varepsilon} \mathbf{P}^2) \rangle.$$

Observe that by the definition of P_k in (??),

$$\dot{P}_k = -\dot{z}_k \cdot \nabla P_k + i(\dot{\beta}_k \cdot (y - z_k) - \frac{\dot{b}}{4} |y - z_k|^2) P_k + \dot{z} a'(z) \rho_k \quad \text{where} \quad \rho_k = [e^{i\Gamma_k} \rho](y - z_k).$$

By (??), (??)–(??) and (??),

$$|\dot{\beta}_k| + |\dot{b}| + |\dot{z} a'(z)| \lesssim s^{-2} \log^{-2}(s).$$

Since

$$\int ||\mathbf{P} + \varepsilon|^2 (\mathbf{P} + \varepsilon) - |\mathbf{P}^2| \mathbf{P} - (2\varepsilon |\mathbf{P}|^2 + \bar{\varepsilon} \mathbf{P}^2)| \lesssim \|\varepsilon\|_{H^1}^2,$$

we obtain

$$\begin{aligned} & \left| \left\langle i(\dot{\beta}_k \cdot (y - z_k) - \frac{\dot{b}}{4} |y - z_k|^2) P_k + \dot{z} a'(z) \rho_k, |\mathbf{P} + \varepsilon|^2 (\mathbf{P} + \varepsilon) - |\mathbf{P}^2| \mathbf{P} - (2\varepsilon |\mathbf{P}|^2 + \bar{\varepsilon} \mathbf{P}^2) \right\rangle \right| \\ & \lesssim s^{-2} \log^{-2}(s) \|\varepsilon\|_{H^1}^2. \end{aligned}$$

Next, note that

$$|\mathbf{P} + \varepsilon|^2 (\mathbf{P} + \varepsilon) - |\mathbf{P}^2| \mathbf{P} - (2\varepsilon |\mathbf{P}|^2 + \bar{\varepsilon} \mathbf{P}^2) = 2|\varepsilon|^2 \mathbf{P} + \varepsilon^2 \bar{\mathbf{P}} + |\varepsilon|^2 \varepsilon.$$

By (??) and (??)–(??), $|\dot{z}| \lesssim s^{-1}$ and thus by (??)–(??),

$$|\langle \dot{z}_k \cdot \nabla P_k, |\varepsilon|^3 \rangle| \lesssim s^{-1} \|\varepsilon\|_{H^1}^3 \lesssim s^{-2} \log^{-\frac{3}{2}}(s) \|\varepsilon\|_{H^1}^2.$$

For $j \neq k$, since $e^{-\kappa z} \lesssim s^{-2}$ by (??)–(??) and the decay properties of P_k, P_j ,

$$|\langle \dot{z}_k \cdot \nabla P_k, 2|\varepsilon|^2 P_j + \varepsilon^2 \bar{P}_j \rangle| \lesssim |s|^{-3} \|\varepsilon\|_{H^1}^2.$$

Gathering these computations, we have obtained

$$D_s \mathbf{H}(s, \varepsilon) = \sum_k \langle \dot{z}_k \cdot \nabla P_k, 2|\varepsilon|^2 P_k + \varepsilon^2 \bar{P}_k \rangle + O(s^{-2} \|\varepsilon\|_{H^1}^2). \quad (3.34)$$

Second,

$$D_\varepsilon \mathbf{H}(s, \varepsilon) = -\Delta \varepsilon + \varepsilon - (|\mathbf{P} + \varepsilon|^2 (\mathbf{P} + \varepsilon) - |\mathbf{P}|^2 \mathbf{P}),$$

so that the equation (??) of ε rewrites

$$i\dot{\varepsilon} - D_\varepsilon \mathbf{H}(s, \varepsilon) - i \frac{\dot{\lambda}}{\lambda} \Lambda \varepsilon + (1 - \dot{\gamma}) \varepsilon + \mathcal{E}_{\mathbf{P}} = 0.$$

In particular,

$$\begin{aligned} \langle D_\varepsilon \mathbf{H}(s, \varepsilon), \dot{\varepsilon} \rangle &= \langle i D_\varepsilon \mathbf{H}(s, \varepsilon), i \dot{\varepsilon} \rangle \\ &= \frac{\dot{\lambda}}{\lambda} \langle D_\varepsilon \mathbf{H}(s, \varepsilon), \Lambda \varepsilon \rangle - (1 - \dot{\gamma}) \langle i D_\varepsilon \mathbf{H}(s, \varepsilon), \varepsilon \rangle - \langle i D_\varepsilon \mathbf{H}(s, \varepsilon), \mathcal{E}_{\mathbf{P}} \rangle. \end{aligned}$$

We recall that

$$\langle -\Delta \varepsilon, \Lambda \varepsilon \rangle = \|\nabla \varepsilon\|^2, \quad \langle \varepsilon, \Lambda \varepsilon \rangle = 0, \quad \langle |\varepsilon|^2 \varepsilon, \Lambda \varepsilon \rangle = \frac{1}{2} \int |\varepsilon|^4,$$

and thus, using also (??)–(??), (??), and the decay properties of Q ,

$$|\langle D_\varepsilon \mathbf{H}(s, \varepsilon), \Lambda \varepsilon \rangle| \lesssim \|\varepsilon\|_{H^1}^2 + \|\varepsilon\|_{H^1}^4 \lesssim \|\varepsilon\|_{H^1}^2.$$

In particular, from (??) and (??)–(??), we deduce

$$\left| \frac{\dot{\lambda}}{\lambda} \langle D_\varepsilon \mathbf{H}(s, \varepsilon), \Lambda \varepsilon \rangle \right| \lesssim s^{-1} \log^{-1}(s) \|\varepsilon\|_{H^1}^2.$$

Note that the estimate on b in (??)–(??) implies $|b| \lesssim s^{-1} \log^{-1}(s) \ll s^{-1}$ which avoids the use of virial localized identities (as in [?, ?]) to control the above term. By (??) and (??)–(??), we estimate

$$|(1 - \dot{\gamma}) \langle i D_\varepsilon \mathbf{H}(s, \varepsilon), \varepsilon \rangle| \lesssim s^{-2} \|\varepsilon\|_{H^1}^2.$$

Finally, integrating by parts, using (??) and (??)–(??), we have

$$|\langle i D_\varepsilon \mathbf{H}(s, \varepsilon), \mathcal{E}_{\mathbf{P}} \rangle| \lesssim \langle |\nabla \varepsilon|, |\nabla \mathcal{E}_{\mathbf{P}}| \rangle + \langle |\varepsilon| + |\varepsilon|^3, |\mathcal{E}_{\mathbf{P}}| \rangle \lesssim s^{-2} \log^{-2}(s) \|\varepsilon\|_{H^1}.$$

The collection of above estimates finishes the proof of (??).

step 3 Variation of the localized momentum. We now claim: for all $s \in [s^*, s^{in}]$,

$$\begin{aligned} \left| \frac{d}{ds} [\mathbf{J}(s, \varepsilon(s))] - b \sum_k \langle z_k \cdot \nabla P_k, 2|\varepsilon|^2 P_k + \varepsilon^2 \bar{P}_k \rangle \right| \\ \lesssim s^{-2} \log^{-2}(s) \|\varepsilon(s)\|_{H^1} + s^{-1} \log^{-1}(s) \|\varepsilon(s)\|_{H^1}^2. \end{aligned} \quad (3.35)$$

Indeed, we compute, for any k ,

$$\begin{aligned} \frac{d}{ds}[J_k(s, \varepsilon(s))] &= \dot{b} \operatorname{Im} \int (z_k \cdot \nabla \varepsilon) \bar{\varepsilon} \chi_k + b \operatorname{Im} \int (\dot{z}_k \cdot \nabla \varepsilon) \bar{\varepsilon} \chi_k + b \operatorname{Im} \int (z_k \cdot \nabla \varepsilon) \bar{\varepsilon} \dot{\chi}_k \\ &\quad + b \langle i \dot{\varepsilon}, z_k \cdot (2\chi_k \nabla \varepsilon + \varepsilon \nabla \chi_k) \rangle. \end{aligned}$$

By (??) and (??)–(??), we have

$$\left| \dot{b} \operatorname{Im} \int (z_k \cdot \nabla \varepsilon) \bar{\varepsilon} \chi_k \right| + \left| b \operatorname{Im} \int (\dot{z}_k \cdot \nabla \varepsilon) \bar{\varepsilon} \chi_k \right| \lesssim s^{-2} \|\varepsilon\|_{H^1}^2.$$

Note that by direct computations, (??) and (??)–(??),

$$|\dot{\chi}_k| \lesssim (s^{-1} \log^{-1}(s)|y - z_k| + |\dot{z}_k|) \log^{-1}(s) |\chi'(\log^{-1}(s)(y - z_k(s)))| \lesssim s^{-1} \log^{-1}(s)$$

and so, by (??)–(??),

$$\left| b \operatorname{Im} \int (z_k \cdot \nabla \varepsilon) \bar{\varepsilon} \dot{\chi}_k \right| \lesssim s^{-2} \log^{-2}(s) \|\varepsilon\|_{H^1}^2.$$

Now, we use the equation (??) of ε to estimate $b \langle i \dot{\varepsilon}, z_k \cdot (2\chi_k \nabla \varepsilon + \varepsilon \nabla \chi_k) \rangle$. By integration by parts, we check the following

$$\begin{aligned} \langle \Delta \varepsilon, 2(z_k \cdot \nabla \varepsilon) \chi_k \rangle &= \int |\nabla \varepsilon|^2 (z_k \cdot \nabla \chi_k) - 2 \langle (\nabla \varepsilon \cdot \nabla \chi_k), (z_k \cdot \nabla \varepsilon) \rangle, \\ \langle \Delta \varepsilon, \varepsilon (z_k \cdot \nabla \chi_k) \rangle &= - \int |\nabla \varepsilon|^2 (z_k \cdot \nabla \chi_k) + \frac{1}{2} \int |\varepsilon|^2 (z_k \cdot \nabla (\Delta \chi_k)). \end{aligned}$$

Thus,

$$\langle \Delta \varepsilon, z_k \cdot (2\chi_k \nabla \varepsilon + \varepsilon \nabla \chi_k) \rangle = -2 \langle (\nabla \varepsilon \cdot \nabla \chi_k), (z_k \cdot \nabla \varepsilon) \rangle + \frac{1}{2} \int |\varepsilon|^2 (z_k \cdot \nabla (\Delta \chi_k)).$$

By (??)–(??), $|b| \lesssim s^{-1} \log^{-1}(s)$ and $|z_k| \lesssim \log(s)$. Moreover, $|\nabla \chi_k| \lesssim \log^{-1}(s)$. Therefore,

$$|b \langle (\nabla \varepsilon \cdot \nabla \chi_k), (z_k \cdot \nabla \varepsilon) \rangle| \lesssim s^{-1} \log^{-1}(s) \|\varepsilon\|_{H^1}^2.$$

Similarly, by $|\nabla (\Delta \chi_k)| \lesssim \log^{-3}(s)$, we obtain

$$\left| b \int |\varepsilon|^2 (z_k \cdot \nabla (\Delta \chi_k)) \right| \lesssim s^{-1} \log^{-3}(s) \|\varepsilon\|_{H^1}^2.$$

In conclusion for term $\Delta \varepsilon$ in the equation of ε , we obtain

$$|b \langle \Delta \varepsilon, z_k \cdot (2\chi_k \nabla \varepsilon + \varepsilon \nabla \chi_k) \rangle| \lesssim s^{-1} \log^{-1}(s) \|\varepsilon\|_{H^1}^2.$$

For the mass term in the equation of ε , we simply check by integration by parts that

$$\langle \varepsilon, z_k \cdot (2\chi_k \nabla \varepsilon + \varepsilon \nabla \chi_k) \rangle = 0.$$

We also check that

$$\langle i \Lambda \varepsilon, z_k \cdot (2\chi_k \nabla \varepsilon + \varepsilon \nabla \chi_k) \rangle = 2 \langle i \varepsilon, (z_k \cdot \nabla \varepsilon) \chi_k \rangle + \langle i (y \cdot \nabla \varepsilon), \varepsilon (z_k \cdot \nabla \chi_k) \rangle,$$

and thus, by (??)–(??),

$$|b| \left| \frac{\dot{\lambda}}{\lambda} \right| |\langle i \Lambda \varepsilon, z_k \cdot (2\chi_k \nabla \varepsilon + \varepsilon \nabla \chi_k) \rangle| \lesssim s^{-2} \log^{-1}(s) \|\varepsilon\|_{H^1}^2.$$

Next, from (??),

$$|b \langle \mathcal{E}_{\mathbf{P}}, z_k \cdot (2\chi_k \nabla \varepsilon + \varepsilon \nabla \chi_k) \rangle| \lesssim s^{-3} \log^{-2}(s) \|\varepsilon\|_{H^1}.$$

Now, we only have to deal with the term

$$b\langle |\mathbf{P} + \varepsilon|^2(\mathbf{P} + \varepsilon) - |\mathbf{P}|^2\mathbf{P}, z_k \cdot (2\chi_k \nabla \varepsilon + \varepsilon \nabla \chi_k) \rangle.$$

Recall that $|\mathbf{P} + \varepsilon|^2(\mathbf{P} + \varepsilon) - |\mathbf{P}|^2\mathbf{P} = (2\varepsilon|\mathbf{P}|^2 + \bar{\varepsilon}\mathbf{P}^2) + 2|\varepsilon|^2\mathbf{P} + \varepsilon^2\bar{\mathbf{P}} + |\varepsilon|^2\varepsilon$. First, by (??)–(??), it is clear that

$$|b\langle 2|\varepsilon|^2\mathbf{P} + \varepsilon^2\bar{\mathbf{P}} + |\varepsilon|^2\varepsilon, z_k \cdot (2\chi_k \nabla \varepsilon + \varepsilon \nabla \chi_k) \rangle| \lesssim s^{-1} \|\varepsilon\|_{H^1}^3 \lesssim s^{-2} \log^{-\frac{3}{2}}(s) \|\varepsilon\|_{H^1}^2.$$

Second, since $|b| \lesssim s^{-1} \log^{-1}(s)$, $|z_k| \lesssim \log(s)$ and $|\nabla \chi_k| \lesssim \log^{-1}(s)$,

$$|b\langle 2\varepsilon|\mathbf{P}|^2 + \bar{\varepsilon}\mathbf{P}^2, \varepsilon(z_k \cdot \nabla \chi_k) \rangle| \lesssim s^{-1} \log^{-1}(s) \|\varepsilon\|_{H^1}^2.$$

Third, by the decay property of Q and the definition of χ_k ,

$$|b\langle 2\varepsilon\left(|\mathbf{P}|^2 - \sum_j |P_j|^2\right) + \bar{\varepsilon}\left(\mathbf{P}^2 - \sum_j P_j^2\right), (z_k \cdot \nabla \varepsilon)\chi_k \rangle| \lesssim s^{-2} \|\varepsilon\|_{H^1}^2,$$

and, for $j \neq k$,

$$|b\langle 2\varepsilon|P_j|^2 + \bar{\varepsilon}P_j^2, (z_k \cdot \nabla \varepsilon)\chi_k \rangle| \lesssim s^{-2} \|\varepsilon\|_{H^1}^2.$$

Finally, we compute by integration by parts,

$$\begin{aligned} \langle 2\varepsilon|P_k|^2 + \bar{\varepsilon}P_k^2, (z_k \cdot \nabla \varepsilon)\chi_k \rangle &= -\langle z_k \cdot \nabla P_k, 2|\varepsilon|^2P_k + \varepsilon^2\bar{P}_k \rangle \\ &\quad - \frac{1}{2} \operatorname{Re} \left(\int (2|\varepsilon|^2P_k^2 + \varepsilon^2|P_k|^2) (z_k \cdot \nabla \chi_k) \right). \end{aligned}$$

As before,

$$\left| b \operatorname{Re} \left(\int (2|\varepsilon|^2P_k^2 + \varepsilon^2|P_k|^2) (z_k \cdot \nabla \chi_k) \right) \right| \lesssim s^{-1} \log^{-1}(s) \|\varepsilon\|_{H^1}^2.$$

The collection of above bounds concludes the proof of (??).

step 3 Conclusion. Recall that, by (??), $|\dot{z}_k - bz_k| \lesssim s^{-1} \log^{-1}(s)$, and so

$$|\langle (\dot{z}_k - bz_k) \cdot \nabla P_k, 2|\varepsilon|^2P_k + \varepsilon^2\bar{P}_k \rangle| \lesssim s^{-1} \log^{-1}(s) \|\varepsilon\|_{H^1}^2,$$

and (??) now follows from (??), (??). This concludes the proof of Proposition ?? $\square$

3.4. End of the proof of Proposition ??. We close the bootstrap estimates (??) and prove (??).

step 1 Closing the estimates in ε . By (??) in Proposition ?? and then (??)–(??), we have

$$\left| \frac{d}{ds} [\mathbf{F}(s, \varepsilon(s))] \right| \lesssim s^{-2} \log^{-2}(s) \|\varepsilon\|_{H^1} + s^{-1} \log^{-1}(s) \|\varepsilon\|_{H^1}^2 \lesssim s^{-3} \log^{-\frac{7}{2}}(s).$$

Thus, by integration on $[s, s^{in}]$ for any $s \in [s^*, s^{in}]$, using $\varepsilon(s^{in}) = 0$ (see (??)), we obtain

$$|\mathbf{F}(s, \varepsilon(s))| \lesssim s^{-2} \log^{-\frac{7}{2}}(s).$$

By (??) in Proposition ??, we obtain

$$\|\varepsilon(s)\|_{H^1}^2 \lesssim s^{-2} \log^{-\frac{7}{2}}(s).$$

Therefore, for s_0 large enough, for all $s \in [s^*, s^{in}]$,

$$\|\varepsilon(s)\|_{H^1}^2 \leq \frac{1}{2} s^{-2} \log^{-3}(s),$$

which strictly improves the estimate on $\|\varepsilon\|_{H^1}^2$ in (??).

step 2 Closing the parameter estimates. First, note that from (??), $|\dot{\beta}| \lesssim s^{-2} \log^{-2}(s)$. Together with the choice $\beta(s^{in}) = \beta^{in} = 0$ (see (??)), direct integration in time gives, for all $s \in [s^*, s^{in}]$, $|\beta(s)| \lesssim s^{-1} \log^{-2}(s)$. For s_0 large enough, we obtain, for all $s \in [s^*, s^{in}]$,

$$|\beta(s)| < \frac{1}{2} s^{-1} \log^{-\frac{3}{2}}(s),$$

which strictly improves the estimate on $\beta(s)$ in (??).

Second, recall from (??), (??) and the definition of $a(z)$ in (??), for all $s \in [s^*, s^{in}]$,

$$\left| \dot{b} + c_a z^{\frac{1}{2}} e^{-\kappa z} \right| \lesssim s^{-2} \log^{-2}(s), \quad |\dot{z} z^{-1} - b| \lesssim s^{-1} \log^{-2}(s).$$

Since $|\dot{b}| \lesssim s^{-2} \log^{-1}(s)$ and $|\dot{z} z^{-1}| \lesssim s^{-1} \log^{-1}(s)$, it follows that

$$\left| \dot{b} b + c_a \dot{z} z^{-\frac{1}{2}} e^{-\kappa z} \right| \lesssim s^{-3} \log^{-3}(s).$$

Integrating on $[s, s^{in}]$ for any $s \in [s^*, s^{in})$, using the special relation between b^{in} and z^{in} fixed in (??)

$$b^2(s^{in}) = \frac{2c_a}{\kappa} z^{-\frac{1}{2}}(s^{in}) e^{-\kappa z(s^{in})}, \quad b(s^{in}) > 0,$$

we obtain

$$\left| b^2 - \frac{2c_a}{\kappa} z^{-\frac{1}{2}} e^{-\kappa z} \right| \lesssim s^{-2} \log^{-3}(s) + \int_s^{s^{in}} \left| \dot{z} z^{-\frac{3}{2}} e^{-\kappa z} \right| ds' \lesssim s^{-2} \log^{-3}(s), \quad b(s) > 0. \quad (3.36)$$

From (??)–(??) and (??), we have

$$\left| \frac{2c_a}{\kappa} z^{-\frac{1}{2}} e^{-\kappa z} - s^{-2} \log^{-2}(s) \right| \lesssim s^{-2} \log^{-\frac{5}{2}}(s).$$

Therefore, the following estimate on $b(s)$ follows from (??)

$$|b^2 - s^{-2} \log^{-2}(s)| \lesssim s^{-2} \log^{-\frac{5}{2}}(s).$$

This implies, for all $s \in [s^*, s^{in}]$,

$$|b - s^{-1} \log^{-1}(s)| \lesssim s^{-1} \log^{-\frac{3}{2}}(s), \quad (3.37)$$

which strictly improves the estimate on $b(s)$ in (??).

Finally, we address the estimate on $z(s)$. From (??), (??)–(??) and (??), we have

$$\left| b - \left(\frac{2c_a}{\kappa} \right)^{\frac{1}{2}} z^{-\frac{1}{4}} e^{-\frac{\kappa}{2} z} \right| + \left| \dot{z} z^{-1} - \left(\frac{2c_a}{\kappa} \right)^{\frac{1}{2}} z^{-\frac{1}{4}} e^{-\frac{\kappa}{2} z} \right| \lesssim s^{-1} \log^{-2}(s). \quad (3.38)$$

Using $z \lesssim \log^{-1}(s)$, we obtain

$$\left| \frac{d}{ds} \left(z^{-\frac{3}{4}} e^{\frac{\kappa}{2} z} \right) - \left(\frac{\kappa c_a}{2} \right)^{\frac{1}{2}} \right| \lesssim \log^{-1}(s) + \left| \dot{z} z^{-\frac{7}{4}} e^{\frac{\kappa}{2} z} \right| \lesssim \log^{-1}(s). \quad (3.39)$$

We need to adjust the initial choice of $z(s^{in}) = z^{in}$ through a topological argument (see [?] for a similar argument). We define ζ and ξ the following two functions on $[s^*, s^{in}]$

$$\zeta(s) = \left(\frac{2}{\kappa c_a} \right)^{\frac{1}{2}} z^{-\frac{3}{4}} e^{\frac{\kappa}{2} z}, \quad \xi(s) = (\zeta(s) - s)^2 s^{-2} \log(s). \quad (3.40)$$

Then, (??) writes

$$|\dot{\zeta}(s) - 1| \lesssim \log^{-1}(s). \quad (3.41)$$

According to (??), our objective is to prove that there exists a suitable choice of

$$\zeta(s^{in}) = \zeta^{in} \in [s^{in} - s^{in} \log^{-\frac{1}{2}}(s^{in}), s^{in} + s^{in} \log^{-\frac{1}{2}}(s^{in})],$$

so that $s^* = s_0$. Assume for the sake of contradiction that for all $\zeta^\sharp \in [-1, 1]$, the choice

$$\zeta^{in} = s^{in} + \zeta^\sharp s^{in} \log^{-\frac{1}{2}}(s^{in})$$

leads to $s^* = s^*(\zeta^\sharp) \in (s_0, s^{in})$. Since all estimates in (??) except the one on $z(s)$ have been strictly improved on $[s^*, s^{in}]$, it follows from $s^*(\zeta^\sharp) \in (s_0, s^{in})$ and continuity that

$$|\zeta(s^*(\zeta^\sharp)) - s^*| = s^* \log^{-\frac{1}{2}} s^* \quad \text{i.e.} \quad \zeta(s^*(\zeta^\sharp)) = s^* \pm s^* \log^{-\frac{1}{2}} s^*.$$

We need a transversality condition to reach a contradiction. We compute:

$$\dot{\xi}(s) = 2(\zeta(s) - s)(\dot{\zeta}(s) - 1)s^{-2} \log(s) - (\zeta(s) - s)^2(2s^{-3} \log(s) - s^{-3}).$$

At $s = s^*$, this gives

$$|\dot{\xi}(s^*) + 2(s^*)^{-1}| \lesssim (s^*)^{-1} \log^{-\frac{1}{2}}(s^*).$$

Thus, for s_0 large enough,

$$\dot{\xi}(s^*) < -(s^*)^{-1}. \quad (3.42)$$

Define the function Φ by

$$\Phi : \zeta^\sharp \in [-1, 1] \mapsto (\zeta(s^*) - s^*)(s^*)^{-1} \log^{\frac{1}{2}}(s^*) \in \{-1, 1\}.$$

A standard consequence of the transversality property (??) is the continuity of the function $\zeta^\sharp \in [-1, 1] \mapsto s^*(\zeta^\sharp)$. In particular, the function Φ is also continuous from $[-1, 1]$ to $\{1, -1\}$. Moreover, for $\zeta^\sharp = -1$ and $\zeta^\sharp = 1$, $\xi(s^*) = 1$ and $\dot{\xi}(s^*) < 0$ by (??) and so in these cases $s^* = s^{in}$. Thus, $\Phi(-1) = -1$ and $\Phi(1) = 1$, but this is in contradiction with the continuity.

In conclusion, there exists at least a choice of

$$\zeta(s^{in}) = \zeta^{in} \in (s^{in} - s^{in} \log^{-\frac{1}{2}}(s^{in}), s^{in} + s^{in} \log^{-\frac{1}{2}}(s^{in}))$$

such that $s^* = s_0$.

step 3 Conclusion. To finish proving (??), we only have to prove the estimate on $\lambda(s)$. From (??) and (??), we obtain

$$\left| \frac{\dot{\lambda}}{\lambda} + s^{-1} \log^{-1}(s) \right| \lesssim s^{-1} \log^{-\frac{3}{2}}(s).$$

By integration on $[s, s^{in}]$, for any $s \in [s_0, s^{in}]$, using the value $\lambda(s^{in}) = \lambda^{in} = \log^{-1}(s^{in})$ (see (??)), we have

$$|\log(\lambda(s)) + \log(\log(s))| \lesssim \log^{-\frac{1}{2}}(s),$$

and thus

$$|\lambda(s) - \log^{-1}(s)| \lesssim \log^{-\frac{3}{2}}(s). \quad (3.43)$$

This concludes the proof of Proposition ??.

4. COMPACTNESS ARGUMENTS

The objective of this section is to finish the construction of Theorem ?? by passing to the limit on a sequence of solutions given by Proposition ??.

4.1. Construction of a sequence of backwards solutions. We claim the following consequence of Proposition ??.

Lemma 9. *There exist $t_0 > 1$ and a sequence of solutions $u_n \in \mathcal{C}([t_0 - T_n, 0], \Sigma)$ of (??), where*

$$T_n \rightarrow +\infty \quad \text{as } n \rightarrow +\infty, \quad (4.1)$$

satisfying the following estimates, for all $t \in [t_0 - T_n, 0]$,

$$\begin{aligned} \left| z_n(t) - \frac{2}{\kappa} \log(t + T_n) \right| &\lesssim \log(\log(t + T_n)), \quad |\lambda_n(t) - \log^{-1}(t + T_n)| \lesssim \log^{-\frac{3}{2}}(t + T_n), \\ |b_n(t) - (t + T_n)^{-1} \log^{-3}(t + T_n)| + |\beta_n(t)| + \|\varepsilon_n(t)\|_{H^1} &\lesssim (t + T_n)^{-1} \log^{-\frac{7}{2}}(t + T_n), \\ |a_n(t)| &\lesssim (t + T_n)^{-2} \log^{-1}(t + T_n), \quad \varepsilon_n(0) \equiv 0, \end{aligned} \quad (4.2)$$

where $(\lambda_n, z_n, \gamma_n, \beta_n, b_n)$ are the parameters of the decomposition of u_n given by Lemma ??, i.e.

$$u_n(t, x) = \frac{e^{i\gamma_n(t)}}{\lambda_n(t)} \left(\sum_k [e^{i\Gamma_{k,n}} Q_{a_n}] \left(\frac{x}{\lambda_n(t)} - z_n(t) e_k \right) + \varepsilon_n \left(t, \frac{x}{\lambda_n(t)} \right) \right), \quad (4.3)$$

with $\Gamma_{k,n}(t, y) = \beta_n(t)(e_k \cdot y) - \frac{b_n(t)}{4}|y|^2$ and $Q_{a_n} = Q + a_n \rho$. Moreover, for all $t \in [t_0 - T_n, 0]$,

$$\int |u_n(t, x)|^2 |x|^2 dx \lesssim 1. \quad (4.4)$$

Proof. Applying Proposition ?? with $s^{in} = n$ for any large n , there exists a solution $u_n(t)$ of (??) defined on the time interval $[-T_n, 0]$ where

$$T_n = \int_{s_0}^n \lambda_n^2(s) ds,$$

and whose decomposition satisfies the uniform estimates (??) on $[-T_n, 0]$. First, we see that (??) follows directly from the estimate on $\lambda_n(s)$ in (??).

Proof of (??). From (??) and the definition of the rescaled time s (see (??)), for any $s \in [s_0, n]$, we have

$$t(s) + T_n = \int_{s_0}^s \lambda_n^2(s') ds' \quad \text{where} \quad |\lambda_n^2(s) - \log^{-2}(s)| \lesssim \log^{-\frac{5}{2}}(s).$$

Fix $\bar{s}_0 > s_0$ large enough independent of n so that, for all $\bar{s}_0 < s < n$,

$$\frac{1}{2}s \log^{-2}(s) \leq \int_{s_0}^s \lambda_n^2(s') ds' = s \log^{-2}(s) + O(s \log^{-\frac{5}{2}}(s)) \leq \frac{3}{2}s \log^{-2}(s).$$

Fix $t_0 = \frac{3}{2}\bar{s}_0 \log^{-2}(\bar{s}_0)$. Then, for all $t \in [t_0 - T_n, 0]$,

$$t + T_n = s \log^{-2}(s) \left(1 + O(\log^{-\frac{1}{2}}(s)) \right) \geq \frac{1}{2}s \log^{-2}(s),$$

and

$$s = (t + T_n) \log^2(t + T_n) \left(1 + O(\log^{-\frac{1}{2}}(t + T_n)) \right).$$

Thus, estimates (??) are direct consequences of (??).

Proof of (??). From (??) and $\varepsilon_n(0) \equiv 0$, we have $u_n(0) \in \Sigma$. It is then standard (see e.g. [?], Proposition 6.5.1) that $u_n \in \mathcal{C}([t_0 - T_n, 0], \Sigma)$. We claim the following preliminary estimates. Fix $A = \frac{16}{\kappa} \geq 8$. For any $k \in \{1, \dots, K\}$, for all $t \in [t_0 - T_n, 0]$,

$$\frac{1}{\lambda_n^2(t)} \int \left| Q_{a_n} \left(\frac{x}{\lambda_n(t)} - z_n(t)e_k \right) \right|^2 |x|^2 dx \lesssim 1, \quad (4.5)$$

$$\frac{1}{\lambda_n^2(t)} \int_{|x| > A} \left| \nabla_x \left([e^{i\Gamma_{k,n}} Q_{a_n}] \left(\frac{x}{\lambda_n(t)} - z_n(t)e_k \right) \right) \right|^2 dx \lesssim (t + T_n)^{-4}. \quad (4.6)$$

Indeed, (??) follows from a change of variable and the decay properties of Q and ρ ,

$$\frac{1}{\lambda_n^2(t)} \int \left| Q_{a_n} \left(\frac{x}{\lambda_n(t)} - z_n(t)e_k \right) \right|^2 |x|^2 dx = \int |Q_{a_n}(y)|^2 |\lambda_n(t)y + \lambda_n(t)z_n(t)e_k|^2 dy \lesssim 1.$$

where we have used from (??),

$$\lambda_n(t)z_n(t) \lesssim 1. \quad (4.7)$$

To show (??), we see first that by (??),

$$|\nabla [e^{i\Gamma_{k,n}} Q_{a_n}](y)|^2 \lesssim |\nabla Q_{a_n}(y)|^2 + (|\beta_n|^2 + b_n^2 |y|^2) Q_{a_n}^2(y) \lesssim e^{-\frac{3}{2}|y|}.$$

Thus, by change of variable (using $A \geq 8$),

$$\begin{aligned} \frac{1}{\lambda_n^2(t)} \int_{|x| > A} \left| \nabla_x \left([e^{i\Gamma_{k,n}} Q_{a_n}] \left(\frac{x}{\lambda_n(t)} - z_n(t)e_k \right) \right) \right|^2 dx \\ = \frac{1}{\lambda_n^2(t)} \int_{|y + z_n(t)e_k| > A/\lambda_n(t)} |\nabla [e^{i\Gamma_{k,n}} Q_{a_n}](y)|^2 dy \\ \lesssim \log^2(t + T_n) \int_{|y| > \frac{A}{2} \log(t + T_n)} e^{-\frac{3}{2}|y|} dy \lesssim (t + T_n)^{-\frac{A}{2}} = (t + T_n)^{-4}, \end{aligned}$$

where we have used from (??) (possibly taking a larger t_0),

$$|y + z_n(t)e_k| > \frac{A}{\lambda_n(t)} \Rightarrow |y| > \frac{A}{\lambda_n(t)} - |z_n(t)| > \left(\frac{3A}{4} - \frac{4}{\kappa} \right) \log(t + T_n) \geq \frac{A}{2} \log(t + T_n).$$

Thus (??) is proved. Observe that (??)–(??) and (??) imply

$$\|xu_n(0)\|_{L^2} \lesssim 1, \quad \|\nabla u_n(t)\|_{L^2(|x| > A)} \lesssim (t + T_n)^{-1} \log^{-\frac{5}{2}}(t + T_n). \quad (4.8)$$

Define $\varphi : \mathbb{R}^2 \rightarrow [0, 1]$ by $\varphi(x) = (|x| - A)^2$ for $|x| > A$ and $\varphi(x) = 0$ otherwise. By elementary computations,

$$\frac{d}{dt} \int |u_n|^2 \varphi = 2 \operatorname{Im} \int (\nabla \varphi \cdot \nabla u_n) \bar{u}_n = 4 \int_{|x| > A} \left(\frac{x}{|x|} \cdot \nabla u_n \right) \bar{u}_n \varphi^{\frac{1}{2}}.$$

Thus, by (??),

$$\left| \frac{d}{dt} \int |u_n|^2 \varphi \right| \lesssim \left(\int |u_n|^2 \varphi \right)^{\frac{1}{2}} \left(\int_{|x| > A} |\nabla u_n(t)|^2 \right)^{\frac{1}{2}} \lesssim (t + T_n)^{-1} \log^{-\frac{5}{2}}(t + T_n) \left(\int |u_n|^2 \varphi \right)^{\frac{1}{2}}.$$

By integration and (??), the following uniform bound holds on $[t_0 - T_n, 0]$,

$$\int |u_n(t, x)|^2 \varphi(x) dx \lesssim 1 \text{ and thus } \int |u_n(t, x)|^2 x^2 dx \lesssim 1,$$

which finishes the proof of (??). $\square$

4.2. Compactness argument. By (??)–(??), the sequence $(u_n(t_0 - T_n))$ is bounded in Σ . Therefore, there exists a subsequence of (u_n) (still denoted by (u_n)) and $u_0 \in \Sigma$ such that

$$\begin{aligned} u_n(t_0 - T_n) &\rightharpoonup u_0 \quad \text{weakly in } H^1(\mathbb{R}^2), \\ u_n(t_0 - T_n) &\rightarrow u_0 \quad \text{in } H^\sigma(\mathbb{R}^2) \text{ for } 0 \leq \sigma < 1, \text{ as } n \rightarrow +\infty. \end{aligned}$$

Let u be the solution of (??) corresponding to $u(t_0) = u_0$. By the local Cauchy theory for (??) (see [?] and [?]) and the properties of the sequence $u_n(t)$ (recall that $T_n \rightarrow \infty$), it follows that $u \in \mathcal{C}([t_0, +\infty), \Sigma)$. Moreover, for all $0 \leq \sigma < 1$, for all $t \in [t_0, +\infty)$,

$$u_n(t - T_n) \rightarrow u(t) \quad \text{in } H^\sigma.$$

By weak convergence in H^1 , $u(t)$ satisfies (??) for all $t \geq t_0$. Moreover, the decomposition $(\vec{p}, \varepsilon)$ of u satisfies, for all $t \geq t_0$,

$$\vec{p}_n(t - T_n) \rightarrow \vec{p}(t), \quad \varepsilon_n(t - T_n) \rightarrow \varepsilon(t) \text{ in } H^\sigma, \quad \varepsilon_n(t - T_n) \rightharpoonup \varepsilon(t) \text{ in } H^1 \quad (4.9)$$

(see e.g. [?], Claim p.598). In particular, for all $t \in [t_0, +\infty)$, $u(t)$ decomposes as

$$u(t, x) = \frac{e^{i\gamma(t)}}{\lambda(t)} \left(\sum_k [e^{i\Gamma_k} Q_a] \left(\frac{x - \lambda(t)z(t)e_k}{\lambda(t)} \right) + \varepsilon \left(t, \frac{x}{\lambda(t)} \right) \right), \quad (4.10)$$

where $\Gamma_k(t, y) = \beta(t)(e_k \cdot y) - \frac{b(t)}{4}|y|^2$ and

$$\begin{aligned} \left| z(t) - \frac{2}{\kappa} \log(t) \right| &\lesssim \log(\log(t)), \quad |\lambda(t) - \log^{-1}(t)| \lesssim \log^{-\frac{3}{2}}(t), \\ |b(t) - t^{-1} \log^{-3}(t)| + |\beta(t)| + \|\varepsilon(t)\|_{H^1} &\lesssim t^{-1} \log^{-\frac{7}{2}}(t), \quad |a(t)| \lesssim t^{-2} \log^{-1}(t), \\ \int |u(t, x)|^2 |x|^2 dx &\lesssim 1. \end{aligned} \quad (4.11)$$

Note that by (??), we have for all $k \in \{1, \dots, K\}$,

$$x_k(t) = \lambda(t)z(t)e_k \rightarrow \frac{2}{\kappa}e_k, \quad \text{with} \quad \left| x_k(t) - \frac{2}{\kappa}e_k \right| \lesssim \frac{\log(\log(t))}{\log(t)}.$$

Since $\lambda^{-1}(t)\|\varepsilon(t)\|_{H^1} \lesssim t^{-1} \log^{-\frac{5}{2}}(t)$ and, by (??) and (??),

$$\lambda^{-1}(t) \|e^{i\Gamma_k} Q_a - Q\|_{H^1} \lesssim \lambda^{-1}(t)(|\beta(t)| + |b(t)| + |a(t)|) \lesssim t^{-1} \log^{-2}(t), \quad (4.12)$$

we obtain the following stronger form of (??)

$$\left\| u(t) - e^{i\gamma(t)} \sum_k \frac{1}{\lambda(t)} Q \left(\frac{\cdot - x_k(t)}{\lambda(t)} \right) \right\|_{H^1} \lesssim t^{-1} \log^{-2}(t). \quad (4.13)$$

Next, since for $j \neq k$, for some q ,

$$\lambda^{-2}(t) \int |\nabla Q(y - z(t)e_k) \cdot \nabla Q(y - z(t)e_j)| dy \lesssim |z|^q e^{-\kappa z} \lesssim t^{-1},$$

we also obtain (??). As a final remark, note that by global existence and uniform bound in Σ , the virial identity (??) implies the rigidity $E(u) = 0$. This concludes the proof of Theorem ??.

4.3. Proof of Corollary ??. For $-t_0^{-1} < t < 0$, we set

$$\begin{aligned}\tilde{z}(t) &= z(|t|^{-1}), \quad \tilde{\lambda}(t) = |t|\lambda(|t|^{-1}), \quad \tilde{a}(t) = a(|t|^{-1}), \quad \tilde{b}(t) = b(|t|^{-1}), \\ \tilde{\gamma}(t) &= \gamma(|t|^{-1}), \quad \tilde{\beta}(t) = \beta(|t|^{-1}), \quad \tilde{\varepsilon}(t) = \varepsilon(|t|^{-1}), \quad \tilde{\Gamma}_k(t, y) = \tilde{\beta}(t)(e_k \cdot y) - \frac{\tilde{b}(t)}{4}|y|^2,\end{aligned}$$

so that from (??),

$$\begin{aligned}\left| \tilde{z}(t) - \frac{2}{\kappa} \log |t| \right| &\lesssim \log |\log |t||, \quad \left| \tilde{\lambda}(t) - |t| \log |t|^{-1} \right| \lesssim |\log |t||^{-\frac{3}{2}}, \\ \left| \tilde{b}(t) - |t| \log |t|^{-3} \right| + |\tilde{\beta}(t)| + \|\tilde{\varepsilon}(t)\|_{H^1} &\lesssim |t| \log |t|^{-\frac{7}{2}}, \quad |\tilde{a}(t)| \lesssim |t| \log |t|^{-1}.\end{aligned}\tag{4.14}$$

We see from (??) that the pseudo-conformal transform $v(t)$ of $u(t)$ as defined in (??) satisfies

$$v(t, x) = e^{-i\frac{|x|^2}{4|t|}} w(t, x), \quad w(t, x) = \frac{e^{i\tilde{\gamma}(t)}}{\tilde{\lambda}(t)} \left(\sum_k [e^{i\tilde{\Gamma}_k} Q_{\tilde{a}}] \left(\frac{x}{\tilde{\lambda}(t)} - \tilde{z}(t)e_k \right) + \tilde{\varepsilon} \left(t, \frac{x}{\tilde{\lambda}(t)} \right) \right).$$

Note in particular that $\tilde{\lambda}(t)\tilde{z}(t) \sim \frac{2}{\kappa}|t|$ as $t \uparrow 0$. From this, it follows that

$$|v(t, x)|^2 \rightharpoonup K \|Q\|_{L^2}^2 \delta_0 \quad \text{as } t \uparrow 0.$$

Finally, since $\nabla v(t, x) = e^{-i\frac{|x|^2}{4|t|}} \left(\nabla w - i\frac{x}{2|t|} w \right) (t, x)$, and as $t \uparrow 0$,

$$\begin{aligned}\frac{1}{|t|^2} \int |x|^2 |w(t, x)|^2 dx &\lesssim \left| \frac{\tilde{\lambda}(t)}{t} \right|^2 \int \left| \sum_k [e^{i\tilde{\Gamma}_k} Q_{\tilde{a}}] (y - \tilde{z}(t)e_k) + \tilde{\varepsilon}(t, y) \right|^2 |y|^2 dy \lesssim 1, \\ \int |\nabla w(t, x)|^2 dx &\sim K \|\nabla Q\|_{L^2}^2 |t|^{-2} |\log |t||^2,\end{aligned}$$

we obtain (??). Note that $\int |x|^2 |v(t, x)|^2 \lesssim t^2$ implies by (??) that $\int |x|^2 |v(t, x)|^2 = t^2 E(v)$. Thus, $E(v) > 0$.

REFERENCES

- [1] C. Antonini. Lower bounds for the L^2 minimal periodic blow-up solutions of critical nonlinear Schrödinger equation. *Differential Integral Equations* 15 (2002), no. 6, 749–768.
- [2] V. Banica. Remarks on the blow-up for the Schrödinger equation with critical mass on a plane domain. *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) 3 (2004), no. 1, 139–170.
- [3] V. Banica, R. Carles and T. Duyckaerts. Minimal blow-up solutions to the mass-critical inhomogeneous NLS equation. *Comm. Partial Differential Equations* 36 (2011), no. 3, 487–531.
- [4] H. Berestycki and P.-L. Lions, Nonlinear scalar field equations. I. Existence of a ground state. *Arch. Rational Mech. Anal.* 82: 313–345, 1983.
- [5] T. Boulenger. Minimal mass blow up for NLS on a manifold. *PhD dissertation*, Orsay 2012.
- [6] J. Bourgain and W. Wang, Construction of blowup solutions for the nonlinear Schrödinger equation with critical nonlinearity. *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (4) 25 (1997), no. 1-2, 197–215.
- [7] T. Cazenave. *Semilinear Schrödinger equations*. New York University – Courant Institute, New York, 2003.
- [8] T. Cazenave and F. B. Weissler. The Cauchy problem for the critical nonlinear Schrödinger equation in H^s . *Nonlinear Anal.*, 14(10):807–836, 1990.
- [9] S.-M. Chang, S. Gustafson, K. Nakanishi and T.-P. Tsai. Spectra of linearized operators for NLS solitary waves. *SIAM J. Math. Anal.*, 39(4):1070–1111, 2007/08.
- [10] C. Collot. Type II blow up manifold for the energy super critical wave equation. *arXiv:1407.4525*
- [11] R. Côte, Y. Martel and F. Merle Construction of multi-soliton solutions for the L^2 -supercritical gKdV and NLS equations. *Rev. Mat. Iberoam.* 27 (2011), no. 1, 273–302.

- [12] B. Dodson. Global well-posedness and scattering for the mass critical nonlinear Schrödinger equation with mass below the mass of the ground state. *arXiv:1104.1114*.
- [13] R. Donninger, M. Huang, J. Krieger and W. Schlag. Exotic blowup solutions for the u^5 focusing wave equation in $\mathbb{R}^3$. *Michigan Math. J.* 63 (2014), no. 3, 451–501.
- [14] T. Duyckaerts, C.E. Kenig and F. Merle. Profiles for bounded solutions of dispersive equations, with applications to energy-critical wave and Schrödinger equations. *Commun. Pure Appl. Anal.* 14 (2015), no. 4, 1275–1326.
- [15] T. Duyckaerts and F. Merle. Dynamic of threshold solutions for energy-critical NLS. *Geom. Funct. Anal.* 18 (2009), no. 6, 1787–1840.
- [16] C. Fan. Propagation of regularity and the localization of log-log blow up. *arXiv:1510.00961*
- [17] M. A. Herrero and J. J. L. Velázquez. Flat blow-up in one-dimensional semilinear heat equations. *Differential Integral Equations* 5 (1992), 973–997.
- [18] J. Jendrej. Construction of type II blow-up solutions for the energy-critical wave equation in dimension 5. *arXiv:1503.05024*
- [19] R. Killip and T. Tao and M. Visan. The cubic nonlinear Schrödinger equation in two dimensions with radial data. *J. Eur. Math. Soc.* 11 (2009), no. 6, 1203–1258.
- [20] J. Krieger, E. Lenzmann and P. Raphaël. Nondispersive solutions to the L^2 -critical half-wave equation. *Arch. Ration. Mech. Anal.* 209 (2013), no. 1, 61–129.
- [21] J. Krieger, Y. Martel and P. Raphaël. Two-soliton solutions to the three-dimensional gravitational Hartree equation. *Comm. Pure Appl. Math.* 62 (2009), no. 11, 1501–1550.
- [22] J. Krieger and W. Schlag. Non-generic blow-up solutions for the critical focusing NLS in 1-D. *J. Eur. Math. Soc.* 11 (2009), 1–125.
- [23] J. Krieger and W. Schlag. Full range of blow up exponents for the quintic wave equation in three dimensions. *J. Math. Pures Appl.* (9) 101 (2014), no. 6, 873–900.
- [24] J. Krieger, W. Schlag and D. Tataru. Renormalization and blow up for charge one equivariant critical wave maps. *Invent. Math.* 171 (2008), no. 3, 543–615.
- [25] M. K. Kwong. Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in $\mathbb{R}^n$. *Arch. Rational Mech. Anal.*, 105(3):243–266, 1989.
- [26] S. Le Coz, Y. Martel, P. Raphaël. Minimal mass blow up solutions for a double power nonlinear Schrödinger equations To appear in *Revista Matemática Iberoamericana*.
- [27] Y. Martel. Asymptotic N -soliton-like solutions of the subcritical and critical generalized Korteweg de Vries equations. *Amer. J. Math.* 127 (2005), no. 5, 1103–1140.
- [28] Y. Martel and F. Merle. Nonexistence of blow-up solution with minimal L^2 -mass for the critical gKdV equation. *Duke Math. J.* 115 (2002), no. 2, 385–408.
- [29] Y. Martel and F. Merle. Inelastic interaction of nearly equal solitons for the quartic gKdV equation. *Invent. Math.* 183 (2011), no. 3, 563–648.
- [30] Y. Martel and F. Merle. Stability of blow up profile and lower bound on the blow up rate for the critical generalized KdV equation. *Ann. of Math.*, 127:235–280, 2002.
- [31] Y. Martel and F. Merle. Multi-solitary waves for nonlinear Schrödinger equations. *Annales de l'IHP (C) Non Linear Analysis*, 23 (2006), 849–864.
- [32] Y. Martel, F. Merle and P. Raphaël. Blow up for the critical generalized Korteweg de Vries equation. I: Dynamics near the soliton. *Acta Math.* 212 (2014), no. 1, 59–140.
- [33] Y. Martel, F. Merle and P. Raphaël. Blow up for the critical generalized Korteweg de Vries equation. II: Minimal mass dynamics. *J. of Math. Eur. Soc.* 17, 1855–1925 (2015).
- [34] Y. Martel, F. Merle and P. Raphaël. Blow up for the critical generalized Korteweg de Vries equation. III: Exotic regimes. *Annali della Scuola Normale Superiore di Pisa XIV*, 575–631 (2015).
- [35] Y. Martel, F. Merle and T.-P. Tsai. Stability in H^1 of the sum of K solitary waves for some nonlinear Schrödinger equations. *Duke Math. J.* 133 (2006), 405–466.
- [36] F. Merle. Construction of solutions with exactly k blow-up points for the Schrödinger equation with critical nonlinearity. *Comm. Math. Phys.* 129 (1990), no. 2, 223–240.
- [37] F. Merle. Determination of blow-up solutions with minimal mass for nonlinear Schrödinger equations with critical power. *Duke Math. J.* 9:427–454, 1993.
- [38] F. Merle and P. Raphaël. On universality of blow-up profile for L^2 critical nonlinear Schrödinger equation. *Invent. Math.*, 156(3):565–672, 2004.

- [39] F. Merle and P. Raphaël. The blow-up dynamic and upper bound on the blow-up rate for critical nonlinear Schrödinger equation. *Ann. of Math. (2)*, 161(1):157–222, 2005.
- [40] F. Merle and P. Raphaël. Sharp upper bound on the blow up rate for critical nonlinear Schrödinger equation. *Geom. Funct. Anal.* 13 (2003), 591–642.
- [41] F. Merle and P. Raphaël. On a sharp lower bound on the blow-up rate for the L^2 critical nonlinear Schrödinger equation. *J. Amer. Math. Soc.*, 19(1):37–90 (electronic), 2006.
- [42] F. Merle and P. Raphaël. Profiles and quantization of the blow up mass for critical nonlinear Schrödinger equation. *Comm. Math. Phys.* 253 (2005), no. 3, 675–704.
- [43] F. Merle and P. Raphaël and I. Rodnianski. Type II blow up for the energy supercritical NLS to appear in *Cambridge Math. Jour.*
- [44] F. Merle, P. Raphaël and J. Szeftel. The instability of Bourgain-Wang solutions for the L^2 critical NLS. *Amer. J. Math.* 135 (4), 967–1017, 2013.
- [45] F. Merle, P. Raphaël and J. Szeftel. On collapsing ring blow up solutions to the mass supercritical NLS. *Duke Math. J.* 163 (2), 369–431, 2014.
- [46] F. Merle and H. Zaag. On the stability of the notion of non-characteristic point and blow-up profile for semilinear wave equations. *Comm. Math. Phys.* 333 (2015), no. 3, 1529–1562.
- [47] E. Miot. Dynamique des points vortex dans une équation de Ginzburg-Landau complexe. (French) Séminaire: Equations aux Dérivées Partielles. 2009–2010, Exp. No. XXI, 13 pp. Ecole Polytech., Palaiseau, 2012.
- [48] T. Mizumachi. Weak interaction between solitary waves of the generalized KdV equations. *SIAM J. Math. Anal.* 35 (2003), no. 4, 1042–1080
- [49] M. Musso, F. Pacard and J. Wei. Finite-energy sign-changing solutions with dihedral symmetry for the stationary nonlinear Schrödinger equation. *J. Eur. Math. Soc.* 14 (2012), no. 6, 1923–1953.
- [50] H. Nawa. Asymptotic and limiting profiles of blowup solutions of the nonlinear Schrödinger equation with critical power. *Comm. Pure Appl. Math.* 52 (1999), no. 2, 193–270.
- [51] G. Perelman. On the formation of singularities in solutions of the critical nonlinear Schrödinger equation. *Ann. Henri Poincaré* 2 (2001), no. 4, 605–673.
- [52] G. Perelman. Two soliton collision for nonlinear Schrödinger equations in dimension 1. *Ann. Inst. H. Poincaré, Anal. Non Linéaire* 28 (2011) 357–384.
- [53] F. Planchon and P. Raphaël. Existence and stability of the log-log blow-up dynamics for the L^2 -critical nonlinear Schrödinger equation in a domain. *Ann. Henri Poincaré*, 8(6):1177–1219, 2007.
- [54] P. Raphaël. Stability of the log-log bound for blow up solutions to the critical non linear Schrödinger equation. *Math. Ann.* 331 (2005), no. 3, 577–609.
- [55] P. Raphaël and J. Szeftel. Existence and uniqueness of minimal blow-up solutions to an inhomogeneous mass critical NLS. *J. Amer. Math. Soc.*, 24(2):471–546, 2011.
- [56] M. I. Weinstein. Nonlinear Schrödinger equations and sharp interpolation estimates. *Comm. Math. Phys.*, 87(4):567–576, 1982/83.
- [57] M. I. Weinstein. Modulational stability of ground states of nonlinear Schrödinger equations. *SIAM J. Math. Anal.*, 16 (1985), 472–491.

CMLS, ÉCOLE POLYTECHNIQUE, CNRS, 91128 PALAISEAU, FRANCE

E-mail address: yvan.martel@polytechnique.edu

UNIVERSITÉ DE NICE SOPHIA-ANTIPOLIS, INSTITUT UNIVERSITAIRE DE FRANCE, EUROPEAN RESEARCH COUNCIL, FRANCE

E-mail address: pierre.rafael@unice.fr