La^{III} and Zn^{II} Cooperatively Template a Metal-Organic Capsule

Dong Yang,^{†,‡} Jake L. Greenfield,[†] Tanya K. Ronson,[†] Larissa K. S. von Krbek,^{†,×} Le Yu,[‡] Jonathan R. Nitschke^{†*}

[†] Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, United Kingdom

[‡] Key Laboratory of Synthetic and Natural Functional Molecule of the Ministry of Education, College of Chemistry and Materials Science, Northwest University, Xi'an 710069, China

ABSTRACT: An organic subcomponent was designed with 2-formyl-8-aminoquinoline and triazole-pyridine ends. The relative orientations and geometries of these two ends enabled this subcomponent to assemble together with Zn^{II} and La^{III} cations to generate a heterobimetallic tetrahedral capsule. The La^{III} cations each template three imine bonds that hold together a threefold-symmetric metallo-ligand, defining the center of each tetrahedron face. The Zn^{II} cations occupy the other ends of these C₃ axes, defining the vertices of the tetrahedron. This is the first example where subcomponent self-assembly brought into being the faces of a polyhedron, as opposed to the vertices. Host-guest studies show positively cooperative binding towards ReO₄[−], the encapsulation of which also resulted in the quenching of capsule fluorescence.

Self-assembly brings the subunits together of the most intricate biological structures,¹ enabling them to function.² Natural assemblies have inspired the construction of synthetic supramolecular structures, which have found uses in stimuli responsive behavior,³ recognition,⁴ adsorption,⁵ transport⁶ and catalysis.⁷

D-block metal ions, particularly those from the first row, have proven very useful in bringing together organic-ligand building blocks into complex assemblies.⁸ Among these are macrocycles,⁹ helicates,¹⁰ catenanes,¹¹ knots,¹² and hollow capsules.¹³ The functions associated with these capsules, in particular, have made them a topic of considerable current interest.¹⁴

Lanthanide-containing architectures are more challenging to design, due to the variable coordination numbers and geometries of these cations. These complexes are nonetheless of great technological interest, as a result of their optical, chiroptical and magnetic properties.¹⁵ In addition to mononuclear complexes and dinuclear helicates,^{15a} research groups led by Hamacek,¹⁶ Duan¹⁷ and Sun¹⁸ have designed intricate three-dimensional functional lanthanide assemblies.

The combination of lanthanides and *d*-block metals within a single heterometallic metal-organic structure could enable functions derived from both, in fields that include luminescence,^{15b, 19} catalysis,²⁰ chemosensors²¹ and magnets.^{20, 22} The design of such heterometallic architectures is challenging, however, not only due to the coordinative flexibility of the lanthanides, but also because ligands

for *d*-block metals often coordinate well to lanthanides. Complex mixtures, or homometallic complexes, thus often result from mixtures of ligands with both kinds of metal ions.²¹

With the aim of expanding the scope of subcomponent self-assembly^{8c} to encompass the formation of new heterometallic architectures, we designed subcomponent **A** (Scheme 1) to simultaneously coordinate lanthanide and *d*-block ions, selectively. The 2-formyl-8-aminoquinoline moiety of **A** presents an awkward fit for hexacoordinate zinc(II). Whereas this group condensed to form double-helical poly(imine)s around Cu^I templates.²³ We hypothesized that imine condensation around a larger lanthanide template might generate a trimeric macrocycle, however, which would array the triazole(pyridine) sites so as to coordinate to zinc(II) centers at the vertices of a face-capped tetrahedron. This hypothesis was confirmed in the synthesis of capsule **1**.

Scheme 1. Subcomponent self-assembly of heterobimetallic Zn^{II}–La^{III} capsule **1**, with the aza-18-crown-6 moiety templated by La^{III} highlighted in red.

The reaction of subcomponent **A** (12 equiv) with Zn(OTf)₂ (4.0 equiv) and La(OTf)₃ (4.0 equiv) in acetonitrile gave a light-yellow solution of heterobimetallic capsule **1**, which could also be prepared as the perchlorate from Zn(ClO₄)₂ and La(ClO₄)₃. Pure **1** was isolated by precipitation with diethyl ether. A single imine peak was observed in the ¹H NMR spectra regardless of whether TfO[−] (Figures S11, S17) or ClO₄[−] (Figures 1, S12, S22) was used as the counter anion, consistent with *T* point symmetry. However, only broad signals were observed in the ¹H NMR spectra when tetrakis(3,5-bis(trifluoromethyl)phenyl)borate (BARF[−]) or bis(trifluoromethanesulfonyl)imide (Tf₂N[−]) was used as counter anion (Figures S26, S27). Likewise, treatment of subcomponent **A** (12 equiv) with only one of the two metal

templates, either $\text{Zn}(\text{OTf})_2$ (8.0 equiv) or $\text{La}(\text{OTf})_3$ (8.0 equiv), did not result in capsule formation (Figure S14). Treatment of **1** with EDTA in order to remove La^{III} resulted in destruction of the capsule framework (Figure S25). These experiments together indicate that La^{III} , Zn^{II} and the counter anion all play important roles in the stabilization of capsule **1**.

ESI-MS provided further evidence for the formulation of capsule **1** (Figure S24). DOSY NMR indicated that the ^1H peaks assigned to **1** exhibit the same diffusion rate, consistent with the formation of only a single species (Figures 1, S18, S23) with a diffusion coefficient of $3.2 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$ for both $\mathbf{1} \cdot (\text{ClO}_4)_2$ and $\mathbf{1} \cdot (\text{OTf})_2$, which corresponds to an approximate solvodynamic radius of 2.0 nm.²⁴

Figure 1. ^1H NMR and DOSY spectra (CD_3CN , 400 MHz, 25 °C) of $\mathbf{1} \cdot (\text{ClO}_4)_{20}$.

Single crystals of $\mathbf{1} \cdot (\text{ClO}_4)_{20}$ were obtained by slow vapor diffusion of diethyl ether into an acetonitrile solution. The ClO_4^- salt of capsule **1** crystallized in the cubic space group, $I23$, with two tetrahedra in one unit cell. $\text{Zn}^{\text{II}}\text{--La}^{\text{III}}$ heterobimetallic tetrahedral capsule **1** (Figure 2) possesses faces and vertices of a single handedness, giving rise to T point symmetry. The condensation of three equivalents of sub-component **A** formed macrocyclic faces that contain alternating imine and quinoline donors, all six of which coordinate equatorially to a La^{III} center.

Figure 2. a) Crystal structure of $\mathbf{1} \cdot (\text{ClO}_4)_{20}$. b) and c) are orthogonal views of the facial Ln^{III} center of **1** (Zn^{II} = yellow, La^{III} = pink C = gray, N = blue, O = red, Cl = green, H = white).

The bowl-shaped macrocyclic faces pucker outwards, with the La^{III} center 0.64 Å out of the center of the macrocyclic plane toward the outside of the capsule and a bend angle of 154.4°, measured as the N-La-N angle between opposite imine and quinoline donors.

In addition to the six nitrogen donors of the macrocyclic capsule face, each La^{III} center is also coordinated by four

water molecules (Figure 2b, 2c). Three of these water molecules are outside the capsule, while the fourth is located inside the capsule, forming a capped trigonal cupola coordination geometry around La^{III} (Figure S52).²⁵

The terminal triazole-pyridine groups from each residue of **A** coordinate to octahedral Zn^{II} centers at the four vertices of the tetrahedron. The $\text{Zn}^{\text{II}}\cdots\text{Zn}^{\text{II}}$ distances are 19.759(1) Å. A cavity volume of 617 Å³ was determined using the VOIDOO program²⁶ (Figure S55). The distance measured from the center of the capsule to the most distant atoms of the terminal benzyl groups is approximately 1.69 nm, consistent with the DOSY measurement in solution.

The ClO_4^- counter anions were found both inside and outside the cavity. Four ClO_4^- anions are encapsulated at the four corners of the tetrahedral capsule, engaging in anion- π interactions with the electron-deficient quinoline rings. Six well-resolved ClO_4^- anions lie on the six edges (Figure S53), each having six C-H $\cdots$ O close contacts with capsule hydrogen atoms, with distances in the range 2.5–2.7 Å. A further ten ClO_4^- anions could not be resolved due to the weak intensity diffraction data and their disorder.

NOESY spectra (Figures S16 and S21) indicated that **1** adopted the same configuration in solution as in the crystal. Cross-peaks were observed between H1 and H4, indicating the formation of cyclic ligand with covalent (N=C) bonds. The through-space interaction between H2 and H4 on separate faces is consistent with the solid-state configuration shown in Figure 2. The interactions observed between H5-H6 and H3-H7 are also consistent with this structure, but preclude the adoption of a different diastereomeric configuration.²⁷

Based on the crystal structure of $\mathbf{1} \cdot (\text{ClO}_4)_{20}$, we inferred that **1** would be capable of binding four different anions inside its cavity in place of the perchlorates. Hence, we investigated the interactions between **1** and different anions (Figure S31) using ^1H NMR and fluorescence spectroscopy. While only slight changes to the NMR spectra were observed upon addition of 20 equivalents of BF_4^- , HSO_4^- , PF_6^- , CN^- , and H_2PO_4^- to $\mathbf{1} \cdot (\text{ClO}_4)_{20}$ (12 μM), addition of ReO_4^- and TfO^- caused more significant changes, which we infer to have been due to the displacement of bound ClO_4^- by ReO_4^- or TfO^- . ReO_4^- caused a significant downfield shift ($\Delta\delta = 0.48$ ppm) of the imine signal H1 (Figure S28), and TfO^- induced a smaller upfield shift of H1 ($\Delta\delta = -0.03$ ppm). The chemical shift changes for H1-H5 in the presence of various anions are summarized in Table S1. These observations are consistent with the conclusions that the affinity of **1** for ReO_4^- and TfO^- was larger than for ClO_4^- , while BF_4^- , HSO_4^- , PF_6^- , CN^- , and H_2PO_4^- compete only weakly with ClO_4^- .

Figure 3. Diagram showing the different affinities and modes of binding of ClO_4^- , TfO^- and ReO_4^- to capsule **1**.

The binding behavior of capsule **1** was studied by competitive ^1H NMR titration of ReO_4^- , ClO_4^- and TfO^- into solutions of $\mathbf{1}\cdot(\text{ClO}_4)_{20}$ and $\mathbf{1}\cdot(\text{OTf})_{20}$, respectively (Figures S37-S45). TBAREO_4 was added to a solution of $\mathbf{1}\cdot(\text{ClO}_4)_{20}$ (11 μM in CD_3CN) or $\mathbf{1}\cdot(\text{OTf})_{20}$ (70 μM) and the concentrations of **1** and $\mathbf{1}\cdot\text{ReO}_4^-$ were plotted and fitted with a Hill function.²⁸ In all ^1H NMR titrations, new imine signals appeared with increasing relative concentration of the added anions, consistent with anion exchange inside the heterobimetallic tetrahedron. ESI-MS and 2D NMR further supported the anion exchange (Figures S32-S36). The apparent association constants for ReO_4^- towards $\mathbf{1}\cdot(\text{ClO}_4)_{20}$ and $\mathbf{1}\cdot(\text{OTf})_{20}$ were determined to be $(8.88 \pm 0.01) \times 10^3$ and $(3.42 \pm 0.01) \times 10^3$,²⁹ respectively. These values indicate that ReO_4^- binds more strongly to **1** than either ClO_4^- or TfO^- . The Hill coefficients (n)²⁸ were determined to be 5.59 and 4.01, respectively, indicating positively cooperative binding of ReO_4^- in the cases of both $\mathbf{1}\cdot(\text{ClO}_4)_{20}$ and $\mathbf{1}\cdot(\text{OTf})_{20}$.

Binding of TfO^- to $\mathbf{1}\cdot(\text{ClO}_4)_{20}$ was also explored, and an apparent association constant of $(4.09 \pm 0.01) \times 10^3$ was determined (Figures S41, S42). Noncooperative binding was observed in the case of TfO^- , with a Hill coefficient of $n = 1.09$. The cooperative inclusion behavior in the case of perrhenate may be a consequence of the shape and size of the cavity: when one tetrahedral ReO_4^- anion is bound inside a corner of the cavity of **1**, the shape of the cavity may adapt to fit this guest. This change in cavity shape may make it easier to bind the second, third, and fourth perrhenates.

Upon titration of ClO_4^- into an acetonitrile solution of $\mathbf{1}\cdot(\text{OTf})_{20}$ (48 μM , Figures S43-S45), the phenyl protons H2, H3 and H5 were first observed to shift upfield (0-15.7 equiv ClO_4^-), and then downfield (15.7-690.8 equiv ClO_4^-). As there are six ClO_4^- anions located at the edges of **1** in the crystal structure (Figure S53), we hypothesize that ClO_4^- interacts initially with the edges of capsule **1**, and then exchanges with the encapsulated TfO^- when an excess of ClO_4^- is added.

Low-temperature ^{19}F NMR further elucidated the interactions between ClO_4^- and $\mathbf{1}\cdot(\text{OTf})_{20}$. Upon addition of ClO_4^- to $\mathbf{1}\cdot(\text{OTf})_{20}$, (Figure S46), we observed the ^{19}F signals corresponding to encapsulated TfO^- to disappear only after the addition of a large excess (601.7 equiv) of ClO_4^- . This replacement of TfO^- to ClO_4^- in $\mathbf{1}\cdot(\text{OTf})_{20}$ only occurs at high

ClO_4^- concentrations, as expected from the lack of thermodynamic favorability of the reverse process (binding of TfO^- to $\mathbf{1}\cdot(\text{ClO}_4)_{20}$ $(4.09 \pm 0.01) \times 10^3$).

By treating the lower- $[\text{ClO}_4^-]$ and higher- $[\text{ClO}_4^-]$ regimes independently (Figure S45), the binding affinities of ClO_4^- for $\mathbf{1}\cdot(\text{OTf})_{20}$ were calculated to be $(5.85 \pm 0.01) \times 10^3$ (periphery) and 168.9 ± 1.2 (cavity). We infer the greater binding constant for *exo*- binding over *endo*- binding arises from the TfO^- anions already present competing more effectively with ClO_4^- for binding within the cavity of **1**, in comparison with the external binding sites. Comparing the binding affinities of the three anions in CD_3CN , a hierarchy of $\text{ReO}_4^- > \text{TfO}^- > \text{ClO}_4^-$ was observed.

Figure 4. a) Fluorescence emission spectra of $\mathbf{1}\cdot(\text{ClO}_4)_{20}$ (2 μM , $\lambda_{\text{ex}} = 375$ nm, $\lambda_{\text{em}} = 543$ nm) and subcomponent **A** (24 μM , $\lambda_{\text{ex}} = 422$ nm, $\lambda_{\text{em}} = 611$ nm) in MeCN; b) Fluorescence emission spectra during the titration of 20 equiv of ReO_4^- into $\mathbf{1}\cdot(\text{ClO}_4)_{20}$ (2 μM in MeCN, $\lambda_{\text{ex}} = 375$ nm). Inset: The decrease of fluorescence intensity at $\lambda = 543$ nm as a function of perrhenate concentration.

Fluorescence measurements were carried out to study the luminescent behavior of **1** during guest encapsulation. Subcomponent **A** exhibited fluorescent emission centered at 611 nm (Figure 4a). This emission shows an enhancement in intensity and a blue shift of 68 nm after the formation of **1**. These changes in emission were attributed to formation of the imine bonds and to the incorporation of La^{III} .

Upon titration of ReO_4^- (20 equiv) into a solution of $1 \cdot (\text{ClO}_4)_2$, the fluorescence intensity was progressively quenched, reaching a plateau once 16 equivalents of ReO_4^- had been added (Figure 4b). Separate additions of HSO_4^- and CN^- (20 equiv in each case) resulted in quenching of 41% and 34% of the fluorescence intensity of **1** (Figure S49), respectively. Other anions either induced no obvious change in fluorescence (PF_6^- and BF_4^-) or only minimal quenching (H_2PO_4^- and OTf^-). Luminescence measurements may thus provide a more sensitive probe of the peripheral binding of these anions than does ^1H NMR.

Heterobimetallic capsule **1** thus presents an initial use of subcomponent self-assembly to construct the faces, as opposed to the vertices, of a three-dimensional structure, and the introduction of a lanthanide template ion. This capsule shows rich and varied anion-binding behavior, and paves the way for the creation of analogs using first-row transition metals in place of zinc, together with f-block metals in place of lanthanum. Such new heterobimetallic capsules could possess novel functions linked to the magnetism of both d- and f-block metals, together with luminescent behavior of the heavier lanthanides.

ASSOCIATED CONTENT

The Supporting Information is available free of charge on the ACS Publications website at DOI: xxxxxxxx.
Complete experimental details (PDF)
X-ray data for **1** (CCDC 2024202) (CIF)

AUTHOR INFORMATION

Corresponding Author

*jrn34@cam.ac.uk

ORCID

Dong Yang: 0000-0002-1971-7349
Jake L. Greenfield: 0000-0002-7650-5414
Tanya K. Ronson: 0000-0002-6917-3685
Larissa K. S. von Krbek: 0000-0002-4278-5235
Le Yu: 0000-0002-0360-290X
Jonathan R. Nitschke: 0000-0002-4060-5122

Present Addresses

× Kekulé-Institut für Organische Chemie und Biochemie der Rheinischen Friedrich-Wilhelms-Universität Bonn, Gerhard-Domagk-Str. 1, 53121 Bonn, Germany.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

This work was supported by the European Research Council (695009) and the UK Engineering and Physical Sciences Research Council (EPSRC EP/P027067/1). The authors thank Diamond Light Source (UK) for synchrotron beamtime on I19 (CY21497). D.Y. acknowledges the National Natural Science Foundation of China (No. 21971210), Natural Science Foundation of Shaanxi Province (2019KJXX-062) and the Chinese Scholarship Council (CSC). L.K.S.v.K. acknowledges the

Alexander von Humboldt Foundation for a Feodor Lynen Research Fellowship. The authors thank Professor Biao Wu (Northwest University) for useful discussions and suggestions and Dr Felix J. Rizzuto for preliminary crystallographic work.

REFERENCES

- Whitesides, G. M.; Mathias, J. P.; Seto, C. T., Molecular Self-Assembly and Nanochemistry: A Chemical Strategy for the Synthesis of Nanostructures. *Science* **1991**, *254*, 1312-1319.
- (a) Mendes, A. C.; Baran, E. T.; Reis, R. L.; Azevedo, H. S., Self-Assembly in Nature: Using the Principles of Nature to Create Complex Nanobiomaterials. *Wiley Interdiscip. Rev.: Nanomed. Nanobiotechnol.* **2013**, *5*, 582-612; (b) Seeman, N. C., DNA in a Material World. *Nature* **2003**, *421*, 427-431.
- (a) Kim, H.-J.; Kim, T.; Lee, M., Responsive Nanostructures from Aqueous Assembly of Rigid-Flexible Block Molecules. *Acc. Chem. Res.* **2011**, *44*, 72-82; (b) Aliprandi, A.; Mauro, M.; De Cola, L., Controlling and Imaging Biomimetic Self-Assembly. *Nat. Chem.* **2016**, *8*, 10-15.
- (a) Jie, K.; Zhou, Y.; Li, E.; Huang, F., Nonporous Adaptive Crystals of Pillararenes. *Acc. Chem. Res.* **2018**, *51*, 2064-2072; (b) Yang, D.; Zhao, J.; Zhao, Y.; Lei, Y.; Cao, L.; Yang, X.-J.; Davi, M.; de Sousa Amadeu, N.; Janiak, C.; Zhang, Z.; Wang, Y.-Y.; Wu, B., Encapsulation of Halocarbons in a Tetrahedral Anion Cage. *Angew. Chem., Int. Ed.* **2015**, *54*, 8658-8661; (c) Yang, D.; Zhao, J.; Yu, L.; Lin, X.; Zhang, W.; Ma, H.; Gogoll, A.; Zhang, Z.; Wang, Y.; Yang, X.-J.; Wu, B., Air- and Light-Stable P4 and As4 within an Anion-Coordination-Based Tetrahedral Cage. *J. Am. Chem. Soc.* **2017**, *139*, 5946-5951; (d) Xu, Y.; Gsäenger, S.; Minameyer, M. B.; Imaz, I.; Maspoche, D.; Shyshov, O.; Schwer, F.; Ribas, X.; Drewello, T.; Meyer, B.; von Delius, M., Highly Strained, Radially π -Conjugated Porphyrinylene Nanohoops. *J. Am. Chem. Soc.* **2019**, *141*, 18500-18507.
- Suh, M. P.; Cheon, Y. E.; Lee, E. Y., Syntheses and Functions of Porous Metallosupramolecular Networks. *Coord. Chem. Rev.* **2008**, *252*, 1007-1026.
- Gale, P. A.; Davis, J. T.; Quesada, R., Anion Transport and Supramolecular Medicinal Chemistry. *Chem. Soc. Rev.* **2017**, *46*, 2497-2519.
- (a) Brown, C. J.; Toste, F. D.; Bergman, R. G.; Raymond, K. N.; Bergman, R. G.; Raymond, K. N., Supramolecular Catalysis in Metal-Ligand Cluster Hosts. *Chem Rev* **2015**, *115*, 3012-35; (b) Hao, M.; Sun, G.; Zuo, M.; Xu, Z.; Chen, Y.; Hu, X.-Y.; Wang, L., A Supramolecular Artificial Light-Harvesting System with Two-Step Sequential Energy Transfer for Photochemical Catalysis. *Angew. Chem., Int. Ed.* **2020**, *59*, 10095-10100; (c) Zhang, Q.; Catti, L.; Tiefenbacher, K., Catalysis inside the Hexameric Resorcinarene Capsule. *Acc. Chem. Res.* **2018**, *51*, 2107-2114.
- (a) Clever, G. H.; Punt, P., Cation-Anion Arrangement Patterns in Self-Assembled Pd_2L_4 and Pd_4L_8 Coordination Cages. *Acc. Chem. Res.* **2017**, *50*, 2233-2243; (b) Grommet, A. B.; Feller, M.; Klajn, R., Chemical Reactivity under Nanoconfinement. *Nat. Nanotechnol.* **2020**, *15*, 256-271; (c) Zhang, D.; Ronson, T. K.; Nitschke, J. R., Functional Capsules via Subcomponent Self-Assembly. *Acc. Chem. Res.* **2018**, *51*, 2423-2436.
- (a) Debata, N. B.; Tripathy, D.; Chand, D. K., Self-Assembled Coordination Complexes from Various Palladium(II) Components and Bidentate or Polydentate Ligands. *Coord. Chem. Rev.* **2012**, *256*, 1831-1945; (b) Chen, L.-J.; Yang, H.-B., Construction of Stimuli-Responsive Functional Materials via Hierarchical Self-Assembly Involving Coordination Interactions. *Acc. Chem. Res.* **2018**, *51*, 2699-2710; (c) Zhang, Y.-Y.; Gao, W.-X.; Lin, L.; Jin, G.-X., Recent Advances in the Construction and Applications of Heterometallic Macrocycles and Cages. *Coord. Chem. Rev.* **2017**, *344*, 323-344.

10. Albrecht, M., "Let's Twist Again"-Double-Stranded, Triple-Stranded, and Circular Helicates. *Chem. Rev.* **2001**, *101*, 3457-3497.
11. Lewis, J. E. M.; Beer, P. D.; Loeb, S. J.; Goldup, S. M., Metal Ions in the Synthesis of Interlocked Molecules and Materials. *Chem. Soc. Rev.* **2017**, *46*, 2577-2591.
12. (a) Forgan, R. S.; Sauvage, J.-P.; Stoddart, J. F., Chemical Topology: Complex Molecular Knots, Links, and Entanglements. *Chem. Rev.* **2011**, *111*, 5434-5464; (b) Leigh, D. A.; Schaefelberger, F.; Pirvu, L.; Stenlid, J. H.; August, D. P.; Segard, J., Tying different knots in a molecular strand. *Nature* **2020**, *584*, 562-568; (c) Sauvage, J.-P., From Chemical Topology to Molecular Machines (Nobel Lecture). *Angew. Chem., Int. Ed.* **2017**, *56*, 11080-11093.
13. (a) Seidel, S. R.; Stang, P. J., High-Symmetry Coordination Cages via Self-Assembly. *Acc. Chem. Res.* **2002**, *35*, 972-983; (b) Shi, Q.; Zhou, X.; Yuan, W.; Su, X.; Neniškis, A.; Wei, X.; Taujenis, L.; Snarskis, G.; Ward, J. S.; Rissanen, K.; de Mendoza, J.; Orentas, E., Selective Formation of S_4 - and T -Symmetric Supramolecular Tetrahedral Cages and Helicates in Polar Media Assembled via Cooperative Action of Coordination and Hydrogen Bonds. *J. Am. Chem. Soc.* **2020**, *142*, 3658-3670; (c) Chakraborty, S.; Newkome, G. R., Terpyridine-Based Metallosupramolecular Constructs: Tailored Monomers to Precise 2D-Motifs and 3D-Metallocages. *Chem. Soc. Rev.* **2018**, *47*, 3991-4016; (d) Li, K.; Zhang, L.-Y.; Yan, C.; Wei, S.-C.; Pan, M.; Zhang, L.; Su, C.-Y., Stepwise Assembly of $\text{Pd}_6(\text{RuL}_3)_8$ Nanoscale Rhombododecahedral Metal-Organic Cages via Metalloligand Strategy for Guest Trapping and Protection. *J. Am. Chem. Soc.* **2014**, *136*, 4456-4459; (e) Ramsay, W. J.; Ronson, T. K.; Clegg, J. K.; Nitschke, J. R., Bidirectional Regulation of Halide Binding in a Heterometallic Supramolecular Cube. *Angew. Chem., Int. Ed.* **2013**, *52*, 13439-13443; (f) Ramsay, W. J.; Nitschke, J. R., Two Distinct Allosteric Active Sites Regulate Guest Binding Within a $\text{Fe}_8\text{Mo}_{12}^{16+}$ Cubic Receptor. *J. Am. Chem. Soc.* **2014**, *136*, 7038-7043; (g) Yang, Y.; Jia, J.-H.; Pei, X.-L.; Zheng, H.; Nan, Z.-A.; Wang, Q.-M., Diastereoselective Synthesis of O Symmetric Heterometallic Cubic Cages. *Chem. Commun.* **2015**, *51*, 3804-3807; (h) Hardy, M.; Lützen, A., Better Together: Functional Heterobimetallic Macrocyclic and Cage-like Assemblies. *Chem. - Eur. J.* **2020**, *26*, 13332-13346.
14. Although the terms 'capsule' and 'cage' are used interchangeably in the literature, we favor the former as it is associated more with protection and less with imprisonment. (a) Preston, D.; Sutton, J. J.; Gordon, K. C.; Crowley, J. D., A Nona-nuclear Heterometallic Pd_3Pt_6 "Donut"-Shaped Cage: Molecular Recognition and Photocatalysis. *Angew. Chem., Int. Ed.* **2018**, *57*, 8659-8663; (b) Spicer, R. L.; Stergiou, A. D.; Young, T. A.; Duarte, F.; Symes, M. D.; Lusby, P. J., Host-Guest-Induced Electron Transfer Triggers Radical-Cation Catalysis. *J. Am. Chem. Soc.* **2020**, *142*, 2134-2139; (c) Saha, R.; Devaraj, A.; Bhattacharyya, S.; Das, S.; Zangrando, E.; Mukherjee, P. S., Unusual Behavior of Donor-Acceptor Stenhouse Adducts in Confined Space of a Water-Soluble Pd^{II}_8 Molecular Vessel. *J. Am. Chem. Soc.* **2019**, *141*, 8638-8645; (d) Yoshizawa, M.; Catti, L., Bent Anthracene Dimers as Versatile Building Blocks for Supramolecular Capsules. *Acc. Chem. Res.* **2019**, *52*, 2392-2404; (e) Cullen, W.; Misuraca, M. C.; Hunter, C. A.; Williams, N. H.; Ward, M. D., Highly Efficient Catalysis of the Kemp Elimination in the Cavity of a Cubic Coordination Cage. *Nat. Chem.* **2016**, *8*, 231-236; (f) Li, Y.; An, Y.-Y.; Fan, J.-Z.; Liu, X.-X.; Li, X.; Hahn, F. E.; Wang, Y.-Y.; Han, Y.-F., Strategy for the Construction of Diverse Poly-NHC-Derived Assemblies and Their Photoinduced Transformations. *Angew. Chem., Int. Ed.* **2020**, *59*, 10073-10080.
15. (a) Büenzli, J.-C. G.; Piguet, C., Lanthanide-Containing Molecular and Supramolecular Polymetallic Functional Assemblies. *Chem. Rev.* **2002**, *102*, 1897-1928; (b) Barry, D. E.; Caffrey, D. F.; Gunnlaugsson, T., Lanthanide-Directed Synthesis of Luminescent Self-Assembly Supramolecular Structures and Mechanically Bonded Systems from Acyclic Coordinating Organic Ligands. *Chem. Soc. Rev.* **2016**, *45*, 3244-3274; (c) Zhang, G.; Gil-Ramírez, G.; Markevicius, A.; Browne, C.; Vitorica-Yrezabal, I. J.; Leigh, D. A., Lanthanide Template Synthesis of Trefoil Knots of Single Handedness. *J. Am. Chem. Soc.* **2015**, *137*, 10437-10442; (d) Zinna, F.; Di Bari, L., Lanthanide Circularly Polarized Luminescence: Bases and Applications. *Chirality* **2015**, *27*, 1-13.
16. (a) El Aroussi, B.; Zebret, S.; Besnard, C.; Perrotet, P.; Hamacek, J., Rational Design of a Ternary Supramolecular System: Self-Assembly of Pentanuclear Lanthanide Helicates. *J. Am. Chem. Soc.* **2011**, *133*, 10764-10767; (b) Hamacek, J.; Poggiali, D.; Zebret, S.; El Aroussi, B.; Schneider, M. W.; Mastalerz, M., Building Large Supramolecular Nanocapsules with Europium Cations. *Chem. Commun.* **2012**, *48*, 1281-1283.
17. (a) Wang, J.; He, C.; Wu, P.; Wang, J.; Duan, C., An Amide-Containing Metal-Organic Tetrahedron Responding to a Spin-Trapping Reaction in a Fluorescent Enhancement Manner for Biological Imaging of NO in Living Cells. *J. Am. Chem. Soc.* **2011**, *133*, 12402-12405; (b) Zhao, L.; Qu, S.; He, C.; Zhang, R.; Duan, C., Face-Driven Octanuclear Cerium(IV) Luminescence Polyhedra: Synthesis and Luminescent Sensing Natural Saccharides. *Chem. Commun.* **2011**, *47*, 9387-9389.
18. (a) Li, X.-Z.; Zhou, L.-P.; Yan, L.-L.; Yuan, D.-Q.; Lin, C.-S.; Sun, Q.-F., Evolution of Luminescent Supramolecular Lanthanide M_{2n}Ln_3 Complexes from Helicates and Tetrahedra to Cubes. *J. Am. Chem. Soc.* **2017**, *139*, 8237-8244; (b) Yan, L.-L.; Tan, C.-H.; Zhang, G.-L.; Zhou, L.-P.; Bünzli, J.-C.; Sun, Q.-F., Stereocontrolled Self-Assembly and Self-Sorting of Luminescent Europium Tetrahedral Cages. *J. Am. Chem. Soc.* **2015**, *137*, 8550-8555; (c) Li, X.-Z.; Zhou, L.-P.; Yan, L.-L.; Dong, Y.-M.; Bai, Z.-L.; Sun, X.-Q.; Diwu, J.; Wang, S.; Bünzli, J.-C.; Sun, Q.-F., A Supramolecular Lanthanide Separation Approach Based on Multivalent Cooperative Enhancement of Metal Ion Selectivity. *Nat. Commun.* **2018**, *9*, 1-10; (d) Wang, Z.; He, L.; Liu, B.; Zhou, L.-P.; Cai, L.-X.; Hu, S.-J.; Li, X.-Z.; Li, Z.; Chen, T.; Li, X.; Sun, Q.-F., Coordination-Assembled Water-Soluble Anionic Lanthanide Organic Polyhedra for Luminescent Labeling and Magnetic Resonance Imaging. *J. Am. Chem. Soc.* **2020**, *142*, 16409-16419.
19. (a) Aboshyan-Sorgho, L.; Nozary, H.; Aebischer, A.; Bünzli, J.-C. G.; Morgantini, P.-Y.; Kittilstved, K. R.; Hauser, A.; Eliseeva, S. V.; Petoud, S.; Piguet, C., Optimizing Millisecond Time Scale Near-Infrared Emission in Polynuclear Chrome(III)-Lanthanide(III) Complexes. *J. Am. Chem. Soc.* **2012**, *134*, 12675-12684; (b) Glover, P. B.; Ashton, P. R.; Childs, L. J.; Rodger, A.; Kercher, M.; Williams, R. M.; De Cola, L.; Pikramenou, Z., Hairpin-Shaped Heterometallic Luminescent Lanthanide Complexes for DNA Intercalative Recognition. *J. Am. Chem. Soc.* **2003**, *125*, 9918-9919; (c) Wang, Z.; Zhou, L.-P.; Zhao, T.-H.; Cai, L.-X.; Guo, X.-Q.; Duan, P.-F.; Sun, Q.-F., Hierarchical Self-Assembly and Chiroptical Studies of Luminescent 4d-4f Cages. *Inorg. Chem.* **2018**, *57*, 7982-7992.
20. Hu, X.-L.; Sun, C.-Y.; Qin, C.; Wang, X.-L.; Wang, H.-N.; Zhou, E.-L.; Li, W.-E.; Su, Z.-M., Iodine-Templated Assembly of Unprecedented 3d-4f Metal-Organic Frameworks as Photocatalysts for Hydrogen Generation. *Chem. Commun.* **2013**, *49*, 3564-3566.
21. Wu, Z.-L.; Dong, J.; Ni, W.-Y.; Zhang, B.-W.; Cui, J.-Z.; Zhao, B., Unique Chiral Interpenetrating d-f Heterometallic MOFs as Luminescent Sensors. *Inorg. Chem.* **2015**, *54*, 5266-5272.
22. (a) Peng, J.-B.; Zhang, Q.-C.; Kong, X.-J.; Zheng, Y.-Z.; Ren, Y.-P.; Long, L.-S.; Huang, R.-B.; Zheng, L.-S.; Zheng, Z., High-Nuclearity 3d-4f Clusters as Enhanced Magnetic Coolers and Molecular Magnets. *J. Am. Chem. Soc.* **2012**, *134*, 3314-3317; (b) Shi, P.-F.; Xiong, G.; Zhao, B.; Zhang, Z.-Y.; Cheng, P., Anion-Induced Changes of Structure Interpenetration and Magnetic Properties in 3D Dy-Cu Metal-Organic Frameworks. *Chem. Commun.* **2013**, *49*, 2338-2340.
23. (a) Greenfield, J. L.; Rizzuto, F. J.; Goldberga, I.; Nitschke, J. R., Self-Assembly of Conjugated Metallopolymers with Tunable Length and Controlled Regiochemistry. *Angew. Chem., Int. Ed.* **2017**,

- 56, 7541-7545; (b) Greenfield, J. L.; Di, A. M.; Nitschke, J. R.; Evans, E. W.; Di, N. D.; Friend, R. H., Unraveling Mechanisms of Chiral Induction in Double-Helical Metallopolymers. *J. Am. Chem. Soc.* **2018**, *140*, 10344-10353.
24. (a) Avram, L.; Cohen, Y., Diffusion NMR of Molecular Cages and Capsules. *Chem. Soc. Rev.* **2015**, *44*, 586-602; (b) Macchioni, A.; Ciancaleoni, G.; Zuccaccia, C.; Zuccaccia, D., Determining Accurate Molecular Sizes in Solution Through NMR Diffusion Spectroscopy. *Chem. Soc. Rev.* **2008**, *37*, 479-489.
25. Ruiz-Martínez, A.; Alvarez, S., Stereochemistry of Compounds with Coordination Number Ten. *Chem. Eur. J.* **2009**, *15*, 7470-7480.
26. Kleywegt, G. J.; Jones, T. A., Detection, Delineation, Measurement and Display of Cavities in Macromolecular Structures. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* **1994**, *D50*, 178-185.
27. The ligands capping the faces of **1** might adopt clockwise (C) or anticlockwise (A) orientations, which are diastereotopic with respect to the Λ or Δ handedness of the Zn^{II} vertices. The combination of these two stereochemical elements can thus produce diastereomeric pairs of enantiomers ($\text{C}_4\Lambda_4/\text{A}_4\Delta_4$ and $\Lambda_4/\text{C}_4\Delta_4$), as observed previously in the case of a triazatruxene-based ligand. Both solution (NOESY) and solid-state (X-ray) data are consistent with the formation of only the enantiomers $\text{C}_4\Lambda_4/\text{A}_4\Delta_4$, however. (a) Zhang, D.; Ronson, T. K.; Güryel, S.; Thoburn, J. D.; Wales, D. J.; Nitschke, J. R., Temperature Controls Guest Uptake and Release from Zn_4L_4 Tetrahedra. *J. Am. Chem. Soc.* **2019**, *141*, 14534-14538; (b) Zhang, D.; Ronson, T. K.; Greenfield, J. L.; Brotin, T.; Berthault, P.; Leonce, E.; Zhu, J.-L.; Xu, L.; Nitschke, J. R., Enantiopure $[\text{Cs}^+/\text{Xe}^- \text{Cryptophane}]^- \text{Fe}^{\text{II}}_4\text{L}_4$ Hierarchical Superstructures. *J. Am. Chem. Soc.* **2019**, *141*, 8339-8345.
28. (a) Takezawa, H.; Murase, T.; Resnati, G.; Metrangolo, P.; Fujita, M., Recognition of Polyfluorinated Compounds Through Self-Aggregation in a Cavity. *J. Am. Chem. Soc.* **2014**, *136*, 1786-1788; (b) Ercolani, G., Assessment of Cooperativity in Self-Assembly. *J. Am. Chem. Soc.* **2003**, *125*, 16097-16103.
29. These association constants are unitless, because each guest is exchanged for another. Hence, the number of species is the same on both sides of the equilibrium, resulting in a dimensionless K_a .