

**Association between Childhood Psychiatric Disorders and Psychotic Experiences in  
Adolescence: a population-based longitudinal study**

**Caroline Siebald<sup>1\*</sup>, Golam M Khandaker<sup>1,2\*</sup>, Stanley Zammit<sup>3,4</sup>, Glyn Lewis<sup>5</sup>, Peter B  
Jones<sup>1,2</sup>**

\* Joint first authors

<sup>1</sup> Department of Psychiatry, University of Cambridge, UK

<sup>2</sup> Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge, UK

<sup>3</sup> Centre for Mental Health, Addiction and Suicide Research, School of Social and  
Community Medicine, University of Bristol, UK

<sup>4</sup> Institute of Psychological Medicine and Clinical Neurosciences, MRC Centre for  
Neuropsychiatric Genetics and Genomics, Cardiff University, UK

<sup>5</sup> Division of Psychiatry, University College London, London, UK

**Corresponding Author:** Dr Golam Khandaker, Department of Psychiatry, Box 189,  
Cambridge Biomedical Campus, Cambridge CB2 2QQ, UK. Email:

[gmk24@medschl.cam.ac.uk](mailto:gmk24@medschl.cam.ac.uk)

**Word count:**

**Abstract: 248**

**Article: 3,286**

**Tables and Figures: 4+1**

## **Abstract**

**Background:** Adolescent psychotic experiences (PEs) are common, and are associated with both psychotic and non-psychotic illnesses. In order to examine psychopathological and cognitive antecedents of adolescent PEs, we have conducted a longitudinal study of common childhood psychiatric disorders and subsequent adolescent PEs in the population-based prospective ALSPAC birth cohort.

**Method:** Depression, anxiety, attention deficit hyperactivity disorder, oppositional defiant or conduct disorder, and pervasive developmental disorder were diagnosed according to DSM-IV criteria in 8,253 participants at age 8 years. IQ was assessed by WISC-III also at 8 years. PEs, depressive and anxiety symptoms were assessed at 13 years. Logistic regression calculated odds ratio (OR) for PEs at 13 years associated with psychiatric disorders at 8 years. Linear regression calculated mean difference in IQ between groups with and without psychiatric disorder. Mediating effects of IQ, mood and anxiety symptoms on the psychiatric disorder-PEs relationship were examined.

**Results:** In total, 599 children were assessed to have a DSM-IV psychiatric disorder at 8 years (7.2%). These children compared with those without any psychiatric disorder performed worse on all measures of IQ; adjusted mean difference in total IQ -6.17 (95% CI, -7.86, -4.48). Childhood psychiatric disorders were associated with PEs subsequently in adolescence; adjusted OR 1.96 (95% CI, 1.47-2.68). The association between psychiatric disorder and subsequent PEs was partly mediated by, independently, IQ deficit at 8 years and depressive and anxiety symptoms at 13 years.

**Conclusions:** The findings indicate adolescent PEs are associated with general cognitive ability and past and present psychopathological factors.

**Key words:** Psychotic Experiences; Psychotic Symptoms; Childhood; Adolescence; Mental Disorder; Intelligence; IQ; Attention Deficit Hyperactivity Disorder, Oppositional Defiant Disorder; Conduct Disorder; Pervasive Developmental Disorder; Neurodevelopmental Disorders; Depression; Anxiety; Cohort Study

## **1. Introduction**

Psychotic experiences (PEs) are common in the general population [1], especially during childhood and adolescence [2]. Early-life PEs are associated with increased risk of psychosis in adulthood [3, 4] as well as a number of established risk factors for schizophrenia including family history of psychosis, cannabis use, maltreatment, advanced paternal age, IQ deficit, prenatal and childhood infection, and other immunological factors [5-15]. However, recent studies suggest PEs can no longer be regarded as having predictive specificity for psychotic disorders subsequently in adulthood [16, 17]. PEs are associated with a range of non-psychotic common mental disorders both cross-sectionally and longitudinally. PEs in adolescence or adulthood are associated with concurrent anxiety, depression, obsessive compulsive disorder, borderline personality disorder, poor functioning, self-harm and suicidal behaviour [18-23]. These findings are consistent with statistical modelling of the underlying structure of psychiatric symptoms in two birth cohorts which reveal that psychotic phenomena co-occur with depression and anxiety [24, 25], and may be a marker of the severity of mental ill health in a single, unitary dimension of common mental distress in young people [25].

Longitudinal studies suggest PEs are associated with a range of psychiatric disorders subsequently in adulthood including obsessive compulsive disorder, social phobia, dysthymia, bipolar disorder, post-traumatic stress disorder, and suicidality [16, 17]. Persistent childhood PEs are associated with subsequent internalising and externalising psychopathology in the general population [26]. Reports from the Avon Longitudinal Study of Parents and Children (ALSPAC) birth cohort [27, 28], including our own [5], have shown an association between childhood neurodevelopmental disorders and risk of PEs in early-adolescence, which is partly mediated by childhood IQ deficit [5]. However, to our

knowledge no population-based longitudinal study has examined the relationship between common mental disorders of childhood diagnosed according to clinical criteria and risk of PEs subsequently in adolescence. Building on our previous work we have examined psychopathological and cognitive antecedents of adolescent PEs using a broader range of childhood psychiatric disorders as predictors of subsequent PEs in early-adolescence in the ALSPAC birth cohort. We hypothesised psychiatric disorders at age 8 years would be associated with increased risk of PEs at age 13 years. The use of clinical diagnoses of depression, anxiety, oppositional defiant/conduct disorder, attention deficit hyperactivity disorder and pervasive developmental disorder at age 8 years defined according to DSM-IV criteria as predictors of PEs at age 13 years – as opposed to parent-reported neurodevelopmental disorders at age 10 years used in the previous study [5] – makes the current analysis unique. We have also examined mediating effects of IQ at 8 years, and depressive and anxiety symptoms at 13 years on the relationship between psychiatric disorder at 8 years and PEs at 13 years.

## **2. Method**

### **2.1 Sample**

The ALSPAC birth cohort is based on all pregnant women resident in the county of Avon, a geographically defined region in the southwest of England, with expected dates of delivery between April 1991 and December 1992 (<http://www.bristol.ac.uk/alspac/>). The initial ALSPAC cohort consisted of 14,062 live births and 13,988 infants still alive at 12 months [29, 30]. Avon included both urban and rural areas, and the population was broadly representative of all children in the UK. The parents completed regular postal questionnaires about all aspects of their child's health and development since birth. Since the age of 7 years the children attended an annual assessment clinic during which they participated in a range of

face-to-face interviews and physical tests. The current study is based on 8,253 individuals who were assessed for psychiatric disorders at age 8 years. The numbers of individuals with data on IQ at 8 years and psychiatric symptoms at 13 years vary as these tests were completed by different numbers of people.

Ethical approval for the study was obtained from ALSPAC Ethics and Law Committee and the Local Research Ethics Committees.

## **2.2 Assessment of psychiatric disorders at 8 years**

Psychiatric disorders were assessed at age 8 years using the parent version of the Development and Well-Being Assessment (DAWBA) [31] and were coded according to DSM-IV criteria by two experienced psychiatrists [32]. The DAWBA consists of a package of questionnaires that are aimed at establishing the presence of relatively common emotional, behavioural, and hyperactivity disorders in children. It has been validated for use in epidemiological studies in the UK, where it has been used for the child and adolescent mental health survey [33], and internationally [34, 35].

A postal questionnaire containing the parent-version of the DAWBA was sent to mothers when the study child was on average about 8 years old. The parent-completed DAWBA-questionnaire constituted the primary source of information for the diagnosis of psychiatric disorders. Where available, complementary sources of information on a child's emotional and behavioural characteristics were consulted, which included: (i) the teacher version of the DAWBA that addressed potential hyperactivity and conduct disorder at 8 years; (ii) the teacher version of the Strengths and Difficulties Questionnaire (SDQ) at 8 years [36]; (iii) parent completed SDQ at 7 years; (iv) basic reading and spelling competency of the child at

age 7 years; (v) IQ at age 8 years using the Wechsler Intelligence Scale for Children (WISC III, 3rd UK edition) [37]. Psychiatric diagnoses were thus based on a wealth of developmental data gathered from different sources. Diagnoses were made for attention deficit hyperactivity disorder (ADHD), oppositional or conduct disorder (OpCD), pervasive developmental disorder (PDD), anxiety disorder, and depressive disorder. The presence of any DSM-IV psychiatric disorder at 8 years, coded as a single binary variable, was used as the main predictor. Those without a diagnosis were included in the comparison group. Additional analyses were carried out using individual diagnostic categories as predictors, which included all participants without the particular diagnosis in the comparison group. For example, children with depression who did not have ADHD were included in the comparison group for the analyses of ADHD, and so on.

### **2.3 Assessment of IQ at 8 years**

Full scale, verbal, and performance IQ were measured by the WISC III, 3rd UK edition [37]. A shortened version of the test was applied by trained psychologists, whereby alternate items (always starting with item number 1 in the standard form) were used for all ten subtests with the exception of the coding subtest which was administered in its standard form. Use of the shortened version reduced the length of assessment so the children were less likely to tire. This approach has been successfully used in other studies [5, 38, 39]. IQ data obtained using this method have shown robust correlations with neurodevelopmental disorders, and other concurrent neurocognitive measures such as working memory, short-term memory, and sociodemographic factors such as social class [5].

## **2.4 Assessment of psychotic experiences at 13 years**

Psychotic experiences (PEs) were identified through the face-to-face, semi-structured Psychosis-Like Symptom Interview (PLIKSi) conducted by trained psychology graduates in assessment clinics. The PLIKSi has good inter-rater reliability ( $\kappa=0.7$ ) [9]. It comprised 12 'core' questions derived from the Diagnostic Interview Schedule for Children-IV (DISC-IV) [40], and coded according to the rating rules for the Schedules for Clinical Assessment in Neuropsychiatry version 2.0 (SCAN 2.0) [41]. It included key symptoms covering the three main domains of positive psychotic symptoms: hallucinations (visual and auditory); delusions (spied on, persecution, thought being read, reference, control, grandiose ability, and other unspecified); and experiences of thought interference (insertion, withdrawal, and broadcasting). The observer based rating for the presence of any PEs (suspected or definite) in the past six months was used as the outcome. The group with PEs was compared with the rest of the cohort.

## **2.5 Assessment for depressive and anxiety symptoms at 13 years**

Symptoms of depression and anxiety were measured at age 13 years using the short version of the mood and feelings questionnaire (MFQ) completed by the participants [36]. The short MFQ is a validated tool widely used in epidemiological studies [42, 43]. It includes 13 items covering core symptoms of depression and anxiety experienced in the past two weeks. Each item is scored zero (not true), one (sometimes true) or two (true) giving a total score of 0-26.

## **2.6 Statistical analysis**

Baseline characteristics between the groups with and without any psychiatric disorder at 8 years were compared by independent sample t-test (continuous data) and chi-squared test (categorical data). Linear regression calculated the mean difference (95% CI) in IQ scores at



8 years between those with and without any psychiatric disorder. Binary logistic regression calculated the odds ratio (OR) for PEs at 13 years in the group with any psychiatric disorder at 8 years compared with those without. All regression models were controlled for age at assessment, sex, mother's social class, and highest educational level.

Mediating effect of IQ at 8 years on the association between psychiatric disorders at 8 years and PEs at 13 years was examined two ways. First, separate regression models assessed the associations between: (1) exposure (any psychiatric disorder) and outcome (PEs); (2) exposure and mediator (total IQ score); (3) mediator and outcome; (4) exposure and outcome controlling for mediator. We expected, in the final step, the exposure-outcome relationship would be attenuated (partial mediation) or eliminated (complete mediation). In case of partial mediation, the extent to which the psychiatric disorder-PEs estimate was attenuated after inclusion of total IQ (the mediator) was calculated. Secondly, mediation analyses were conducted using the STATA package 'paramed'. It calculated the controlled direct effect (i.e. the effect of any psychiatric disorder (exposure) on the risk of PEs (outcome) after adjusting for IQ); natural direct effect (i.e. the effect on the risk of PEs of a change in IQ among unexposed individuals compared with exposed individuals); total effect (the product of controlled direct effect and natural direct effect). Mediating effect of total MFQ score at 13 years on the association between psychiatric disorders at 8 years and subsequent PEs at 13 years was examined using the same approach for IQ described above. The 'paramed' approach was omitted as MFQ scores were not normally distributed. Instead, the association between psychiatric disorders at 8 years and PEs at 13 years were examined by stratifying the sample according to tertiles of MFQ score.

### **3. Results**

#### **3.1 Psychiatric disorders at 8 years and baseline characteristics**

At 8 years, 599 out of total 8,253 children (7.2%) were assessed to have a DSM-IV psychiatric disorder (depression, anxiety, oppositional defiant/conduct disorder, ADHD, PDD). Out of these, 465 children had one (5.6%), 136 children had two (1.3%), and 28 children had three or more psychiatric disorders (0.3%). Oppositional/conduct disorder was the most common (Figure 1). Psychiatric disorders were more common in boys (Table 1).

--- Please insert Table 1 and Figure 1 here ---

#### **3.2 Association with IQ at 8 years**

Compared with children with no psychiatric disorder, children with a psychiatric disorder as a group performed worse on all measures of IQ at 8 years (Table 2). The results remained similar after adjusting for a number of potential confounders. We also examined total IQ at 8 years separately in five specific psychiatric disorders. Compared with the rest of the cohort, mean total IQ was lower in all disorders except depression. The largest deficit in total IQ was observed for pervasive developmental disorder.

--- Please insert Table 2 here ---

#### **3.3 Association with psychotic experiences at 13 years**

Data on psychiatric disorders at 8 years and the outcome of PEs at 13 years were available for 5,528 individuals. In the group with psychiatric disorders 75 developed PEs (20.8%), while in the group with no psychiatric disorder 673 developed PEs (13.0%). There was nearly two-fold increased risk of PEs at 13 years for individuals with any psychiatric disorder at 8 years

compared with those without. The association persisted after adjusting for potential confounders (Table 3). Separate analyses using specific psychiatric disorders showed all childhood disorders were individually associated with PEs in adolescence; the highest OR was observed for pervasive developmental disorder (Table 3).

--- Please insert Table 3 here ---

### **3.4 Effect of IQ at 8 years**

Step by step regression analyses showed about a fifth (21%) of the association between psychiatric disorders at 8 years and PEs at 13 years could be accounted for by deficit in total IQ at 8 years among children with psychiatric disorders. The association between psychiatric disorders at 8 years and PEs at 13 years (OR 1.76; 95% CI, 1.35-2.30) was attenuated after including total IQ score at 8 years as a predictor (OR 1.56; 95% CI, 1.16-2.10). This finding was consistent with additional mediation analyses, which revealed that a small but significant part of the relationship between psychiatric disorders and subsequent PEs could be accounted for by IQ. The overall adjusted OR for PEs at 13 years for psychiatric disorders at 8 years was 1.66 (95% CI, 1.19-2.33) after taking into account sociodemographic factors as well as total MFQ score at 13 years (i.e. total effect), which was composed in part by IQ deficit among children with a psychiatric disorder; adjusted OR 1.05 (95% CI, 1.01-1.10) (Table 4).

--- Please insert Table 4 here ---

### **3.5 Effect of MFQ score at 13 years**

Step by step regression analyses showed about 13% of the association between psychiatric disorders at 8 years and PEs at 13 years could be accounted for by higher MFQ scores at 13

years among children with psychiatric disorders. The association between psychiatric disorders at 8 years and PEs at 13 years (OR 1.76; 95% CI, 1.35-2.30) was attenuated after including total MFQ score at 13 years as a predictor (OR 1.63; 95% CI, 1.23-2.17). Further examination of the association between psychiatric disorders at 8 years and PEs at 13 years was carried out dividing the sample into thirds according to tertiles of MFQ score. It showed that the association between psychiatric disorders at 8 years and PEs at 13 years was present only in participants in the top third of MFQ scores (OR 1.65; 95% CI, 1.14-2.40). The ORs for those in the bottom (1.74; 95% CI, 0.95-3.16) and middle (1.53; 95% CI, 0.82-2.84) thirds of MFQ scores were not statistically significant.

#### **4. Discussion**

The findings suggest that childhood psychiatric disorders diagnosed according to DSM-IV criteria at 8 years are associated with increased risk of psychotic experiences at 13 years, which persists after controlling for potential confounders such as age, sex, ethnicity, maternal social class and educational level. IQ at 8 years and depressive and anxiety symptoms at 13 years both independently mediated part of the relationship between childhood psychiatric disorders and subsequent PEs in adolescence. Although all childhood psychiatric disorders were associated with adolescent PEs, the highest OR was observed for PDD. Over half of the children with PDD developed PEs at 13 years compared with about an eighth in the comparison group. However, multivariable regression analyses for PDD lacked statistical power as reflected by wide confidence interval around the ORs because only a small number of participants met the criteria for this diagnosis. As expected, this group also had the highest deficit in IQ. These findings are consistent with the neurodevelopmental hypothesis of schizophrenia which posits abnormal neurodevelopment as a cause of the illness [44, 45]. Population-based longitudinal studies reporting impaired motor, cognitive, language and

social development during infancy, childhood and adolescence in future cases of schizophrenia support the neurodevelopmental hypothesis [46-53]. Meta-analysis of population-based longitudinal studies have confirmed a premorbid IQ deficit of about half a standard deviation in future cases of schizophrenia compared with healthy controls [54]. The findings are also consistent with three previous studies from the ALSPAC cohort reporting about two-fold increased risk of PEs in adolescence for autism, dyslexia and other neurodevelopmental disorders in childhood [5, 27, 28], although lack of an association between autistic traits and PEs has also been reported [55]. The use of clinical diagnosis of PDD as defined by DSM-IV criteria as exposure, as opposed to neurodevelopmental symptoms [27, 28] or parent-reported disorder [5] used in the previous studies, could explain a larger effect observed in the current study.

Looking at the underlying mechanism for PEs, the study shows cognitive deficit (as measured by IQ) and increased depressive and anxiety symptoms (as measured by MFQ score) both independently, in part explain the risk of adolescent PEs associated with past psychopathology. The findings are consistent with the idea that adolescent PEs have a cognitive facet to their origin. Indeed, adolescent PEs assessed at 13 years have been previously reported to be associated with lower childhood IQ at 8 years in the ALSPAC cohort [9]. Similarly, PEs at 12 years have been reported to be associated with lower IQ at 5 years in another cohort [10]. The findings are also consistent with previous population-based studies showing an association between PEs and other psychiatric symptoms such as depression and anxiety. Population-based studies have previously shown that adolescent PEs may be a marker of multiple concurrent depressive or anxiety disorders (i.e. severity of mental ill-health) or of the likelihood of clinically concerning behaviours such as suicidal thoughts or self-harm [18-21]. Moreover, item response theory analysis of the underlying

structure of psychiatric symptoms in the ALSPAC birth cohort have revealed that psychotic phenomena co-occur with depression and anxiety where they represent severity of mental ill-health in a single, unitary dimension of common mental distress in young people [25]. Thus, the current study supports the previous findings by showing that concurrent depressive and anxiety symptoms partly account for adolescent PEs.

The use of a general population-based birth cohort, large sample size, objective assessments for IQ, and PEs are particular strengths of this study. Diagnoses of psychiatric disorders at 8 years were based on primarily parent completed DAWBA questionnaire. Although where available complementary sources of information on a child's emotional and behavioural characteristics were consulted (see Methods) the use of parent reported data as opposed to direct assessment to establish a diagnosis is a limitation of this study. We did not have data on family psychiatric history and substance use to examine whether these variables confound the association between childhood psychiatric disorder and subsequent PE in early-adolescence, which is a potential limitation of this study. Another limitation is missing data. A third of participants assessed for psychiatric diagnoses at 8 years did not attend assessment for PEs at 13 years. Presence of a psychiatric disorder in childhood was associated with missing data for PEs at follow-up. If participants with psychiatric disorder at 8 years who did not develop PEs at 13 years were more likely to be missing at follow up this would lead to a spurious over estimation of the association between childhood psychiatric disorder and subsequent PEs. However, this is unlikely. Missing data was also associated with lower maternal education and lower maternal socio-economic status. In the ALSPAC cohort both of these factors are associated with higher PEs, lower IQ, and adverse health and social outcomes. PEs were not assessed at baseline. Although we cannot be certain it's likely that PEs at 13 years were new onset phenomena in majority of subjects since baseline assessment

at age 8 years. This is because early-life PEs are transitory phenomena in most children; a 6-year follow-up of the ALSPAC birth cohort suggests about 80% of subjects with PEs in early-adolescence do not have PEs at age 18 years [3].

To our knowledge, this is one of the first population-based longitudinal studies of common childhood psychiatric disorders and subsequent PEs in early-adolescence. The findings implicate past and present psychopathological factors as well as general cognitive ability in the origin of adolescent PEs. In future, studies should use repeated assessment of PEs over the early life course in order to elucidate factors associated with persistence and remission of these symptoms. Individuals with childhood psychiatric disorders are known to be at higher risk of developing psychotic and non-psychotic illnesses during adult life. In future studies should also examine whether adolescent PEs can account for the association between childhood and adult psychiatric disorders.

## **Acknowledgment**

We are grateful to all families who took part in this study, midwives for their help in recruitment, and the whole ALSPAC team, including interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. Prof Jones acknowledges support from the Wellcome Trust (095844/Z/11/Z & 088869/Z/09/Z) and NIHR (RP-PG-0606-1335). The UK Medical Research Council and the Wellcome Trust grant ref 092731 and the University of Bristol provide core support for the ALSPAC cohort.

## **Declaration of interest**

None of the authors has any conflicts of interest to declare. Prof Jones received an honorarium that he donated to his department from Roche for taking part in an advisory board to advise on education about schizophrenia to psychiatrists. He directs the National Institute for Health Research Collaborations for Leadership in Applied Health Research and Care for Cambridgeshire and Peterborough (CLAHRC-CP) of which this work forms part.



## Reference

1. van Os, J., et al., *A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness-persistence-impairment model of psychotic disorder*. Psychol Med, 2009. **39**(2): p. 179-95.
2. Kelleher, I., et al., *Prevalence of psychotic symptoms in childhood and adolescence: a systematic review and meta-analysis of population-based studies*. Psychol Med, 2012. **42**(9): p. 1857-1863.
3. Zammit, S., et al., *Psychotic Experiences and Psychotic Disorders at Age 18 in Relation to Psychotic Experiences at Age 12 in a Longitudinal Population-Based Cohort Study*. Am J Psychiatry, 2013. **170**(7): p. 742-750.
4. Poulton, R., et al., *Children's self-reported psychotic symptoms and adult schizophreniform disorder: a 15-year longitudinal study*. Arch Gen Psychiatry, 2000. **57**(11): p. 1053-8.
5. Khandaker, G.M., et al., *A population-based longitudinal study of childhood neurodevelopmental disorders, IQ and subsequent risk of psychotic experiences in adolescence*. Psychol Med, 2014. **44**(15): p. 3229-3238.
6. Khandaker, G.M., et al., *Childhood Epstein-Barr Virus infection and subsequent risk of psychotic experiences in adolescence: A population-based prospective serological study*. Schizophr Res, 2014. **158**(1-3): p. 19-24.
7. Khandaker, G.M., et al., *A population-based prospective birth cohort study of childhood neurocognitive and psychological functioning in healthy survivors of early life meningitis*. Ann Epidemiol, 2015. **25**(4): p. 236-42.
8. Zammit, S., et al., *Investigating whether adverse prenatal and perinatal events are associated with non-clinical psychotic symptoms at age 12 years in the ALSPAC birth cohort*. Psychol Med, 2009. **39**(9): p. 1457-67.


9. Horwood, J., et al., *IQ and non-clinical psychotic symptoms in 12-year-olds: results from the ALSPAC birth cohort*. Br J Psychiatry, 2008. **193**(3): p. 185-91.
10. Polanczyk, G., et al., *Etiological and clinical features of childhood psychotic symptoms: results from a birth cohort*. Arch Gen Psychiatry, 2010. **67**(4): p. 328-38.
11. Morgan, C., et al., *Adversity, cannabis use and psychotic experiences: evidence of cumulative and synergistic effects*. Br J Psychiatry, 2014. **204**: p. 346-53.
12. Wigman, J.T., et al., *Early trauma and familial risk in the development of the extended psychosis phenotype in adolescence*. Acta Psychiatr Scand, 2012. **126**(4): p. 266-73.
13. Janssen, I., et al., *Childhood abuse as a risk factor for psychotic experiences*. Acta Psychiatr Scand, 2004. **109**(1): p. 38-45.
14. Foutz, J. and B. Mezuk, *Advanced paternal age and risk of psychotic-like symptoms in adult offspring*. Schizophr Res, 2015. **165**(2-3): p. 123-7.
15. Khandaker, G.M., et al., *Association of serum interleukin 6 and C-reactive protein in childhood with depression and psychosis in young adult life: a population-based longitudinal study*. JAMA Psychiatry, 2014. **71**(10): p. 1121-8.
16. Fisher, H.L., et al., *Specificity of childhood psychotic symptoms for predicting schizophrenia by 38 years of age: a birth cohort study*. Psychol Med, 2013. **43**(10): p. 2077-86.
17. Rossler, W., et al., *Sub-clinical psychosis symptoms in young adults are risk factors for subsequent common mental disorders*. Schizophr Res, 2011. **131**(1-3): p. 18-23.
18. Kelleher, I., et al., *Clinicopathological significance of psychotic experiences in non-psychotic young people: evidence from four population-based studies*. Br J Psychiatry, 2012. **201**(1): p. 26-32.

19. Nishida, A., et al., *Associations between psychotic-like experiences and mental health status and other psychopathologies among Japanese early teens*. Schizophr Res, 2008. **99**(1-3): p. 125-33.
20. Wigman, J.T., et al., *Evidence that psychotic symptoms are prevalent in disorders of anxiety and depression, impacting on illness onset, risk, and severity--implications for diagnosis and ultra-high risk research*. Schizophr Bull, 2012. **38**(2): p. 247-57.
21. Hui, C., et al., *Psychiatric morbidity, functioning and quality of life in young people at clinical high risk for psychosis*. Schizophr Res, 2013. **148**(1-3): p. 175-80.
22. Bortolon, C. and S. Raffard, *Self-reported psychotic-like experiences in individuals with obsessive-compulsive disorder versus schizophrenia patients: characteristics and moderation role of trait anxiety*. Compr Psychiatry, 2015. **57**: p. 97-105.
23. Pearse, L.J., et al., *A study of psychotic symptoms in borderline personality disorder*. J Nerv Ment Dis, 2014. **202**(5): p. 368-71.
24. Caspi, A., et al., *The p Factor: One General Psychopathology Factor in the Structure of Psychiatric Disorders?* Clin Psychol Sci, 2014. **2**(2): p. 119-137.
25. Stochl, J., et al., *Mood, anxiety and psychotic phenomena measure a common psychopathological factor*. Psychol Med, 2015. **45**(7): p. 1483-93.
26. Downs, J.M., et al., *Persisting psychotic-like experiences are associated with both externalising and internalising psychopathology in a longitudinal general population child cohort*. Schizophr Res, 2013. **144**(1-3): p. 99-104.
27. Sullivan, S., et al., *The Association Between Autism Spectrum Disorder and Psychotic Experiences in the Avon Longitudinal Study of Parents and Children (ALSPAC) Birth Cohort*. J Am Acad Child Adolesc Psychiatry, 2013. **52**(8): p. 806-814.
28. Bevan Jones, R., et al., *The association between early autistic traits and psychotic experiences in adolescence*. Schizophr Res, 2012. **135**(1-3): p. 164-9.

29. Boyd, A., et al., *Cohort Profile: The 'Children of the 90s'--the index offspring of the Avon Longitudinal Study of Parents and Children*. *Int J Epidemiol*, 2013. **42**(1): p. 111-127.
30. Fraser, A., et al., *Cohort Profile: The Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort*. *Int J Epidemiol*, 2013. **42**(1): p. 97-110.
31. Goodman, R., et al., *The Development and Well-Being Assessment: description and initial validation of an integrated assessment of child and adolescent psychopathology*. *J Child Psychol Psychiatry*, 2000. **41**(5): p. 645-55.
32. APA, *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV. 4th ed.*, 1995, American Psychiatric Association: Washitngton D.C.
33. Ford, T., R. Goodman, and H. Meltzer, *The British Child and Adolescent Mental Health Survey 1999: the prevalence of DSM-IV disorders*. *J Am Acad Child Adolesc Psychiatry*, 2003. **42**(10): p. 1203-11.
34. Fleitlich-Bilyk, B. and R. Goodman, *Prevalence of child and adolescent psychiatric disorders in southeast Brazil*. *J Am Acad Child Adolesc Psychiatry*, 2004. **43**(6): p. 727-34.
35. Heiervang, E., A. Goodman, and R. Goodman, *The Nordic advantage in child mental health: separating health differences from reporting style in a cross-cultural comparison of psychopathology*. *J Child Psychol Psychiatry*, 2008. **49**(6): p. 678-85.
36. Angold, A., E.J. Costello, and S.C. Messer, *Development of a short questionnaire for use in epidemiological studies of depression in children and adolescents*. *International Journal of Methods in Psychiatric Research*, 1995. **5**: p. 237- 249.
37. Wechsler, D., S. Golombok, and J. Rust, *Wechsler Intelligence Scale for Children (3rd Edition) (WISC-III UK)*. 1992: The Psychological Corporation.

38. Finch, A.J. and W.B. Chihldress, *A comparison of WISC selected subtest short forms with MR children*. *Mental Retardation*, 1975. **13**: p. 20- 21.
39. Stricker, G., M. Merbaum, and P. Tangeman, *WAIS short forms, information transmission and approximation of full scale IQ*. *Journal of Clinical Psychology*, 1968. **25**: p. 170- 172.
40. Shaffer, D., et al., *NIMH Diagnostic Interview Schedule for Children Version IV (NIMH DISC-IV): description, differences from previous versions, and reliability of some common diagnoses*. *J Am Acad Child Adolesc Psychiatry*, 2000. **39**(1): p. 28-38.
41. WHO, *SCAN: Schedules for Clinical Assessment in Neuropsychiatry Version 2.0*. 1994, Geneva, Switzerland: Psychiatric Publishers International/American Psychiatric Press Inc.
42. Sharp, C., I.M. Goodyer, and T.J. Croudace, *The Short Mood and Feelings Questionnaire (SMFQ): a unidimensional item response theory and categorical data factor analysis of self-report ratings from a community sample of 7-through 11-year-old children*. *J Abnorm Child Psychol*, 2006. **34**(3): p. 379-91.
43. Thapar, A. and P. McGuffin, *Validity of the shortened Mood and Feelings Questionnaire in a community sample of children and adolescents: a preliminary research note*. *Psychiatry Res*, 1998. **81**(2): p. 259-68.
44. Weinberger, D.R., *Implications of normal brain development for the pathogenesis of schizophrenia*. *Arch Gen Psychiatry*, 1987. **44**(7): p. 660-9.
45. Murray, R.M. and S.W. Lewis, *Is schizophrenia a neurodevelopmental disorder?* *BMJ (Clinical Research Edition)*, 1987. **295**(6600): p. 681- 682.
46. Jones, P., et al., *Child development risk factors for adult schizophrenia in the British 1946 birth cohort*. *Lancet*, 1994. **344**(8934): p. 1398-402.

47. Crow, T.J., D.J. Done, and A. Sacker, *Childhood precursors of psychosis as clues to its evolutionary origins*. Eur Arch Psychiatry Clin Neurosci, 1995. **245**(2): p. 61-9.
48. Cannon, M., et al., *Evidence for early-childhood, pan-developmental impairment specific to schizophreniform disorder: results from a longitudinal birth cohort*. Arch Gen Psychiatry, 2002. **59**(5): p. 449-56.
49. Cannon, T.D., et al., *Childhood cognitive functioning in schizophrenia patients and their unaffected siblings: a prospective cohort study*. Schizophr Bull, 2000. **26**(2): p. 379-93.
50. David, A.S., et al., *IQ and risk for schizophrenia: a population-based cohort study*. Psychol Med, 1997. **27**(6): p. 1311-23.
51. Davidson, M., et al., *Behavioral and intellectual markers for schizophrenia in apparently healthy male adolescents*. Am J Psychiatry, 1999. **156**(9): p. 1328-35.
52. Isohanni, M., et al., *Early developmental milestones in adult schizophrenia and other psychoses. A 31-year follow-up of the Northern Finland 1966 Birth Cohort*. Schizophrenia Res, 2001. **52**(1-2): p. 1-19.
53. Bearden, C.E., et al., *A prospective cohort study of childhood behavioral deviance and language abnormalities as predictors of adult schizophrenia*. Schizophr Bull, 2000. **26**(2): p. 395-410.
54. Khandaker, G.M., et al., *A quantitative meta-analysis of population-based studies of premorbid intelligence and schizophrenia*. Schizophr Res, 2011. **132**(2-3): p. 220-7.
55. Taylor, M.J., et al., *A longitudinal twin study of the association between childhood autistic traits and psychotic experiences in adolescence*. Mol Autism, 2015. **6**: p. 44.



OpCD=Oppositional/ conduct disorder; ADHD=Attention deficit hyperactivity disorder; PDD=Pervasive developmental disorder

**Figure 1: Prevalence of common psychiatric disorders (DSM-IV) at 8 years in the ALSPAC birth cohort**

**Table 1: Baseline characteristics of individuals with psychiatric disorders at 8 years**

Characteristic	Psychiatric disorder	No disorder	P-value <sup>a</sup>
Total No.	599	7654	-
Male sex, no. (%)	413 (68.9)	3832 (50.1)	<0.001
Age at assessment, mean (SD), years	7.69 (0.15)	7.68 (0.14)	0.400
British white ethnicity, no. (%)	565 (98.6)	7253 (98.2)	0.819
Mother's social class, no. (%)			0.017
I (professional)	20 (4.20)	462 (7.26)	
II (intermediate)	154 (32.3)	2177 (34.2)	
III nm (skilled, non-manual)	208 (43.8)	2705 (42.5)	
III m (skilled, manual)	33 (6.93)	422 (6.63)	
IV (partly skilled)	47 (9.87)	501 (7.87)	
V (unskilled)	14 (2.94)	100 (1.57)	
Maternal highest education, no. (%)			0.643
Secondary school	93 (16.2)	1053 (14.2)	
Vocational	52 (9.10)	660 (8.90)	
O level	196 (34.2)	2607 (35.2)	
A level	149 (26.0)	1903 (25.6)	
Degree	83 (14.5)	1189 (16.04)	

<sup>a</sup> Independent sample t-test for continuous data (age), Chi-squared test for proportions (sex, ethnicity, maternal social class, and education)



**Table 2: IQ at 8 years in children with and without psychiatric disorders**

Psychiatric disorder/ IQ	Psychiatric disorder		No disorder		Mean difference (95% CI)	
	No.	Mean (SD)	No.	Mean (SD)	Unadjusted	Adjusted <sup>a</sup>
<i>Any psychiatric disorder</i>						
Total IQ	400	98.50 (18.63)	5665	105.57 (16.03)	-7.06 (-8.71, -5.42)	-6.17 (-7.86, -4.48)
Verbal IQ	402	101.96 (19.10)	5691	108.44 (16.40)	-6.48 (-8.18, -4.80)	-5.94 (-7.66, -4.21)
Performance IQ	401	94.82 (18.75)	5683	100.79 (16.76)	-5.97 (-7.68, -4.26)	-4.85 (-6.69, -3.01)
<i>Individual disorder (total IQ)</i>						
Oppositional/ conduct	175	100.22 (17.42)	5880	105.28 (16.22)	-5.07 (-7.51, -2.62)	-3.03 (-6.08, -0.98)
Anxiety	174	100.73 (17.29)	5891	105.23 (16.26)	-4.50 (-6.95, -2.04)	-4.13 (-6.63, -1.64)
ADHD	116	93.84 (21.34)	5939	105.36 (16.08)	-11.51 (-14.49, -8.54)	-9.63 (-12.73, -6.53)
Depression	27	97.63 (19.09)	6038	105.13 (16.28)	-7.50 (-13.67, -1.34)	-5.09 (-11.58, 1.39)
PDD	10	82.00 (18.48)	6055	105.14 (16.28)	-23.14 (-33.24, -13.04)	-24.47 (-35.68, -13.27)

<sup>a</sup> Adjusted for age at IQ testing, sex, ethnicity, maternal social class and highest education

**Table 3: ORs for PEs at 13 years for psychiatric disorders at 8 years**

Psychiatric disorder at 8 years	Total No.	PEs, No. (%)	OR (95% CI) for PEs at 13 years		
			Unadjusted	Model 1 <sup>a</sup>	Model 3 <sup>c</sup>
Any disorder	5528	673 (13.02)	1.76 (1.35-2.30)	1.82 (1.39-2.38)	1.96 (1.47-2.68)
Oppositional/ conduct	5519	710 (13.25)	1.70 (1.15-2.52)	1.78 (1.20-2.64)	1.81 (1.17-2.80)
Anxiety	5528	717 (13.35)	1.60 (1.07-2.38)	1.62 (1.08-2.41)	1.70 (1.10-2.61)
ADHD	5519	723 (13.35)	1.55 (0.94-2.53)	1.65 (1.00-2.72)	1.91 (1.22-3.26)
Depression	5528	741 (13.46)	3.22 (1.29-7.99)	3.21 (1.29-7.99)	3.00 (1.13-7.96)
PDD	5528	743 (13.46)	8.03 (2.15-29.99)	8.31 (2.22-31.05)	11.79 (2.77-50.13)

<sup>a</sup> Adjusted for sex

<sup>b</sup> Adjusted for age at PEs, sex, maternal social class and highest education

**Table 4: Mediation of the psychiatric disorder-PEs relationship by IQ**

Mediator	OR (95% CI) for PEs
IQ at 8 years	
Controlled direct effect <sup>a</sup>	1.58 (1.13-2.22)
Natural direct effect <sup>b</sup>	1.05 (1.01-1.10)
Total effect <sup>c</sup>	1.66 (1.19-2.33)

<sup>a</sup> The effect of any psychiatric disorder (exposure) on the risk of PEs (outcome) after adjusting for IQ (mediator)

<sup>b</sup> The effect on the risk of PEs of a change in IQ among unexposed individuals compared with exposed individuals

<sup>c</sup> The product of controlled direct effect and natural direct effect