

# Human METTL12 is a mitochondrial methyltransferase that modifies citrate synthase

Virginie F. Rhein\*, Joe Carroll\*, Shujing Ding, Ian M. Fearnley and John E. Walker

Medical Research Council Mitochondrial Biology Unit, University of Cambridge, UK

## Correspondence

J. E. Walker, Medical Research Council Mitochondrial Biology Unit, University of Cambridge, Wellcome Trust/MRC Building, Cambridge Biomedical Campus, Hills Road, Cambridge CB2 0XY, UK  
Fax: +44 1223 252715  
Tel: +44 1223 252701  
E-mail: walker@mrc-mbu.cam.ac.uk

\*These authors contributed equally

(Received 28 February 2017, revised 31 March 2017, accepted 4 April 2017, available online 27 April 2017)

doi:10.1002/1873-3468.12649

Edited by Stuart Ferguson

**The protein methylome in mammalian mitochondria has been little studied until recently. Here, we describe that lysine-368 of human citrate synthase is methylated and that the modifying enzyme, localized in the mitochondrial matrix, is methyltransferase-like protein 12 (METTL12), a member of the family of 7 $\beta$ -strand methyltransferases. Lysine-368 is near the active site of citrate synthase, but removal of methylation has no effect on its activity. In mitochondria, it is possible that some or all of the enzymes of the citric acid cycle, including citrate synthase, are organized in metabolons to facilitate the channelling of substrates between participating enzymes. Thus, possible roles for the methylation of Lys-368 are in controlling substrate channelling itself, or in influencing protein–protein interactions in the metabolon.**

**Keywords:** citrate synthase; metabolons; METTL12; mitochondria; protein methylation; substrate channelling

The methylation of lysine and arginine residues in non-histone proteins is being associated increasingly with the regulation of many different cellular activities [1,2]. Lysine residues can also be ubiquitinated and sumoylated, and the interplay between methylation, acetylation and these other post-translational modifications, at the same and neighbouring sites, adds complexity to the regulation of biological processes by post-translational modification [3–5]. Trimethylation of a lysine residue in mitochondrial apocytochrome c by Ctm1p in the

cytoplasm of *Saccharomyces cerevisiae* is well known [6,7], although the mammalian orthologue is unmodified. Methylation of yeast cytochrome c promotes its import into the mitochondrial intermembrane space of yeast mitochondria, and modulates its interaction with other proteins [8,9]. Four other major mitochondrial proteins have been reported to contain trimethylated lysines. They are citrate synthase [10] and the  $\beta$ -subunit of the electron transfer flavoprotein (ETF $\beta$ ) [11], both found in the mitochondrial matrix, the ADP/ATP

## Abbreviations

ASMTL, acetylserotonin O-methyltransferase-like; COMTD1, catechol O-methyltransferase domain-containing protein 1; Ctm1p, cytochrome c methyltransferase; ECE2, Endothelin-converting enzyme 2; EFM4, elongation factor methyltransferase 4; ETD, electron transfer dissociation; ETF $\beta$ ,  $\beta$ -subunit of the electron transfer flavoprotein; HAP1, human haploid cell line 1; HEK, human embryonic kidney; HEMK1, heme biosynthesis gene K methyltransferase family member 1; KMT2A, histone-lysine N-methyltransferase 2A; METTL10, methyltransferase-like protein 10; METTL12, methyltransferase-like protein 12; METTL13, methyltransferase-like protein 13; METTL9, methyltransferase-like protein 9; MTRF1L, mitochondrial translation release factor 1-like; MTS, mitochondrial targeting sequence; NDUFAF7, NADH dehydrogenase (ubiquinone) complex I assembly factor 7; NDUF3, NADH dehydrogenase (ubiquinone) 1 $\beta$  subcomplex subunit 3; NDUF52, NADH dehydrogenase (ubiquinone) iron-sulphur protein 2; PRDM15, positive regulatory domain-containing protein 15; PRMT8, protein arginine methyltransferase 8; RRAD1, ribosomal RNA adenine demethylase domain-containing protein 1; RSAD1, radical S-adenosyl methionine domain-containing protein 1; SAM, S-adenosyl-methionine; SET, SET domain (*Drosophila* Su(var)3-9, Enhancer of zeste and Trithorax); SETD4, SET domain-containing protein 4; TOM and TIM, translocases of the outer and inner mitochondrial membranes respectively.

translocase [12] and the c-subunit in the rotor of the ATP synthase [13], both major components of the inner mitochondrial membrane. Two additional proteins are methylated in complex I. An arginine residue in the NDUFS2 subunit is symmetrically dimethylated [14], and three histidine residues near to the N terminus of the NDUFB3 subunit are methylated to varying extents [15]. In addition to these sites characterized in proteins purified from mitochondria, other human proteins that locate to mitochondria have been reported in cell-wide methylome studies to contain methylated lysine and arginine residues [16–18].

The methylation of proteins and RNA molecules is catalysed by methyltransferases with *S*-adenosyl-methionine as the methyl donor [19,20]. Until recently, the only human methyltransferases known to be located in mitochondria were HEMK1, which methylates a glutamine residue in the mitochondrial translation release factor MTRF1L [21], and NDUFAF7, an assembly factor for complex I [22]. In a continuing systematic study of protein methylases in human mitochondria, we have shown that NDUFAF7 symmetrically dimethylates Arg-85 in the NDUFS2 subunit of complex I, and that this modification is an essential step in the assembly of the complex [23]. We have also demonstrated that METTL20 methylates lysine residues adjacent to the recognition loop of ETF $\beta$  [11]. HEMK1, NDUFAF7 and METTL20 are all members of the cell-wide protein family of 7 $\beta$ -strand methyltransferases. This family is characterized by a common core similar to a Rossmann fold with a twisted seven-stranded  $\beta$ -sheet structure and usually six associated  $\alpha$ -helices, although the number of  $\alpha$ -helices varies [19,20]. Another member of the family, NDUFAF5, also located in the matrix of mitochondria, hydroxylates Arg-73 in subunit NDUFS7 of complex I [24]. This modification is an essential early step in the assembly of the complex. Previously, we have confirmed that another 7 $\beta$ -strand methyltransferase, METTL12, is found in the matrix of the organelle [11]. Here, we demonstrate that METTL12 modifies Lys-368 of citrate synthase in an external surface region close to its catalytic site.

## Materials and methods

### Bioinformatic analyses

The presence of N-terminal mitochondrial targeting sequences (MTSs) in the 208 known and putative methyltransferases [25] was examined with MitoProt, iPSORT, TargetP and MitoFates [26–29]. The secondary structures of proteins were predicted with Jpred [30] and PSIPRED [31].

### Cell culture

Parental human embryonic kidney cells (HEK293T) and the same cells overexpressing METTL12 were grown under an atmosphere of 5% CO<sub>2</sub> at 37 °C in high glucose (25 mM) Dulbecco's modified Eagle medium (DMEM) supplemented with tetracycline-free FBS (10% v/v), penicillin (100 units·mL<sup>-1</sup>) and streptomycin (0.1 mg·mL<sup>-1</sup>). The medium also included blasticidin (15  $\mu$ g·mL<sup>-1</sup>) and Zeocin (100  $\mu$ g·mL<sup>-1</sup>), or blasticidin (15  $\mu$ g·mL<sup>-1</sup>) and hygromycin (100  $\mu$ g·mL<sup>-1</sup>) respectively. A human haploid cell line, HAP1 (Horizon Discovery), derived from a chronic myelogenous leukaemia, was grown at 37 °C in the presence of 5% CO<sub>2</sub> in Iscove's modified Dulbecco's medium (IMDM) containing 10% (v/v) FBS, penicillin (100 units·mL<sup>-1</sup>) and streptomycin (0.1 mg·mL<sup>-1</sup>). A cell line, HAP1- $\Delta$ METTL12 (Horizon Discovery HZGHC000536c011), derived from the parental cells by disruption of *METTL12* by CRISPR-Cas9 technology was grown under the same conditions. The proliferation of cells was assayed at 37 °C in a humidified atmosphere containing 5% CO<sub>2</sub> with an IncuCyte HD instrument (Essen Bioscience, Welwyn Garden City, UK). Initially,  $1 \times 10^5$  parental HAP1 cells or HAP1- $\Delta$ METTL12 cells were seeded into three wells in a six-well plate. Cell proliferation was followed by measuring confluence over a period of time and quantified with INCUCYTE software (version 2011A; Essen Bioscience).

### Expression of tagged METTL12

The cDNA for human METTL12 (Source BioScience, Cambridge, UK) was amplified by PCR with the forward and reverse primers 5'-ACCGAGCTCGGATCCGGAATGGCCGCGCTGCGTCGAAT-3' and 5'-ACCGGTCCCTCGAGATGAGAGCCTTGAATCAAGTAAG-3', respectively, which introduced Flag and Strep II tags at its C terminus. The amplified sequence was cloned into plasmid pcDNA5/FRT/TO. This plasmid and plasmid pOG44 were cotransfected in the presence of lipofectamine 2000 (Life Technologies, Paisley, UK) into human HEK293T Flp-In T-Rex cells (total DNA 1  $\mu$ g; pOG44:pcDNA5/FRT/TO, 7 : 1 by weight) (Life Technologies) [32]. After 24 h, the cells were transferred into selective medium containing blasticidin (15  $\mu$ g·mL<sup>-1</sup>) and hygromycin (100  $\mu$ g·mL<sup>-1</sup>), and inducible cell lines expressing the recombinant protein were selected. Expression of tagged METTL12 was induced for 24 h with doxycycline (20 ng·mL<sup>-1</sup>), then mitochondria [32] or mitoplasts [11,33] were prepared from the cells.

### Protein analyses

Samples of proteins were fractionated by SDS/PAGE on Novex 10–20% acrylamide gradient gels (Life Technologies) and then transferred by electrophoresis to Immobilon P

membranes (Millipore, Watford, UK). Membranes were washed with a solution (PBST) made from PBS containing Tween-20 (0.01%, v/v), and then blocked with a solution of dried skimmed milk in PBST, or, when a methyl-specific antibody was used, in PBST plus bovine serum albumin. Then, the membrane was exposed to the primary antibody. The primary antibodies were specific for methyllysine and trimethyllysine (Abcam, Cambridge, UK; ab23366 and ab76118), human mitochondrial malate dehydrogenase, citrate synthase and ETF $\beta$  (Proteintech, Manchester, UK; 15462-1-AP, 16131-1-AP and 17925-1-AP respectively). Bound primary antibodies were detected by chemiluminescence generated with an anti-rabbit secondary antibody conjugated to horse radish peroxidase (Thermo Fisher Scientific, Loughborough, UK) and ECL prime reagents (GE Healthcare, Little Chalfont, UK). Bound proteins were stained with Coomassie Blue R250 0.08% (w/v) dye dissolved in an aqueous solution containing 20% methanol (v/v) and 2.8% acetic acid (v/v). The membrane was washed with 50% methanol (v/v) and dried in air. Then it was wetted with 100% methanol and reprobed with another antibody.

### Mass spectrometry

Samples were fractionated by SDS/PAGE as above, and proteins were detected by staining with Coomassie blue dye. The stained bands were excised, digested 'in-gel' [34] with trypsin or AspN protease, and the peptides were analysed in an LTQ Orbitrap XL-electron transfer dissociation (ETD) mass spectrometer (Thermo Scientific, Hemel Hempstead, UK) coupled with a Proxeon Easy-nLC (Thermo Scientific) for nanoscale reverse-phase peptide separation, as described before [11,23]. Proteins were identified by interrogation of a human protein sequence database with the peptide mass and fragmentation data, with the aid of Proteome Discoverer 1.3 (Thermo Scientific) and the Mascot search engine, version 2.4.0 [35]. The identities of peptides containing methylated lysine residues, and the specific site of methylation, were determined by manual interpretation of ETD fragmentation data. Fragment ions generated by ETD are labelled as c and z-ions, with the related z+H-ions included with z-ions [36]. The relative abundances of peptides bearing one, two, three or no methyl groups were determined with Xcalibur data analysis software from calculated peak areas of Gaussian smoothed extracted ion chromatograms, applying a 5 p.p.m.  $m/z$  tolerance and overlapping peptide retention times.

### Assay of citrate synthase

The activity of citrate synthase in parental HAP1 cells and HAP1- $\Delta$ METTL12 cells was determined by an immunocapture-based microplate assay (Abcam) requiring the measurement of thionitrobenzoic acid released by the reaction of 5,5-dithiobis-(2-nitrobenzoic acid) with CoA-SH

produced by the enzymic generation of citrate from oxaloacetate and acetyl-CoA [37]. The concentration of citrate synthase in cell lysates was determined with an ELISA assay kit (Elabscience, Wuhan, China). Both assays were performed in a Spectramax Plus 384 microplate reader (Molecular Devices, Wokingham, UK), and analysed with SOFTMAX PRO software (Molecular Devices).

### Immunocapture of citrate synthase

Mitochondria from wild-type HAP1 cells and HAP1- $\Delta$ METTL12 cells, were solubilized at a protein concentration of 5 mg·mL<sup>-1</sup> with 1% (w/v) dodecylmaltoside prepared in PBS containing complete protease inhibitors (Roche, Mannheim, Germany). An anti-citrate synthase monoclonal antibody (Abcam ab128564) was added (5  $\mu$ L per 100  $\mu$ L of mitochondrial solution), and the sample was kept at 4 °C for 4 h. A 50% (w/v) slurry of protein-G agarose (10  $\mu$ L; Abcam) was added, and the sample was mixed for 4 h at 4 °C. Unbound material was removed, and the agarose was washed four times with PBS containing 0.1% dodecylmaltoside. Bound proteins were eluted with a solution of 0.2 M glycine-HCl, pH 2.5, and neutralized with a 1 M Tris solution to a final concentration of ca. 130 mM.

## Results

### Mitochondrial protein methyltransferases

Previously, a catalogue of known and predicted methyltransferases in human mitochondria has been presented [23]. The entries were selected from a larger catalogue of 208 known and putative methyltransferases [25] on the basis of their having an N-terminal sequence with the characteristic features of a mitochondrial targeting signal that would be removed by proteolysis during import of the protein into the organelle, to generate the mature protein in the matrix of the mitochondrion [38]. This past catalogue contained not only known protein methyltransferases but also methyltransferases that modify RNA, and others involved in the biogenesis of small molecules. The new catalogue presented here (Table 1) is restricted to known protein methyltransferases and unknown predicted methyltransferases. Thus, potentially, some unknown entries could modify proteins, and others either RNA or small molecules. Four proteins, PRMT8, PRMT9, PRDM15 and KMT2A, are all predicted by two programs to be mitochondrial proteins, but PRMT8 is a plasma membrane protein [39] and the others are found in the nucleus and/or have been shown to methylate nuclear proteins [40–43]. Therefore, they have been excluded from the new catalogue. Another potential methyltransferase, SETD4, has been

**Table 1.** Known and possible protein methyltransferases in human mitochondria.

Protein	Family <sup>a</sup>	MTS <sup>b</sup>	Function <sup>c</sup>	Target <sup>d</sup>	Location <sup>e</sup>	References
HEMK1	7β	1	Gln-Me	PRF1-L	M	[21,63]
METTL12	7β	3	Lys-Me	CS	M	This work
METTL20	7β	3	Lys-Me	ETFβ	M	[11,45]
NDUFAF5	7β	4	Arg-OH	NDUFS7	M	[24,61,63]
NDUFAF7	7β	4	Arg-Me	NDUFS2	M	[23,62,63]
SETD9	SET	4	Unknown	Unknown	M	[64]
RSAD1	R-SAM	4	Unknown	Unknown	M	[64]
COMTD1	7β	1	Unknown	Unknown	M	[63]
SETD4	SET	2	Unknown	Unknown	N, C	[44]
METTL9	7β	4	Unknown	Unknown	Unknown	
ASMTL	7β	3	Unknown	Unknown	Unknown	
RRNAD1	7β	4	Unknown	Unknown	Unknown	

<sup>a</sup>7β, 7β-strand; SET, SET domain (*Drosophila* Su(var)3-9, Enhancer of zeste and Trithorax); R-SAM, radical S-adenosylmethionine. <sup>b</sup>MTS, number of independent predictions of the presence of a N-terminal mitochondrial targeting sequence; unless there is experimental evidence for their localization in mitochondria, singly predicted proteins have been excluded. <sup>c</sup>Gln-Me, Lys-Me and Arg-Me, glutamine, lysine and arginine methyltransferases respectively; Arg-OH, arginine hydroxylase. <sup>d</sup>PRF1-L, peptide release factor 1-like; ETFβ, the β-subunit of the electron transfer flavoprotein; CS, citrate synthase. <sup>e</sup>Subcellular location of methyltransferase; M, mitochondria; N, nucleus; C, cytoplasm.

described as being a component of both the nucleus and the cytoplasm [44]. As the cytoplasmic attribution could conceivably represent the mitochondria, SETD4 remains in this new restricted catalogue.


### Overexpression of METTL12 and methylation of human citrate synthase

METTL12, was overexpressed in HEK293T cells, and, in comparison to an equivalent cell line overexpressing a second mitochondrial methyltransferase, METTL20, that modifies two lysine residues in ETFβ [11,45], the overexpression of METTL12 was accompanied by an increase in the methylation of a lysine residue (or residues) in citrate synthase (Fig. 1). In order to identify the methylated site or sites, peptides in AspN and trypsin digests of human citrate synthase were analysed by mass spectrometry from the samples where METTL12 and METTL20 had been overexpressed (Tables S1–S4). The numbering of residues in the sequence of citrate synthase refers to the mature protein produced by removal of the N-terminal 27 amino acid MTS [10]. The fragmentation spectra of two triply charged ions of an AspN peptide corresponding to residues 363–374, with *m/z* values of 470.59 and 456.57, correspond to trimethylated and unmethylated states respectively. These fragmentation spectra contained the ions *c*<sub>5</sub>–*c*<sub>6</sub> and *z*<sub>6</sub>–*z*<sub>7</sub>, which show that residue Lys-368 is trimethylated (Fig. 1), whereas the ions *c*<sub>3</sub>–*c*<sub>4</sub> and *z*<sub>8</sub>–*z*<sub>9</sub> show Lys-366 is unmodified. From the relative abundances of the mono-, di- and trimethylated forms of the peptide (Table S5), it was calculated that 99% of this residue was trimethylated in cells

overexpressing METTL12, in contrast to 14.7% of the residue being trimethylated in control cells. In the same control cells, the abundances of dimethylated, monomethylated and unmethylated states were 14.6%, 26.4% and 44.3% respectively (Fig. 1). Similar levels of partial methylation of this site were observed in cells overexpressing METTL20 (Fig. 1), which has no effect on citrate synthase, but instead modifies two lysine residues in ETFβ [11,45].

### Impact of disruption of METTL12

The 240 amino acids of the mitochondrial precursor of human METTL12 are encoded by a single human gene on chromosome 11 with three exons interrupted by two introns. In the near-haploid human cell line HAP1-ΔMETTL12, exon III has been disrupted by the insertion of a single base. This insertion introduces a frame-shift, leading to a truncated version of METTL12 with the N-terminal mitochondrial import sequence from residues 1–21 [29] and residues 1–30 of the mature protein, plus unrelated protein sequence terminated 66 amino acids later. Analysis of the mitochondria from wild-type HAP1 cells, revealed the expected presence of the three prominent mitochondrial proteins containing trimethyllysine residues, namely citrate synthase, the ADP/ATP translocase and the c-subunit of ATP synthase. Similarly, the mitochondria from HAP1-ΔMETTL12 cells also contained the trimethylated ADP/ATP translocase and the c-subunit of ATP synthase, and the same level of citrate synthase as wild-type cells, but in contrast to the wild-type cells, the HAP1-ΔMETTL12 cells were devoid of the trimethylated form of citrate synthase (Fig. 2A).




**Fig. 1.** The methylation of human citrate synthase accompanying overexpression of METTL12. (A) Fractionation by SDS/PAGE of mitochondria from HEK293T cells overexpressing, in track a, METTL12, and in tracks b and c, METTL20. Proteins in tracks a and b were detected with an antibody recognizing methylated lysine residues, and track c is track b, stripped of bound antibodies and reprobred with antibodies against citrate synthase (CS) and ETF $\beta$ . The positions of the proteins containing trimethylated lysine residues are indicated on the left; HSP70, 70-kDa heat shock protein; EF1 $\alpha$ , elongation factor 1 $\alpha$ ; ANT, ADP/ATP translocase. (B) The relative peak areas taken from the extracted ion chromatograms of the AspN peptide EQGKAKNPWPNV (residues 363–374) of citrate synthase in mitoplast samples, containing unmethylated, and mono-, di- and trimethylated lysine-368 residues (white, light grey, dark grey and black areas respectively). Column a, control cells; b and c, cells overexpressing METTL20 and METTL12 respectively. (C,D) Fragmentation spectra of the AspN peptide produced by ETD of triply charged ions with  $m/z$  470.59 and 456.57 respectively. The asterisks indicate ions arising from neutral losses of CH<sub>3</sub>NO and C<sub>2</sub>H<sub>5</sub>NO from Gln and Asn residues. In the insets, fragment ions have been mapped onto the amino acid sequence.

Both samples also contained additional bands, corresponding to contaminant nonmitochondrial HSP70 proteins and elongation factor 1 $\alpha$ , which also contain trimethylated lysines [46,47].

The methylation of Lys-368 of citrate synthase in wild-type HAP1 cells was verified by mass spectrometric analysis of the AspN peptide from residues 363–374. This analysis revealed that 74% of the peptide was trimethylated, 10% was dimethylated, 9% was

monomethylated and 7% was unmethylated (Fig. 2 and Table S6). In contrast, in HAP1- $\Delta$ METTL12 cells, the level of the unmethylated peptide was 97% (Fig. 2 and Table S6). The residual 3% was predominantly monomethylated, probably arising from artefactual methyl esterification of residue Glu-363 by acidic methanol used in the detection of the protein in the SDS/PAGE gel [48]. Spectra arising from the fragmentation of triply charged ions of this peptide with  $m/z$



**Fig. 2.** Effect of disruption of *METTL12* on the methylation of human citrate synthase. (A) Mitoplasts from a, wild-type HAP1 cells and b, HAP1- $\Delta$ METTL12 cells, fractionated by SDS/PAGE and western blotted with antibodies against trimethyllysine (TML) and citrate synthase (CS). An antibody against malate dehydrogenase (MDH2), which is found in the matrix of mitochondria, provided a loading control. On the right-hand side, the same membrane stained with Coomassie Blue dye. (B) The relative peak areas obtained from the extracted ion chromatograms of the various methylation states of the AspN peptide representing residues 363–374 of citrate synthase. The peptide was produced by digestion of the appropriate band in the fractionated mitoplast samples. Columns a and b, respectively, wild-type and HAP1- $\Delta$ METTL12 cells. For the meaning of the various shades, see the legend to Fig. 1. (C,D) ETD fragmentation spectra of the AspN peptide from residues 363–374 from citrate synthase immunocaptured from mitochondria from HAP1 wild-type and HAP1- $\Delta$ METTL12 cells; they represent the trimethylated and unmethylated forms of the peptide respectively. The asterisks indicate ions arising from neutral losses of  $\text{CH}_3\text{NO}$  and  $\text{C}_2\text{H}_5\text{NO}$  from Gln and Asn residues. In the insets, fragment ions have been mapped onto the amino acid sequence.

values of 456.57 and 470.59, confirmed the presence of unmethylated forms of Lys-368 in HAP1- $\Delta$ METTL12 cells and of trimethylated forms in HAP1 parental cells (Fig. 2).


### Impact of the absence of METTL12 on HAP1 cells

Under the limited range of conditions that were investigated, the removal of METTL12 had no impact on the proliferation of HAP1 cells (Fig. 3A). Additionally, neither the amount (wild-type,  $0.13 \pm 0.03$  and HAP1- $\Delta$ METTL12  $0.14 \pm 0.02$   $\text{ng}\cdot\mu\text{g}^{-1}$  of cell lysate), nor the enzymic activity of citrate synthase (Fig. 3) was affected.


## Discussion

### Subcellular site of methylation of mitochondrial proteins

Nuclear-encoded mitochondrial proteins are synthesized in the cytoplasm and are transferred to the organelle via TOM and TIM complexes in the outer and inner membranes respectively [38]. Thus, it is conceivable that the post-translational introduction of methyl groups into specific mitochondrial proteins by methyltransferases could occur at any stage during their transit to their final destinations in the organelle.



**Fig. 3.** Comparison of properties of wild-type HAP1 cells and HAP1-ΔMETTL12 cells. (A) Growth rates. Initially, ca.  $10^5$  HAP1 wild-type cells (●) and  $10^5$  HAP1-ΔMETTL12 cells (□) were seeded into six-well plates, and their confluence was measured over 96 h. The data points represent the mean values  $\pm$  SD ( $n = 3$ ). (B) Activity of citrate synthase in cell lysates from wild-type cells (●) and HAP1-ΔMETTL12 cells (□) ( $n = 3$ , SD values are smaller than the symbols). (C) Activity of citrate synthase in  $6.25 \mu\text{g}\cdot\text{mL}^{-1}$  of cell lysates from wild-type cells (a) and HAP1-ΔMETTL12 cells (b) ( $n = 3$ ,  $\pm$ SD). The concentration of citrate synthase in wild-type and mutant cells did not differ significantly.



**Fig. 4.** The predicted structure of human METTL12. (A) Predicted secondary structure of mature METTL12. The locations of  $\alpha$ -helices indicated above the sequence by red or green bars, and  $\beta$ -sheets by blue arrows. Helix Z,  $\beta$ -sheets 1-7 and intervening  $\alpha$ -helices A-E characterize the  $7\beta$ -strand methyltransferase fold. Helix C, intervening between strands 3 and 4 in the canonical fold for this family, is not present in METTL12. The additional predicted helices I and II are found in the structure of the related methyltransferase domain of endothelin-converting enzyme-2 [65]. Helix I is positioned over the fold. Conserved motifs I, post-I, II and III are also found in  $7\beta$ -strand methyltransferases [66]. The post-II motif sequence (underlined) is highly conserved in other related methyltransferases METTL10, METTL13 and the methyltransferase domain of endothelin-converting enzyme-2. (B) Topology of the  $7\beta$ -strand fold and flanking  $\alpha$ -helices. Triangles, circles and lines represent  $\beta$ -strands,  $\alpha$ -helices and joining loops respectively. The positions of the substrate binding sites are shown.

Previously, in human mitochondria, three methyltransferases (HEMK1, NDUFAF7, METTL20), with SAM as the methyl donor, have been found uniquely in the

matrix space (Table 1) [11,21,23]. In addition, NDUFAF5, which probably uses SAM as a cofactor in the introduction of a hydroxyl group into the side chain

of Arg-73 in the NDUFS7 subunit of complex I, is found in the same subcellular location [24]. METTL12, also located in the mitochondrial matrix, adds a fourth protein-modifying methyltransferase to this growing list. Its predicted secondary structure indicates that it is a 7 $\beta$ -strand methyltransferase with five of the six associated canonical  $\alpha$ -helices (Fig. 4). METTL12 has been placed into a subfamily of 7 $\beta$ -strand methyltransferases, with human METTL10, METTL13 and ECE2, and yeast EFM4 [25]. They share regions of sequence homology including a postmotif-II sequence that interacts with the specific substrate [49]. Like METTL12, both METTL10 and EFM4 are lysine methyltransferases [50,51]. Therefore, it is likely that members of this subfamily all catalyse the methylation of lysine residues.


### The site of methylation in citrate synthase

The three-dimensional structure of human citrate synthase has not been determined, but the sequences of the human and porcine enzymes are 98% identical, and therefore the structure of the porcine enzyme [52] provides an excellent model for the human orthologue. It is known that, as in the human enzyme, Lys-368 in the porcine protein is trimethylated [10]. The active enzyme is a homodimer, and Lys-368 is exposed on the surface of the enzyme in a loop in an extended structure that forms part of the binding pocket for CoA (Fig. 5). Lys-366, nearby, is unmodified. Despite the modified residue being close to the active site of the enzyme, removal of the methyl group had no effect on the activity of the human enzyme *in vitro* (Fig. 3), and recombinant porcine citrate synthase lacking the methylation had a similar specific activity to the methylated enzyme isolated from a natural source [53]. However, this is unlikely to reflect the *in vivo* situation in the mitochondrial matrix, where citrate synthase does not function in isolation, but as a component of the TCA cycle of enzymes. Moreover, while the removal of the methyl groups in HAP1- $\Delta$ METTL12 cells had no effect on cellular growth under the conditions that were investigated (Fig. 3); in the high-glucose IMDM media, HAP1 cells can use aerobic glycolysis to generate the ATP and metabolites needed for growth. Therefore, under these conditions, the impact of any altered activity of citrate synthase on cell growth will be minimal.

In contrast to the observed methylation, there was no evidence for the acetylation of lysine residues of citrate synthase in the mass spectrometric analysis of AspN or tryptic peptides (Tables S1–S4 and S7–S10). The equivalent concentrations of citrate synthase

measured in HAP1 cells, despite there being significantly different levels of methylation, suggest that the modification does not influence the stability of the enzyme. However, the central location of the methylated site, its conservation throughout metazoans, and also in fungi and plants (Fig. 6), and the conservation of METTL12 across metazoans suggests that the modification is likely to be biologically significant.

Citrate synthase catalyses the first step of the tricarboxylic acid cycle, and there is extensive evidence that it forms a metabolon or metabolons with other citric acid cycle enzymes to facilitate the channelling of substrates between enzymes in the same pathway [54–57]. One proposal is that all of the enzymes that perform the cycle, plus aspartate aminotransferase and nucleoside diphosphate kinase, are organized around the  $\alpha$ -ketoglutarate dehydrogenase complex [58]. Therefore, one possibility is that under particular cellular conditions, as yet undefined, the methylation of Lys-368 regulates contacts with other enzymes in a metabolon of part or all of the citric acid cycle.



**Fig. 5.** Location of methylated lysine-368 in citrate synthase. The image, made with PyMol and shown in cartoon representation, displays the active site of one monomer of the dimeric porcine enzyme (PDB 2CTS). The CoA molecule is pink, and the side chain of the trimethylated Lys-368 is red. The side chain of unmodified Lys-366 is blue. In addition to CoA, the active site also contains citrate (CIT, green). Amino acids H274, H320 and D375 participate in catalysis.

Hs	358	PNVLLEQGKAK	NPWP	NVDAHS	378					
Ss	358	PNVLLEQGKAK	NPWP	NVDAHS	378					
Rn	358	PNILLEQGKAK	NPWP	NVDAHS	378					
Sv	388	PNVLLEQGKAK	NPWP	NVDAHS	408					
Ii	388	PNVLLEQGKAK	NPWP	NVDAHS	408					
Xl	387	PNILLEQGKAK	NPWP	NVDAHS	407					
Dr	387	PNVLLEQGKAK	NPWP	NVDAHS	407					
Dm	386	PPILTETGKVK	NPWP	NVDAHS	406					
Dp	384	PDILLQTGKVK	NPWP	NVDAHS	404					
Cb	387	PGILLEQGKAK	NPWP	NVDSHS	407					
Lg	387	PDILLEQGKAK	NPWP	NVDAHS	407					
Ep	393	PDVLLLEQGK	AKNPWP	NVDAHS	413					
Ta	402	PDVLLLEHGK	AKNPWP	NVDAHS	422					
Sc	396	PGVLTKHGK	TKNPWP	NVDSHS	416					
At	392	PPVLTELGK	VKNPWP	NVDAHS	412					
		*	:	*	:	**	.	*****	:	**

**Fig. 6.** Conservation of lysine-368 in citrate synthase. Alignment of sequences adjacent to human Lys-368 from representative species across metazoans, in yeast and in a plant. The trimethyl-lysines in the human and porcine proteins are red, and the grey box highlights a conserved residue in the catalytic triad of the enzyme. Hs, *Homo sapiens*; Ss, *Sus scrofa*; Rn, *Rattus norvegicus*; Sv, *Sturnus vulgaris*; Ii, *Iguana iguana*; Xl, *Xenopus laevis*; Dr, *Danio rerio*; Dm, *Drosophila melanogaster*; Dp, *Daphnia pulex*; Cb, *Caenorhabditis briggsae*; Lg, *Lottia gigantea*; Ep, *Exaoptasia pallida*; Ta, *Trichoplax adhaerens*; Sc, *Saccharomyces cerevisiae*; At, *Arabidopsis thaliana*. The alignment was made with Clustal Omega. The conserved human, porcine and rat proteins are numbered according to the mature porcine protein, after removal on the N-terminal 27 amino acid mitochondrial targeting peptide [10]. The numbering of all other proteins is based on the full-length sequence. The symbols \*, : and . denote identical, strongly conserved and weakly conserved residues respectively.

### Roles of methylation of mitochondrial proteins

Methylated proteins in mitochondria fall into two broad categories. In the first are ETF $\beta$ , where lysine residues 199 and 202 are methylated [11,45], and citrate synthase described here. A characteristic of these proteins is that the extent of methylation is partial, and in the case of at least ETF $\beta$ , it varies according to growth conditions. In ETF $\beta$ , the modified site is adjacent to the recognition loop of the ETF, which has the remarkable property of recognizing and binding to as many as 13 different acyl-CoA dehydrogenases so as to allow electron transfer between the dehydrogenases and the ETF [59,60]. It has been postulated that methylation of ETF $\beta$  helps to regulate these protein–protein interactions [11,45], similar to one of the proposals being made here to explain the methylation of citrate synthase. Unlike regulation of pathways in other subcellular locations of human cells, there is no evidence in mitochondria that methylation is a reversible process, and so far, no mitochondrial

protein demethylase has been identified. The second category consists of proteins where the site of methylation is modified completely, as in Arg-85 of the NDUFS2 subunit of complex I [14], ADP/ATP translocase [12] and the c-subunit of the ATP synthase complex [13]. To this category, the complex I subunit, NDUFS7, can be added, where Arg-73 is completely hydroxylated by NDUFAF5 [14,24]. In the cases of NDUFS2 and NDUFS7, the modifications are absolutely required in the process of assembly of complex I, and, in each case, the absence of the modification stalls the assembly process at specific intermediate stages [23,24,61,62]. Therefore, it is possible that, in the ADP/ATP translocase and the c-subunit of the ATP synthase, the complete methylation of the modified residues is also an obligatory step in the biogenesis and/or assembly of the proteins. The identification of the modifying enzymes would help in the testing of this hypothesis.

### Acknowledgements

We thank M. G. Montgomery for preparing Fig. 5. This work was supported by the Medical Research Council of the United Kingdom by grant MC\_U1065663150 and by Programme Grant MR/M009858/1, both to JEW and a Fellowship from the Swiss Novartis Foundation to VFR.

### Author contributions

JEW designed research and supervised project and prepared the manuscript; VFR, JC, SD and IMF performed research; VFR, JC, SD, IMF and JEW analysed data.

### References

- Huang J and Berger SL (2008) The emerging field of dynamic lysine methylation of non-histone proteins. *Curr Opin Genet Dev* **18**, 152–158.
- Clarke SG (2013) Protein methylation at the surface and buried deep: thinking outside the histone box. *Trends Biochem Sci* **38**, 243–252.
- Erce MA, Pang CN, Hart-Smith G and Wilkins MR (2012) The methylproteome and the intracellular methylation network. *Proteomics* **12**, 564–586.
- Lu Z, Cheng Z, Zhao Y and Volchenboum SL (2011) Bioinformatic analysis and post-translational modification crosstalk prediction of lysine acetylation. *PLoS One* **6**, e28228.
- Lothrop AP, Torres MP and Fuchs SM (2013) Deciphering post-translational modification codes. *FEBS Lett* **587**, 1247–1257.

- 6 DeLange RJ, Glazer AN and Smith EL (1969) Presence and location of an unusual amino acid, epsilon-N-trimethyllysine, in cytochrome c of wheat germ and *Neurospora*. *J Biol Chem* **244**, 1385–1388.
- 7 Polevoda B, Martzen MR, Das B, Phizicky EM and Sherman F (2000) Cytochrome c methyltransferase, Ctm1p, of yeast. *J Biol Chem* **275**, 20508–20513.
- 8 Park KS, Frost B, Tuck M, Ho LL, Kim S and Paik WK (1987) Enzymatic methylation of *in vitro* synthesized apocytochrome c enhances its transport into mitochondria. *J Biol Chem* **262**, 14702–14708.
- 9 Winter DL, Abeygunawardena D, Hart-Smith G, Erce MA and Wilkins MR (2015) Lysine methylation modulates the protein-protein interactions of yeast cytochrome C Cyc1p. *Proteomics* **15**, 2166–2176.
- 10 Bloxham DP, Parmelee DC, Kumar S, Walsh KA and Titani K (1982) Complete amino acid sequence of porcine heart citrate synthase. *Biochemistry* **21**, 2028–2036.
- 11 Rhein VF, Carroll J, He J, Ding S, Fearnley IM and Walker JE (2014) Human METTL20 methylates lysine residues adjacent to the recognition loop of the electron transfer flavoprotein in mitochondria. *J Biol Chem* **289**, 24640–24651.
- 12 Aquila H, Bogner W and Klingenberg M (1982) ADP/ATP-translocator from beef heart mitochondria: amino acid sequence and surface labelling. *Hoppe-Seyler's Z Physiol Chem* **363**, 894.
- 13 Chen R, Fearnley IM, Palmer DN and Walker JE (2004) Lysine 43 is trimethylated in subunit c from bovine mitochondrial ATP synthase and in storage bodies associated with Batten disease. *J Biol Chem* **279**, 21883–21887.
- 14 Carroll J, Ding S, Fearnley IM and Walker JE (2013) Post-translational modifications near the quinone binding site of mammalian complex I. *J Biol Chem* **288**, 24799–24808.
- 15 Carroll J, Fearnley IM, Skehel JM, Runswick MJ, Shannon RJ, Hirst J and Walker JE (2005) The post-translational modifications of the nuclear encoded subunits of complex I from bovine heart mitochondria. *Mol Cell Proteomics* **4**, 693–699.
- 16 Cao XJ, Arnaudo AM and Garcia BA (2013) Large-scale global identification of protein lysine methylation *in vivo*. *Epigenetics* **8**, 477–485.
- 17 Guo A, Gu H, Zhou J, Mulhern D, Wang Y, Lee KA, Yang V, Aguiar M, Kornhauser J, Jia X *et al.* (2014) Immunoaffinity enrichment and mass spectrometry analysis of protein methylation. *Mol Cell Proteomics* **13**, 372–387.
- 18 Larsen SC, Sylvestersen KB, Mund A, Lyon D, Mullari M, Madsen MV, Daniel JA, Jensen LJ & Nielsen ML (2016) Proteome-wide analysis of arginine monomethylation reveals widespread occurrence in human cells. *Sci Signal* **9**, rs9.
- 19 Martin JL and McMillan FM (2002) SAM (dependent) I AM: the S-adenosylmethionine-dependent methyltransferase fold. *Curr Opin Struct Biol* **12**, 783–793.
- 20 Schubert HL, Blumenthal RM and Cheng X (2003) Many paths to methyltransfer: a chronicle of convergence. *Trends Biochem Sci* **28**, 329–335.
- 21 Ishizawa T, Nozaki Y, Ueda T and Takeuchi N (2008) The human mitochondrial translation release factor HMRFL is methylated in the GGQ motif by the methyltransferase HMPPrmC. *Biochem Biophys Res Commun* **373**, 99–103.
- 22 Carilla-Latorre S, Gallardo ME, Annesley SJ, Calvo-Garrido J, Graña O, Accari SL, Smith PK, Valencia A, Garesse R, Fisher PR *et al.* (2010) MidA is a putative methyltransferase that is required for mitochondrial complex I function. *J Cell Sci* **123**, 1674–1683.
- 23 Rhein VF, Carroll J, Ding S, Fearnley IM and Walker JE (2013) NDUFAF7 methylates arginine 85 in the NDUFS2 subunit of human complex I. *J Biol Chem* **288**, 33016–33026.
- 24 Rhein VF, Carroll J, Ding S, Fearnley IM and Walker JE (2016) NDUFAF5 hydroxylates NDUFS7 at an early stage in the assembly of human complex I. *J Biol Chem* **291**, 14851–14860.
- 25 Petrossian TC and Clarke SG (2011) Uncovering the human methyltransferasome. *Mol Cell Proteomics* **10**, M110.000976.
- 26 Claros MG and Vincens P (1996) Computational method to predict mitochondrially imported proteins and their targeting sequences. *Eur J Biochem* **241**, 779–786.
- 27 Bannai H, Tamada Y, Maruyama O, Nakai K and Miyano S (2002) Extensive feature detection of N-terminal protein sorting signals. *Bioinformatics* **18**, 298–305.
- 28 Emanuelsson O, Nielsen H, Brunak S and von Heijne G (2000) Predicting subcellular localization of proteins based on their N-terminal amino acid sequence. *J Mol Biol* **300**, 1005–1016.
- 29 Fukasawa Y, Tsuji J, Fu SC, Tomii K, Horton P and Imai K (2015) MitoFates: improved prediction of mitochondrial targeting sequences and their cleavage sites. *Mol Cell Proteomics* **14**, 1113–1126.
- 30 Cole C, Barber JD and Barton GJ (2008) The Jpred 3 secondary structure prediction server. *Nucleic Acids Res* **36**, W197–W201.
- 31 Jones DT (1999) Protein secondary structure prediction based on position-specific scoring matrices. *J Mol Biol* **292**, 195–202.
- 32 He J, Cooper HM, Reyes A, Di Re M, Kazak L, Wood SR, Mao CC, Fearnley IM, Walker JE and Holt IJ (2012) Human C4orf14 interacts with the mitochondrial nucleoid and is involved in the biogenesis

- of the small mitochondrial ribosomal subunit. *Nucleic Acids Res* **40**, 6097–6108.
- 33 Klement P, Nijtmans LG, Van den Bogert C and Houstek J (1995) Analysis of oxidative phosphorylation complexes in cultured human fibroblasts and amniocytes by blue-native-electrophoresis using mitoplasts isolated with the help of digitonin. *Anal Biochem* **231**, 218–224.
- 34 Wilm M, Shevchenko A, Houthaeve T, Breit S, Schweigerer L, Fotsis T and Mann M (1996) Femtomole sequencing of proteins from polyacrylamide gels by nano-electrospray mass spectrometry. *Nature* **379**, 466–469.
- 35 Perkins DN, Pappin DJ, Creasy DM and Cottrell JS (1999) Probability-based protein identification by searching sequence databases using mass spectrometry data. *Electrophoresis* **20**, 3551–3567.
- 36 Sun RX, Dong MQ, Song CQ, Chi H, Yang B, Xiu LY, Tao L, Jing ZY, Liu C, Wang LH *et al.* (2010) Improved peptide identification for proteomic analysis based on comprehensive characterization of electron transfer dissociation spectra. *J Proteome Res* **9**, 6354–6367.
- 37 Srere PA, Brazil H and Gonen L (1963) The citrate condensing enzyme of pigeon breast muscle and moth flight muscle. *Acta Chem Scand* **17**, S129–S134.
- 38 Mossmann D, Meisinger C and Vögtle FN (2012) Processing of mitochondrial presequences. *Biochim Biophys Acta* **1819**, 1098–1106.
- 39 Lee J, Sayegh J, Daniel J, Clarke S and Bedford MT (2005) PRMT8, a new membrane-bound tissue-specific member of the protein arginine methyltransferase family. *J Biol Chem* **280**, 32890–32896.
- 40 Hsieh JJ, Ernst P, Erdjument-Bromage H, Tempst P and Korsmeyer SJ (2003) Proteolytic cleavage of MLL generates a complex of N- and C-terminal fragments that confers protein stability and subnuclear localization. *Mol Cell Biol* **23**, 186–194.
- 41 Shibuya K, Kudoh J, Okui M and Shimizu N (2005) Identification of a novel zinc finger protein gene (ZNF298) in the GAP2 of human chromosome 21q. *Biochem Biophys Res Commun* **332**, 557–568.
- 42 Yang Y, Hadjikyriacou A, Xia Z, Gayatri S, Kim D, Zurita-Lopez C, Kelly R, Guo A, Li W, Clarke SG *et al.* (2015) PRMT9 is a type II methyltransferase that methylates the splicing factor SAP145. *Nat Commun* **6**, 6428.
- 43 Hadjikyriacou A, Yang Y, Espejo A, Bedford MT and Clarke SG (2015) Unique features of human protein arginine methyltransferase 9 (PRMT9) and its substrate RNA splicing factor SF3B2. *J Biol Chem* **290**, 16723–16743.
- 44 Faria JA, Corrêa NC, de Andrade C, de Angelis Campos AC, Dos Santos Samuel de Almeida R, Rodrigues TS, de Goes AM, Gomes DA and Silva FP (2013) SET domain-containing protein 4 (SETD4) is a newly identified cytosolic and nuclear lysine methyltransferase involved in breast cancer cell proliferation. *J Cancer Sci Ther* **5**, 58–65.
- 45 Malecki J, Ho AY, Moen A, Dahl HA and Falnes PØ (2015) Human METTL20 is a mitochondrial lysine methyltransferase that targets the  $\beta$  subunit of electron transfer flavoprotein (ETF $\beta$ ) and modulates its activity. *J Biol Chem* **290**, 423–434.
- 46 Dever TE, Costello CE, Owens CL, Rosenberry TL and Merrick WC (1989) Location of seven post-translational modifications in rabbit elongation factor 1 alpha including dimethyllysine, trimethyllysine, and glycerylphosphorylethanolamine. *J Biol Chem* **264**, 20518–20525.
- 47 Wang C, Lin JM and Lazarides E (1992) Methylations of 70,000-Da heat shock proteins in 3T3 cells: alterations by arsenite treatment, by different stages of growth and by virus transformation. *Arch Biochem Biophys* **297**, 169–175.
- 48 Chen G, Liu H, Wang X and Li Z (2010) *In vitro* methylation by methanol: proteomic screening and prevalence investigation. *Anal Chim Acta* **661**, 67–75.
- 49 Falnes PØ, Jakobsson ME, Davydova E, Ho A and Malecki J (2016) Protein lysine methylation by seven- $\beta$ -strand methyltransferases. *Biochem J* **473**, 1995–2009.
- 50 Lipson RS, Webb KJ and Clarke SG (2010) Two novel methyltransferases acting upon eukaryotic elongation factor 1A in *Saccharomyces cerevisiae*. *Arch Biochem Biophys* **500**, 137–143.
- 51 Shimazu T, Barjau J, Sohtome Y, Sodeoka M and Shinkai Y (2014) Selenium-based *S*-adenosylmethionine analog reveals the mammalian seven-beta-strand methyltransferase METTL10 to be an EF1A1 lysine methyltransferase. *PLoS One* **9**, e105394.
- 52 Remington S, Wiegand G and Huber R (1982) Crystallographic refinement and atomic models of two different forms of citrate synthase at 2.7 and 1.7 Å resolution. *J Mol Biol* **158**, 111–152.
- 53 Evans CT, Owens DD, Slaughter CA and Srere PA (1988) Characterization of mutant TMK368K pig citrate synthase expressed in and isolated from *Escherichia coli*. *Biochem Biophys Res Commun* **157**, 1231–1238.
- 54 Morgunov I and Srere PA (1998) Interaction between citrate synthase and malate dehydrogenase. Substrate channeling of oxaloacetate. *J Biol Chem* **273**, 29540–29544.
- 55 Shatalin K, Lebreton S, Rault-Leonardon M, Vélot C and Srere PA (1999) Electrostatic channeling of oxaloacetate in a fusion protein of porcine citrate synthase and porcine mitochondrial malate dehydrogenase. *Biochemistry* **38**, 881–889.
- 56 Vélot C and Srere PA (2000) Reversible transdominant inhibition of a metabolic pathway. *In vivo* evidence of

- interaction between two sequential tricarboxylic acid cycle enzymes in yeast. *J Biol Chem* **275**, 12926–12933.
- 57 Wu F and Minter S (2015) Krebs cycle metabolon: structural evidence of substrate channeling revealed by cross-linking and mass spectrometry. *Angew Chem Int Ed Engl* **54**, 1851–1854.
- 58 Lyubarev AE and Kurganov BI (1989) Supramolecular organization of tricarboxylic acid cycle enzymes. *Biosystems* **22**, 91–102.
- 59 Toogood HS, van Thiel A, Basran J, Sutcliffe MJ, Scrutton NS and Leys D (2004) Extensive domain motion and electron transfer in the human electron transferring flavoprotein/medium chain Acyl-CoA dehydrogenase complex. *J Biol Chem* **279**, 32904–32912.
- 60 Toogood HS, Leys D and Scrutton NS (2007) Dynamics driving function: new insights from electron transferring flavoproteins and partner complexes. *FEBS J* **274**, 5481–5504.
- 61 Sugiana C, Pagliarini DJ, McKenzie M, Kirby DM, Salemi R, Abu-Amero KK, Dahl HH, Hutchison WM, Vascotto KA, Smith SM *et al.* (2008) Mutation of C20orf7 disrupts complex I assembly and causes lethal neonatal mitochondrial disease. *Am J Hum Genet* **83**, 468–478.
- 62 Zurita Rendón O, Silva Neiva L, Sasarman F and Shoubridge EA (2014) The arginine methyltransferase NDUFAF7 is essential for complex I assembly and early vertebrate embryogenesis. *Hum Mol Genet* **23**, 5159–5170.
- 63 Pagliarini DJ, Calvo SE, Chang B, Sheth SA, Vafai SB, Ong SE, Walford GA, Sugiana C, Boneh A, Chen WK *et al.* (2008) A mitochondrial protein compendium elucidates complex I disease biology. *Cell* **134**, 112–123.
- 64 Rhee HW, Zou P, Udeshi ND, Martell JD, Mootha VK, Carr SA and Ting AY (2013) Proteomic mapping of mitochondria in living cells via spatially restricted enzymatic tagging. *Science* **339**, 1328–1331.
- 65 Tempel W, Wu H, Dombrovsky L, Zeng H, Loppnau P, Zhu H, Plotnikov AN and Bochkarev A (2009) An intact SAM-dependent methyltransferase fold is encoded by the human endothelin-converting enzyme-2 gene. *Proteins* **74**, 789–793.
- 66 Petrossian TC and Clarke SG (2009) Multiple Motif Scanning to identify methyltransferases from the yeast proteome. *Mol Cell Proteomics* **8**, 1516–1526.

## Supporting information

Additional Supporting Information may be found online in the supporting information tab for this article:

**Table S1.** Protein identification of citrate synthase from HEK293T cells overexpressing METTL12.

**Table S2.** Protein identification of citrate synthase from HEK293T cells overexpressing METTL20.

**Table S3.** Protein identification of citrate synthase from HEK293T cells overexpressing METTL12.

**Table S4.** Protein identification of citrate synthase from HEK293T cells overexpressing METTL20.

**Table S5.** Quantitation of lysine 368 methylation levels of citrate synthase in HEK293T cells.

**Table S6.** Quantitation of lysine 368 methylation levels of citrate synthase in HAP1 cells.

**Table S7.** Protein identification of citrate synthase from HAP1 wild-type cells.

**Table S8.** Protein identification of citrate synthase from HAP1 wild-type cells.

**Table S9.** Protein identification of citrate synthase from HAP1- $\Delta$ METTL12 cells.

**Table S10.** Protein identification of citrate synthase from HAP1- $\Delta$ METTL12 cells.