

Regulation of neuronal development and function by ROS

Matthew C. W. Oswald¹, Nathan Garnham², Sean T. Sweeney² and Matthias Landgraf¹

¹ Department of Zoology, University of Cambridge, UK

² Department of Biology, University of York, Heslington York, UK

Correspondence

M. C. W. Oswald, M. Landgraf, Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3EJ, UK
Fax: +44 1223 336 676
Tel: +44 1223 769 348
E-mails: mo364@cam.ac.uk (MCWO); ml10006@cam.ac.uk (ML)
and

S. T. Sweeney, Department of Biology, University of York, Heslington York YO10 5DD, UK

Fax: +44 1904 328505

Tel: +44 1904 328537

E-mail: sean.sweeney@york.ac.uk

(Received 12 December 2017, accepted 2 January 2018, available online 26 January 2018)

doi:10.1002/1873-3468.12972

Edited by Wilhelm Just

Reactive oxygen species (ROS) have long been studied as destructive agents in the context of nervous system ageing, disease and degeneration. Their roles as signalling molecules under normal physiological conditions is less well understood. Recent studies have provided ample evidence of ROS-regulating neuronal development and function, from the establishment of neuronal polarity to growth cone pathfinding; from the regulation of connectivity and synaptic transmission to the tuning of neuronal networks. Appreciation of the varied processes that are subject to regulation by ROS might help us understand how changes in ROS metabolism and buffering could progressively impact on neuronal networks with age and disease.

Keywords: axon; cytoskeleton; dendrite; NADPH oxidase; nervous system; neuronal polarity; pathfinding; plasticity; reactive oxygen species; synapse

The increase in atmospheric oxygenation is presumed to have set the pace of evolutionary change. The symbiotic acquisition of mitochondria 1.45 billion years ago generating the eukaryota further allowed diversification *via* the efficient metabolic use of diatomic oxygen. Reactive oxygen species (ROS), highly reactive molecules and free radicals derived from molecular oxygen are produced as natural by-products of normal respiratory metabolism, with the major source being the mitochondria. Mitochondria leak bursts of ROS as a function of respiration [1] pointing to a link between metabolism and ageing-related damage. Other

subcellular locations of ROS production continue to be identified and include the endoplasmic reticulum (ER) [2], peroxisome [3], the cytosol [4], plasma membrane [5] and extracellular space [6]. However, there is a growing opinion and body of evidence that ROS can act as physiological signalling molecules. In this review we will focus on the role of ROS as a physiological signal in the nervous system during development and as a regulator of neuronal function.

By their highly reactive nature, ROS are damaging to the cell, oxidising proteins, lipids and DNA and are normally regarded as detrimental to cell function.

Abbreviations

BH₄, tetrahydrobiopterin; CamKII, calcium/calmodulin-dependent kinase II; CRMP2, collapsin response mediator protein 2; ER, endoplasmic reticulum; ERK, extracellular signal-regulated kinase; Grx1, glutaredoxin 1; HFS, high-frequency stimulation; IP3Rs, inositol-3-phosphate receptors; JNK, Jun-N-terminal Kinase; LTD, long-term depression; LTP, long-term potentiation; MICAL, molecule interacting with CasL; MsrB, methionine sulfoxide reductase; NGF, nerve growth factor; PKC, protein kinase C; PP, protein phosphatase; PTEN, phosphatase and tensin homolog; PVIs, parvalbumin-expressing inhibitory interneurons; redox, reduction-oxidation; ROS, reactive oxygen species; RyRs, ryanodine receptors.

Indeed, an overwhelming of the defences against ROS is termed oxidative stress and is commonly associated with cellular damage seen in neurodegenerative disorders, including Parkinson's [7] and Alzheimer's disease [8]. ROS and the accumulation of ROS-related damage are also associated with ageing; oxidised lipids, DNA damage and the accumulation of lipofuscin (the 'aging pigment', autofluorescent material found in endosomes consisting of oxidised lipids, proteins, transition metals and senescent mitochondria) [9]. Increasing evolutionary complexity and concomitant demand for oxygen-dependent energy production *via* reduction–oxidation (redox) reactions also produced a diversification in defence mechanisms against ROS. This is particularly notable in energy demanding tissues such as heart and liver. The nervous system is anomalous in this framework as nerve cells are very energy demanding yet at the same time inadequately equipped with antioxidant defence [10,11]. Interestingly, much of the ROS defence within the nervous system occurs in glia [12]. It is becoming increasingly apparent, however, that evolution may have made a virtue out of a necessity and co-opted ROS for cellular signalling mechanisms. For such a framework lowered antioxidant defence or ROS buffering in neurons would be permissive and necessary.

Regulation of ROS in the nervous system, the case for the defence

Studies of ROS in the nervous system have primarily focused on ROS as damaging agents and the defence against ROS. Protection against ROS is mediated by multilayered constitutive and adaptive forms of defence. In the brain, static defences against ROS generally are seen to be (a) constitutive and enzymatic as seen in the standing high concentrations of enzymes such as superoxide dismutases, catalases, thioredoxin reductases and glutathione peroxidases or (b) constitutive and nonenzymatic as mediated by defence molecules (termed ROS scavengers) present in the cell, *via* synthesis or diet, such as alpha-tocopherol (vitamin E), ascorbic acid, β -carotene and tetrahydrobiopterin (BH_4). BH_4 is a molecule of particular interest. For example, BH_4 is a highly sensitive scavenger of H_2O_2 and Hydroxyl ions, while also important for the synthesis of the neurotransmitters dopamine, serotonin and noradrenaline. This suggests a functional link between ROS abundance, buffered by BH_4 levels, and neurotransmitter function (for review see [13]). A second set of defence mechanisms is adaptive and mediated by transcription factors [14]. Two pathways are prominent: (a) the NRF2/Keap system, predominant

in glial cells [12] and (b) the Jun-N-terminal Kinase (JNK)/AP-1 system, seen to be a major protective mechanism in neurons. Both promote the transcription of genes encoding antioxidant response proteins. For example, NRF2 promotes the expression of glutathione-S-transferases in glia [15] while in neurons AP-1 activation upregulates sulfiredoxin [16]. NRF2 activity in glia mediates neuronal protection in a nonautonomous manner, partly through the ensheathing nature of symbiotic glial–neuron interactions. Indeed, keeping NRF2 function low in neurons allows dendritic and synaptic development and their regulation *via* redox-sensitive signalling pathways, such as JNK/AP-1 and WNT [17].

Neuronal polarity

Most neuronal cell types are explicitly polarised, endowed with a major axonal neurite that mediates long-range connectivity and is primarily presynaptic, dedicated to passing on information, while the somato-dendritic compartment of the cell is composed of branched smaller diameter neurites that are largely postsynaptic. The establishment of neuronal polarity has until recently mostly been studied *in vitro*, using low-density cultures of cortical and hippocampal neurons. Under such conditions neuronal polarity first manifests with the emergence of a primary neurite, which extends more rapidly than others and develops into the axon, while the other minor neurites adopt postsynaptic dendritic characteristics. Several signalling pathways, including TGF- β , growth factors (e.g. BDNF), LKB and PI3 kinases, have been implicated in bringing about and maintaining asymmetries of the cytoskeleton. Characteristically, axons contain microtubules whose plus ends face away from the cell body, while dendrites have microtubules of mixed (mammals) or opposite polarity (e.g. *Drosophila* and *Caenorhabditis elegans*; for reviews see [18,19]). Perhaps inspired by work from other systems that have associated NADPH oxidase-generated ROS with the regulation of cell polarisation and growth, for example, *Arabidopsis* hair cell outgrowth [20] and the enforcement of apical dominance in *Aspergillus* hyphae [21], ROS have been investigated as signals regulating the polarisation of neurons. Indeed, *in vitro* studies suggest that ROS produced by NADPH oxidases were required alongside growth factors for the differentiation of neuronal characteristics by PC12 and SH-SY5Y cells, such as axonal outgrowth [22–25]. *In vivo*, gene expression profiles show that subunits of the NOX2 NADPH oxidase complex are present at the right place and time in mouse and rat embryonic hippocampal neurons

[26,27]. In the context of neuronal differentiation mediated by nerve growth factor (NGF), Neuregulin or Retinoic Acid ROS appear to be permissive, acting in parallel to these growth factors. As documented for several growth factors, ROS can enhance downstream kinase signalling by inhibition of phosphatases, whose active sites contain cysteines that are susceptible to oxidation [28]. The activity of protein kinases is critical during cellular polarisation, mediating positive feedback loops and signal amplification. Multiple pathways converge onto PI3 kinase, which is enriched at the tip of the future axon as neurons adopt a polarised morphology [29,30] (reviewed in [18]). Interestingly, PI3 kinase signalling can be regulated by ROS, by redox-mediated inhibition of the phosphatase and tensin homolog (PTEN), which antagonises the PI3 kinase product phosphoinositide-3-phosphate [31,32].

Studies on embryonic hippocampal neurons and cerebellar granule cells in culture have underpinned the idea that ROS contribute to the establishment of neuronal polarity. As cerebellar granule cells differentiate during the first 3 days *in vitro* overall ROS levels increase and become specifically enriched at growth cones and other cytoskeletally dynamic sites. This is paralleled by a ROS-dependent rise in Tau and MAP2 expression levels, which are representative of axonal and dendritic cytoskeletal specialisation [33]. Underlining a requirement for ROS in these processes, neurons derived from NOX2 knockout mice showed reduced neurite length compared to control cells [33]. Similarly, in embryonic hippocampal neurons pharmacological or genetic reduction in NADPH oxidase activity, e.g. by expression of the dominant negative Dnp22^{phox} regulatory subunit, led to delayed and reduced axon outgrowth [26]. Conversely, overactivation of NADPH oxidase by overexpression of the accessory protein p47^{phox} promoted axonal growth [34]. These studies suggest that NOX2 activity could indeed contribute to neuronal polarisation and promote neurite outgrowth.

The mechanisms by which physiological levels of ROS support neurite outgrowth and axon specification appear to involve release of calcium from intracellular stores, a potent second messenger regulating cytoskeletal organisation and dynamics (reviewed in [35,36]). Short respiratory bursts of NADPH oxidase-generated ROS promote calcium release by redox modification of ryanodine (RyRs) and inositol-3-phosphate receptors (IP3Rs). These in turn lead to increased expression of Rac1, an activator of the NOX2 complex, thus generating a positive feedback loop that can convert initially transient ROS bursts into sustained ROS activation and high levels of intracellular calcium [34]; (Fig. 1). However, the extent to which ROS signalling

promotes the establishment of neuronal polarity *in vivo* remains to be determined. Phenotypes of NADPH oxidase knockout and RNAi knockdown experiments in several experimental animal models suggest that removal of any one NADPH oxidase still allows the nervous systems to form and function remarkably well, at least to a level sufficient for survival under laboratory conditions [37,38]. This could indicate a degree of functional redundancy. It also suggests that ROS might have a modulatory role rather than being critically required for the establishment of neuronal polarity. This would not be entirely surprising, as during nervous system development neural progenitor cells and their progeny are invariably located within asymmetric environments where local enrichment of numerous cues, including cell adhesion and extracellular matrix proteins, can contribute to cellular polarisation [39–41] (for review see [18]).

Cytoskeletal modifications by ROS and growth cone pathfinding

Reactive oxygen species can regulate cytoskeletal change at multiple levels; directly *via* redox modification of structural cytoskeletal proteins and indirectly by modification of proteins or signalling pathways that regulate cytoskeletal dynamics. For example, actin and tubulin monomers contain multiple cysteine and methionine residues exposed to the cytoplasm that are subject to redox modifications, notably glutathionylation, nitrosylation and carbonylation [42–49]. Indeed, all major cytoskeletal elements and many cytoskeleton-associated proteins are subject to direct redox modifications of some kind [50–52], with substantial fractions of actin, tubulin and neurofilaments found glutathionylated under normal physiological conditions [53]. The question of which residues are modified, under what conditions and how this impacts on protein function and dynamics remains live. For alpha-actin purified from rabbit muscles, *in vitro* and cell culture studies have shown Cys 374 sensitive to glutathionylation, leading to a reduced rate of actin polymerisation and altered actin dynamics [44,54]. Actin Cys 374 glutathionylation has been proposed to occur in response to growth factor and integrin-stimulated signalling following interactions with the extracellular matrix [54]. This actin modification is thought to regulate the disassembly of the actinomyosin complex during cell spreading [52]. Along the same lines, disruption of protein deglutathionylation by mutation of glutaredoxin 1 (Grx1), the gene coding for an enzyme that catalyses actin deglutathionylation, led to reduced actin polymerisation and impaired

polarisation, chemotaxis, adhesion, and phagocytosis by neutrophils. Conversely, blocking NOX activity led to increased formation of filamentous actin [55]. Thus, ROS have increasingly been recognised as important regulators of actin dynamics.

One of the principal ROS sources in neurons is NADPH oxidases. Their activity is highly regulated, making them prime candidates as regulators of growth cone cytoskeletal dynamics (for reviews see [56] and [57]). How NADPH oxidase-generated ROS regulate cytoskeletal dynamics in neuronal growth cones remains poorly understood. Pharmacological inhibition of NADPH oxidase activity or lowering cytosolic ROS levels led to reduced F-actin content, retrograde flow and neurite outgrowth. Few studies have documented the localisation of NOX complex components in neuronal growth cones *in vitro*, none as yet *in vivo* [58,59]. In cultured *Aplysia* bag cell neurons the main enzymatic subunit of a NOX2-type NADPH oxidase, NOX2/gp91^{phox}, was seen localised to the plasma membrane, largely distinct from the regulatory subunit p40^{phox} found associated with filopodial actin bundles (Fig. 1). Interestingly, a local stimulus of growth cone interaction with apCAM-coated beads triggered colocalisation of both subunits to the site of growth cone–substrate interaction. Thus, in growth cones NADPH oxidase subunits are localised to the periphery at sites of actin assembly [58]. Their activity is regulated by multiple pathways, including activation following translocation of regulatory subunits, such as p40^{phox} or Rac1, but also by other signalling pathways, such as protein kinase C (PKC) [59]. The extent to which local NADPH oxidase activation regulates growth cone dynamics directly, *via* oxidation of actin and tubulin, or indirectly, through modulation of other signalling pathways, remains to be seen. Both are likely. A study using cultured marsupial kidney cells demonstrated that hydrogen peroxide-induced chemotaxis and filopodial dynamics were in large part indirectly regulated by ROS *via* local extracellular signal-regulated kinase (ERK) pathway activation, which promoted actin retrograde flow by differential activation and recruitment of cofilin and the Arp2/3 nucleator at the leading edge [60]. It is conceivable that ROS signalling in the nervous system might similarly be utilised for noncell autonomous communication, either during the development of synaptic connections or their subsequent adjustment. For example, in a mouse model for multiple sclerosis, persistent activation of NOX2/gp91^{phox} in microglia leads to the impairment of synaptic plasticity in adjacent hippocampal neurons [61].

Thus far, the clearest evidence for post-translational redox modification of cytoskeletal proteins directing growth cone pathfinding derives from studies of Semaphorin-Plexin signalling. The cytoplasmic tail of Plexin interacts with the NADPH-dependent monooxygenase, molecule interacting with CasL (MICAL), which is activated upon binding of Semaphorin guidance cues [62–67] (for review see [68]; Fig. 1). Elegant experiments by the Terman laboratory and collaborators demonstrated that the amino-terminal NADPH-dependent redox domain of MICAL-1 binds F-actin and directly oxidises the methionine residues Met44 and Met47 [64,65]. These redox modifications are regulated and can be reversed by a methionine sulfoxide reductase, (MsrB/SelR), which catalyses the reduction of methionine sulfoxide to methionine [66]. MICAL-1-mediated oxidation of actin weakens inter-actin contacts while simultaneously increasing the binding affinity of the F-actin severing protein cofilin by more than an order of magnitude. This synergistic effect of actin destabilisation by oxidation and concomitantly increased binding of cofilin promotes F-actin disassembly. In addition, MICAL-mediated oxidation decreases the capacity of actin for repolymerisation, thus further impacting on the dynamics of the actin cytoskeleton [67]. Thus, Semaphorin-Plexin guidance cue–receptor interactions are directly transduced to redox modification of the actin cytoskeleton, leading to F-actin disassembly and altered reassembly dynamics.

In addition, MICAL activation also impacts on the microtubule cytoskeleton *via* another binding partner, collapsin response mediator protein 2 (CRMP2). CRMP2 is responsible for Semaphorin-induced growth cone collapse [69] and interacts with tubulin heterodimers regulating microtubule dynamics 1 [62,70]. Current data suggest that MICAL-1 binds CRMP2, and that upon MICAL monooxygenase activation (e.g. by Semaphorin binding to Plexin), hydrogen peroxide is produced that oxidises CRMP2. Oxidation of CRMP2 promotes formation of disulfide-linked CRMP2 homodimers, which interact with thioredoxin, which in turn promotes their phosphorylation by GSK-3 β , microtubule disassembly and growth cone collapse [71]; (Fig. 1). In summary, direct redox modifications of cytoskeletal elements, notably actin and tubulin, as well as of regulators of cytoskeletal dynamics, such as CRMP2, lie at the heart of Semaphorin-Plexin growth cone guidance. It is conceivable that redox modification of cytoskeletal proteins extends to other guidance cue signalling pathways, either directly or in a modulatory capacity where the cellular redox state determines growth cone responses to extracellular cues.




Fig. 1. Regulation of the growth cone cytoskeleton by ROS. (1) Semaphorin binding to Plexin activates MICAL monoxygenase. MICAL interactions with F-actin lead to oxidation of the conserved amino acid Met44 [64]. Oxidation of actin destabilises F-actin filaments and promotes binding of the actin severing protein Profilin, thus promoting F-actin disassembly [67]. Oxidised actin monomers have a reduced propensity for polymerisation. MICAL redox activity is opposed by the MsrB enzyme SelR, which specifically reduces Met-44-R-sulfoxide [66]. MICAL activation *via* Semaphorin-Plexin binding also generates H₂O₂ that can oxidise CRMP2, enabling it to form a disulfide-linked homodimer and to transiently interact with Thioredoxin, which stimulates phosphorylation of CRMP2 by glycogen synthase kinase-3, promoting CRMP2 modification of microtubules and growth cone collapse [71]. (2) Superoxide produced by the NOX2 NADPH oxidase catalytic subunit (gp91^{phox}) is regulated by translocation of the p40^{phox} regulatory subunit from its association with F-actin to the plasma membrane upon growth cone engagement with a substrate or guidance cue [58]. Converted into hydrogen peroxide by extracellular superoxide dismutase (EC-SOD), H₂O₂ enters the cytoplasm *via* aquaporins and can oxidise cytoskeletal proteins (e.g. F-actin) [132,133]. (3) H₂O₂ also modifies RyRs and IP3Rs triggering release of calcium from internal stores in the ER. (4) Changes in intracellular calcium modify activities of cytoskeletal regulatory proteins, directly or indirectly, e.g. *via* the regulation of Calcium/calmodulin-dependent kinase II (CamKII) or the phosphatase calcineurin or activation of the protease calpain. Elevated calcium levels also lead to expression of the cytoskeletal and NADPH oxidase regulator Rac1, thus generating a positive feedback loop that can amplify and sustain transient respiratory bursts [34].

Connectivity and structural plasticity

For decades ROS have been implicated in neurodegenerative conditions, largely thought of as destructive agents [72,73]. Increasingly ROS have also been viewed as regulators and modulators of signalling pathways and gene expression, of which many are known to regulate neuronal growth and plasticity (reviewed in [56,57,74]). Several years ago we provided the first direct *in vivo* evidence of ROS as regulators of synaptic terminal growth, under pathological conditions in an experimental animal model (*Drosophila*) for lysosomal storage disease [75]. The study demonstrated that oxidative stress resulting from lysosomal storage dysfunction led to activation of the JNK cascade and activation of the immediate early genes c-Jun and c-Fos (AP-1) which in turn led to altered growth of neuromuscular junction terminals [75]. Along with a prior study by Sanyal and colleagues [76], this work identified AP-1 as the major adaptive response to ROS in

neurons. The JNK/AP-1 signalling pathway has long been known to be critical in many neuronal functions and is a well-known mediator of synaptic and oxidative stress responses (reviewed in [74]). AP-1 is a heterodimer composed of the leucine-zipper transcription factors Fos and Jun. Fos is one of the major immediate early transcription factors mediating long-term synaptic changes during long-term potentiation (LTP) [77,78], although the actual mechanism inducing JNK phosphorylation and subsequent AP-1 activation in this process remains obscure. Activation of JNK/AP-1 by oxidative stress is thought to reinforce autophagy, and many genes encoding autophagy proteins are direct transcriptional targets of AP-1 [79]. Some evidence suggests that activation of autophagy *via* the oxidative stress-induced JNK/AP-1 pathway can regulate synaptic terminal size and strength at the *Drosophila* larval neuromuscular synapse [75,76,80]. The data point to the importance of the JNK/AP-1

pathway as regulating synaptic function and to ROS as critical upstream signals during normal physiological conditions as well as under oxidative stress.

In a follow-up study we since asked whether ROS also act as regulators of synaptic terminal growth and plasticity under normal physiological conditions, which until now has remained largely unexplored. Indeed, we found that ROS, in particular hydrogen peroxide, are necessary for activity-induced synaptic terminal growth and are sufficient to drive synaptic terminal growth [81]. Specifically, overactivation of motoneurons leads to increased mitochondrial ROS levels at the presynaptic neuromuscular junction of *Drosophila* larvae, previously also reported for cultured hippocampal neurons [82,83]. In *Drosophila* larvae, activity-generated ROS promote altered synaptic terminal growth, generating more, albeit smaller synaptic varicosities (boutons) along with a reduction in the number of synaptic release sites. Postsynaptic dendrites similarly undergo homeostatic structural adjustments in response to activity-generated ROS, leading to smaller dendritic arbours, which we previously showed equates to reduced synaptic input sites and reduced synaptic drive [84]. The signalling and downstream effector pathways of neuronal activity-generated ROS are only just being sketched out. We found that neuronal ROS signal *via* the conserved redox-sensitive protein DJ-1 β , a homologue of vertebrate DJ-1 (PARK7) [85,86], which appears to act as a neuronal redox sensor. Oxidation of DJ-1 β increases inhibitory interactions with the phosphatase PTEN thus leading to disinhibition of PI3kinase signalling, a known regulator of synaptic terminal growth [87–90]. Whether changes in dendritic growth are also regulated by PTEN-PI3kinase as a ROS effector pathway, and how synaptic terminal growth might be coupled to synaptic connectivity remain to be determined. Importantly, ROS are obligate signals for activity-regulated structural plasticity of synaptic terminals, either instructive or permissive. Targeted abrogation of neuronal ROS signalling in *Drosophila* larval motoneurons, by expression of a modified form of the DJ-1 β redox sensor (the conserved Cysteine 106 mutated to a nonoxidisable Alanine [86,91]) prevented the locomotor network from adjusting homeostatically in response to increased levels of network activity, resulting in abnormal motor output [81].

Precisely how neuronal activation leads to the generation of ROS signals is not clear and is likely context specific. In general, ROS are formed as obligate by-products of mitochondrial respiratory ATP synthesis, by ‘leakage’ of the electron transport chain [72]. Thus, mitochondrial ROS could potentially provide neurons

with a readout of their energetic demand. In addition, NMDA receptor stimulation has been shown to trigger ROS generation by either mitochondria [92,93] or NADPH oxidases [94]. NADPH oxidase activity is subject to complex regulatory pathways, of which many are associated with neuronal activation, such as elevated intracellular calcium levels, Protein kinases C and A, as well as calmodulin and calcium/calmodulin-dependent kinase II (CamKII) [34,56,95–98].

It has become increasingly evident that during normal nervous system development and function ROS act as second messengers that regulate multiple aspects, from neuronal polarity and axon growth cone behaviour to structural plasticity. As such ROS have more recently been investigated as potentially involved in the aetiology of psychiatric disorders with neurodevelopmental origins. For example, in mammalian brains cortical parvalbumin-expressing inhibitory interneurons (PVI) are critical to the excitation–inhibition balance and therefore function of many cortical networks. These fast-spiking neurons are arguably highly susceptible to ROS generated by mitochondrial ATP metabolism and it is thought that their surrounding perineuronal nets confer protection of the PVI to oxidative challenges [99]. Imbalances in ROS metabolism and buffering are thought of as potentially impacting on the development of cortical networks during so-called critical periods, leading to sub-optimal network performance and neuropsychiatric disorders [100]. Supporting this hypothesis are reports of *post mortem* brain tissue from patients with schizophrenia, bipolar or autism spectrum disorder exhibiting reductions in cortical PVI [101–103]. Moreover, a recent study did indeed find that conditions of elevated oxidative stress preceded and correlated with reduced integrity of PVI in mouse models for these disorders [104].

Synaptic transmission and plasticity

Synaptic plasticity describes the ability of synapses to adjust their strength, connectivity and structure in response to previously experienced activity. The inherent plasticity of neurons is key to neuronal network development and in networks allows for adaptation, memory and learning. Synaptic strength may be enhanced or reduced depending upon the neuronal context and the nature of stimulation, the best-studied examples being LTP and long-term depression (LTD). LTP was originally described following repetitive stimulation of the perforant path fibres to the dentate area of the hippocampus [105]. The high-frequency stimulation (HFS) used for induction of LTP results in opening of NMDA receptors and thus elevated intracellular

Ca²⁺. This leads to the adjustment of synaptic strength *via* direct and transcriptionally regulated modification of synaptic proteins, and changes in the composition of synaptic protein complexes (for review see [57,106]).

A role for ROS in synaptic plasticity has been demonstrated in various model systems and areas of the nervous system (Fig. 2). ROS production is elevated in hippocampal slice preparations following increased neuronal activity, NMDA receptor activation and subsequent LTP [92]. In mouse hippocampus NMDA receptor activation triggers ROS generation through the NOX2 NADPH oxidase, regulated by PKC [94]. Importantly, acute application of cell permeable superoxide scavengers can block HFS-induced LTP in hippocampal slices. Dysregulation of ROS *via* transgenic mis-expression of SOD1 or Catalase similarly blocked LTP, suggesting that LTP requires ROS and at the same time is sensitive to the cellular redox state [107–110]. Conversely, bath applied elevation of ROS in hippocampal slices can be sufficient to induce LTP in the CA1 region [111]. ROS are also required and sufficient for the induction and maintenance of spinal cord LTP, contributing to central sensitisation and chronic neuropathic pain [112]. Interestingly, in cerebellar Purkinje neurons superoxide is required for LTD, although in these cells synaptic depression (as opposed to potentiation) requires elevated intracellular calcium concentration [113].

The specific protein targets of ROS for regulating synaptic plasticity and the interplay between Ca²⁺ and

ROS-regulated pathways are exciting and thriving research fields [57]. In particular, the link between ROS and Ca²⁺-regulated plasticity signalling pathways raises the possibility that synaptic redox state provides a permissive environment that, depending upon context, positively or negatively tunes neuronal plasticity in response to Ca²⁺ signalling. One of the recurrent challenges is to identify the mechanisms through which ROS act, whether these are direct redox modification of effector proteins, or whether ROS impact on cellular function indirectly by modulating other signalling pathways. For example, several protein phosphatases (PP), including PP1 and PP2, PTEN and calcineurin (PP 2B), are regulated by ROS, perhaps suggesting a general role for ROS as negative regulators of kinase cascade pathways [114–119]. In the capacity of phosphatase inhibitors ROS have modulatory access to pathways known to regulate synaptic plasticity. Indeed, in response to H₂O₂ application the phosphorylation state of ERK is upregulated in hippocampal slices, cortical neurons and PC12 cells [120,121]. Similarly, activation of PKC, which is required for hippocampal LTP, is triggered by ROS [111] (Fig. 2).

Direct regulation of synaptic function by redox modification of synaptic proteins also appears likely. For example, exposure of frog NMJs to H₂O₂ revealed their capability to directly regulate synaptic release probabilities. While synaptic strength remained unchanged, the authors observed altered quantal release synchronicity when comparing proximal vs. distal parts of the NMJ, suggesting a modulatory role for




Fig. 2. ROS regulation of Synaptic Plasticity. (1) Presynaptic ROS, derived from mitochondria or NADPH oxidase activity regulates vesicle release *via* oxidation of SNAP-25 [124]. ROS regulate release probabilities with subsynaptic terminal resolution [122,123]. (2) Increases in ROS lead to activation of JNK and AP-1, which promote expression of antioxidant encoding genes [14] and others required for autophagy [79]. AP-1 mediates neuronal adaptive responses to ROS [75,76]. (3) Postsynaptic LTP, in response to HFS, drives recruitment and opening of NMDA receptors and consequent elevation of intracellular Ca²⁺ concentration. HFS causes elevated ROS production and a shift towards an oxidative environment in the synaptic terminal (4) [83,92]. HFS-induced LTP requires ROS (5) [108] and exogenous application of ROS (6) is sufficient to induce LTP in the absence of HFS [111]. ROS regulate canonical synaptic plasticity pathways *via* direct oxidative modification, and inhibition of phosphatases PP1, PP2, PTEN and Calcineurin resulting in increased kinase signalling including ERK and PKC (7) [114–121]. Also, ROS-activated PKC stimulates NADPH-oxidase activation and exacerbated ROS production (8) [107,111].

ROS or redox state with subsynaptic resolution [122,123]. In a follow-up study the same group identified SNAP-25, a component of the SNARE complex and regulator of synaptic vesicle fusion, as a direct ROS target, whose oxidation by H₂O₂ mediated down-regulation of release synchronicity [124]. Excitingly, a recent study demonstrated a role for targeted redox modifications in regulating presynaptic homeostatic adjustments of quantal content. Postsynaptic Semaphorin to presynaptic Plexin signalling activates the MICAL monooxygenase. This is known to specifically modify the actin cytoskeleton, which might impact on the presynaptic cytoskeleton in general and actin-mediated vesicle tethering in particular [125] (Fig. 2).

In summary, current evidence suggests that synaptic plasticity is regulated by both direct and indirect modes of ROS action. It will almost certainly be determined by context, such as local redox state of pre- or postsynaptic sites, the nature (e.g. hydrogen peroxide vs. superoxide vs. nitric oxide), source and subcellular localisation of ROS (e.g. mitochondrial vs. plasma membrane localised NADPH oxidase vs. cytoskeleton associated monooxygenase). In addition, different protein modifications might be subject to distinct ROS concentration thresholds, which could endow ROS signalling with a further degree of flexibility and complexity.

Conclusions

A large body of work has demonstrated the importance of ROS as signals that regulate numerous processes during neuronal development and nervous system function. With the majority of work to date having been carried out in cultured cells, it will be important to verify these by studying ROS signalling *in vivo*. While whole animal knockout mutants, for example, of NADPH oxidases, are informative, future work would ideally use cell-specific mosaic manipulations with which to unambiguously determine cell type specificity of ROS requirements. One central aspect will be to determine how ROS generation is regulated, for example, the subcellular localisation of NADPH oxidases and other ROS generators. Previous studies have reported cue-dependent changes in regulatory subunits as mechanisms for controlling ROS production [58,126]. Equally important for our understanding will be genetically encoded tools for visualising ROS and cellular redox potential, ideally with specificity for different ROS species [127–130]. Under normal physiological conditions ROS signals are expected to be confined in space and time. In the nervous system, in particular, there is a need for new reagents that will

allow genetically targeted *in vivo* manipulation of ROS generators and scavengers with spatial and temporal control appropriate for studying neuronal cell behaviour and synaptic transmission. Following the success of optogenetic modulators of neuronal excitability, this might take the route for engineering optogenetic solutions for ROS generation and sequestration [131].

Acknowledgements

We apologise to all colleagues whose work we were unable to include due to space limitations. This work was supported by BBSRC studentship to NG and BBSRC research grants to ML (BB/IO1179X/1, BB/M002934/1) and to STS (BB/I012273/1, BB/M002322/1).

References

- 1 Wang W, Fang H, Groom L, Cheng A, Zhang W, Liu J, Wang X, Li K, Han P, Zheng M *et al.* (2008) Superoxide flashes in single mitochondria. *Cell* **134**, 279–290.
- 2 Gross E, Sevier CS, Heldman N, Vitu E, Bentzur M, Kaiser CA, Thorpe C and Fass D (2006) Generating disulfides enzymatically: reaction products and electron acceptors of the endoplasmic reticulum thiol oxidase Ero1p. *Proc Natl Acad Sci USA* **103**, 299–304.
- 3 Boveris A and Chance B (1973) The mitochondrial generation of hydrogen peroxide. General properties and effect of hyperbaric oxygen. *Biochem J* **134**, 707–716.
- 4 Kukreja RC, Kontos HA, Hess ML and Ellis EF (1986) PGH synthase and lipoxygenase generate superoxide in the presence of NADH or NADPH. *Circ Res* **59**, 612–619.
- 5 O'Donnell VB and Azzi A (1996) High rates of extracellular superoxide generation by cultured human fibroblasts: involvement of a lipid-metabolizing enzyme. *Biochem J* **318** (Pt 3), 805–812.
- 6 McNally JS, Davis ME, Giddens DP, Saha A, Hwang J, Dikalov S, Jo H and Harrison DG (2003) Role of xanthine oxidoreductase and NAD(P)H oxidase in endothelial superoxide production in response to oscillatory shear stress. *Am J Physiol Heart Circ Physiol* **285**, H2290–H2297.
- 7 Spina MB and Cohen G (1989) Dopamine turnover and glutathione oxidation: implications for Parkinson disease. *Proc Natl Acad Sci USA* **86**, 1398–1400.
- 8 Martins RN, Harper CG, Stokes GB and Masters CL (1986) Increased cerebral glucose-6-phosphate dehydrogenase activity in Alzheimer's disease may reflect oxidative stress. *J Neurochem* **46**, 1042–1045.
- 9 Höhn A and Grune T (2013) Lipofuscin: formation, effects and role of macroautophagy. *Redox Biol* **1**, 140–144.

- 10 Ahlgren-Beckendorf JA, Reising AM, Schander MA, Herdler JW and Johnson JA (1999) Coordinate regulation of NAD(P)H:quinone oxidoreductase and glutathione-S-transferases in primary cultures of rat neurons and glia: role of the antioxidant/electrophile responsive element. *Glia* **25**, 131–142.
- 11 Shih AY, Johnson DA, Wong G, Kraft AD, Jiang L, Erb H, Johnson JA and Murphy TH (2003) Coordinate regulation of glutathione biosynthesis and release by Nrf2-expressing glia potently protects neurons from oxidative stress. *J Neurosci* **23**, 3394–3406.
- 12 Jimenez-Blasco D, Santofimia-Castaño P, González A, Almeida A and Bolaños JP (2015) Astrocyte NMDA receptors' activity sustains neuronal survival through a Cdk5-Nrf2 pathway. *Cell Death Differ* **22**, 1877–1889.
- 13 Longo N (2009) Disorders of bipterin metabolism. *J Inherit Metab Dis* **32**, 333–342.
- 14 Baxter PS and Hardingham GE (2016) Adaptive regulation of the brain's antioxidant defences by neurons and astrocytes. *Free Radic Biol Med* **100**, 147–152.
- 15 Baxter PS, Bell KFS, Hasel P, Kaindl AM, Fricker M, Thomson D, Cregan SP, Gillingwater TH and Hardingham GE (2015) Synaptic NMDA receptor activity is coupled to the transcriptional control of the glutathione system. *Nat Commun* **6**, 6761.
- 16 Papadia S, Soriano FX, Léveillé F, Martel M-A, Dakin KA, Hansen HH, Kaindl A, Sifringer M, Fowler J, Stefovská V *et al.* (2008) Synaptic NMDA receptor activity boosts intrinsic antioxidant defenses. *Nat Neurosci* **11**, 476–487.
- 17 Bell KFS, Al-Mubarak B, Martel M-A, McKay S, Wheelan N, Hasel P, Márkus NM, Baxter P, Deighton RF, Serio A *et al.* (2015) Neuronal development is promoted by weakened intrinsic antioxidant defences due to epigenetic repression of Nrf2. *Nat Commun* **6**, 7066.
- 18 Yogev S and Shen K (2017) Establishing neuronal polarity with environmental and intrinsic mechanisms. *Neuron* **96**, 638–650.
- 19 Schelski M and Bradke F (2017) Neuronal polarization: from spatiotemporal signaling to cytoskeletal dynamics. *Mol Cell Neurosci* **84**, 11–28.
- 20 Foreman J, Demidchik V, Bothwell JHF, Mylona P, Miedema H, Torres MA, Linstead P, Costa S, Brownlee C, Jones JDG *et al.* (2003) Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. *Nature* **422**, 442–446.
- 21 Semighini CP and Harris SD (2008) Regulation of apical dominance in *Aspergillus nidulans* hyphae by reactive oxygen species. *Genetics* **179**, 1919–1932.
- 22 Suzukawa K, Miura K, Mitsushita J, Resau J, Hirose K, Crystal R and Kamata T (2000) Nerve growth factor-induced neuronal differentiation requires generation of Rac1-regulated reactive oxygen species. *J Biol Chem* **275**, 13175–13178.
- 23 Goldsmit Y, Erlich S and Pinkas-Kramarski R (2001) Neuregulin induces sustained reactive oxygen species generation to mediate neuronal differentiation. *Cell Mol Neurobiol* **21**, 753–769.
- 24 Kamata H, Oka S-I, Shibukawa Y, Kakuta J and Hirata H (2005) Redox regulation of nerve growth factor-induced neuronal differentiation of PC12 cells through modulation of the nerve growth factor receptor, TrkA. *Arch Biochem Biophys* **434**, 16–25.
- 25 Nitti M, Furfaro AL, Cevasco C, Traverso N, Marinari UM, Pronzato MA and Domenicotti C (2010) PKC delta and NADPH oxidase in retinoic acid-induced neuroblastoma cell differentiation. *Cell Signal* **22**, 828–835.
- 26 Wilson C, Núñez MT and González-Billault C (2015) Contribution of NADPH oxidase to the establishment of hippocampal neuronal polarity in culture. *J Cell Sci* **128**, 2989–2995.
- 27 Tejada-Simon MV, Serrano F, Villasana LE, Kanterewicz BI, Wu GY, Quinn MT and Klann E (2005) Synaptic localization of a functional NADPH oxidase in the mouse hippocampus. *Mol Cell Neurosci* **29**, 97–106.
- 28 Tonks NK (2005) Redox redux: revisiting PTPs and the control of cell signaling. *Cell* **121**, 667–670.
- 29 Ménager C, Arimura N, Fukata Y and Kaibuchi K (2004) PIP3 is involved in neuronal polarization and axon formation. *J Neurochem* **89**, 109–118.
- 30 Shi S-H, Jan LY and Jan YN (2003) Hippocampal neuronal polarity specified by spatially localized mPar3/mPar6 and PI 3-kinase activity. *Cell* **112**, 63–75.
- 31 Kim RH, Peters M, Jang Y, Shi W, Pintilie M, Fletcher GC, DeLuca C, Liepa J, Zhou L, Snow B *et al.* (2005) DJ-1, a novel regulator of the tumor suppressor PTEN. *Cancer Cell* **7**, 263–273.
- 32 Kim Y-C, Kitaura H, Taira T, Iguchi-Arigo SMM and Ariga H (2009) Oxidation of DJ-1-dependent cell transformation through direct binding of DJ-1 to PTEN. *Int J Oncol* **35**, 1331–1341.
- 33 Olgún-Albuérne M and Morán J (2015) ROS produced by NOX2 control *in vitro* development of cerebellar granule neurons development. *ASN Neuro* **7**, 1–28.
- 34 Wilson C, Muñoz-Palma E, Henríquez DR, Palmisano I, Núñez MT, Di Giovanni S and González-Billault C (2016) A feed-forward mechanism involving the NOX complex and RyR-mediated Ca²⁺ release during axonal specification. *J Neurosci* **36**, 11107–11119.
- 35 Akiyama H and Kamiguchi H (2013) Second messenger networks for accurate growth cone guidance. *Dev Neurobiol* **75**, 411–422.
- 36 Gasperini RJ, Pavez M, Thompson AC, Mitchell CB, Hardy H, Young KM, Chilton JK and Foa L (2017)

- How does calcium interact with the cytoskeleton to regulate growth cone motility during axon pathfinding? *Mol Cell Neurosci* **84**, 29–35.
- 37 Aguirre J and Lambeth JD (2010) Nox enzymes from fungus to fly to fish and what they tell us about Nox function in mammals. *Free Radic Biol Med* **49**, 1342–1353.
- 38 Sirokmány G, Donkó Á and Geiszt M (2016) Nox/Duox family of NADPH oxidases: lessons from knockout mouse models. *Trends Pharmacol Sci* **37**, 318–327.
- 39 Randlett O, Poggi L, Zolessi FR and Harris WA (2011) The oriented emergence of axons from retinal ganglion cells is directed by laminin contact in vivo. *Neuron* **70**, 266–280.
- 40 Desai RA, Gao L, Raghavan S, Liu WF and Chen CS (2009) Cell polarity triggered by cell-cell adhesion via E-cadherin. *J Cell Sci* **122**, 905–911.
- 41 McNeill H, Ozawa M, Kemler R and Nelson WJ (1990) Novel function of the cell adhesion molecule uvomorulin as an inducer of cell surface polarity. *Cell* **62**, 309–316.
- 42 Dalle-Donne I, Rossi R, Giustarini D, Gagliano N, Lusini L, Milzani A, Di Simplicio P and Colombo R (2001) Actin carbonylation: from a simple marker of protein oxidation to relevant signs of severe functional impairment. *Free Radic Biol Med* **31**, 1075–1083.
- 43 Dalle-Donne I, Rossi R, Giustarini D, Gagliano N, Di Simplicio P, Colombo R and Milzani A (2002) Methionine oxidation as a major cause of the functional impairment of oxidized actin. *Free Radic Biol Med* **32**, 927–937.
- 44 Dalle-Donne I, Giustarini D, Rossi R, Colombo R and Milzani A (2003) Reversible S-glutathionylation of Cys 374 regulates actin filament formation by inducing structural changes in the actin molecule. *Free Radic Biol Med* **34**, 23–32.
- 45 Dalle-Donne I, Rossi R, Milzani A, Di Simplicio P and Colombo R (2001) The actin cytoskeleton response to oxidants: from small heat shock protein phosphorylation to changes in the redox state of actin itself. *Free Radic Biol Med* **31**, 1624–1632.
- 46 Chai YC, Ashraf SS, Rokutan K, Johnston RB and Thomas JA (1994) S-thiolation of individual human neutrophil proteins including actin by stimulation of the respiratory burst: evidence against a role for glutathione disulfide. *Arch Biochem Biophys* **310**, 273–281.
- 47 Landino LM, Hasan R, McGaw A, Cooley S, Smith AW, Masselam K and Kim G (2002) Peroxynitrite oxidation of tubulin sulfhydryls inhibits microtubule polymerization. *Arch Biochem Biophys* **398**, 213–220.
- 48 Landino LM, Moynihan KL, Todd JV and Kennett KL (2004) Modulation of the redox state of tubulin by the glutathione/glutaredoxin reductase system. *Biochem Biophys Res Commun* **314**, 555–560.
- 49 Lassing I, Schmitzberger F, Björnstedt M, Holmgren A, Nordlund P, Schutt CE and Lindberg U (2007) Molecular and structural basis for redox regulation of beta-actin. *J Mol Biol* **370**, 331–348.
- 50 Lind C, Gerdes R, Hammell Y, Schuppe-Koistinen I, von Löwenhielm HB, Holmgren A and Cotgreave IA (2002) Identification of S-glutathionylated cellular proteins during oxidative stress and constitutive metabolism by affinity purification and proteomic analysis. *Arch Biochem Biophys* **406**, 229–240.
- 51 Fratelli M, Demol H, Puype M, Casagrande S, Eberini I, Salmona M, Bonetto V, Mengozzi M, Duffieux F, Miclet E *et al.* (2002) Identification by redox proteomics of glutathionylated proteins in oxidatively stressed human T lymphocytes. *Proc Natl Acad Sci USA* **99**, 3505–3510.
- 52 Fiaschi T, Cozzi G, Raugei G, Formigli L, Ramponi G and Chiarugi P (2006) Redox regulation of beta-actin during integrin-mediated cell adhesion. *J Biol Chem* **281**, 22983–22991.
- 53 Sparaco M, Gaeta LM, Tozzi G, Bertini E, Pastore A, Simonati A, Santorelli FM and Piemonte F (2006) Protein glutathionylation in human central nervous system: potential role in redox regulation of neuronal defense against free radicals. *J Neurosci Res* **83**, 256–263.
- 54 Wang J, Boja ES, Tan W, Tekle E, Fales HM, English S, Mieyal JJ and Chock PB (2001) Reversible glutathionylation regulates actin polymerization in A431 cells. *J Biol Chem* **276**, 47763–47766.
- 55 Sakai J, Li J, Subramanian KK, Mondal S, Bajrami B, Hattori H, Jia Y, Dickinson BC, Zhong J, Ye K *et al.* (2012) Reactive oxygen species-induced actin glutathionylation controls actin dynamics in neutrophils. *Immunity* **37**, 1037–1049.
- 56 Massaad CA and Klann E (2011) Reactive oxygen species in the regulation of synaptic plasticity and memory. *Antioxid Redox Signal* **14**, 2013–2054.
- 57 Hidalgo C and Arias-Cavieres A (2016) Calcium, reactive oxygen species, and synaptic plasticity. *Physiology (Bethesda)* **31**, 201–215.
- 58 Munnamalai V, Weaver CJ, Weisheit CE, Venkatraman P, Agim ZS, Quinn MT and Suter DM (2014) Bidirectional interactions between NOX2-type NADPH oxidase and the F-actin cytoskeleton in neuronal growth cones. *J Neurochem* **130**, 526–540.
- 59 Munnamalai V and Suter DM (2009) Reactive oxygen species regulate F-actin dynamics in neuronal growth cones and neurite outgrowth. *J Neurochem* **108**, 644–661.
- 60 Taulet N, Delorme-Walker VD and DerMardirossian C (2012) Reactive oxygen species regulate protrusion efficiency by controlling actin dynamics. *PLoS ONE* **7**, e41342.
- 61 Di Filippo M, de Iure A, Giampà C, Chiasserini D, Tozzi A, Orvietani PL, Ghiglieri V, Tantucci M,

- Durante V, Quiroga-Varela A *et al.* (2016) Persistent activation of microglia and NADPH oxidase [corrected] drive hippocampal dysfunction in experimental multiple sclerosis. *Sci Rep* **6**, 20926.
- 62 Schmidt EF, Shim S-O and Strittmatter SM (2008) Release of MICAL autoinhibition by semaphorinplexin signaling promotes interaction with collapsin response mediator protein. *J Neurosci* **28**, 2287–2297.
- 63 Terman JR, Mao T, Pasterkamp RJ, Yu H-H and Kolodkin AL (2002) MICALs, a family of conserved flavoprotein oxidoreductases, function in plexin-mediated axonal repulsion. *Cell* **109**, 887–900.
- 64 Hung R-J, Pak CW and Terman JR (2011) Direct redox regulation of F-actin assembly and disassembly by Mical. *Science* **334**, 1710–1713.
- 65 Hung R-J, Yazdani U, Yoon J, Wu H, Yang T, Gupta N, Huang Z, van Berkel WJH and Terman JR (2010) Mical links semaphorins to F-actin disassembly. *Nature* **463**, 823–827.
- 66 Hung R-J, Spaeth CS, Yesilyurt HG and Terman JR (2013) SelR reverses Mical-mediated oxidation of actin to regulate F-actin dynamics. *Nat Cell Biol* **15**, 1445–1454.
- 67 Grintsevich EE, Yesilyurt HG, Rich SK, Hung R-J, Terman JR and Reisler E (2016) F-actin dismantling through a redox-driven synergy between Mical and cofilin. *Nat Cell Biol* **18**, 876–885.
- 68 Zhou Y, Gunput R-AF, Adolfs Y and Pasterkamp RJ (2011) MICALs in control of the cytoskeleton, exocytosis, and cell death. *Cell Mol Life Sci* **68**, 4033–4044.
- 69 Goshima Y, Nakamura F, Strittmatter P and Strittmatter SM (1995) Collapsin-induced growth cone collapse mediated by an intracellular protein related to UNC-33. *Nature* **376**, 509–514.
- 70 Fukata Y, Itoh TJ, Kimura T, Ménager C, Nishimura T, Shiromizu T, Watanabe H, Inagaki N, Iwamatsu A, Hotani H *et al.* (2002) CRMP-2 binds to tubulin heterodimers to promote microtubule assembly. *Nat Cell Biol* **4**, 583–591.
- 71 Morinaka A, Yamada M, Itofusa R, Funato Y, Yoshimura Y, Nakamura F, Yoshimura T, Kaibuchi K, Goshima Y, Hoshino M *et al.* (2011) Thioredoxin mediates oxidation-dependent phosphorylation of CRMP2 and growth cone collapse. *Sci Signal* **4**, ra26.
- 72 Halliwell B (1992) Reactive oxygen species and the central nervous system. *J Neurochem* **59**, 1609–1623.
- 73 Wang X and Michaelis EK (2010) Selective neuronal vulnerability to oxidative stress in the brain. *Front Aging Neurosci* **2**, 12.
- 74 Milton VJ and Sweeney ST (2012) Oxidative stress in synapse development and function. *Dev Neurobiol* **72**, 100–110.
- 75 Milton VJ, Jarrett HE, Gowers K, Chalak S, Briggs L, Robinson IM and Sweeney ST (2011) Oxidative stress induces overgrowth of the *Drosophila* neuromuscular junction. *Proc Natl Acad Sci USA* **108**, 17521–17526.
- 76 Sanyal S, Sandstrom DJ, Hoefler CA and Ramaswami M (2002) AP-1 functions upstream of CREB to control synaptic plasticity in *Drosophila*. *Nature* **416**, 870–874.
- 77 Fleischmann A, Hvalby O, Jensen V, Strekalova T, Zacher C, Layer LE, Kvello A, Reschke M, Spanagel R, Sprengel R *et al.* (2003) Impaired long-term memory and NR2A-type NMDA receptor-dependent synaptic plasticity in mice lacking c-Fos in the CNS. *J Neurosci* **23**, 9116–9122.
- 78 Tischmeyer W and Grimm R (1999) Activation of immediate early genes and memory formation. *CMLS. Cell Mol Life Sci* **55**, 564–574.
- 79 Wu H, Wang MC and Bohmann D (2009) JNK protects *Drosophila* from oxidative stress by transcriptionally activating autophagy. *Mech Dev* **126**, 624–637.
- 80 Berke B, Wittnam J, McNeill E, Van Vactor DL and Keshishian H (2013) Retrograde BMP signaling at the synapse: a permissive signal for synapse maturation and activity-dependent plasticity. *J Neurosci* **33**, 17937–17950.
- 81 Oswald M, Brooks PS, Zwart MF and Mukherjee A (2016) Reactive oxygen species regulate activity-dependent neuronal structural plasticity|bioRxiv. *bioRxiv* [Preprint].
- 82 Hongpaisan J, Winters CA and Andrews SB (2003) Calcium-dependent mitochondrial superoxide modulates nuclear CREB phosphorylation in hippocampal neurons. *Mol Cell Neurosci* **24**, 1103–1115.
- 83 Hongpaisan J, Winters CA and Andrews SB (2004) Strong calcium entry activates mitochondrial superoxide generation, upregulating kinase signaling in hippocampal neurons. *J Neurosci* **24**, 10878–10887.
- 84 Zwart MF, Randlett O, Evers JF and Landgraf M (2013) Dendritic growth gated by a steroid hormone receptor underlies increases in activity in the developing *Drosophila* locomotor system. *Proc Natl Acad Sci USA* **110**, E3878–E3887.
- 85 Meulener M, Whitworth AJ, Armstrong-Gold CE, Rizzu P, Heutink P, Wes PD, Pallanck LJ and Bonini NM (2005) *Drosophila* DJ-1 mutants are selectively sensitive to environmental toxins associated with Parkinson's disease. *Curr Biol* **15**, 1572–1577.
- 86 Wilson MA (2011) The role of cysteine oxidation in DJ-1 function and dysfunction. *Antioxid Redox Signal* **15**, 111–122.
- 87 Jordán-Álvarez S, Fouquet W, Sigrist SJ and Acebes A (2012) Presynaptic PI3K activity triggers the formation of glutamate receptors at neuromuscular terminals of *Drosophila*. *J Cell Sci* **125**, 3621–3629.
- 88 Cuesto G, Enriquez-Barreto L, Caramés C, Cantarero M, Gasull X, Sandi C, Ferrús A, Acebes A and

- Morales M (2011) Phosphoinositide-3-kinase activation controls synaptogenesis and spinogenesis in hippocampal neurons. *J Neurosci* **31**, 2721–2733.
- 89 Martín-Peña A, Acebes A, Rodríguez J-R, Sorribes A, de Polavieja GG, Fernández-Fúnez P and Ferrús A (2006) Age-independent synaptogenesis by phosphoinositide 3 kinase. *J Neurosci* **26**, 10199–10208.
- 90 Kumar V, Zhang M-X, Swank MW, Kunz J and Wu GY (2005) Regulation of dendritic morphogenesis by Ras-PI3K-Akt-mTOR and Ras-MAPK signaling pathways. *J Neurosci* **25**, 11288–11299.
- 91 Meulener MC, Xu K, Thomson L, Thompson L, Ischiropoulos H and Bonini NM (2006) Mutational analysis of DJ-1 in *Drosophila* implicates functional inactivation by oxidative damage and aging. *Proc Natl Acad Sci USA* **103**, 12517–12522.
- 92 Bindokas VP, Jordán J, Lee CC and Miller RJ (1996) Superoxide production in rat hippocampal neurons: selective imaging with hydroethidine. *J Neurosci* **16**, 1324–1336.
- 93 Dugan LL, Sensi SL, Canzoniero LM, Handran SD, Rothman SM, Lin TS, Goldberg MP and Choi DW (1995) Mitochondrial production of reactive oxygen species in cortical neurons following exposure to N-methyl-D-aspartate. *J Neurosci* **15**, 6377–6388.
- 94 Brennan AM, Won Suh S, Joon Won S, Narasimhan P, Kauppinen TM, Lee H, Edling Y, Chan PH and Swanson RA (2009) NADPH oxidase is the primary source of superoxide induced by NMDA receptor activation. *Nat Neurosci* **12**, 857–863.
- 95 Sorce S, Stocker R, Seredenina T, Holmdahl R, Aguzzi A, Chio A, Depaulis A, Heitz F, Olofsson P, Olsson T *et al.* (2017) NADPH oxidases as drug targets and biomarkers in neurodegenerative diseases: what is the evidence? *Free Radic Biol Med* **112**, 387–396.
- 96 Bánfi B, Tirone F, Durussel I, Knisz J, Moskwa P, Molnár GZ, Krause K-H and Cox JA (2004) Mechanism of Ca²⁺ activation of the NADPH oxidase 5 (NOX5). *J Biol Chem* **279**, 18583–18591.
- 97 Pandey D, Gratton J-P, Rafikov R, Black SM and Fulton DJR (2011) Calcium/calmodulin-dependent kinase II mediates the phosphorylation and activation of NADPH oxidase 5. *Mol Pharmacol* **80**, 407–415.
- 98 Tirone F and Cox JA (2007) NADPH oxidase 5 (NOX5) interacts with and is regulated by calmodulin. *FEBS Lett* **581**, 1202–1208.
- 99 Cabungcal J-H, Steullet P, Morishita H, Kraftsik R, Cuenod M, Hensch TK and Do KQ (2013) Perineuronal nets protect fast-spiking interneurons against oxidative stress. *Proc Natl Acad Sci USA* **110**, 9130–9135.
- 100 Do KQ, Cuenod M and Hensch TK (2015) Targeting oxidative stress and aberrant critical period plasticity in the developmental trajectory to schizophrenia. *Schizophr Bull* **41**, 835–846.
- 101 Beasley CL and Reynolds GP (1997) Parvalbumin-immunoreactive neurons are reduced in the prefrontal cortex of schizophrenics. *Schizophr Res* **24**, 349–355.
- 102 Hashemi E, Ariza J, Rogers H, Noctor SC and Martínez-Cerdeño V (2017) The number of parvalbumin-expressing interneurons is decreased in the medial prefrontal cortex in autism. *Cereb Cortex* **27**, 1931–1943.
- 103 Lewis DA, Curley AA, Glausier JR and Volk DW (2012) Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia. *Trends Neurosci* **35**, 57–67.
- 104 Steullet P, Cabungcal J-H, Coyle J, Didriksen M, Gill K, Grace AA, Hensch TK, LaMantia A-S, Lindemann L, Maynard TM *et al.* (2017) Oxidative stress-driven parvalbumin interneuron impairment as a common mechanism in models of schizophrenia. *Mol Psychiatry* **22**, 936–943.
- 105 Bliss TV and Lomo T (1973) Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J Physiol (Lond)* **232**, 331–356.
- 106 Herring BE and Nicoll RA (2016) Long-term potentiation: from CaMKII to AMPA receptor trafficking. *Annu Rev Physiol* **78**, 351–365.
- 107 Klann E, Roberson ED, Knapp LT and Sweatt JD (1998) A role for superoxide in protein kinase C activation and induction of long-term potentiation. *J Biol Chem* **273**, 4516–4522.
- 108 Klann E (1998) Cell-permeable scavengers of superoxide prevent long-term potentiation in hippocampal area CA1. *J Neurophysiol* **80**, 452–457.
- 109 Thiels E, Urban NN, Gonzalez-Burgos GR, Kanterewicz BI, Barrionuevo G, Chu CT, Oury TD and Klann E (2000) Impairment of long-term potentiation and associative memory in mice that overexpress extracellular superoxide dismutase. *J Neurosci* **20**, 7631–7639.
- 110 Gahtan E, Auerbach JM, Groner Y and Segal M (1998) Reversible impairment of long-term potentiation in transgenic Cu/Zn-SOD mice. *Eur J Neurosci* **10**, 538–544.
- 111 Knapp LT and Klann E (2002) Potentiation of hippocampal synaptic transmission by superoxide requires the oxidative activation of protein kinase C. *J Neurosci* **22**, 674–683.
- 112 Lee KY, Chung K and Chung JM (2010) Involvement of reactive oxygen species in long-term potentiation in the spinal cord dorsal horn. *J Neurophysiol* **103**, 382–391.
- 113 Fujii H and Hirano T (2002) Calcineurin regulates induction of late phase of cerebellar long-term depression in rat cultured Purkinje neurons. *Eur J Neurosci* **16**, 1777–1788.

- 114 Ferri A, Gabbianelli R, Casciati A, Paolucci E, Rotilio G and Carri MT (2000) Calcineurin activity is regulated both by redox compounds and by mutant familial amyotrophic lateral sclerosis-superoxide dismutase. *J Neurochem* **75**, 606–613.
- 115 Ferri A, Gabbianelli R, Casciati A, Celsi F, Rotilio G and Carri MT (2001) Oxidative inactivation of calcineurin by Cu, Zn superoxide dismutase G93A, a mutant typical of familial amyotrophic lateral sclerosis. *J Neurochem* **79**, 531–538.
- 116 Namgaladze D, Hofer HW and Ullrich V (2002) Redox control of calcineurin by targeting the binuclear Fe(2+)-Zn(2+) center at the enzyme active site. *J Biol Chem* **277**, 5962–5969.
- 117 Wu KLH, Wu C-A, Wu C-W, Chan SHH, Chang AYW and Chan JYH (2013) Redox-sensitive oxidation and phosphorylation of PTEN contribute to enhanced activation of PI3K/Akt signaling in rostral ventrolateral medulla and neurogenic hypertension in spontaneously hypertensive rats. *Antioxid Redox Signal* **18**, 36–50.
- 118 Kim W, Youn H, Kang C and Youn B (2015) Inflammation-induced radioresistance is mediated by ROS-dependent inactivation of protein phosphatase 1 in non-small cell lung cancer cells. *Apoptosis* **20**, 1242–1252.
- 119 O’Loughlen A, Pérez-Morgado MI, Salinas M and Martín ME (2003) Reversible inhibition of the protein phosphatase 1 by hydrogen peroxide. Potential regulation of eIF2 alpha phosphorylation in differentiated PC12 cells. *Arch Biochem Biophys* **417**, 194–202.
- 120 Crossthwaite AJ, Hasan S and Williams RJ (2002) Hydrogen peroxide-mediated phosphorylation of ERK1/2, Akt/PKB and JNK in cortical neurones: dependence on Ca(2+) and PI3-kinase. *J Neurochem* **80**, 24–35.
- 121 Zhang L and Jope RS (1999) Oxidative stress differentially modulates phosphorylation of ERK, p38 and CREB induced by NGF or EGF in PC12 cells. *Neurobiol Aging* **20**, 271–278.
- 122 Giniatullin AR and Giniatullin RA (2003) Dual action of hydrogen peroxide on synaptic transmission at the frog neuromuscular junction. *J Physiol (Lond)* **552**, 283–293.
- 123 Tsentsevitsky A, Nikolsky E, Giniatullin R and Bukharaeva E (2011) Opposite modulation of time course of quantal release in two parts of the same synapse by reactive oxygen species. *Neuroscience* **189**, 93–99.
- 124 Giniatullin AR, Darios F, Shakirzyanova A, Davletov B and Giniatullin R (2006) SNAP25 is a pre-synaptic target for the depressant action of reactive oxygen species on transmitter release. *J Neurochem* **98**, 1789–1797.
- 125 Orr BO, Fetter RD and Davis GW (2017) Retrograde semaphorin-plexin signalling drives homeostatic synaptic plasticity. *Nature* **550**, 109–113.
- 126 Khan SJ, Abidi SNF, Skinner A, Tian Y and Smith-Bolton RK (2017) The *Drosophila* Duox maturation factor is a key component of a positive feedback loop that sustains regeneration signaling. *PLoS Genet* **13**, e1006937.
- 127 Breckwoldt MO, Armoundas AA, Aon MA, Bendszus M, O’Rourke B, Schwarzländer M, Dick TP and Kurz FT (2016) Mitochondrial redox and pH signaling occurs in axonal and synaptic organelle clusters. *Sci Rep* **6**, 23251.
- 128 Morgan B, Van Laer K, Owusu TNE, Ezeriņa D, Pastor-Flores D, Amponsah PS, Tursch A and Dick TP (2016) Real-time monitoring of basal H₂O₂ levels with peroxiredoxin-based probes. *Nat Chem Biol* **12**, 437–443.
- 129 Zhao Y, Hu Q, Cheng F, Su N, Wang A, Zou Y, Hu H, Chen X, Zhou H-M, Huang X *et al.* (2015) SoNar, a highly responsive NAD⁺/NADH sensor, allows high-throughput metabolic screening of anti-tumor agents. *Cell Metab* **21**, 777–789.
- 130 Bilan DS, Pase L, Joosen L, Gorokhovatsky AY, Ermakova YG, Gadella TWJ, Grabher C, Schultz C, Lukyanov S and Belousov VV (2013) HyPer-3: a genetically encoded H₂O₂ probe with improved performance for ratiometric and fluorescence lifetime imaging. *ACS Chem Biol* **8**, 535–542.
- 131 Kim CK, Adhikari A and Deisseroth K (2017) Integration of optogenetics with complementary methodologies in systems neuroscience. *Nat Rev Neurosci* **18**, 222–235.
- 132 Bienert GP, Møller ALB, Kristiansen KA, Schulz A, Møller IM, Schjoerring JK and Jahn TP (2007) Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. *J Biol Chem* **282**, 1183–1192.
- 133 Viceli Dalla Sega F, Zamboni L, Fiorentini D, Rizzo B, Caliceti C, Landi L, Hrelia S and Prata C (2014) Specific aquaporins facilitate Nox-produced hydrogen peroxide transport through plasma membrane in leukaemia cells. *Biochim Biophys Acta* **1843**, 806–814.