

Feasting on terrestrial organic matter: Dining in a dark lake changes microbial decomposition

Amelia Fitch¹  | Chloe Orland¹ | David Willer¹ | Erik J. S. Emilson^{1,2}  | Andrew J. Tanentzap¹

¹Department of Plant Sciences, University of Cambridge, Cambridge, UK

²Natural Resources Canada, Great Lakes Forestry Centre, Sault Ste. Marie, Ontario,

Correspondence

Amelia Fitch, Ecosystems and Global Change Group, Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EA, UK.
Email: afitch016@gmail.com

Funding information

Gates Cambridge Trust; Natural Environment Research Council, Grant/Award Number: NE/L006561/1; Biotechnology and Biological Sciences Research Council, Grant/Award Number: BB/M011194/1

Abstract

Boreal lakes are major components of the global carbon cycle, partly because of sediment-bound heterotrophic microorganisms that decompose within-lake and terrestrially derived organic matter (t-OM). The ability for sediment bacteria to break down and alter t-OM may depend on environmental characteristics and community composition. However, the connection between these two potential drivers of decomposition is poorly understood. We tested how bacterial activity changed along experimental gradients in the quality and quantity of t-OM inputs into littoral sediments of two small boreal lakes, a dark and a clear lake, and measured the abundance of operational taxonomic units and functional genes to identify mechanisms underlying bacterial responses. We found that bacterial production (BP) decreased across lakes with aromatic dissolved organic matter (DOM) in sediment pore water, but the process underlying this pattern differed between lakes. Bacteria in the dark lake invested in the energetically costly production of extracellular enzymes as aromatic DOM increased in availability in the sediments. By contrast, bacteria in the clear lake may have lacked the nutrients and/or genetic potential to degrade aromatic DOM and instead mineralized photo-degraded OM into CO₂. The two lakes differed in community composition, with concentrations of dissolved organic carbon and pH differentiating microbial assemblages. Furthermore, functional genes relating to t-OM degradation were relatively higher in the dark lake. Our results suggest that future changes in t-OM inputs to lake sediments will have different effects on carbon cycling depending on the potential for photo-degradation of OM and composition of resident bacterial communities.

KEYWORDS

bacterial production, boreal, carbon cycling, enzyme activity, lake sediments, photo-oxidation

1 | INTRODUCTION

Inputs of terrestrially derived organic matter (t-OM) are a major driver in lake ecosystems, but the ecological consequences of changes in their quantity and quality remain poorly understood (Solomon et al., 2007). Nearly a third of all the terrestrial carbon (C) input into inland waters is buried into lake sediments (Tranvik, Downing, &

Cotner, 2016). Much of this burial, especially of larger particulate material, occurs in nearshore environments of small lakes through mechanisms including sedimentation and flocculation, where heavier matter and coagulated dissolved organic matter (DOM) fall out of suspension and accumulate (von Wachenfeldt & Tranvik, 2016; Wurtsbaugh et al., 2015). However, littoral sediments remain

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. *Global Change Biology* Published by John Wiley & Sons Ltd

understudied relative to pelagic environments, especially in the context of future changes to t-OM exports. Terrestrial organic matter is increasingly exported into receiving waters, due to mechanisms including declines in anthropogenic sulfur deposition (Evans, Monteith, & Cooper, 2005; Monteith, Stoddard, & Vesely, 1997) and increases in terrestrial primary production (Freeman, Ostle, Fenner, & Kang, 2004) and soil decomposition (Findlay, 2005), and should thereby enhance sedimentation (Kritzberg et al., 2006; von Wachenfeldt, Sobek, Bastviken, & Tranvik, 2008). Additionally, northward shifts in deciduous forests (Boisvert-Marsh, Perie, & Blois, 2014), and the spread of fires and insect outbreaks (Schindler & Lee, 2004), are expected to change the composition of t-OM that will be buried into receiving waters (e.g., Jaffé et al., 2012).

Although heterotrophic decomposition, particularly by bacteria, is the primary mechanism by which t-OM is assimilated into aquatic ecosystems (Kirchman, ; Solomon et al., 2007), relatively little is known about how and why bacterial activity changes along gradients in the quantity and quality of buried t-OM. For example, bacterial production (BP) has been found to vary by 70% across freshwater sediments with different DOM concentrations (Cole, Findlay, & Pace, 1988). One explanation for this is that bacteria assimilate low molecular weight (LMW) labile DOM quickly and easily so their production can be elevated in the presence of LMW compounds (Breggren et al., 2010; Kirchman, ; Smith & Prairie, 2009). By contrast, high molecular weight (HMW) compounds, including lignin, humic acid, and aromatic molecules, require the production of enzymes that hydrolyze or oxidize complex structures into LMW compounds before they can be assimilated (Burns et al., 2013; Fuchs, Mattias, & Heider, 2011; Kirchman, ; Sinsabaugh, Findlay, Franchini, & Fischer, 2003). Therefore, while HMW compounds are thought to be less amenable to microbial processing, the production of extracellular enzymes can allow bacteria to utilize these substrates and access both C and nutrients associated with terrestrially derived OM (Guillemette, McCallister, & Giorgio, 2016; Judd, Crump, & Kling, 2006; Kritzberg et al., 2006; Tranvik, 2009). Extracellular enzyme activity (EEA) should increase with concentrations of t-OM in lakes with limited LMW exudates (DeAngelis, Allgaier, & Chavarría, 2011; Sinsabaugh, 1997), providing a source of C and nutrients, but may result in less bacterial growth as it comes with a high metabolic cost (Chróst & Siuda, 2002; Sieczko, Maschek, & Peduzzi, 2010). Because extracellular enzymes are energetically costly to produce, bacterial growth may even be stagnant or negative with increasing concentrations of HMW OM (Chróst, 1991).

Environmental conditions, such as sediment light exposure and nutrient availability, will also interact with t-OM supply to dictate how bacterial communities will respond to future increases in the sedimentation of t-OM. For example, high light levels at the sediment–water interface can increase the availability of LMW phytoplankton exudates and photo-oxidize up to 70%–95% of the HMW t-OM found in lakes into LMW compounds (Hunting et al., 2013; Kirchman, ; Ward & Cory, 2003). This process can therefore negate the reliance of bacterial communities on EEAs. By contrast, lakes with lower sediment light exposure will have reduced oxidation of dissolved compounds at the sediment–water interface, increasing

their availability to bacteria and their ability to settle into sediments by forming organic particles (von Wachenfeldt et al., 2008). Bacteria may consequently rely on EEAs to acquire LMW C from HMW t-OM (Kirchman,). Because EEA is the rate-limiting step for making HMW C available to bacteria, the total pool of available LMW C could also be smaller in these cases (Azam & Cho, 1987; Jørgensen, 1987; Münster, 2012). Enhanced BP from LMW compounds will also depend on an adequate supply of nutrients for building proteins and cells (Goldman, Caron, & Dennett, 1987; Reche, Pace, & Cole, 2010; Russell, 2014; Smith & Prairie, 2009; Sterner, Clasen, Lampert, & Weisse, 1997).

Changes in bacterial functions involved in heterotrophic decomposition, including EEA, may also arise because the ability to degrade HMW compounds is taxonomically restricted (Bertilsson, Eiler, Nordqvist, & Jørgensen, 2007; DeAngelis et al., 2011; Ruiz-Gonzalez, Nino-García, Lapierre, & Giorgio, 2007; Zifcakova, Vetrovsky, Howe, & Baldrian, 2016). Consequently, shifts in bacterial community composition with increasing t-OM inputs will alter bacterial function (Logue et al., 2016). The functioning of whole communities can also change independently of composition if there is a high degree of redundancy among taxonomically distinct lineages (Ruiz-Gonzalez et al., 2007). However, irrespective of the underlying cause of change, no study has yet connected actual and potential functioning of sediment bacterial communities to the quantity and quality of t-OM input into lake sediments. Understanding this connection can help improve predictions of future changes in diverse ecosystem processes such as C cycling, food web production, and water quality (Solomon et al., 2007).

Here, our aim was to characterize how bacterial activity changed along fine-scale gradients in the quantity and quality of t-OM in the littoral sediments of small boreal lakes. We addressed this aim by analyzing OM from pore water collected at the interface of the sediment and overlying lake water, where t-OM is primarily deposited (Sobek et al., 2007). We focused on the effect of terrestrially sourced particulate OM as this is the primary way in which t-OM enters the littoral zone and benthic food web (Scharnweber et al., 2010; Wetzel, 2002). Boreal lakes also have varying water clarity and dissolved organic carbon (DOC) concentrations, ranging fivefold from approximately 4 to 20 mg C L⁻¹ (Sobek, Tranvik, Prairie, Kortelainen, & Cole, 1996; von Wachenfeldt & Tranvik, 2016), so can have very different responses to changes in t-OM additions because of these differences in overlying water quality (Ask, Karlsson, & Jansson, 2012). We predicted that sediment bacteria would allocate carbon differently, either to enzyme production or CO₂ production, in a clear and dark lake with increasing additions of terrestrially derived OM, and we used metagenomics to reveal the underlying mechanisms for these changes. We tested our prediction in sediment mesocosms that mirror natural ecosystems in their biogeochemical dynamics and provide a controlled way to replicate t-OM inputs across lakes with contrasting water quality and clarity (Tanentzap et al., 2014). Microbial community composition and the environment are closely linked, and a major challenge is to decouple these two effects on microbial function (Logue et al., 2016).

2 | MATERIALS AND METHODS

2.1 | Study site

Our experiment was deployed in the nearshore region of two lakes near Sudbury, Ontario Canada: Lake Laurentian (46°27'9.74"N, 80°56'35.42"W) and Swan Lake (46°21'58.96"N, 81°3'48.58"W). The sites have minimal human disturbance and are surrounded by similar early-successional forest, but differ in their overlying water quality. Swan Lake is more oligotrophic than Lake Laurentian, with mean (\pm SE) total phosphorus concentrations (mid-lake surface grabs taken during our sampling) of $9.3 \pm 0.4 \mu\text{g/L}$ vs. $35.2 \pm 2.5 \mu\text{g/L}$ respectively. Swan also had more than four times higher light levels at the sediment surface. Mean (\pm SE) light levels in Swan were $6,378 \pm 149 \text{ lx}$ vs. $1,482 \pm 28 \text{ lx}$ in Laurentian, as measured every 60 min over the duration of our experiment using Hobo UA-002-64 light loggers. The loggers were installed on the sediment surface to measure light intensity reaching the sediment–water interface of 24 mesocosms per lake. These differences were consistent with the overlying lake water, as DOC concentrations were 2 mg/L in Swan vs. 7 mg/L in Laurentian. We hereafter refer to the two lakes as the “clear” and “dark” lake, respectively.

2.2 | Experimental design

We submerged sediment amended with different types of t-OM on the bottom of the study lakes in the nearshore environment (0.30–0.75 m depth) during July 2015 after Tanentzap et al. (2014). Submergence exposed the experimental sediments to natural overlying water conditions. Briefly, sediments composed of 0%, 5%, 25%, 35%, and 50% t-OM (dry-weight basis) and locally sourced inorganic material were mixed with particle sizes and vertical structuring of all material mimicking natural lake sediments (Tanentzap et al., 2014). We then filled 17.5 L, $50.8 \times 38.1 \times 12.7$ (height) cm HDPE containers with 8 cm of sediments (total sediment volume \sim 15 L). For each t-OM quantity, material was added in a 1:2, 1:1, or 2:1 dry–mass ratio of deciduous to coniferous litterfall collected from nearby forests. Each of the 5 quantities \times 3 qualities combinations was replicated three times in each lake. Mesocosms were arranged in a block design between two sampling bays, submerged in rows at increasing distance from the shoreline, and covered with a 1×1 mm nylon mesh screen to standardize shading and resuspension within lakes. Importantly, sediment pore water samples taken from our mesocosms have been found to reflect the biogeochemistry of natural lake sediments (Tanentzap et al., 2014), allowing us to extrapolate our findings to field conditions.

The sediment manipulations interacted with lake conditions to produce experimental gradients in pore water OM quality and quantity within each lake, which we directly measured from optical properties and DOC concentrations, respectively (Supporting Information Figure S1). These pore water dissolved pools were distinct from the overlying lake water and represented the outcomes from mineralization of the added (i.e., sedimented) terrestrial OM. We then followed

recommendations to analyze our responses in relation to these continuous gradients rather than the original factorial levels (Cottingham, Lennon, & Brown, 2005). Avoiding the use of categorical variables to represent predictors that are clearly continuous, for example, levels of DOM released from sediment additions, has the added benefit of allowing us to develop predictive models that can demonstrate scientific understanding (see Houlihan, McKinney, Anderson, & McGill, 2017 for further discussion).

2.3 | Water chemistry and gas sampling

We collected pore water samples from our mesocosms during three sampling periods in June, July and August of 2016. A 3 ml polypropylene syringe was secured horizontally immediately beneath the sediment surface along one side of the HDPE container prior to submergence in the lake. The wall of the syringe that faced the sediment was removed and covered in ca. 250 μm nylon mesh. Each sampling syringe was then connected to 122 cm of nylon tubing that was purged of water before any sample collection.

On each sampling period, we extracted 45 ml of pore water into an airtight 60-ml syringe. pH was immediately measured with a handheld meter (HI 9126, Hanna Instruments, Woonsocket, RI, USA), before filtering 25 ml of each sample through a 0.5- μm glass fiber filter (Macherey-Nagel MN 85/90) and into a 20-ml glass scintillation vial. Glass vials were preacidified for a sample pH of approximately 2–3 to counter the well-documented effects of metal quenching of DOM fluorescence that becomes negligible below a pH of 3.0 (Poulin, Ryan, & Aiken, 2004; Spencer, Bolton, & Baker, 1998). In the lab, we measured two widely used DOM metrics using a Cary 60 UV Vis spectrophotometer and a Cary Eclipse fluorescent spectrophotometer (Agilent Technologies, Santa Clara, CA, USA), after analyzing the samples for DOC concentration on a Shimadzu TOC-5000A (Shimadzu Co, Columbia, MD, USA). The first DOM metric was the specific UV254 absorbance (SUVA), an indication of the average aromatic fraction of DOM per unit DOC. Higher values of SUVA indicate a more reduced state due to intact ring structures that have yet to be oxidized and tend to be HMW DOM (Lavonen et al., 2016). All UV absorbance values were corrected for total iron concentrations, because iron absorbs UV at a similar wavelength to SUVA, and can artificially increase measured SUVA values (O'Donnell, Aiken, Walvoord, & Butler, 2018; Weishaar et al., 1992). Iron was measured using the FerroVer method (Hach Company, 2014) with a Hach DR3900 spectrophotometer (HACH, Loveland, CO, USA). The second metric we measured was the humification index (HIX), for which higher values correspond to longer wavelengths of fluorescing molecules as humification of DOM proceeds (Fellman, Hood, & Spence, 2010). In addition to DOC and DOM, we also measured total dissolved nitrogen (TDN) and total dissolved phosphorus (TDP) using persulfate digestion and ascorbic acid methods, respectively (Hach Company, 2014), with a Hach DR3900 spectrophotometer. TDP and TDN values that were below the detection limit were set to the minimum detection value (0.05 and 1.0 mg/L, respectively).

We also inferred dissolved CO₂ concentration from the total inorganic carbon concentration, pH and temperature on the day of sample collection from a 45 ml water sample. 2 ml of 0.5 M HCL were injected into the sample before drawing in 15 ml of ambient air, closing the stopcock, and shaking for 2 min to equilibrate the air and acidified sample. 10 ml of the headspace air was then collected in a gas syringe for measurement of CO₂ concentration on a SRI 8610C gas chromatograph within 48 hr of collection (SRI Instruments, Torrance, CA, USA). Concentration of CO₂ in the ambient air was measured from a volume of 10 ml and used to correct the headspace measurements. We calculated final pore water concentrations of CO₂ following methods from Åberg and Wallin (2014) by applying the Bunsen solubility coefficient and ideal gas law, accounting for pH and the ambient air concentration of CO₂.

2.4 | Bacterial production

Bacterial production (BP) was measured using ³H leucine incorporation after Pace, Giorgio, Fischer, Condon, and Malcom (2014). Duplicate 1.5 ml pore water samples were collected on ice in each of June, July and August 2016 approximately two weeks after sampling water chemistry to analyze how bacteria responded to antecedent DOM conditions. In the laboratory, we added 100 µl of 17.5 nM L-[4,5-³H] leucine (1 mCi/ml, PerkinElmer, Waltham, MA, USA) to each sample. Duplicate “kill” samples were prepared for each combination of OM quality and quantity in each lake. These samples immediately received 0.3 ml of 50% trichloroacetic acid (TCA), killing any live cells, and providing a measure of background ³H leucine incorporation that was later subtracted from the incubated values. After incubating for 1 hr, each sample received 0.3 ml of 50% TCA to stop ³H leucine uptake. Samples were subsequently centrifuged at 14,243 g for 10 min and the supernatant was discarded. Each sample was then washed twice with 5% TCA, dried using a vacuum centrifuge, and stored at -20°C. ³H incorporation was later measured using a Beckman Coulter LS6500 liquid scintillation counter (Beckman Coulter, Brea, CA, USA). Prior to counting, each sample received 1 ml of Optiphase Hisafe 2 Scintillation fluid and was vortexed for 30 s. This step was repeated twice more for a total sample volume of 3 ml. Decays per minute measured on the scintillation counter were then converted to BP using standard conversion factors (Bade, Houser, & Scanga, 1998).

2.5 | Extracellular enzyme activity

We assayed five hydrolytic enzymes (β -1,4-glucosidase, β -D-1,4-cellobiosidase, β -1,4-xylosidase, leucine aminopeptidase, and phosphatase) and two oxidative enzymes (peroxidase and phenol oxidase) in pore water collected in July and August of 2016. Samples were taken one week after water chemistry sampling to test how bacterial enzyme exudation responded to DOM, and one week before BP sampling to test how enzyme activities affected subsequent bacterial C utilization. Pore water samples were assayed directly without dilution (Sinsabaugh et al., 2003), and without buffer to capture the

in situ pH (German et al., 2011). Optimal incubation times for hydrolytic enzymes were determined by standard time trials as recommended and described in German et al. (2011). We found these times to be 2 hr for β -1,4-glucosidase and leucine aminopeptidase and 1 hr for β -D-1,4-cellobiosidase, β -1,4-xylosidase, and phosphatase. Timings of oxidative incubations were performed as suggested in Sinsabaugh et al. (2004). Fluorescence and absorbance were read on a Synergy H1 microplate reader (BioTek, Winooski, VT, USA). β -D-1,4-cellobiosidase, β -1,4-xylosidase, and β -1,4-glucosidase were summed as total hydrolase activity, as these enzymes collectively target structural components of plant-derived OM (Emilsson, Kreuzweiser, Gunn, & Mykytczuk, 2016). Similarly, peroxidase and phenol oxidase activity were expressed as total oxidase activity because they have a similar functional role (Sinsabaugh, 1997; Sinsabaugh et al., 2004).

2.6 | Metagenomics shotgun sequencing

DNA was extracted in duplicate from each mesocosm in August 2016 using a PowerSoil DNA Isolation Kit (MoBio Laboratories Inc, Carlsbad, CA, USA), and following the manufacturer's protocol. Sequencing libraries were prepared with 1 ng of genomic DNA per sample using the Nextera XT DNA Sample Preparation and dual-barcoding with Nextera XT Indexes (Illumina, San Diego, CA, USA) following the manufacturer's instructions. Libraries were quantified on a Qubit 3.0 Fluorometer (ThermoFisher Scientific, Waltham, MA, USA) and on a Bioanalyzer HS DNA chip (Agilent, Santa Clara, CA) and pooled in equimolar concentrations into a single sample. Samples were sequenced on an Illumina platform using a NextSeq 500/550 Mid Output Kit v2 (300 cycles, paired-end).

Raw sequences were processed at a read depth of approximately 3.3 million reads per sample following the European Molecular Biology Laboratory-European Bioinformatics Institute (EBI) pipeline version 3.0 (Mitchell et al., 2007). In brief, the SeqPrep tool (<https://github.com/jstjohn/SeqPrep>, version 1.1) was used to merge paired-end overlapping reads, Trimmomatic (Bolger, Lohse, & Usadel, 2014, version 0.35) was used to trim low quality ends and sequences with >10% undetermined nucleotides, and sequences <100 nucleotides were removed using Biopython (Cock et al., 2009, version 1.65). Sequences were functionally annotated by predicting coding sequences (pCDS) above 60 nucleotides with FragGeneScan (Rho, Tang, & Ye, 2017, version 1.20). Read matches were then generated against pCDS using a subset of databases from InterProScan (Jones et al., 2014, version 5.19-58.0) and summarized using the Gene Ontology terms. Sequences were taxonomically annotated with QIIME (Caporaso et al., 2010, version 1.9.1). Representative 16S sequences were classified using the SILVA reference database (Quast et al., 2017, release 128) at 97% sequence identity following the open-reference Operational Taxonomic Unit (OTU) picking method with reverse strand matching enabled (Caporaso et al., 2010). The sequences were deposited in EBI Metagenomics under the project accession number ERP019980. We normalized both the functional and OTU datasets by transforming the abundance of annotated

sequences to abundances relative to the per sample total to account for sequencing bias (McMurdie & Holmes, ; Vincent, Derome, Boyle, Culley, & Charette, 2008). We focused our analysis on the subset of functional genes that matched the groupings of hydrolytic (“cellulase activity,” “glucosidase activity,” and “xylan 1,4-beta-xylosidase activity”) and oxidative enzymes (“peroxidase activity” and “catechol 1–2-dioxygenase activity”) measured for extracellular activity, as well as genes required to catabolize aromatic DOM (“aromatic compound catabolic process”) and genes for oxidoreductase activity involving oxygen (“oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen”), an important step in the decomposition of aromatics (Sinsabaugh, 1997).

2.7 | Statistical analysis

We tested the effects of DOM quantity and quality on bacterial responses using the linear mixed-effects models with the *lmer* function in the *lme4* package in R v3.2 (R Core Team, 1998). Bacterial responses included BP, CO₂ concentrations, hydrolase activity, oxidase activity, phosphatase activity, and leucine amino peptidase activity. We transformed responses with a quarter-power rather than logarithmically because small values in some of the responses resulted in heavy left-tail-distributions with the latter. Each response was then predicted as a function of DOC, SUVA and HIX, in addition to sampling month and environmental variables pH, TDN and TDP. The model for CO₂ also included BP as a predictor because CO₂ is a combined product of bacterial respiration and photo-mineralization, which can account for up to 50% and 90% of total production respectively, depending on environmental conditions (Kirchman, ; Ward & Cory, 2003). We accounted for the blocking design of our experiment, whereby the mesocosms were distributed around two sampling bays, by including bay as a fixed factor. We also accounted for repeated measurements of the same mesocosm and blocking row to account for proximity to shoreline and decreasing depth by including these factors as random effects. Finally, we allowed mean responses to vary between the dark and light lake (i.e., lake fixed effect) and tested whether the effects of DOC, DOM quality, and BP (for CO₂ response) differed between lakes with contrasting light penetration (i.e., a statistical interaction). Temperature was excluded from the models because it did not vary among mesocosms within lakes (mean ± SE of 15.85 ± 0.03C and 14.0 ± 0.02C in the dark and clear lake, respectively).

We determined the best-fitting model that explained our responses using backwards stepwise selection. Models were compared using the Akaike information criterion (AIC) and we sequentially removed predictors that did not increase AIC by more than 2, ensuring the more parsimonious models also never had AICs that were more than 2 values higher than the lowest observed value across the entire model set. Main effects were only dropped after their interactions. We assessed the significance of factors in the best supported model with *p*-values calculated from a *t*-distribution using the R package *lmerTest* (Kuznetsova, Brockhoff, & Bojesen, 2014). Degrees of freedom were adjusted with the Kenward–Roger

method, which reduces bias in small sample settings (Kenward & Roger, 1997).

We also analyzed the relative abundances of OTUs and functional genes relating to t-OM degradation to identify potential reasons for lake-dependent bacterial responses. First, we tested for a compositional difference between lake bacterial communities using a permutational multivariate analysis of variance (perMANOVA) with Bray–Curtis distances between samples. We also tested whether compositional differences were associated with sediment pore water DOC, SUVA, HIX, and pH within mesocosms of each lake by including these variables in the perMANOVA. Significance of marginal effects was assessed with 1,000 permutations of the raw data, which were restricted to each lake for the covariates, using the *adonis2* function in the R package *vegan* version 2.5-2 (Oksanen et al., 2004). We did not include TDN and TDP in these analyses to minimize collinearity; this issue was not a concern for linear mixed-effects models as none of the parameter estimates were strongly intercorrelated ($r > 0.75$). Differences in OTU relative abundance between lakes were visualized with a nonmetric multidimensional scaling (NMDS) ordination. We also used a perMANOVA with Bray–Curtis distances to test whether the relative abundances of the subset of functional genes could be explained by environmental variables and lake identity. Again, we visualized statistically significant factors ($p < 0.05$) using a NMDS ordination. In a separate perMANOVA, we tested whether the original manipulations of t-OM quantity and quality were also correlated with the composition of pore water bacterial communities and functional genes. Finally, we tested if the relative abundance of the subset of functional genes differed on average between lakes using paired *t* tests. We paired mesocosms that were in the same position in our experimental block design but located in different lakes.

Finally, we quantified the difference in bacterial community composition between the natural sediments of the two study lakes at the start of the experiment. We calculated the Bray–Curtis dissimilarity index between six pooled samples from each lake, where a value of 0 indicated communities were identical and a value closer to 1 indicated communities were entirely different, that is, no overlapping OTUs (Bloom, 1981).

3 | RESULTS

3.1 | Bacterial responses to t-OM addition depended on lake clarity

We found that BP decreased with the aromatic fraction of DOM (i.e., SUVA) across both lakes ($t = -2.04$, $df = 156$, $p = 0.042$), such as if the bacteria expended more energy to break down aromatic DOM than to grow. For example, a 25% increase above the mean aromaticity of 3.37 L mg⁻¹ cm⁻¹ decreased BP below its mean of 0.54 g C L⁻¹ day⁻¹ by 14.5% (Figure 1a). The decrease in BP with an increasing fraction of aromatic pore water DOM in the dark lake was associated with a small increase in CO₂ production ($t = -2.68$, $df = 158$, $p = 0.008$, Figure 1b), suggesting that bacteria in the dark

lake were incurring a higher metabolic and respiratory cost associated breaking down aromatic OM. Specifically, CO₂ increased by 2% above the mean value of 8.59 mg-CO₂ L⁻¹ with a 25% decrease in BP. In contrast to the dark lake, greater UV exposure at the sediment–water interface in the clear lake may have led to photo-oxidation of DOC that increased CO₂ production (lake effect: $t = 13.38$, $df = 155$, $p < 0.001$). Greater photo-degradation in the clear lake may have also resulted in a lower concentration of aromatic DOM relative to the dark lake (two-sample t test: $t = -2.64$, $df = 143$, $p = 0.009$, Figure 2a). Despite the higher pore water DOC concentration in the clear lake, of which less was aromatic and presumably more was LMW C (Figure 2a,b), BP was lower on average ($t = -6.54$, $df = 120$, $p < 0.001$). The lower BP likely arose because of nutrient limitation that was inseparable from the lake-level effect (Figure 2d–e). On average, the clear lake always had 13.6% and 15.0% less available N and P, respectively, relative to C (i.e., higher pore water C:N and C:P ratios) when comparing the same mesocosms between lakes ($t = 7.59$, $df = 118$, $p < 0.001$; $t = 16.74$, $df = 100$, $p < 0.001$). Full results of model selection are given in Supporting Information Table S1.

3.2 | Extracellular enzyme production enabled aromatic OM decomposition in the dark lake

The decrease in bacterial productivity in the dark lake corresponded with a positive association between hydrolytic and oxidative enzyme activity and the fraction of aromatic pore water DOM. Bacteria were presumably limited for LMW substrates in the dark lake and subsequently produced more hydrolytic and oxidative enzymes with higher fractions of SUVA ($t = 2.09$, $df = 99$, $p = 0.039$ and $t = 4.27$, $df = 111$, $p < 0.001$, respectively; Figure 3). For example, a 25% increase in the mean aromatic fraction of 3.37 L mg-C⁻¹ cm⁻¹ increased oxidative and hydrolytic enzyme activity by 14.6% and 11.4% above the mean values of 0.05 μmol ml⁻¹ hr⁻¹ and 230.0 ηmol ml⁻¹ hr⁻¹, respectively. There was no response to aromaticity in the clear lake for oxidative ($t = -1.33$, $df = 112$,

$p = 0.100$) or hydrolytic enzymes ($t = 0.66$, $df = 106$, $p = 0.419$), suggesting that bacteria were utilizing less terrestrially derived OM for growth. Both phosphatase and leucine aminopeptidase had no response to DOM quality or DOC concentration (Supporting Information Table S2).

3.3 | Different bacterial communities underlie varying responses to t-OM addition

We found evidence that, in addition to differences in available nutrients, bacterial activity might have also differed between the two lakes because of underlying differences in community composition and subsequent functional potential to decompose t-OM. The relative abundance of pore water bacterial OTUs differed significantly between the two lakes across all mesocosms ($F_{1,49} = 2.01$, $p = 0.001$, Figure 4a). Correspondingly, we found that genes relating to oxidative enzymes ($t = 4.42$, $df = 24$, $p < 0.001$), aromatic catabolism ($t = 9.22$, $df = 24$, $p < 0.001$) and oxidoreductase activity ($t = 18.71$, $df = 24$, $p < 0.001$) were on average 1.1–3.0 more times abundant in the darker lake where microbial activity responded more strongly to increasing fractions of aromatic OM (Figure 5a–c). There were slightly higher abundances of hydrolytic genes in the clear lake ($t = -2.38$, $df = 24$, $p = 0.025$, Figure 5d), but because enzyme expression and exudation may be more strongly induced by low concentrations of LMW substrates than the presence of HMW DOM, the clear lake bacterial communities did not produce more enzymes in response to increasing DOC quantity (Supporting Information Table S2).

The observed differences in bacterial community composition and function may have arisen because in-lake processes, including photo-oxidation at the sediment–water interface, caused the added t-OM to diverge between lakes. Communities varied with pore water DOC ($F_{1,49} = 1.42$, $p = 0.037$) and pH gradients in each lake ($F_{1,49} = 1.48$, $p = 0.009$; Figure 4a), which themselves differed between lakes (Figure 2). Functional genes capable of degrading inputs of t-OM also differed between the two lakes ($F_{1,49} = 4.13$,




FIGURE 1 Energetic tradeoffs in the dark but not the clear lake. (a) BP decreased with specific ultraviolet absorbance (SUVA) across both lakes, resulting in (b) more CO₂ production where BP was lower, but only in the dark lake. Solid lines are mean effects ±95% confidence intervals




FIGURE 2 Pore water differed between mesocosms in the dark (LAU) and clear (SWA) lake. The dark lake had (a) more SUVA, (b) lower DOC concentrations, (c) slightly higher pH, (d) lower C:P and (e) C:N ratios, (f) more BP, and (g) less CO₂ production. Nonoverlapping notches indicate differences in the two medians based on 95% confidence intervals (Chambers, Cleveland, Kleiner, & Tukey, 1983). The upper and lower whiskers extended 1.5 times the interquartile range, with points outside of this range plotted




FIGURE 3 Enzyme activities increased with aromaticity of DOM in the dark lake. (a) Oxidase and (b) hydrolase activity increased with specific UV254 absorbance (SUVA). Solid lines are mean effects \pm 95% confidence intervals

$p = 0.035$) and varied along gradients of humic DOM ($F_{1,49} = 4.63$, $p = 0.030$) and pH in sediment pore water of each lake ($F_{1,49} = 5.24$, $p = 0.031$; Figure 4b). Consistent with these results, we found that the composition of pore water bacterial communities changed with the original quantity and quality of t-OM added in each lake ($F_{1,50} = 2.75$, $p = 0.016$; $F_{3,50} = 3.48$, $p = 0.001$, respectively). The presence of functional genes mirrored these results ($F_{1,50} = 2.22$, $p = 0.001$; $F_{3,50} = 1.80$, $p = 0.001$, respectively). Both lakes also had very different resident bacterial communities at the start of our

experiment, as expected if the existing environment was responsible for altering the quality of t-OM received by the sediment bacteria (Bray–Curtis dissimilarity between lakes = 0.90).

4 | DISCUSSION

Our analysis of sediment pore water at the interface between sediments and the overlying water column in a dark and a clear lake shows that environmental conditions interact with community




FIGURE 4 Bacterial community and functional composition differ between the dark (LAU) and clear (SWA) lake. (a) Nonmetric multidimensional scaling (NMDS) of OTU abundance in individual mesocosms (stress = 0.13), where lake, pH, and dissolved organic carbon (DOC) were correlated with OTU relative abundances within mesocosms (for all, $p < 0.05$ with perMANOVA). (b) NMDS of relative abundance of functional genes related to t-OM degradation (stress = 0.02), where lake, pH, and HIX and were correlated with functional gene composition within mesocosms (for all, $p < 0.05$ with perMANOVA)




FIGURE 5 Functional genes involved in OM degradation were relatively more abundant in the dark than clear lake. Genes considered were those involved in (a) oxidase activity, which includes the sum of peroxidase and catechol, 1-2-dioxygenase gene abundances, (b) aromatic catabolism, (c) oxidoreductase activity, and (d) hydrolase activity, which is the sum of glucosidase, cellobiosidase, and xylosidase gene abundances. The line indicates a 1:1 ratio. Points above or below the line indicate a higher abundance of genes in the clear and dark lake, respectively

composition to control the functioning of bacterial communities and ultimately their responses to t-OM additions. Despite the same t-OM being added to the experimental sediments in each lake, biogeochemical processing differed. More DOC and CO₂ and less aromatic DOM was produced in the sediment pore water of the clear lake than in the dark lake. The bacterial community in the dark lake had higher relative abundances of functional genes related to t-OM decomposition, enabling greater enzyme activity. A relatively lower abundance of these functional genes limited the bacterial response to aromatic DOM in the clear lake. The overall higher amount of CO₂ production in the clear lake suggested that the bacterial community instead mineralized the photo-oxidized, LMW DOM (Hessen, 1992; Moran & Zepp, 1985). Mineralization likely arose because of nutrient limitation relative to the dark lake, where terrestrial nutrient inputs were higher and bacterial production could be sustained at a higher overall level (Figure 2f; López-Urrutia & Morán, 2012; Reche et al., 2010). As future influxes of t-OM increase across boreal lakes (Creed et al., 2018; Sobek et al., 2007), these water clarity-dependent responses suggest CO₂ release may be greater in clear lakes with high levels of photo-oxidation at the sediment surface (Lapierre, Guillemette, Breggren, & del Giorgio, 2013). Dark lakes may instead experience a decrease in primary production and shift toward retaining rather than mineralizing terrestrial carbon, such as by burying it in sediment (Gudasz et al., 2017; Isidorova et al., 2015; Seekell et al., 2014). However, over the longer term, the darkening of clear lakes will reduce light exposure on the sediment surface, increasing OM burial and encouraging bacterial communities to develop in sediment pore water that can utilize this material (Judd, Crump, & Kling, 2007; Rofner et al., 2016).

The ability for bacteria in the dark lake sediments to break down increasing fractions of terrestrially derived organic matter was possible because they had the genes to produce hydrolyzing and oxidizing enzymes (Judd et al., 2007; Ward & Cory, 2003). Although this may have allowed the dark lake bacteria to access nutrients and bioavailable growth-promoting amino acids associated with t-OM, the metabolic cost of producing enzymes likely resulted in less growth (Logue et al., 2016; López-Urrutia & Morán, 2012; Roiha, Matheiu, & Rautio, 2015; Yamashita, Fichot, Shen, Jaffé, & Benner, 2016). Additionally, bacteria in the dark lake had a limited availability of LMW substrates from photosynthetically derived and photo-oxidized OM (Bertilsson & Tranvik, 1998; Karlsson et al., 2009). The availability of LMW substrates was further limited by the rate at which extracellular enzymes could oxidize and hydrolyze HMW t-OM. As there was no resulting pool of excess LMW substrates to facilitate metabolic cycling (Chróst, 1991; Hessen, 1992), the sediment bacterial community in the dark lake consequently did not produce excess CO₂ in addition to the respiration required for basal metabolic function, and instead produced extracellular enzymes to degrade DOM with a greater terrestrial signature (Chróst & Rai, 1993; Russell, 2014; Sinsabaugh, 1997).

In the clear lake, bacterial metabolic cycling of a LMW DOC pool and direct photo-mineralization at the sediment surface likely accounted for higher CO₂ production. Previous work has shown that

sunlight exposure photo-degrades aromatic DOM and leads to higher concentrations of LMW compounds (Anesio, Graneli, Aiken, Kieber, & Mopper, 2007; Bertilsson & Tranvik, 1998; Sulzberger & Durisch-Kaiser, 2017). This process could lead to more bioavailable carbon and elevated CO₂ production (Lapierre et al., 2013; Lapierre & del Giorgio, 2014). However, growth of sediment bacteria in the clear lake may have been limited by the lower availability of N and P relative to C (per mass of DOC) (Figure 2d,e; Goldman et al., 1987; Reche et al., 2010; Smith & Prairie, 2009; Sterner, Elser, Fee, Guildford, & Chrzanowski, 2009; Sterner et al., 1997), despite the available LMW compounds. This nutrient limitation can explain why communities would have metabolically cycled C instead of allocating it to biomass, thus producing more CO₂ than the dark lake (del Giorgio & Cole, 1998; Hessen, 1992; Karlsson, Jansson, & Jonsson, 2002; Reche et al., 2010; Smith & Prairie, 2009; Zwart et al., 2016). Therefore, increasing t-OM influx may have varied effects on in-lake carbon cycling (Lapierre & del Giorgio, 2014). Photo-mineralization can also produce up to 90% of CO₂ in aquatic systems, and may have accounted for some of the CO₂ observed in the clear lake pore water (Ward & Cory, 2003). This high rate of photo-oxidation in the clear lake as opposed to the dark lake also altered the mesocosm pore water DOM quality, which may have shifted the composition of the clear lake bacterial community to taxa that were better able to degrade photoproducts, whereas the dark lake community may have been predisposed to degrade t-OM (Judd et al., 2007; Lønborg et al., 2007; Sharpless et al., 2015; Ward & Cory, 2003).

Our results add to the growing evidence that terrestrially derived OM is not recalcitrant (Judd et al., 2006; Kellerman, Dittmar, Kothawala, & Tranvik, 2014; Kritzbeg, Langenheder, & Lindstrom, 2016; McCallister & del Giorgio, 2014; Roiha et al., 2015). Rather, t-OM can undergo photo-induced transformations and be degraded by EEAs, leading to CO₂ production in lake sediments (Karlsson et al., 2012). Although bacterial activity was not directly stimulated by aromatic OM, it was relatively higher in lake sediments dominated by aromatic OM, thereby suggesting that the terrestrially sourced fraction of DOM is relatively bioavailable and has a potentially high turn-over rate (Karlsson et al., 2009; Kleber et al.,). These findings challenge the paradigm that t-OM is highly degraded, recalcitrant, and otherwise unavailable to microbial degradation due to its chemical composition (Judd et al., 2007; McCallister & del Giorgio, 2014; Sollins, Homann, & Caldwell, 2015; Zwart et al., 2016). Consequently, we suggest that t-OM inputs to lakes will not remain unchanged in lake sediments, but will be altered by microorganisms and photo-processes, which may change carbon budgets and/or transfer heterotrophic productivity to higher trophic levels (Cole, Caraco, Kling, & Kratz, 1994; Forsström, Roiha, & Rautio, 2013; McCallister & del Giorgio, 2014; Tanentzap et al., 1988). More broadly, our results advance previous work by showing how the fate of t-OM in lake sediments will ultimately depend on local environmental conditions and the bacterial community's genetic potential to degrade different OM qualities.

We found that variation in the functional genes of sediment bacteria and nutrient availability could explain differences in t-OM

utilization between two lakes of contrasting water clarity. This finding suggests that future inputs of t-OM, such as those observed across the Boreal region (Monteith et al., 1997; Pregitzer, Zak, Burton, Ashby, & Macdonald, 2013; Solomon et al., 2007), can have dramatically different outcomes for whole-lake C cycles depending on lake-specific characteristics (Ruiz-Gonzalez et al., 2007; Traving et al., 2017; Zwart et al., 2016). In clear lakes, increased inputs of t-OM may induce a positive feedback loop, whereby increased CO₂ production stimulates primary production, leading to additional inputs of LMW compounds from algae (Hare et al., 2007; Schippers, Lürling, & Scheffer, 2015). In contrast, elevated inputs of t-OM in dark lakes could lead to an increase in organic matter burial and an increasingly heterotrophic food web (Forsström et al., 2013; Gudasz et al., 2017). Future predictions of how t-OM will impact whole-lake processes clearly need to consider lake-specific characteristics, such as water clarity and nutrient availability.

ACKNOWLEDGEMENTS

Funding for this research came from a Natural Environment Research Council grant NE/L006561/1 to AJT, a Biotechnology and Biological Sciences Research Council grant BB/M011194/1 to DW, and a scholarship from the Gates Cambridge Trust to AF. We thank John Anderson, Jean-François Lapierre and an anonymous reviewer for comments that improved an earlier draft, Laura Tessier, Jim McGeer, and Michael Wilkie for help measuring bacterial production, Cyndy Desjardins, Beth Smith, Timo Rossberg, Mélanie Münch, Cassia Michel, and Jasmine Louste-Fillion for field sampling, and The Living with Lakes Centre at Laurentian University for logistical support. We are also grateful to Conservation Sudbury for allowing us to carry out our experiment in Lake Laurentian.

ORCID

Amelia Fitch  <http://orcid.org/0000-0003-1757-2219>

Erik J. S. Emilson  <http://orcid.org/0000-0002-1516-9728>

REFERENCES

- Åberg, J., & Wallin, M. B. (2014). Evaluation a fast headspace method for measuring DIC and subsequent calculation of pCO₂ in freshwater systems. *Inland Waters*, 4, 157–166. <https://doi.org/10.5268/IW-4.2.694>
- Anesio, A. M., Graneli, W., Aiken, G. R., Kieber, D. J., & Mopper, K. (2007). Effect of humic substance photodegradation on bacterial growth and respiration in lake water. *Applied and Environmental Microbiology*, 71, 6267–6275. <https://doi.org/10.1128/AEM.71.10.6267-6275.2005>
- Ask, J., Karlsson, J., & Jansson, M. (2012). Net ecosystem production in clear-water and brown-water lakes. *Global Biogeochemistry Cycles*, 26, GB1017. <https://doi.org/10.1029/2010GB003951>
- Azam, F., & Cho, B. C. (1987). Bacterial utilization of organic matter in the sea. M. Fletcher (Ed.), *Ecology of microbial communities* (pp. 261–281). Cambridge, UK: Cambridge University Press.
- Bade, D., Houser, J., & Scanga, S. (1998). *Methods of the cascading trophic interactions project*, 5th ed. Madison, WI: Center for Limnology: University of Wisconsin.
- Berggren, M., Ström, L., Laudon, H., Karlsson, J., Jonsson, A., Giesler, R., ... Jansson, M. (2010). Lake secondary production fueled by rapid transfer of low molecular weight organic carbon from terrestrial sources to aquatic consumers. *Ecology Letters*, 13, 870–880. <https://doi.org/10.1111/j.1461-0248.2010.01483.x>
- Bertilsson, S., Eiler, A., Nordqvist, A., & Jorgensen, N. O. G. (2007). Links between bacterial production, amino-acid utilization and community composition in productive lakes. *The ISME Journal*, 1, 532–544. <https://doi.org/10.1038/ismej.2007.64>
- Bertilsson, S., & Tranvik, L. J. (1998). Photochemically produced carboxylic acids as substrates for freshwater bacterioplankton. *Limnology and Oceanography*, 43, 885–895.
- Bloom, S. A. (1981). Similarity indices in community studies: Potential pitfalls. *Marine Ecology Progress Series*, 5, 125–128. <https://doi.org/10.4319/lo.1998.43.5.0885>
- Boisvert-Marsh, L., Perie, C., & De Blois, S. (2014). Shifting with climate? Evidence for recent changes in tree species distribution at high latitudes. *Ecosphere*, 5, 1–33. <https://doi.org/10.1890/ES14-00111.1>
- Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina Sequence Data. *Bioinformatics*, 30, 114–120. <https://doi.org/10.1093/bioinformatics/btu170>
- Burns, R. G., DeForest, J. L., Marxsen, J., Sinsabaugh, R. L., Stromberger, M. E., Wallenstein, M. D., ... Zoppini, A. (2013). Soil enzymes in a changing environment: Current knowledge and future directions. *Soil Biology and Biochemistry*, 58, 216–234. <https://doi.org/10.1016/j.soilbio.2012.11.009>
- Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., ... Knight, R. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nature Methods*, 7, 335–336. <https://doi.org/10.1038/nmeth.f.303>
- Chambers, J. M., Cleveland, W. S., Kleiner, B., & Tukey, P. A. (1983). Comparing data distributions. In *Graphical methods for data analysis*. Belmont, CA: Wadsworth International Group, 62 pp.
- Chróst, R. J. (1991). Environmental control of the synthesis and activity of aquatic microbial ectoenzymes. In R. J. Chróst (Ed.), *Microbial Enzymes in Aquatic Environments* (pp. 29–59). Berlin, Germany: Springer.
- Chróst, R., & Rai, H. (1993). Ectoenzyme activity and bacterial secondary production in nutrient-impooverished and nutrient-enriched freshwater mesocosms. *Microbial Ecology*, 25, 131–150. <https://doi.org/10.1007/BF00177191>
- Chróst, R. J., & Siuda, W. (2002). Ecology and Microbial Enzymes in Lake Ecosystems. In R. G. Burns, & R. P. Dick (Eds.), *Enzymes in the Environment. Activity, Ecology and Applications* (pp. 35–72). Boca Raton, FL: CRC Press.
- Cock, P. A., Antao, T., Chang, J. T., Chapman, B. A., Cox, C. J., Dalke, A., ... de Hoon, M. J. L. (2009). Biopython: Freely available Python tools for computational molecular biology and bioinformatics. *Bioinformatics*, 25, 1422–1423. <https://doi.org/10.1093/bioinformatics/btp163>
- Cole, J. J., Caraco, N. F., Kling, G. W., & Kratz, T. K. (1994). Carbon dioxide supersaturation in the surface waters of lakes. *Science*, 265, 1568–1570. <https://doi.org/10.1126/science.265.5178.1568>
- Cole, J. J., Findlay, S., & Pace, M. L. (1988). Bacterial production in fresh and saltwater ecosystems: A cross-system overview. *Marine Ecology Progress Series*, 43, 1–10. <https://doi.org/10.3354/meps043001>
- Cottingham, K. L., Lennon, J. T., & Brown, B. L. (2005). Knowing when to draw the line: Designing more informative ecological experiments. *Frontiers in Ecology and the Environment*, 3, 145–152. [https://doi.org/10.1890/1540-9295\(2005\)003\[0145:KWTDTL\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2005)003[0145:KWTDTL]2.0.CO;2)
- Creed, I. F., Bergström, A., Trick, C. G., Grimm, N. B., Hessen, D. O., Karlsson, J., ... Weyhenmeyer, G. A. (2018). Global change-driven effects on dissolved organic matter composition: Implications for food webs of northern lakes. *Global Change Biology*, 24. <https://doi.org/10.1111/gcb.14129>

- DeAngelis, K. M., Allgaier, M., & Chavarria, Y. (2011). Characterization of trapped lignin-degrading microbes in tropical forest soil. *PLOS One*, *6*, e19306. <https://doi.org/10.1371/journal.pone.0019306>
- del Giorgio, P. A., & Cole, J. J. (1998). Bacterial growth efficiency in natural aquatic systems. *Annual Review of Ecology and Systematics*, *29*, 503–541. <https://doi.org/10.1146/annurev.ecolsys.29.1.503>
- Emilsson, C. E., Kreuzweiser, D. P., Gunn, J. M., & Mykytczuk, N. C. S. (2016). Effects of land use on the structure and function of leaf-litter microbial communities in boreal streams. *Freshwater Biology*, *61*, 1049–1061. <https://doi.org/10.1111/fwb.12765>
- Evans, C. D., Monteith, D. T., & Cooper, D. M. (2005). Long-term increases in surface water dissolved organic carbon: Observations, possible causes and environmental impacts. *Environmental Pollution*, *137*, 55–71. <https://doi.org/10.1016/j.envpol.2004.12.031>
- Fellman, J. B., Hood, E., & Spence, R. G. M. (2010). Fluorescence spectroscopy opens new windows into dissolved organic matter dynamics in freshwater ecosystems: A review. *Limnology and Oceanography*, *55*, 2452–2462. <https://doi.org/10.4319/lo.2010.55.2452>
- Findlay, S. E. G. (2005). Increased carbon transport in the Hudson River: Unexpected consequence of nitrogen deposition? *Frontiers in Ecology and the Environment*, *3*, 133–137. [https://doi.org/10.1890/1540-9295\(2005\)003\[0133:ICTITH\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2005)003[0133:ICTITH]2.0.CO;2)
- Forsström, L., Roiha, T., & Rautio, M. (2013). Responses of microbial food web to increased allochthonous DOM in an oligotrophic subarctic lake. *Aquatic Microbial Ecology*, *68*, 171–184. <https://doi.org/10.3354/ame01614>
- Freeman, C., Ostle, N. J., Fenner, N., & Kang, H. (2004). A regulatory role for phenol oxidase during decomposition in peatlands. *Soil Biology and Biochemistry*, *36*, 1663–1667. <https://doi.org/10.1016/j.soilbio.2004.07.012>
- Fuchs, G., Mattias, B., & Heider, J. (2011). Microbial degradation of aromatic compounds – from one strategy to four. *Nature Reviews Microbiology*, *9*, 803–816. <https://doi.org/10.1038/nrmicro2652>
- German, D. P., Weintraub, M. N., Grandy, A. S., Lauber, C. L., Rinkes, Z. L., & Allison, S. D. (2011). Optimization of hydrolytic and oxidative enzyme methods for ecosystem studies. *Soil Biology and Biochemistry*, *43*, 1387–1397. <https://doi.org/10.1016/j.soilbio.2011.03.017>
- Goldman, J. C., Caron, D. A., & Dennett, M. R. (1987). Regulation of gross growth efficiency and ammonium regeneration in bacteria by substrate C: N ratio. *Limnology and Oceanography*, *32*, 1239–1252. <https://doi.org/10.4319/lo.1987.32.6.1239>
- Gudasz, C., Ruppenthal, M., Kalbiz, K., Cerli, C., Fiedler, S., Oelmann, Y., ... Karlsson, J. (2017). Contributions of terrestrial organic carbon to northern lake sediments. *Limnology and Oceanography*, *2*, 218–227. <https://doi.org/10.1002/lo.2.10051>
- Guillemette, F., McCallister, S. L., & del Giorgio, P. A. (2016). Selective consumption and metabolic allocation of terrestrial and algal carbon determine allochthony in lake bacteria. *The ISME Journal*, *10*, 1373–1382. <https://doi.org/10.1038/ismej.2015.215>
- Hach Company (2014). *Water Analysis Handbook*, 7th ed. Loveland, CO: Hach Company.
- Hare, C. E., Leblanc, K., DiTullio, G. R., Kudela, R. M., Zhang, Y., Lee, P. A., ... Hutchins, D. A. (2007). Consequences of increased temperature and CO₂ for phytoplankton community structure in the Bering Sea. *Marine Ecology Progress Series*, *352*, 9–16. <https://doi.org/10.3354/meps07182>
- Hessen, D. O. (1992). Dissolved organic carbon in a humic lake: Effects on bacterial production and respiration. *Hydrobiologia*, *229*, 115–123. <https://doi.org/10.1007/BF00006995>
- Houlahan, J. E., McKinney, S. T., Anderson, M. T., & McGill, B. J. (2017). The priority of prediction in ecological understanding. *Oikos*, *126*, 1–7. <https://doi.org/10.1111/oik.03726>
- Hunting, E. R., White, C. M., van Gemert, M., Mes, D., Stam, E., van der Geest, H. G., ... Admiraal, W. (2013). UV radiation and organic matter composition shape bacterial functional diversity in sediments. *Frontiers in Microbiology*, *4*, 317. <https://doi.org/10.3389/fmicb.2013.00317>
- Isidorova, A., Bravo, A. G., Riise, G., Bouchet, S., Björn, E., & Sobek, S. (2015). The effect of lake browning and respiration mode on the burial and fate of carbon and mercury in the sediment of two boreal lakes. *Journal of Geophysical Research Biosciences*, *121*, 233–245. <https://doi.org/10.1002/2015JG003086>
- Jaffé, R., Yamashita, Y., Maie, N., Cooper, W. T., Dittmar, T., Dodds, W. K., ... Watanabe, A. (2012). Dissolved organic matter in headwater streams: Compositional variability across climatic regions of North America. *Geochimica (Beijing) Et Cosmochimica Acta*, *94*, 95–108. <https://doi.org/10.1016/j.gca.2012.06.031>
- Jones, P., Binns, D., Chang, H. Y., Fraser, M., Li, W., McAnulla, C., ... Hunter, S. (2014). InterProScan 5: Genome-scale protein function classification. *Bioinformatics*, *30*, 1236–1240. <https://doi.org/10.1093/bioinformatics/btu031>
- Jørgensen, N. O. G. (1987). Free amino acids in lakes: Concentrations and assimilation rates in relation to phytoplankton and bacterial production. *Limnology and Oceanography*, *32*, 97–111. <https://doi.org/10.4319/lo.1987.32.1.0097>
- Judd, K. E., Crump, B. C., & Kling, G. W. (2006). Variation in dissolved organic matter controls bacterial production and community composition. *Ecology*, *87*, 2068–2079. [https://doi.org/10.1890/0012-9658\(2006\)87\[2068:VIDOMC\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[2068:VIDOMC]2.0.CO;2)
- Judd, K. E., Crump, B. C., & Kling, G. C. (2007). Bacterial responses in activity and community composition to photo-oxidation of dissolved organic matter from soil and surface waters. *Aquatic Sciences*, *69*, 96–107. <https://doi.org/10.1007/s00027-006-0908-4>
- Karlsson, J., Berggren, M., Ask, J., Byström, P., Jonsson, A., Laudon, H., & Jansson, M. (2012). Terrestrial organic matter support of lake food webs: Evidence from lake metabolism and stable hydrogen isotopes of consumers. *Limnology and Oceanography*, *57*, 1042–1048. <https://doi.org/10.4319/lo.2012.57.4.1042>
- Karlsson, J., Byström, P., Ask, J., Persson, L., & Jansson, M. (2009). Light limitation of nutrient-poor lake ecosystems. *Nature*, *460*, 506–509. <https://doi.org/10.1038/nature08179>
- Karlsson, J., Jansson, M., & Jonsson, A. (2002). Similar relationships between pelagic primary and bacterial production in clearwater and humic lakes. *Ecology*, *83*, 2902–2910. <https://doi.org/10.2307/3072025>
- Kellerman, A. M., Dittmar, T., Kothawala, D. N., & Tranvik, L. J. (2014). Chemodiversity of dissolved organic matter in lakes driven by climate and hydrology. *Nature Communications*, *5*, 3804. <https://doi.org/10.1038/ncomms4804>
- Kenward, M. G., & Roger, J. H. (1997). Small sample inference for fixed effects from restricted maximum likelihood. *Biometrics*, *53*, 983–997. <https://doi.org/10.2307/2533558>
- Kirchman, D. L. (2013). Degradation of organic material. In D. L. Kirchman (Ed.), *Processes in microbial ecology* (pp. 79–98). New York, NY: Oxford University Press.
- Kleber, M., Nico, P. S., Plante, A., Filley, T., Kramer, M., Swanston, C., & Sollins, P. (2011). Old and stable soil organic matter is not necessarily chemically recalcitrant: Implications for modeling concepts and temperature sensitivity. *Global Change Biology*, *17*, 1097–1107. <https://doi.org/10.1111/j.1365-2486.2010.02278.x>
- Kritzberg, E. S., Graneli, W., Björk, J., Bronmark, C., Hallgren, P., Nicolle, A., ... Hansson, L. A. (2014). Warming and browning of lakes: Consequences for pelagic carbon metabolism and sediment delivery. *Freshwater Biology*, *59*, 325–336. <https://doi.org/10.1111/fwb.12267>
- Kritzberg, E., Langenheder, S., & Lindström, E. (2006). Influence of dissolved organic matter source on lake bacterioplankton structure and function—implications for seasonal dynamics of community composition. *FEMS Microbial Ecology*, *56*, 406–417. <https://doi.org/10.1111/j.1574-6941.2006.00084.x>

- Kuznetsova, A., Brockhoff, P. B., & Bojesen, R. H. (2016) *lmerTest: Tests in linear mixed effects models. R package version 2.0-33*. Retrieved from <https://CRAN.R-project.org/package=lmerTest>
- Lapierre, J.-F., & del Giorgio, P. A. (2014). Partial coupling and differential regulation of biologically and photochemically labile dissolved organic carbon across boreal aquatic networks. *Biogeosciences*, *11*, 5969–5985. <https://doi.org/10.5194/bg-11-5969-2014>
- Lapierre, J.-F., Guillemette, F., Breggen, M., & del Giorgio, P. A. (2013). Increases in terrestrially derived carbon stimulate organic carbon processing and CO₂ emission in boreal aquatic ecosystems. *Nature Communications*, *4*. <https://doi.org/10.1038/ncomms3972>
- Lavonen, E. E., Kothawala, D. N., Tranvik, L. J., Gonsior, M., Schmitt-Kopplin, P., & Kohler, S. J. (2015). Tracking changes in the optical properties and molecular composition of dissolved organic matter during drinking water production. *Water Resources*, *85*, 286–294. <https://doi.org/10.1016/j.watres.2015.08.024>
- Logue, J. B., Stedmon, C. A., Kellerman, A. M., Nielsen, N. J., Andersson, A. F., Laudon, H., ... Kritzbeg, E. S. (2016). Experimental insights into the importance of aquatic bacterial community composition to the degradation of dissolved organic matter. *The ISME Journal*, *10*, 533–545. <https://doi.org/10.1038/ismej.2015.131>
- Lønborg, C., Nieto-Cid, M., Hernando-Morales, V., Ruiz-Hernández, M., Tiera, E., & Álvarez-Salgado, X. A. (2016) Photochemical alteration of dissolved organic matter and the subsequent effects on bacterial carbon cycling and diversity. *FEMS Microbial Ecology*, *95*, fiw048. <https://doi.org/10.1093/femsec/fiw048>
- López-Urrutia, Á., & Morán, X. A. G. (2007). Resource limitation of bacterial production distorts the temperature dependence of oceanic carbon cycling. *Ecology*, *88*, 817–822. <https://doi.org/10.1093/femsec/fiw048>
- McCallister, S. L., & del Giorgio, P. A. (2012). Evidence for the respiration of ancient terrestrial organic C in northern temperate lakes and streams. *Proceedings of the National Academy of Sciences of the United States of America*, *109*, 16963–16968. <https://doi.org/10.1073/pnas.1207305109>
- McMurdie, P. J., & Holmes, S. (2014). Waste not, want not: Why rarefying microbiome data is inadmissible. *PLOS Computational Biology*, *10*, e1003531. <https://doi.org/10.1371/journal.pcbi.1003531>
- Mitchell, A., Bucchini, F., Cochrane, G., Denise, H., Hoopen, P., Fraser, M., ... Finn, R. D. (2016). EBI metagenomics in 2016 - an expanding and evolving resource for the analysis and archiving of metagenomic data. *Nucleic Acids Research*, *44*, D595–D603. <https://doi.org/10.1093/nar/gkv1195>
- Monteith, D. T., Stoddard, J. L., & Vesely, J. (2007). Dissolved organic carbon trends resulting from changes in atmospheric deposition chemistry. *Nature*, *450*, 537–540. <https://doi.org/10.1038/nature06316>
- Moran, M. A., & Zepp, R. (1997). Role of photoreactions in the formation of biologically labile compounds from dissolved organic matter. *Limnology and Oceanography*, *42*, 1307–1316. <https://doi.org/10.4319/lo.1997.42.6.1307>
- Münster, U. (1985). Investigations about structure, distribution and dynamics of different organic substrates in the DOM of lake Plußsee. *Archiv Für Hydrobiologie Supplements.*, *70*, 429–480.
- O'Donnell, J. A., Aiken, G. R., Walvoord, M. A., & Butler, K. D. (2012). Dissolved organic matter composition winter flow in the Yukon River basin: Implications of permafrost thaw and increased groundwater discharge. *Global Biogeochemical Cycles*, *26*, GB0E06. <https://doi.org/10.1029/2012GB004341>
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., ... Wagner, H. (2018). *Vegan: Community ecology package. R package version 2.5-2*. Received from <https://CRAN.R-project.org/package=vegan>
- Pace, M. L., del Giorgio, P., Fischer, D., Condon, R., & Malcom, H. (2004). Estimates of bacterial production using the leucine incorporation method are influenced by differences in protein retention of microcentrifuge tubes. *Limnology and Oceanography, Methods*, *2*, 55–61. <https://doi.org/10.4319/lom.2004.2.55>
- Poulin, B. A., Ryan, J. N., & Aiken, G. R. (2014). Effects of iron on optical properties of dissolved organic matter. *Environmental Science and Technology*, *48*, 10098–10106. <https://doi.org/10.1021/es502670r>
- Pregitzer, K. S., Zak, D. R., Burton, A. J., Ashby, J. A., & Macdonald, N. W. (2004). Chronic nitrate additions dramatically increase the export of carbon and nitrogen from northern hardwood ecosystems. *Biogeochemistry*, *68*, 179–197. <https://doi.org/10.1023/B:BIOG.0000025737.29546.f0>
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., ... Glöckner, F. O. (2013). The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Research*, *41*, D590–D596. <https://doi.org/10.1093/nar/gks1219>
- R Core Team (2017). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- Reche, I., Pace, M. L., & Cole, J. J. (1998). Interactions of photobleaching and inorganic nutrients in determining bacterial growth on colored dissolved organic carbon. *Microbial Ecology*, *36*, 270–280. <https://doi.org/10.1007/s002489900>
- Rho, M., Tang, H., & Ye, Y. (2010). FragGeneScan: Predicting genes in short and error-prone Reads. *Nucleic Acids Research*, *38*, e191. <https://doi.org/10.1093/nar/gkq747>
- Rofner, C., Peter, H., Catalán, N., Drewes, F., Sommaruga, R., & Pérez, M. T. (2017). Climate-related changes of soil characteristics affect bacterial community composition and function of high altitude and latitude lakes. *Global Change Biology*, *23*, 2331–2344. <https://doi.org/10.1111/gcb.13545>
- Roiha, T., Matheiu, C., & Rautio, M. (2016). Allochthonous carbon is a major regulator to bacterial growth and community composition in subarctic freshwaters. *Nature Scientific Reports*, *6*, 34456. <https://doi.org/10.1038/srep34456>
- Ruiz-Gonzalez, C., Nino-Garcia, J. P., Lapierre, J. F., & Giorgio, P. A. (2015). The quality of organic matter shapes the functional biogeography of bacterioplankton across boreal freshwater ecosystems. *Global Ecology and Biogeography*, *24*, 1487–1498. <https://doi.org/10.1111/geb.12356>
- Russell, J. (2007). The energy spilling reactions of bacteria and other organisms. *Journal of Molecular Microbiology and Biotechnology*, *13*, 1–11. <https://doi.org/10.1159/000103591>
- Scharnweber, K., Syväranta, J., Hilt, S., Brauns, M., Vanni, M. j., Brothens, S., ... Mehner, T. (2014). Whole-lake experiments reveal the fate of terrestrial particulate organic carbon in benthic food webs of shallow lakes. *Ecology*, *95*, 1496–1505. <https://doi.org/10.1890/13-0390.1>
- Schindler, D. W., & Lee, P. G. (2010). Comprehensive conservation planning to protect biodiversity and ecosystem services in Canadian boreal regions under a warming climate and increasing exploitation. *Biological Conservation*, *143*, 1571–1586. <https://doi.org/10.1016/j.biocon.2010.04.003>
- Schippers, P., Lüring, M., & Scheffer, M. (2004). Increase of atmospheric CO₂ promotes phytoplankton productivity. *Ecology Letters*, *7*, 446–451. <https://doi.org/10.1111/j.1461-0248.2004.00597.x>
- Seekell, D. A., Lapierre, J. F., Ask, J., Bergström, A. K., Deininger, A., & Rodríguez, P. (2015). Karlsson J (2015) The influence of dissolved organic carbon on primary production in northern lakes. *Limnology and Oceanography*, *60*, 1276–1285. <https://doi.org/10.1002/lno.10096>
- Sharpless, C. M., Aeschbacher, M., Page, S. E., Wenk, J., Sander, M., & Kristopher, M. (2014). Photooxidation-induced changes in optical, electrochemical, and photochemical properties of humic substances. *Environmental Science and Technology*, *48*, 2688–2696. <https://doi.org/10.1021/es403925g>

- Sieczko, A., Maschek, M., & Peduzzi, P. (2015). Algal extracellular release in river-floodplain dissolved organic matter: Response of extracellular enzymatic activity during a post-flood period. *Frontiers in Microbiology*, 6, 80. <https://doi.org/10.3389/fmicb.2015.00080>
- Sinsabaugh, R. L. (2010). Phenol oxidase, peroxidase and organic matter dynamics of soil. *Soil Biology and Biochemistry*, 42, 391–404. <https://doi.org/10.1016/j.soilbio.2009.10.014>
- Sinsabaugh, R. L., Findlay, S., Franchini, P., & Fischer, D. (1997). Enzymatic analysis of riverine bacterioplankton production. *Limnology and Oceanography*, 42, 29–38. <https://doi.org/10.4319/lo.1997.42.1.0029>
- Sinsabaugh, R., Saiya-Cork, K., Long, T., Osgood, M., Neher, D., Zak, D., & Norby, R. (2003). Soil microbial activity in a Liquidambar plantation unresponsive to CO₂-driven increases in primary production. *Applied Soil Ecology*, 24, 263–271. [https://doi.org/10.1016/S0929-1393\(03\)00002-7](https://doi.org/10.1016/S0929-1393(03)00002-7)
- Smith, E. M., & Prairie, Y. T. (2004). Bacterial metabolism and growth efficiency in lakes: The importance of phosphorus availability. *Limnology and Oceanography*, 49, 137–147. <https://doi.org/10.4319/lo.2004.49.1.0137>
- Sobek, S., Durisch-Kaiser, E., Zurbrugg, R., Wongfun, N., Wessels, M., Pasche, N., & Wehrli, B. (2009). Organic carbon burial efficiency in lake sediments controlled by oxygen exposure time and sediment source. *Limnology and Oceanography*, 54, 2243–2254. <https://doi.org/10.4319/lo.2009.54.6.2243>
- Sobek, S., Tranvik, L. J., Prairie, Y. T., Kortelainen, P., & Cole, J. J. (2007). Patterns and regulation of dissolved organic carbon: An analysis of 7,500 widely distributed lakes. *Limnology and Oceanography*, 52, 1208–1219. <https://doi.org/10.4319/lo.2007.52.3.1208>
- Sollins, P., Homann, P., & Caldwell, B. A. (1996). Stabilization and destabilization of soil organic matter: Mechanisms and controls. *Geoderma*, 74, 65–105. [https://doi.org/10.1016/S0016-7061\(96\)00036-5](https://doi.org/10.1016/S0016-7061(96)00036-5)
- Solomon, C. T., Jones, S. E., Weidel, B. C., Buffam, I., Fork, M. L., Karlsson, J., ... Saros, J. E. (2015). Ecosystem consequences of changing inputs of terrestrial dissolved organic matter to lakes: Current knowledge and future challenges. *Ecosystems*, 18, 376–389. <https://doi.org/10.1007/s10021-015-9848-y>
- Spencer, G. M., Bolton, L., & Baker, A. (2007). Freeze/thaw and pH effects of freshwater dissolved organic matter fluorescence and absorbance properties from a number of UK locations. *Water Resources*, 41, 2941–2950. <https://doi.org/10.1016/j.watres.2007.04.012>
- Sterner, R. W., Clasen, J., Lampert, W., & Weisse, T. (1998). Carbon: Phosphorus stoichiometry and food chain production. *Ecology Letters*, 1, 146–150. <https://doi.org/10.1046/j.1461-0248.1998.00030.x>
- Sterner, R. W., Elser, J. J., Fee, E. J., Guildford, S. J., & Chrzanowski, T. H. (1997). The light: Nutrient ratio in lakes: The balance of energy and materials affects ecosystem structure and process. *The American Naturalist*, 150, 633–684. <https://doi.org/10.1086/286088>
- Sulzberger, B., & Durisch-Kaiser, E. (2009). Chemical characterization of dissolved organic matter (DOM): A prerequisite for understanding UV-induced changes of DOM absorption properties and bioavailability. *Aquatic Sciences*, 71, 104–126. <https://doi.org/10.1007/s00027-008-8082-5>
- Tanentzap, A. J., Szkokan-Emilson, E. J., Desjardins, C. M., Orland, C., Yakimovich, K., Dirszowsky, R., ... Gunn, J. (2017). Bridging between litterbags and whole-ecosystem experiments: A new approach for studying lake sediments. *Journal of Limnology*, 76, 431–437. <https://doi.org/10.4081/jlimnol.2017.1588>
- Tanentzap, A. J., Szkokan-Emilson, E. J., Keilstra, B. W., Arts, M. T., Yan, N. D., & Gunn, J. M. (2014). Forests fuel fish growth in freshwater deltas. *Nature Communications*, 5, 4077. <https://doi.org/10.1038/ncomms5077>
- Tranvik, L. J. (1988). Availability of dissolved organic carbon for planktonic bacteria in oligotrophic lakes of differing humic content. *Microbial Ecology*, 16, 311–322. <https://doi.org/10.1007/BF02011702>
- Tranvik, L. J., Downing, J. A., & Cotner, J. B. (2009). Lakes and reservoirs as regulators of carbon cycling and climate. *Limnology and Oceanography*, 54, 2298–2314. https://doi.org/10.4319/lo.2009.54.6_part_2.2298
- Traving, S. J., Bentzon-Tilia, M., Knudsen-Leerbeck, H., Mantikci, M., Hansen, J. L. S., Stedmon, C. A., ... Riemann, L. (2016). Coupling bacterioplankton populations and environment to community function in coastal temperate waters. *Frontiers in Microbiology*, 7, 1533. <https://doi.org/10.3389/fmicb.2016.01533>
- Vincent, A. T., Derome, N., Boyle, B., Culley, A. I., & Charette, S. J. (2017). Next-generation sequencing (NGS) in the microbiological world: How to make the most of your money. *Journal of Microbiological Methods*, 138, 60–71. <https://doi.org/10.1016/j.mimet.2016.02.016>
- von Wachenfeldt, E., Sobek, S., Bastviken, D., & Tranvik, L. J. (2008). Linking allochthonous dissolved organic matter and boreal lake sediment carbon sequestration: The role of light-mediated flocculation. *Limnology and Oceanography*, 53, 2416–2426. <https://doi.org/10.4319/lo.2008.53.6.2416>
- von Wachenfeldt, E., & Tranvik, L. J. (2008). Sedimentation in boreal lakes – The role of flocculation of allochthonous dissolved organic matter in the water column. *Ecosystems*, 11, 803–814. <https://doi.org/10.1007/s10021-008-9162-z>
- Ward, C. P., & Cory, R. M. (2016). Complete and partial photo-oxidation of dissolved organic matter draining permafrost soils. *Environmental Science and Technology*, 50, 3545–3553. <https://doi.org/10.1021/acs.est.5b05354>
- Weishaar, J. L., Aiken, G. R., Bergamaschi, B. A., Fram, M. S., Fujii, R., & Mopper, K. (2003). Evaluation of Specific Ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon. *Environmental Science and Technology*, 37(4703), 4708. <https://doi.org/10.1021/es030360x>
- Wetzel, R. (1992). Gradient-dominated ecosystems: Sources and regulatory functions of dissolved organic matter in fresh water ecosystems. *Hydrobiologia*, 229, 181–198. <https://doi.org/10.1007/BF00007000>
- Wurtsbaugh, W., Cole, J., McKnight, D., Brezonik, P., MacIntyre, S., & Potter, K. (2002). Report of The Workshop on Emerging Research Issues for Limnology: The study of Inland Waters. American Society of Limnological Oceanography.
- Yamashita, Y., Ficht, C. G., Shen, Y., Jaffé, R., & Benner, R. (2015). Linkages among fluorescent dissolved organic matter, dissolved amino acids and lignin-derived phenols in a river-influenced ocean margin. *Frontiers in Marine Science*, 2, 92. <https://doi.org/10.3389/fmars.2015.00092>
- Zifcakova, L., Vetrovsky, T., Howe, A., & Baldrian, P. (2016). Microbial activity in forest soil reflects the changes in ecosystem properties between summer and winter. *Environmental Microbiology*, 18, 288–301. <https://doi.org/10.1111/1462-2920.13026>
- Zwart, J. A., Craig, N., Kelly, P. T., Sebestyen, S. D., Solomon, C. T., Weidel, B. C., & Jones, S. E. (2016). Metabolic and physiochemical responses to a whole-lake experimental increase in dissolved organic carbon in a north-temperate lake. *Limnology and Oceanography*, 61, 723–734. <https://doi.org/10.1002/lno.10248>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Fitch A, Orland C, Willer D, Emilson EJS, Tanentzap AJ. Feasting on terrestrial organic matter: Dining in a dark lake changes microbial decomposition. *Glob Change Biol*. 2018;00:1–14. <https://doi.org/10.1111/gcb.14391>