

1 **Dietary intake using 7-day diet diaries in British men and women in the EPIC-Norfolk Study:**
2 **a focus on methodological issues**

3 Marleen AH Lentjes¹, Alison McTaggart¹, Angela A Mulligan¹, Natasha A Powell¹, David Parry-
4 Smith¹, Robert N Luben¹, Amit Bhaniani¹, Ailsa A Welch², Kay-Tee Khaw³

5

6 ¹ University of Cambridge, Department of Public Health and Primary Care, Strangeways Research
7 Laboratories, 2 Worts Causeway, Cambridge, CB1 8RN, United Kingdom.

8 ² University of East Anglia, Medical School, Diet & Health Group, Norwich Research Park,
9 Norwich, NR4 7TJ, United Kingdom.

10 ³ University of Cambridge School of Clinical Medicine, Clinical Gerontology Unit, Addenbrooke's
11 Hospital, Cambridge, CB2 2QQ, United Kingdom.

12

13 Corresponding author:

14 Marleen Lentjes

15 University of Cambridge

16 Department of Public Health and Primary Care

17 Strangeways Research Laboratories

18 2 Worts Causeway

19 Cambridge

20 CB1 8RN

21 United Kingdom

22 Telephone number: 01223 - 740 179

23 Fax: 01223 - 740 177

24 Email: marleen.lentjes@phpc.cam.ac.uk

25

26 Short title: Diet measured with 7dDD in EPIC-Norfolk

27 Keywords: Diet diary, comparison food groups, energy intake, EPIC-Norfolk

1 **Abstract**

2 We aimed to describe the energy, nutrient and crude vs. disaggregated food intake measured by 7-
3 day diet diaries (7dDD) for the full baseline Norfolk cohort recruited for the European Prospective
4 Investigation Into Cancer (EPIC-Norfolk), with emphasis on methodological issues. First data
5 collection phase took place between 1993 and 1998 in Norfolk, East-Anglia (UK). The study asked
6 25,639 men and women, aged 40-79 years, registered with a general practitioner to take part in a
7 health examination and complete a 7dDD. Diary data with at least one day was obtained for 99% of
8 the cohort; 10,354 (89.8%) of the men and 12,779 (91.5%) of the women completed all seven days.
9 Mean (SD) energy intake was 9.44 (2.22) MJ/d and 7.15 (1.66) MJ/d respectively. Energy intake
10 remained approximately stable across the days, but there was apparent underreporting among
11 participants, especially those with BMI >25 kg/m². Micronutrient density was higher among men
12 than women. We conclude that underreporting is an issue, but not more so than in national surveys.
13 How foods were grouped (crude or disaggregated) made a difference to the estimates obtained and
14 comparison of intakes showed wide limits of agreement. The choice of variables influences
15 estimates from the food group data; while this may not alter ranking of individuals within studies,
16 this issue may be relevant when comparing absolute food intakes between studies.

17

1 **Introduction**

2 The EPIC cohort is a ten country, half a million participant collaboration. Its primary purpose was
3 to elucidate diet cancer associations, however the aims of the study have broadened to incorporate
4 other exposures and health outcomes. Prior to recruitment of participants for the European
5 Prospective Investigation into Cancer in Norfolk (EPIC-Norfolk), comparisons of dietary
6 assessment methods were undertaken to establish their relative validity⁽¹⁾, as well as their
7 associations with recovery biomarkers⁽²⁾. Although it was acknowledged that the food frequency
8 questionnaire (FFQ) has its place within the wider EPIC study⁽³⁾, Norfolk participants were asked to
9 complete both an FFQ and a 7-day diet diary (7dDD) because of higher correlations with biomarker
10 data⁽³⁾, its flexibility with respect to hypotheses generation⁽²⁾ and the growing food supply⁽³⁾. The
11 open-ended prospective recording of dietary intake in the 7dDD reduces recall bias⁽⁴⁾, but is
12 substantially more laborious in terms of data entry because of data interpretation. The intakes
13 obtained from 7dDD can be used in an absolute way, but like other observational measures of
14 dietary assessment, are prone to energy underreporting and require vigilance when interpreting
15 results^(3,5). After sixteen years, the 7dDD of the full cohort of over 25,500 participants taking part
16 in the EPIC-Norfolk study are available to study diet-disease associations.

17 Health advice to the general public is given in terms of quantities of foods to be consumed, rather
18 than nutrient intake⁽⁶⁾. This brings another layer of data interpretation, *i.e.* grouping the foods
19 chosen during data-entry into substantial enough groups for statistical analysis, while maintaining
20 consistency within food groups to aid clear guidelines for public health messages. We
21 disaggregated composite dishes into their constituent parts, which in the case of meat dishes has
22 already been shown to improve precision in the estimation of the amount of meat consumed in
23 national survey^(7,8) and cohort data⁽⁹⁾. Disaggregation is important in the formulation of
24 recommendations as well as establishing sources of nutrients. However, the extent to which the
25 measures of disaggregated intakes and cruder measures are in agreement for groups such as fruit,
26 vegetables and fish is less well documented. These comparisons are of importance when food
27 group data across studies are compared, or pooled; a process that is further complicated by
28 decisions by researchers on food classification⁽¹⁰⁾ and unclear information provided by researchers
29 regarding the foods that are included or excluded from particular food groups.

30 We aimed to describe the process from data collection to data fit for statistical analysis, and the
31 different interpretation stages that were involved. We analysed population energy and nutrient
32 intakes from the largest prospective cohort who completed 7dDD. Knowing that underreporting
33 takes place, we estimated the proportion of participants who underreported. Finally we analysed

1 amounts of commonly consumed food groups, quantified in the traditional ('label-based') way and
2 using disaggregation, followed by an assessment of agreement between these two ways of data
3 interpretation.

4 **Methods**

5 EPIC-Norfolk

6 The EPIC-Norfolk study started recruitment in 1993⁽¹¹⁾. It invited 30,445 men and women between
7 the ages of 40 and 79 years via 35 general practices based in Norfolk, East-Anglia (UK), of whom
8 25,639 came for a health examination and were asked to complete a 7dDD (Table 1). In the UK
9 National Health Service, where all residents are registered with a general practitioner, practice
10 registers provide a good proxy for population based registers. The study was conducted according
11 to the Declaration of Helsinki and was approved by the Norwich District Health Authority Ethics
12 Committee and all participants gave signed informed consent.

13 Anthropometry and energy requirements

14 During their health examination, participant's weight was measured to the nearest 0.2 kg using a
15 digital scale (Salter, UK). Height was measured to the nearest millimetre using a free-standing
16 stadiometer; for both measures the participant wore light clothing and no shoes. To estimate
17 underreporting, height and weight were used to calculate Basal Metabolic Rate (BMR) using the
18 Henry equation taken from the report of the Scientific Advisory Committee on Nutrition
19 (SACN)⁽¹²⁾. Total Energy Expenditure (TEE) was estimated by multiplying the BMR with three
20 levels of assumed physical activity (PAL) in the EPIC-Norfolk population. The PALs are taken
21 from the same report and based on studies that used doubly labelled water methods to estimate
22 TEE: 1.40 (10th centile), 1.49 (25th centile) and 1.63 (50th centile, representing light physical
23 activity)⁽¹²⁾. PAL values of 1.27 are considered a minimum survival requirement; and PAL levels in
24 healthy, mobile, older adults are considered to be the same as those for adults. We included the 10th
25 centile PAL of 1.40 to represent very low levels of physical activity.

26 The 7-day Diet Diary (7dDD)

27 The 7dDD is an A5-booklet with four pages for each day to record the foods and drinks consumed
28 over seven meal occasions (before breakfast, breakfast, between breakfast and lunch, lunch,
29 between lunch and dinner, dinner and after dinner), based on the diary used in the National Survey
30 of Health and Development⁽¹³⁾. For each day there is a separate area for recipe notation and a
31 checklist of commonly consumed, but often forgotten, foods. The last four pages in the diary are in

1 the style of a general questionnaire where details regarding types of milk, bread and spread are
2 recorded to aid data-entry in case participants did not provide enough detail in their 7dDD.

3 During the participant's health examination, a nurse performed a 24-hour diet recall (24hDR)
4 according to standardised protocol⁽¹⁴⁾ and explained how to complete the booklet and the amount of
5 detail that would be necessary in order to analyse the diaries, avoiding evaluation of a participant's
6 diet and using the aforementioned checklist to aid participant's memory. Participants completed the
7 remaining six days at their home. They were asked to write down the type and amount of foods
8 consumed at the time of consumption. Portions could be estimated by using either household
9 measures (such as teaspoons, mugs etc.), one of the seventeen colour-print photos of commonly
10 consumed foods/dishes or by recording weights from packaging. They returned the diary to the
11 study centre by post, where it was recorded as 'returned' and immediately stored; no contact with
12 the participant was attempted.

13 Data-entry

14 Returned diaries were initially selected for data-entry as a series of nested case-control analyses.
15 The diaries were entered into Data Into Nutrients for Epidemiological Research (DINER), a
16 computer-based coding system developed in-house, described by Welch *et al.*⁽¹⁴⁾. As resources
17 permitted, diaries were entered between 1996 and 2011 by 1-6 trained data-entry clerks, who were
18 blinded to the case or control status of the participant. Over 11,000 food items and nearly 600
19 portions were available to choose from by the time data-entry was completed. To guide the data-
20 entry clerks' work and ensure consistency, a manual was developed and maintained that explained
21 common situations and the decisions to be made.

22 Data cleaning

23 The process of converting handwritten 7dDD into a digital format involved a two-part cleaning
24 process, covering database and diary related errors.

25 *Database checks.* The database checks ensured that food items and their associated data such as
26 portion sizes, nutrient quantities, density, cooking loss, water gain and edible part fractions, were
27 kept consistent across the various source tables.

28 The nutrient data from the 5th edition of McCance & Widdowson Composition of Foods
29 (CoF)⁽¹⁵⁾ and the ten supplement books⁽¹⁶⁻²⁵⁾ were comprehensively checked and missing values
30 were completed for carotenoids, vitamin C, iron, vitamin D, vitamin E and vitamin K. The nutrient
31 data has expanded extensively to include a total of 32 fat fractions, 16 phytoestrogens^(26,27), 6

1 phytosterols⁽²⁸⁾, haem and non-haem iron and 35 distinct flavonoids. For the majority of the added
2 phytochemicals, the nutrient quantity takes the ranges into account, *i.e.* the variety found in
3 published and/or analysed data, by creating separate nutrients for ‘minimum’, ‘median/mean’ and
4 ‘maximum’ nutrient amounts.

5 *Diary checks.* The first checks on a 7dDD were done after entry by the data-enterers where a
6 supplementary program to DINER identified meal times missed or extreme portion sizes. Most
7 checks however were done by the nutritionists’ team using a suite of in-house designed programs.
8 Notes on how to use the different parts were compiled in a single user manual CHEDDAR (Correct
9 Handling of EPIC-Norfolk Data Diminishes Awful Results)⁽²⁹⁾. CHEDDAR ensured similar
10 handling and interpretation of the computer output, as well as an explanation of data management
11 for both paper and digital diary data.

12 The programs for data cleaning and calculation have seen two extensive revisions in the past six
13 years, the first one at the start of the MRC Centre for Nutritional Epidemiology in Cancer
14 Prevention and Survival in 2007⁽³⁰⁾. The original program relied on expensive commercial software
15 and it was thought beneficial to use open-source software. The checking and calculation programs
16 have been called ‘DINERMO’ since its inception (a name reflecting ‘moving onwards’ from data-
17 entry). The checks described by Welch *et al.*⁽¹⁴⁾ were incorporated into DINERMO, and these were
18 extended with date validity checks, whereby diaries containing less than three meal times per day
19 were evaluated for validity; more detailed portion size allocation checks, such as suitability of
20 portion defaults; improvements on day completion checks and a general improvement of interface
21 and data output provided. The second revision of DINERMO took place over the last two years and
22 focussed on making the checking process more efficient by merging elements of checking and
23 calculation together as well as improving computer efficiency. We also incorporated the food
24 group calculation into this version, and revised the user manual to reflect these changes, which is
25 now called EDAM (EPIC-Norfolk Diary All-in-one Method).

26 Output

27 *Nutrients.* The recently revised nutrient calculation program calculates all 208 nutrient quantities
28 (and food group data) for all 25,507 diaries much more rapidly due to parallel processing
29 capabilities. Each food item is calculated for the full range of nutrients. This ensures a high level
30 of flexibility since data can be summed and averaged to provide nutrient variables that can be
31 compared between the seven individual days (or any other time element in the diary, such as per
32 meal) or averaged over the number of days completed. The nutrient intake data can also be

1 combined with the food group data to provide information on sources of nutrients (*e.g.* the amount
2 of vitamin C derived from vegetables).

3 *Food group data.* Food group data enables us to analyse the data in different ways. Most of the
4 food groups are hierarchical in nature. Similar foods are grouped together in the same category, and
5 in the crudest grouping system available, foods are labelled with names such as ‘vegetable’, ‘meat’,
6 ‘dairy’ etc. (Table 2). A detailed food grouping system creates sub categories such as
7 ‘brassicaceae’, ‘beef’, ‘cheese’. Another group focuses on all fruit and vegetable (F&V) varieties
8 (*e.g.* further categorising brassicaceae into vegetables such as Brussels sprouts, cabbage, broccoli)
9 and their respective preparation methods as being either ‘raw/fresh’, ‘cooked’, ‘dried’, ‘sauce/soup’,
10 ‘juice’ or ‘dish’. Specific groups, not fitting into the hierarchy, were created for several projects
11 such as a group which matches the 7dDD food items to their respective FFQ item (if present), dairy
12 food groups and a group for canned products. For dairy, the food items were characterised on three
13 elements: dairy source (milk, cheese, cream, butter, yogurt etc), dairy fat content (skimmed/semi-
14 skimmed/whole; double/single; full fat/reduced fat; categories of the percentage of fat in spreads)
15 and subjective dairy content (100% dairy/high dairy/low dairy/non-dairy).

16 The above crude, qualitative food groupings have in common that the food item (*e.g.* custard) can
17 only be categorised into one category (*e.g.* cereal, other) of the same food group (‘crude’). Another,
18 quantitative set of groupings identifies the fractions of a food item that are fruit, vegetable, red
19 meat, white meat, processed meat, fatty fish and white fish (Table 2). For example, veal stewed in
20 tomato sauce would be classified as 40% red meat *and* 55% vegetable (though in the crude group
21 100% of the weight consumed would have been assigned to ‘meat’). Fractions for disaggregation
22 were mainly obtained by calculating recipes published in the CoF⁽¹⁵⁾ and its supplements^(16–25), as
23 well as collected manufacturers’ data for commercial products; an approach which is similar to
24 methods applied in other food databases^(7,9).

25 Although a distinction was made between the qualitative and quantitative food groups, they can be
26 combined (Table 2). For example, ‘Apple crumble’ contains ‘apple, used in dishes’ according to
27 the F&V variety group, multiplying the portion size consumed with the fruit fraction group (here
28 0.62) will give the amount of cooked apple in this dish. Another example, an estimate of a
29 minimum and maximum likely amount of canned products within (homemade) food items can be
30 obtained by adding the percentages in the latter seven columns in Table 2 (*e.g.* tuna and sweetcorn
31 sandwich filling, though classified as a non-canned food item, could contain as much as 85% of
32 canned products).

1 Statistical analysis

2 All analyses were stratified by sex. We calculated the mean, median, standard deviation, 2.5th and
3 97.5th percentile of energy intake (EI), macronutrients and a selection of micronutrients.
4 Differences between the sexes were tested using Mann-Whitney test; a *P*-value of <0.05 was
5 considered significant. In order to compare energy adjusted intakes with nationally representative
6 data^(31,32) we stratified the results by age (≤ 65 years; > 65 years).

7 The mean of a participant's EI and TEE was plotted against the difference (TEE-EI) to test for
8 agreement using Bland-Altman plots⁽³³⁾. Approximately 95% of the observations are between the
9 limits of agreement ($\text{mean}_{\text{diff}} \pm (1.96 * \text{SD}_{\text{diff}})$). Underreporters were crudely defined as participants
10 with a difference in TEE - EI that resulted in values greater than 0; the degree of underreporting was
11 expressed in three categories (0-1 MJ/d, 1-2 MJ/d, ≥ 2 MJ/d). We stratified the results by body mass
12 index (BMI, ≤ 25 kg/m², > 25 kg/m²), but without any further adjustment or transformation of the
13 data.

14 A similar descriptive analysis was completed for meat, fish, fruit and vegetable consumption,
15 comparing the crude (qualitative) grouping and the disaggregated (quantitative) food groupings
16 (Table 3). The Spearman correlations of these food group data were calculated and Bland-Altman
17 plots were created to assess agreement. All analyses were conducted using SPSS v.19 (IBM).

18 **Results**

19 Response rate 7dDD

20 At their health examination, 25,639 participants were asked about their dietary habits during an
21 interviewed 24hDR (which formed the first day of the 7dDD). Only 132 participants did not
22 complete this. Diary data was obtained for 11,535 (99.4%) of the men, of whom 10,354 (89.8%)
23 returned a fully completed 7dDD (data for seven days). Among women this was 13,972 (99.6%)
24 and 12,779 (91.5%) respectively.

25 Energy intake

26 The mean energy consumed in men was 9.44 MJ/d with a standard deviation (SD) of 2.22 MJ/d and
27 in women 7.15 MJ/d with a SD of 1.66 MJ/d. However, EI was 0.4 MJ/d lower among overweight
28 and obese participants ($P < 0.001$) compared to participants with a BMI ≤ 25 kg/m² (Table 3). The
29 95% confidence interval of the mean difference between EI and TEE (TEE-EI) showed wide limits
30 of agreement and were associated with the mean, with marked differences between sexes and BMI
31 categories (Table 3). For the median PAL-level of 1.63, the non-stratified mean (SD) difference

1 between TEE and EI was 1.79 (1.98) MJ/d; we therefore created three levels of energy
2 disagreement, <1; 1-2; ≥ 2 MJ/d). The estimated proportion of underreporters (assuming all levels
3 of disagreement were correctly identified) could be as large as 39-86% among men and 49-91%
4 among women, depending on the assumed PAL-level. The degree of estimated underreporting was
5 lower in participants with a BMI ≤ 25 kg/m², but 2-3 times as many participants underreported by
6 > 2 MJ/d when overweight/obese. We explored whether the length of the 7dDD contributed to the
7 mean lower EI, but found no evidence of declining EI as diary completion progressed (Figure 1).

8 Nutrient intake

9 There were small, but mostly significant differences in men and women, when comparing the two
10 age groups (≤ 65 vs. > 65 years) in their contribution of macronutrient intake to total energy
11 consumption (Figure 2). Energy was mainly provided by carbohydrates, followed by fat, protein
12 and alcohol. Of the energy providers, only sugars were consumed in a greater proportion by women
13 than by men. Micronutrient intake, with the exception of vitamin C, was significantly higher
14 among men compared to women ($P < 0.001$), although when expressed per MJ of EI, women
15 consumed a more nutrient-dense diet than men (Table 4).

16 Intake of fish, meat, fruit and vegetables

17 Table 5 shows that mean intakes of foods consumed were influenced by aggregation or
18 disaggregation of the data. For F&V, the crude groupings underestimated the amounts consumed;
19 however, for meat and fish the quantities were overestimated with the crude grouping system. The
20 Spearman correlations between the crude and disaggregated data varied between .85 and .99 (Table
21 5). Despite these high correlations, the agreement between the two measures was low (Figure 3).
22 For example, disaggregated meat consumption could be 93 g/d higher or 39 g/d lower than the
23 crude measure of meat consumption. We observed for all food groups that with higher mean
24 consumption, the differences between the two food grouping methods became larger
25 (heteroscedasticity).

26 Discussion

27 We have shown the developments and capacities of the DINERMO nutrient and food group
28 calculation programs on 25,507 participants (99%) who contributed diary data at baseline to the
29 EPIC-Norfolk cohort. Their mean EI remained stable across the diary days. There was
30 underreporting of EI, especially among overweight/obese participants. Micronutrient density was
31 higher among women than men. The food groups compared correlated highly, but they disagreed
32 greatly in absolute quantities; with the biggest differences for meat and vegetables.

1 In nutritional epidemiology, underreporting of EI is well established. Results from the OPEN study
2 have shown energy underreporting to be prevalent in 21% of the men (by reporting 10% lower
3 energy intakes than required) and 22% of the women (14% lower energy intakes than required)⁽⁵⁾.
4 Measurements of urinary nitrogen and potassium excretion in a sub-cohort of EPIC-Norfolk
5 confirmed that underreporting takes place⁽³⁾. When EI was graphed against BMR quintiles, PAL
6 values diminished with increasing BMR and these PAL values were relatively low (1.22-1.33)⁽³⁾. In
7 this analysis, we wanted to be able to express underreporting in energy amounts, and not in PALs as
8 is common with the Goldberg criteria⁽³⁴⁾. Hence we chose to apply a Bland-Altman plot to assess
9 agreement between TEE and EI, and compensated our crude definition of defining even small
10 deviations above zero to mean underreporting, by including three levels of energy disagreement.
11 We used age, weight and height as biomarkers for energy requirement, and calculated the BMR
12 with the Henry equations that are used to estimate energy requirements in the UK⁽¹²⁾. This BMR
13 formula has indicated that 79% can be categorised within $\pm 10\%$ of the measured resting metabolic
14 rate⁽¹²⁾. Results showed that the TEE exceeded EI in at least 40-50% of the EPIC-Norfolk cohort.
15 Although these proportions of underreporters cannot be directly compared against doubly labelled
16 water methods nor the Goldberg cut-offs, since we did not account for error in both the TEE as well
17 as the EI, we saw similar trends: women were more likely to underreport and that underreporting
18 increased in overweight/obese participants. However, EI in the 50-65 age category in the National
19 Diet and Nutrition Survey (NDNS) reports a mean (SD) of 9.55 (2.38) MJ/d for men and 6.91
20 (1.74) MJ/d for women⁽³¹⁾, which is close to the EI in EPIC-Norfolk. A similar comparison can be
21 made with the 65-74 age category in the survey for people over 65 years old where men consumed a
22 mean (SD) of 8.21 (1.97) MJ/d and women 6.07 (1.38) MJ/d⁽³²⁾. The percentage of energy derived
23 from protein and fat were slightly lower and carbohydrate and alcohol were slightly higher than in
24 EPIC-Norfolk⁽³²⁾. Underreporting in EPIC-Norfolk is hence comparable and the small differences
25 in EI with these surveys are likely to be because of sampling (different age distributions) as well as
26 differences in data processing programs.

27 The EI was approximately stable during diary completion. This is encouraging considering others
28 have observed a downward trend with diaries exceeding three days⁽⁴⁾. However, the diary days are
29 (mostly) consecutive and measures of variety could as a result be lower than expected. This has
30 been compensated by requesting participants to complete another 7dDD after 18 months⁽³⁵⁾; a
31 subset of these repeat 7dDD has been used to correct odds ratios for measurement error due to
32 variation in nutrient intake⁽³⁰⁾.

33 The micronutrient data presented here do not include sources of dietary supplements. Supplements
34 are being used by 40% of the cohort⁽³⁶⁾ and have shown to change the nutrient intake distribution⁽³⁷⁾.

1 How this affects the proportions below the Estimated Average Requirements or above the Safe
2 Upper Levels in this cohort is still to be assessed.

3 A food is more than the sum of its nutrients and public health messages are given in types and
4 amounts of foods to consume; hence many studies and surveys tend to analyse food consumption
5 rather than nutrient consumption, but as a result, comparisons of study results become more
6 complicated. We were unable to compare the consumption data of meat and fish due to differences
7 in groupings of foods in the NDNS data; however, the fruit consumption in EPIC-Norfolk appeared
8 to be up to 15% (men) and 18% (women) higher and vegetable consumption ranged from 4% lower
9 (men) up to 13% higher (women)⁽³⁸⁾. These differences became more pronounced when comparing
10 disaggregated weights and reached 27% for fruit and 8% for vegetables in men and 20% for both
11 fruit and vegetables in women. The F&V consumption after disaggregation could add as much as
12 0.5 to 1 portion to a participant's 5-a-Day; however, it is still unclear whether health effects of F&V
13 are similar when used in a dish such as an apple pie or cauliflower cheese.

14 The EPIC-Norfolk 7dDD data are the largest data collection of its kind. Until now, the only dietary
15 data from the full EPIC-Norfolk cohort was based on an FFQ⁽³⁹⁾, which is known to overestimate
16 fruit, vegetable and milk consumption⁽³⁾. Moreover, the 7dDD enables us to study diet variety and
17 meal patterns, which are areas that have shown potential for intervention^(40,41).

18 The computer programs described are still undergoing development and we hope to change the
19 calculation method to an approach that separates all ingredients within dishes, similar as described
20 by Subar *et al.*⁽⁵⁾, providing a fully 'matured' system which we have named PECORINO, (Precision
21 in EPIC-Norfolk: Calculation Of Recipes Improves Nutrient Output). This system will have several
22 advantages, firstly, the reporting of disaggregated amounts would no longer be restricted to the
23 seven groups mentioned in this paper; secondly, any extension of the food database with nutrient or
24 phytochemical data will be a process that can be limited to 'simple/single foods', since only food
25 items such as 'flour', 'apple', 'sugar' rather than dishes such as 'apple crumble', 'apple pie' will
26 need assessment of their nutrient profile; thirdly, it enables modifications and updating of default
27 recipes published in the CoF to make them better suited for EPIC-Norfolk cohort data.

28 **Conclusion**

29 The response rate for 7dDD in EPIC-Norfolk has been extremely good. Underreporting may be an
30 issue, but not more so than found in national surveys; and underreporting is not likely to have been
31 caused by the length of the diary, since EI did not drop during the seven days. Despite this, the
32 association of underreporting with BMI will be important for interpretation of future endpoint

1 analysis. The large number of variables in the EPIC-Norfolk data has made these data highly
2 flexible to test new hypotheses in nutritional epidemiology or even use new approaches such as
3 hypotheses free nutrient wide association studies along the lines of gene wide association studies⁽⁴²⁾.
4 Groupings of food items and/or disaggregation can cause differences in absolute estimates, though
5 ranking of individuals will be less affected. The choice of aggregated or disaggregated variables
6 will influence estimates of food groups and comparison of results between studies.

7 **Acknowledgements**

8 We thank the dedicated team of data-enterers who entered over 6,500,000 food items, and all
9 nutritionists who had a major contribution to data checking and cleaning in the past sixteen years.
10 We especially wish to acknowledge the fundamental contributions of Sheila Rodwell (Bingham) in
11 her work on dietary biomarkers and her development of the 7dDD. The data on the quantitative
12 food groups was partly made possible by collaboration in the MRC Centre for Nutritional
13 Epidemiology in Cancer Prevention and Survival (CNC); we thank the nutritionists who provided
14 us with these data.

15 **Funding**

16 This work was supported by the Medical Research Council (MRC). The MRC had no role in the
17 design, analysis or writing of this article.

18 **Conflict of interest**

19 None.

20 **Contribution of each author to the manuscript**

21 MAHL prepared the manuscript, did the statistical analysis, checked diary data and assisted in the
22 re-writes of the different DINERMO programs. AM, NP, AMcT checked diary data, oversaw data-
23 entry and contributed with their experience to the many program revisions. DPS wrote the last
24 revision of the DINERMO program. RL, AB maintained earlier versions of DINERMO programs
25 as well as the wider range of cohort data available. AW developed the concept of DINER and the
26 early checking programs. KTK is principal investigator of the EPIC-Norfolk study. All co-authors
27 read the manuscript and provided their input.

28 **Additional/Online material**

29 More detailed information on the categorising of foods and food groups will be provided on our
30 website: www.epic-norfolk.org.uk or can be obtained from the corresponding author.

1 **References**

- 2 1. Bingham SA, Gill C, Welch A, et al. (1994) Comparison of dietary assessment methods in
3 nutritional epidemiology: weighed records v. 24 h recalls, food-frequency questionnaires and
4 estimated-diet records. *The Br J Nutr* **72**, 619–43.
- 5 2. Bingham SA, Gill C, Welch A, et al. (1997) Validation of dietary assessment methods in the
6 UK arm of EPIC using weighed records, and 24-hour urinary nitrogen and potassium and
7 serum vitamin C and carotenoids as biomarkers. *Int J Epidemiol* **26 Suppl 1**, S137–S151.
- 8 3. Bingham SA, Welch AA, McTaggart A, et al. (2001) Nutritional methods in the European
9 Prospective Investigation of Cancer in Norfolk. *Public Health Nutr* **4**, 847–58.
- 10 4. Thompson F & Subar A (2008) Dietary assessment methodology. In *Nutrition in the*
11 *prevention and treatment of disease*, 2nd ed., pp. 2–39 [Coulston AM, Boushey CJ, editors].
12 Burlington: Academic Press (AP).
- 13 5. Subar AF, Kipnis V, Troiano RP, et al. (2003) Using intake biomarkers to evaluate the extent
14 of dietary misreporting in a large sample of adults: the OPEN study. *Am J Epidemiol* **158**, 1–
15 13.
- 16 6. World Cancer Research Fund / American Institute for Cancer Research (2007) *Food,*
17 *Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective.*
18 Washington DC: AICR.
- 19 7. Fitt E, Mak TN, Stephen AM, et al. (2010) Disaggregating composite food codes in the UK
20 National Diet and Nutrition Survey food composition databank. *Eur J Clin Nutr* **64 Suppl 3**,
21 S32–6. Nature Publishing Group.
- 22 8. Cosgrove M, Flynn A & Kiely M (2005) Impact of disaggregation of composite foods on
23 estimates of intakes of meat and meat products in Irish adults. *Public Health Nutr* **8**, 327–37.
- 24 9. Prynne CJ, Wagemakers JJMF, Stephen AM, et al. (2009) Meat consumption after
25 disaggregation of meat dishes in a cohort of British adults in 1989 and 1999 in relation to diet
26 quality. *Eur J Clin Nutr* **63**, 660–6.
- 27 10. Subar AF, Krebs-Smith SM, Cook A, et al. (1998) Dietary sources of nutrients among US
28 adults, 1989 to 1991. *J Am Diet Assoc* **98**, 537–47.
- 29 11. Day N, Oakes S, Luben R, et al. (1999) EPIC-Norfolk: study design and characteristics of the
30 cohort. *Br J Cancer* **80 Suppl 1**, 95–103.
- 31 12. Scientific Advisory Committee on Nutrition (SACN) (2011) *Dietary reference values for*
32 *energy*. London: The Stationary Office (TSO).
- 33 13. Prynne CJ, Paul a. a., Mishra GD, et al. (2007) Changes in intake of key nutrients over 17
34 years during adult life of a British birth cohort. *British Journal of Nutrition* **94**, 368.
- 35 14. Welch AA, McTaggart A, Mulligan AA, et al. (2001) DINER (Data Into Nutrients for
36 Epidemiological Research) - a new data-entry program for nutritional analysis in the EPIC-
37 Norfolk cohort and the 7-day diary method. *Public Health Nutr* **4**, 1253–1265.

- 1 15. Holland B, Welch AA, Unwin D, et al. (1991) *McCance and Widdowson's The Composition*
2 *of Foods*. Cambridge: Royal Society of Chemistry (RSC).
- 3 16. Holland B, Unwin I & Buss D (1988) *Cereals and cereal products. The third supplement to*
4 *McCance & Widdowson's The Composition of Foods (4th Edition)*. Cambridge: RSC/MAFF.
- 5 17. Holland B, Unwin I & Buss D (1989) *Milk products and eggs. The fourth supplement to*
6 *McCance & Widdowson's The Composition of Foods (4th Edition)*. Cambridge: RSC/MAFF.
- 7 18. Holland B, Unwin I & Buss D (1991) *Vegetables, herbs and spices. The fifth supplement to*
8 *McCance & Widdowson's The Composition of Foods (4th Edition)*. Cambridge: RSC/MAFF.
- 9 19. Holland B, Unwin I & Buss D (1992) *Fruit and nuts. The first supplement to McCance &*
10 *Widdowson's The Composition of Foods (5th Edition)*. Cambridge: RSC/MAFF.
- 11 20. Holland B, Welch A & Buss D (1992) *Vegetable dishes. The second supplement to McCance*
12 *& Widdowson's The Composition of Foods (5th Edition)*. Cambridge: RSC/MAFF.
- 13 21. Holland B, Brown J & Buss D (1993) *Fish and fish products. The third supplement to*
14 *McCance & Widdowson's The Composition of Foods (5th Edition)*. Cambridge: RSC/MAFF.
- 15 22. Chan W, Brown J & Buss D (1994) *Miscellaneous foods. The fourth supplement to McCance*
16 *& Widdowson's The Composition of Foods (5th Edition)*. Cambridge: RSC/MAFF.
- 17 23. Chan W, Brown J, Lee S, et al. (1995) *Meat, poultry and game. The fifth supplement to*
18 *McCance & Widdowson's The Composition of Foods (5th Edition)*. Cambridge: RSC/MAFF.
- 19 24. Chan W, Brown J, Church S, et al. (1996) *Meat products and dishes. The sixth supplement to*
20 *McCance & Widdowson's The Composition of Foods (5th Edition)*. Cambridge: RSC/MAFF.
- 21 25. Ministry of Agriculture Fisheries and Food (1998) *Fatty acids. The seventh supplement to*
22 *McCance & Widdowson's The Composition of Foods (5th Edition)*. Cambridge: RSC/MAFF.
- 23 26. Mulligan A, Welch A, McTaggart A, et al. (2007) Intakes and sources of soya foods and
24 isoflavones in a UK population cohort study (EPIC-Norfolk). *Eur J Clin Nutr* **61**, 248–54.
- 25 27. Mulligan AA, Kuhnle GG, Lentjes MA, et al. (2012) Intakes and sources of isoflavones,
26 lignans, enterolignans, coumestrol and soya-containing foods in the Norfolk arm of the
27 European Prospective Investigation into Cancer and Nutrition (EPIC-Norfolk), from 7 d food
28 diaries, using a newly updated database. *Public Health Nutr*, 1–9.
- 29 28. Klingberg S, Andersson H, Mulligan A, et al. (2008) Food sources of plant sterols in the
30 EPIC Norfolk population. *Eur J Clin Nutr* **62**, 695–703.
- 31 29. Lentjes MAH, Welch AA, McTaggart A, et al. (2006) The importance of error checking after
32 data-entry of open-ended dietary methods: CHEDDAR. *International Conference on Dietary*
33 *Assessment Methods* **6**, 110.
34 http://orbit.dtu.dk/fedora/objects/orbit:96782/datastreams/file_3501279/content (accessed
35 November 2012).
- 36 30. Dahm CC, Keogh RH, Spencer EA, et al. (2010) Dietary fiber and colorectal cancer risk: a
37 nested case-control study using food diaries. *J Natl Cancer Inst* **102**, 614–26.

- 1 31. Henderson L, Gregory J, Irving K, et al. (2003) *The National Diet & Nutrition Survey :
2 adults aged 19 to 64 years. Energy, protein, carbohydrate, fat and alcohol intake.* vol. 2.
3 London: Her Majesty's Stationary Office (HMSO).
- 4 32. Finch S, Doyle W, Lowe C, et al. (1998) *National Diet and Nutrition Survey: people aged 65
5 years and over.* vol. 1. London: The Stationary Office (TSO).
- 6 33. Bland JM & Altman DG (1986) Statistical methods for assessing agreement between two
7 methods of clinical measurement. *Lancet* **1**, 307–10.
- 8 34. MacIntyre UE (2009) Measuring food intake. In *Introduction to Human Nutrition*, 2nd ed.,
9 pp. 238–275 [Gibney MJ, Lanham-New SA, Cassidy A, et al., editors]. Oxford: Wiley-
10 Blackwell.
- 11 35. Nelson M & Bingham SA (1997) Assessment of food consumption and nutrient intake. In
12 *Design concepts in nutritional epidemiology*, 2nd ed., pp. 123–169 [Margetts BM, Nelson M,
13 editors]. New York: University Press.
- 14 36. Lentjes MA, Bhaniani A, Mulligan AA, et al. (2011) Developing a database of vitamin and
15 mineral supplements (ViMiS) for the Norfolk arm of the European Prospective Investigation
16 into Cancer (EPIC-Norfolk). *Public Health Nutr* **14**, 459–471.
- 17 37. Lentjes MAH, Bhaniani A, Mulligan AA, et al. (2011) Contribution of cod liver oil
18 supplements to intake and associations with biomarkers of fatty acids in the European
19 Prospective Investigation into Cancer (EPIC-Norfolk) Study. *Proc Nutr Soc* **70**, E103.
- 20 38. Henderson L, Gregory J & Swan G (2002) *The National Diet & Nutrition Survey : adults
21 aged 19 to 64 years. Types and quantities of foods consumed.* vol. 1. London: Her Majesty's
22 Stationary Office (HMSO).
- 23 39. Welch AA, Luben R, Khaw KT, et al. (2005) The CAFE computer program for nutritional
24 analysis of the EPIC-Norfolk food frequency questionnaire and identification of extreme
25 nutrient values. *J Hum Nutr Diet* **18**, 99–116.
- 26 40. Purslow LR, Sandhu MS, Forouhi N, et al. (2008) Energy intake at breakfast and weight
27 change: prospective study of 6,764 middle-aged men and women. *Am J Epidemiol* **167**, 188–
28 92.
- 29 41. Cooper AJ, Sharp SJ, Lentjes MAH, et al. (2012) A prospective study of the association
30 between quantity and variety of fruit and vegetable intake and incident type 2 diabetes.
31 *Diabetes care* **35**, 1293–300.
- 32 42. Forman JP & Willett WC (2012) Nutrient-Wide Association Studies: Another Road to the
33 Same Destination. *Circulation*, 2447–2448.
- 34 43. Wareham NJ, Jakes RW, Rennie KL, et al. (2003) Validity and repeatability of a simple
35 index derived from the short physical activity questionnaire used in the European Prospective
36 Investigation into Cancer and Nutrition (EPIC) study. *Public Health Nutr* **6**, 407–413.

37

38 .

Table 1: Socio-demographic characteristics of the participants in the EPIC-Norfolk study (UK) who attended the health examination and completed at least 1 day of their 7dDD between 1993-1998.

		Men			Women		
		N	Mean/Proportion	SD	N	Mean/Proportion	SD
Age (year)		11,535	59.2	9.3	13,972	58.4	9.3
BMI (kg/m²)		11,513	26.5	3.3	13,944	26.2	4.4
Social class (%)	Professional	868	7.7		875	6.4	
	Managerial	4,330	38.2		4,778	35.1	
	Skilled non-manual	1,419	12.5		2,706	19.9	
	Skilled manual	2,864	25.3		2,878	21.1	
	Partially skilled	1,515	13.4		1,832	13.5	
	Non-skilled	335	3.0		543	4.0	
Education (%)	No qualification	3,512	30.5		5,897	42.2	
	O-level	996	8.6		1,604	11.5	
	A-level	5,246	45.5		4,957	35.5	
	Degree/equivalent	1,772	15.4		1,505	10.8	
Smoking (%)	Current	1,395	12.2		1,565	11.3	
	Former	6,243	54.5		4,459	32.2	
	Never	3,817	33.3		7,809	56.5	
Physical activity (%)*	Inactive	3,554	30.8		4,249	30.4	
	Moderately inactive	2,838	24.6		4,479	32.1	
	Moderately active	2,651	23.0		3,101	22.2	
	Active	2,482	21.5		2,134	15.3	

* Physical activity was measured using a questionnaire and included occupational and leisure time activity; there is no algorithm to convert the answers in this questionnaire to a PAL-level⁽⁴³⁾.

Table 2: Examples of several foods and the categories to which they belong, depending on the food group.

Food item	Qualitative food groups			Quantitative / disaggregated food groups								
	Crude group	Detailed group	F&V variety*	Canned products	Dairy	fruit [†]	vegetables	red meat	white meat	processed meat	white fish	fatty fish
Chicken risotto	cereal, other	rice dish	onion-dish	non-canned	non-dairy	0	.15	0	.18	0	0	0
Beef fillet steak, grilled	meat	red meat	no F or V	non-canned	non-dairy	0	0	1	0	0	0	0
Veal in tomato sauce	meat	red meat dish	tomato-cooked	non-canned	non-dairy	0	.55	.40	0	0	0	0
Quiche Lorraine	cereal, other	pie/flan/quiche, red meat	no F or V	non-canned	low dairy-mixed -full fat	0	0	0	0	.14	0	0
Muesli	cereals	mueslis	mixed fruit-dried	non-canned	non-dairy	.20	0	0	0	0	0	0
Apple crumble	cereal, other	cereal pudding	apple-dish	non-canned	non-dairy	.62	0	0	0	0	0	0
Custard, made up with skimmed milk	cereal, other	milk dessert	no F or V	non-canned	high dairy-milk-skimmed	0	0	0	0	0	0	0
Tuna and	miscellaneous	fish dish	sweetcorn-	non-	non-dairy	0	.25	0	0	0	0	.60

* F&V, Fruit & Vegetable

[†]Another version of this quantitative fruit group exists which acknowledges the contribution of fruit juices to fruit (*i.e.* a proportion of '1').

Food item	Qualitative food groups				Quantitative / disaggregated food groups							
	Crude group	Detailed group	F&V variety*	Canned products	Dairy	fruit [†]	vegetables	red meat	white meat	processed meat	white fish	fatty fish
sweetcorn sandwich filling			cooked	canned								
Iceberg lettuce	vegetables	leafy salad	lettuce-raw	non-canned	non-dairy	0	1	0	0	0	0	0
Vegetable soup, canned	soups	soups	mixed veg-soup	canned	non-dairy	0	.29	0	0	0	0	0
Orange juice, commercial	drinks	orange juice	orange-juice	mixed	non-dairy	0	0	0	0	0	0	0
Lemonade fizzy	drinks	soft drink	no F or V	mixed	non-dairy	0	0	0	0	0	0	0

Table 2: Classification of food items into crude groupings and the criteria applied for disaggregating food items in EPIC-Norfolk.

	Crude / qualitative	Disaggregated / quantitative
Fruit	Food items that are completely fruit, eaten raw or cooked. As well as food items with some minimal additions such as sugar when stewed or canned, but excluding fruit juices.	Food items that are 100% fruit, eaten raw or cooked. As well as fruit used in dishes such as stewed fruit, fruit crumble and fruit fool, for which a percentage of fruit content is assigned (fruit juice, fruit puree in yogurt and jams are set to 0% fruit).
Vegetables	Food items that are completely vegetable (excl. potatoes), eaten raw or cooked. As well as the full weight of dishes such as cauliflower cheese, vegetable burgers and vegetable curries, but not vegetables used in dishes such as soups, sauces and meat/fish stews. 'Pure' pulses and lentils are excluded.	Food items that are 100% vegetable, eaten raw or cooked. As well as vegetables used in dishes, for which a percentage of vegetable content was assigned (potatoes, pulses, lentils and quorn are not considered vegetables).
Meat	Food items that are completely meat (red, white or processed), as well as the full weight of meat dishes such as stews/casseroles, bolognese sauce and chilli con carne, but not rice or pasta dishes, meat pies, soups and offal.	Food items that are 100% meat, as well as meat in dishes, for which a percentage of red, white and processed meat content is assigned. Offal is excluded.
Fish	Food items that are completely fish (white or fatty), crustacea or molluscs, as well as fish dishes such as fish cakes, fish in sauce/bake and battered fish, but not rice or pasta dishes or soups.	Food items that are 100% fish, as well as fish in dishes such as coated fish, fish pies, fish bakes, rice/pasta, omelettes, for which a percentage of fish content was assigned. Crustacea and molluscs are excluded.

Table 3: Energy intake (EI) by subcategories of BMI. Total Energy Expenditure (TEE) was calculated using the Henry formula and different levels of assumed activity (PAL) in the EPIC-Norfolk population. EI and TEE were compared using Bland-Altman plots. The limits of agreement and the proportion of underreporters for each level of assumed PAL are shown, stratified by BMI. Analysis only included participants who completed seven days of their 7dDD.

N	EI* (MJ/d)	TEE-EI (PAL set to 1.40 [†])			TEE-EI (PAL set to 1.49 [†])			TEE-EI (PAL set to 1.63 [†])										
		Mean	SD	Lower‡	Upper	Underreporters§ (%)	Lower‡	Upper	Underreporters§ (%)	Lower‡	Upper	Underreporters§ (%)						
				0-1	1-2	≥2			0-1	1-2	≥2			0-1	1-2	≥2		
				MJ/d	MJ/d	MJ/d			MJ/d	MJ/d	MJ/d			MJ/d	MJ/d	MJ/d		
Men																		
BMI (kg/m ²)																		
≤25	3,491	9.75	2.07	-4.57	3.32	18.9	11.7	8.0	-4.00	3.92	20.9	15.9	14.1	-3.11	4.85	20.4	20.3	28.5
>25	6,843	9.35	2.13	-3.54	4.89	18.8	18.2	26.4	-2.92	5.56	17.0	18.9	37.6	-1.97	6.62	12.8	16.9	56.4
Women																		
BMI (kg/m ²)																		
≤25	5,739	7.41	1.57	-3.09	2.95	24.6	16.3	8.2	-2.63	3.43	26.4	20.6	14.6	-1.91	4.19	23.2	25.8	29.1
>25	7,016	7.02	1.60	-2.32	4.23	24.0	22.1	26.2	-1.84	4.77	20.4	23.9	37.4	-1.09	5.61	13.4	22.1	55.9

* EI, Energy intake; TEE, Total Energy Expenditure; PAL, Physical Activity Level; MJ, Megajoule; SD, standard deviation; BMI, Body Mass Index.

† The chosen PAL values represent the 10th, 25th and 50th centile of the distribution used in the SACN report⁽¹²⁾.

‡ The difference in energy was calculated as: estimated TEE - EI. Approximately 95% of the observations are between the limits of agreement (meandiff ± (1.96 * SDdiff))⁽³³⁾. The upper limit of agreement reflects underreporting of EI and the lower limit of agreement reflects overreporting of EI.

§ Underreporters were defined as participants with a difference in TEE - EI that resulted in values greater than 0; the degree of underreporting was expressed in three categories (0-1 MJ/d, 1-2 MJ/d, ≥2 MJ/d).

Table 4: Measures of central tendency and spread in energy, macronutrient and micronutrient intake in the EPIC-Norfolk cohort, stratified by sex (N=25,507).

	Unit	Men (n=11,535)					Women (n=13,972)				
		Median	Mean	SD*	2.5 th	97.5 th	Median	Mean	SD	2.5 th	97.5 th
Energy	MJ/d	9.36	9.45	2.22	5.36	14.14	7.11	7.15	1.66	4.06	10.61
Protein	g/d	80.9	82.2	18.6	49.4	121.6	64.8	65.4	14.2	39.2	95.0
Fat	g/d	83.9	86.1	26.9	40.5	145.0	63.4	64.6	20.9	27.9	109.3
Saturated fat	g/d	31.2	32.6	11.9	13.4	59.5	23.6	24.6	9.3	9.3	45.4
Carbohydrate	g/d	269.7	273.7	73.9	142.4	433.9	210.9	213.2	54.4	112.8	329.6
Sugars	g/d	115.7	119.5	45.9	42.8	221.4	95.2	97.9	34.9	37.1	174.4
Fibre (NSP)	g/d	15.2	16.1	6.0	7.0	30.4	13.5	14.2	5.0	6.2	25.9
	g/MJ	1.6	1.7	0.6	0.9	3.2	1.9	2.0	0.7	1.0	3.7
Alcohol	g/d	9.0	16.6	21.5	0	74.8	2.7	8.0	12.2	0	42.1
Consumers only [†]	g/d	15.6	22.1	22.3	1.2	80.5	8.6	12.9	13.3	0.8	49.4

* SD, standard deviation; 2.5th/97.5th percentile; NSP, Non-starch polysaccharides; TE, Tocopherol equivalents.

† The distribution among consumers of alcoholic beverages only (alcohol (as a nutrient) >0.5 g/d): men n=8,665; women n=8,660.

		Men (n=11,535)				Women (n=13,972)					
Fat-soluble vitamins											
Retinol	mcg/d	391	780	1,571	99	4,816	300	610	1,239	78	3,905
	mcg/MJ	42	85	206	13	526	42	86	177	13	562
Vitamin D	mcg/d	3.09	3.73	2.74	0.74	10.27	2.45	2.93	2.08	0.57	7.81
	mcg/MJ	0.33	0.40	0.29	0.09	1.09	0.35	0.42	0.30	0.10	1.14
Vitamin E	mg TE/d	10.61	11.62	5.24	4.42	24.83	8.67	9.27	3.78	3.79	18.33
	mg/MJ	1.15	1.23	0.46	0.57	2.37	1.23	1.30	0.42	0.67	2.33
Water-soluble vitamins											
Thiamin	mg/d	1.63	1.72	0.90	0.85	2.83	1.34	1.42	0.74	0.72	2.27
	mg/MJ	0.18	0.18	0.09	0.11	0.29	0.19	0.20	0.10	0.12	0.32
Riboflavin	mg/d	1.88	1.97	0.69	0.90	3.51	1.58	1.65	0.58	0.74	2.94
	mg/MJ	0.20	0.21	0.07	0.11	0.37	0.22	0.23	0.08	0.12	0.41
Ascorbic acid	mg/d	73	85	52	22	212	79	89	50	24	210
	mg/MJ	8	9	6	3	24	11	13	8	4	32
Minerals											
Calcium	mg/d	890	920	298	419	1,575	745	767	249	350	1,323
	mg/MJ	95	98	24	56	152	105	108	29	62	173
Iron	mg/d	12.8	13.4	4.3	6.9	23.0	10.4	10.9	3.4	5.6	18.8
	mg/MJ	1.4	1.4	0.4	0.9	2.4	1.5	1.5	0.4	1.0	2.6

Table 5: Consumption of fruit, vegetables, meat and fish in EPIC-Norfolk (N=25,507). Weights are given for two types of food groupings, crude (qualitative groups) and disaggregated foods.

Food group	Type*	Men (n=11,535)						Women (n=13,972)					
		Median	Mean	SD	2.5 th	97.5 th	r _s [†]	Median	Mean	SD	2.5 th	97.5 th	r _s [†]
Fruit (g/d)	C	117	144	132	0	463		150	173	133	0	485	
	D	133	158	134	0	478	.98	161	183	133	0	494	.99
Vegetables (g/d)	C	118	131	81	0	320		122	135	79	17	321	
	D	140	151	79	28	333	.90	140	150	76	36	328	.92
Meat (g/d)	C	117	126	77	0	297		88	95	61	0	229	
Red (g/d)	D	36	41	34	0	122		25	29	26	0	89	
White (g/d)	D	21	27	32	0	99		18	23	25	0	83	
Processed (g/d)	D	24	28	24	0	87		15	18	17	0	58	
Sum of red, white, processed	D	92	96	49	0	203	.85	68	70	38	0	151	.86
Fish (g/d)	C	33	42	41	0	141		30	36	32	0	116	
Fatty (g/d)	D	0	13	23	0	74		3	12	18	0	59	
White (g/d)	D	14	17	21	0	66		11	14	17	0	54	
Sum of fatty, white	D	23	30	31	0	109	.90	21	26	25	0	86	.89

* Type refers to the type of food group, C=crude, D=disaggregated; SD, standard deviation; r_s, spearman correlation; 2.5th/97.5th percentile.

† Spearman correlation between disaggregated variable and crude variable. All correlations were significant at $P < 0.01$.

Figure 1: Boxplot of averaged and seven individual days of energy intake (kJ/d), stratified by sex (N=25,507). The results of the averaged data are included since these data have been used for most publications to date.

Figure 2: Distribution of macronutrients as a percentage of total energy intake, stratified by sex and age (N=25,507). Using Mann-Whitney statistic, all differences between age-groups (within the same sex), were statistically significant apart from energy intake provided by unsaturated fatty acid intake in women.

Figure 3: Comparison of the crude and disaggregated method for measuring fruit, vegetable, meat and fish consumption in EPIC-Norfolk using Bland-Altman plots. Data only included those participants who completed all seven days of their 7dDD (N=23,149).