

1       **Adults with autism spectrum condition (ASC) have atypical**  
2       **perception of ambiguous figures when bottom-up and top-down**  
3               **interactions are incongruous**

4  
5       **Monika Intaitė<sup>2\*</sup>, Alexandra L. Georgescu<sup>3,1</sup>, Valdas Noreika<sup>4</sup>, Marie A. O. von Saldern<sup>1,6</sup>, Kai**

6               **Vogeley<sup>1,7</sup> & Christine M. Falter-Wagner<sup>5,6,8</sup>**

7  
8       1. Department of Psychiatry and Psychotherapy, University Hospital of Cologne, Cologne, Germany

9       2. Visual Neuroscience Laboratory - IBILI, Faculty of Medicine, University of Coimbra, Coimbra, Portugal

10       3. Institute of Cognitive Neuroscience, University College London, London, United Kingdom

11       4. Department of Psychology, University of Cambridge, Cambridge, United Kingdom

12       5. Institute of Medical Psychology, Medical Faculty, LMU Munich, Germany

13       6. Department of Psychology, University of Cologne, Cologne, Germany

14       7. Institute of Neuroscience and Medicine – Cognitive Neuroscience (INM-3), Research Center Juelich,  
15       Juelich, Germany

16       8. Department of Psychiatry, Medical Faculty, LMU Munich, Germany

17  
18       \*Corresponding author:

19       Monika Intaitė

20       Visual Neuroscience Laboratory - IBILI,

21       Faculty of Medicine, University of Coimbra

22       Azinhaga De Santa Comba, Celas

23       3000-548, Coimbra, Portugal

24       Phone: +351239480200

25       Fax: +351239480217

26       E-mail: [mintaite@fmed.uc.pt](mailto:mintaite@fmed.uc.pt)

## Abstract

We examined the perception of an ambiguous squares stimulus evoking bistable perception in a sample of 31 individuals with autistic spectrum condition (ASC) and 22 matched typical adults (TA). The perception of the ambiguous figure was manipulated by adaptation to unambiguous figures and/or by placing the ambiguous figure into a context of unambiguous figures. This resulted in four conditions testing the independent and combined (congruent and incongruent) manipulations of adaptation (bottom-up) and spatial context (top-down) effects. The strength of perception, as measured by perception of the first reported orientation of the ambiguous stimulus was affected comparably between groups. Nevertheless, the strength of perception, as measured by perceptual durations was affected differently between groups: The perceptual effect was strongest for the ASC group when combined bottom-up and top-down conditions were congruent. In contrast, the strength of the perceptual effect in response to the same condition in the TA group was comparable to the adaptation, but stronger than both the context and the incongruent combined bottom-up and top-down conditions. Furthermore, the context condition was stronger than the incongruent combined bottom-up and top-down condition for the TA group. Thus, our findings support the view of stimulus-specific top-down modulation in ASC.

**Keywords:** Autistic Spectrum Condition; Ambiguous figures; Adaptation; Context

1 **Adults with autism spectrum condition (ASC) have atypical perception of ambiguous**  
2 **figures when bottom-up and top-down interactions are incongruous**

3 Autistic spectrum condition (ASC) is a neurodevelopmental condition characterized  
4 by disturbances in social interaction and communication, as well as stereotyped, repetitive  
5 behaviors and interests (WHO, 1992). The condition additionally exhibits abnormalities in  
6 visual perception and attention (Happé & Frith, 2006; Hill, 2004).

7 Anecdotal evidence suggests that ASC individuals are very accurate and fast at  
8 perceiving visual details of a display (bottom-up perception). A growing body of research  
9 confirms that ASC participants are better in tasks, which require predominantly “local” or  
10 detail-focused processing, e.g., embedded figures (Shah & Frith, 1983), block design (Shah  
11 & Frith, 1993), conjunctive visual search (Plaisted, O’Riordan, & Baron-Cohen, 1998), for a  
12 recent meta-analysis on the former two tasks refer to Muth, Hönekopp, and Falter (2014).  
13 Although evidence for an enhanced local processing is consistent, reports on deficits in global  
14 processing are inconclusive (Dakin & Frith, 2005).

15 On a neural level, a decreased top-down neural control might manifest itself in a local  
16 processing style at the expense of processing the global ‘gist’ (Frith, 2003). Frith (2003)  
17 suggests lack of pruning of neuronal feedback projections during brain development in ASC  
18 individuals, which results in malfunctioning of neural feedback control systems. Functional  
19 neuroimaging studies have supported the view of atypical top-down modulation of early  
20 sensory processing in ASC individuals: task-dependent early sensory processing areas  
21 activate or even hyper-activate, while later processing areas hypo-activate in response to  
22 various perceptual and attentional manipulations (Freitag et al., 2008). Loth, Gómez, &  
23 Happé (2010) and Cook, Barbalat, & Blakemore (2012) show that top-down influence of

1 prior knowledge and top-down dominance on task performance is reduced in ASC. However,  
2 Bird, Catmur, Silani, Frith, & Frith (2006) and Greenaway & Plaisted (2005) imply that top-  
3 down influence on task performance can be stimulus-specific. Mitchell, Mottron, Soulières,  
4 and Ropar (2010) discuss whether top-down processing in ASC might be attenuated in the  
5 sense of a general deficit, or top-down processing might be merely stimulus or task-  
6 dependent.

7         We tested this particular question for the first time and explored spontaneous  
8 influences of bottom-up and top-down information on perception and their interaction in  
9 ASC using an ambiguous figures paradigm. Ambiguous figures are such stimuli that make  
10 our perception constantly oscillate between two (or more) alternative interpretations. When  
11 participants merely observe the ambiguous figures, ASC participants and typical adults  
12 experience similar perceptual changes (Ropar, Mitchell, & Ackroyd, 2003; Sobel, Capps, &  
13 Gopnik, 2005). Our ambiguous figures paradigm, however, allows the implementation of  
14 bottom-up and top-down processing on their own, as well as under combined congruent and  
15 incongruent conditions (Intaité et al., 2013). This is achieved by including adaptation (bottom-  
16 up) and context (top-down) effects into the task. An adaptation effect is obtained in studies  
17 where participants are presented with an unambiguous stimulus before the actual ambiguous  
18 stimulus is shown. The unambiguous stimulus represents one of the possible interpretations  
19 of the ambiguous figure, hence biasing the subsequent perception of the ambiguous figure,  
20 i.e. participants tend to first recognize the alternative interpretation of the subsequently  
21 presented ambiguous figure (Long & Moran, 2007). In this study, all participants were  
22 adapted to an unambiguous squares stimulus in either ‘upwards’ or ‘downwards’ orientation  
23 that was followed by the ambiguous squares stimulus. Contrary to the adaptation effect, the

1 context effect is achieved by presenting an ambiguous test figure within the spatial context  
2 of unambiguous figures representing one of the possible perceptual interpretations. In this  
3 case, the ambiguous test figure is more likely to be perceived in the same orientation as the  
4 context figures (Intaitè et al., 2013; Sundareswara & Schrater, 2008). In the current study, the  
5 ambiguous squares stimulus was presented in the spatial context of unambiguous squares  
6 stimuli. Adaptation and context conditions were presented either in separate trials or within  
7 the same trials. In the latter case, the orientation of the context stimuli either matched the  
8 formerly presented adapting stimulus or not. This resulted in four conditions: an adaptation  
9 only condition, a context only condition, an adaptation different from context condition  
10 (combined congruent), and an adaptation identical to context condition (combined  
11 incongruent) (see Fig. 1). Previously Happé (1996) and Happé & Frith (2006) suggested that  
12 perception in ASC individuals might be less influenced by context in a top-down manner.  
13 Yet, Greenaway & Plaisted (2005) and Mitchell et al. (2010) indicate that top-down  
14 attentional modulation might be stimulus-specific or task-dependent rather than generally  
15 attenuated. Hence, we sought to investigate these two hypotheses of either a general deficit  
16 of top-down modulation (i.e., the ASC group would show weaker responses to context effect  
17 alongside to weaker responses to both combined bottom-up and top-down manipulations) or  
18 a stimulus-specific top-down modulation by testing ambiguous figure perception under these  
19 four conditions.

20

## 1 **Methods**

### 2 **Participants**

3 Thirty-six individuals with Autism Spectrum Condition (ASC; twenty-three males)  
4 and twenty-eight typical adults (TA; fourteen males) took part in the experiment. All ASC  
5 participants met international criteria for autism (F84.0) or Asperger's Syndrome (F84.5)  
6 confirmed by clinical interviews according to ICD-10 (WHO, 1992), as judged by two  
7 clinicians specializing in the assessment and diagnosis of the condition. Participants of this  
8 study were recruited from the database at the Outpatient Clinic for Adults with Autism  
9 Spectrum Disorder at the University Hospital Cologne. Eleven participants (five from the  
10 ASC group and six from the TA group) were excluded from analyses because of an inability  
11 to report reversals in one or more experimental conditions in spite of reporting to see both  
12 interpretations of the ambiguous squares stimulus during the practice period. Thus, the data  
13 of thirty-one ASC (twenty males; age = 43:6 years:months,  $SD = 8.6$ ) and twenty-two TA  
14 participants (twelve males; age = 39:3 years:months,  $SD = 8.7$ ) were entered into the analyses.  
15 Each participant had reported normal or corrected-to-normal vision. Written informed  
16 consent was obtained from all participants and the ethics board of the University Hospital  
17 Cologne formally approved the study.

### 18 **Psychometric testing**

19 Participants completed this study as part of a larger battery of three perceptual  
20 experiments. They also completed a neuropsychological test battery including the Beck  
21 Depression Inventory (BDI, Hautzinger, Keller, & Kühner, 2006), Wechsler Intelligence  
22 Scale (WIE, Aster, Neubauer, & Horn, 2006), the Autism-Spectrum Quotient (AQ, Baron-  
23 Cohen, Wheelwright, Skinner, Martin & Clubley, 2001), the Empathy Quotient (EQ, Baron-

1 Cohen, Wheelwright, 2004), the Systemizing Quotient (SQ, Baron-Cohen, Richler, Bisarya,  
2 Gurunathan, & Wheelwright, 2003), Toronto Alexithymia Scale (TAS-20, Bagby, Parker, &  
3 Taylor, 1994), the Intuitive Physics Test (IPT, Baron-Cohen, S, Wheelwright, Scahill,  
4 Lawson, & Spong, 2001) and Theory of Mind - Reading the Mind in the Eyes Test (Baron-  
5 Cohen, S, Wheelwright, Hill, Raste, & Plumb, 2001). ASC and TA groups were matched  
6 with respect to age, verbal, and performance IQ (Table 1; *largest t* = 1.82). TA participants  
7 were screened for not having a prior history of neurological or psychiatric conditions. Eleven  
8 ASC participants were taking antidepressants at the time of the study. Taking into account  
9 the altered excitation/inhibition balance in the autistic brain (Nelson & Valakh, 2015), we  
10 have rerun our statistics excluding these ASC participants and the results were comparable  
11 to the results obtained with a complete sample. Results excluding ASC participants taking  
12 antidepressants are available online (<https://osf.io/nhrxk/>).

13 <<<Table 1 about here>>>

## 14 **Design and Procedure**

15  
16 An ambiguous squares figure was used as the main experimental stimulus. Each trial  
17 started with a 120 s adaptation period followed by a blank screen presented for 1 s.  
18 Afterwards, participants were presented with an ambiguous squares stimulus (in the four  
19 different conditions described below) for 30 s and had to report the perceived changes of the  
20 square's orientation by pressing one of the two keys (upwards or downwards) on the response  
21 keyboard. An inter-trial interval of 120 s was presented after each trial. A central fixation  
22 point was always present in the middle of the screen/figure except during the intertrial  
23 intervals. The participants were asked to keep their eyes focused on this point throughout the  
24 experiment. The four conditions had the following characteristics:

1           1) The adaptation condition (hereafter AC): During the adaptation period, an  
2 unambiguous ‘upwards’ or ‘downwards’ oriented squares stimulus was presented. During  
3 the test period an ambiguous squares stimulus (i.e., no context) was presented (Fig. 1A).

4           2) The context condition (hereafter CC): During the adaptation period, only the  
5 fixation point was used as a stimulus (i.e., no adaptation). During the test period an  
6 ambiguous squares stimulus was presented in the context of surrounding unambiguous  
7 squares stimuli in either ‘upwards’ or ‘downwards’ orientation (Fig. 1B).

8           3) The adaptation different from context (hereafter ADC) condition: During the  
9 adaptation period, an unambiguous squares stimulus oriented either ‘upwards’ or  
10 ‘downwards’ was presented. During the test period an ambiguous squares stimulus was  
11 presented in the context of surrounding unambiguous squares stimuli in either a ‘downwards’  
12 or ‘upwards’ orientation, respectively (Fig. 1C).

13           4) The adaptation identical to context (hereafter AIC) condition: During the  
14 adaptation period, the unambiguous squares stimulus oriented either ‘upwards’ or  
15 ‘downwards’. During the test period an ambiguous squares stimulus was presented in the  
16 context of surrounding unambiguous squares stimuli in either ‘upwards’ or ‘downwards’  
17 orientation, respectively (Fig. 1D).

18           All stimuli were drawn in black ( $\sim 12.78 \text{ cd/m}^2$ ), presented on a white background ( $\sim$   
19  $14.98 \text{ cd/m}^2$ ), and viewed binocularly. A single figure subtended a visual angle of  $1.51^\circ \times$   
20  $1.51^\circ$ , the fixation point was  $0.02^\circ$  large, and line thickness in degrees of visual angle was  
21 around  $0.07^\circ$ . The entire display consisting of ambiguous squares in the context of  
22 unambiguous squares subtended  $5.82^\circ \times 5.82^\circ$  of visual angle. Stimuli were presented on a  
23 24 inch computer screen (frame rate: 59 Hz, mean luminance of the monitor:  $\sim 13.88 \text{ cd/m}^2$ ,

1 Michelson contrast: 0.15, LED monitor) at a viewing distance of ~ 60 cm. The position of  
2 the fixation point on each version of unambiguous squares was adjusted to match the center  
3 of the screen and the center of the subsequently presented ambiguous squares stimulus.

4 Each participant took part in a 90 min individual testing session. Ambient lighting  
5 was kept constant throughout each testing session. Before testing commenced, the ambiguous  
6 squares stimulus was shown to each participant, and they were instructed to watch it until  
7 reversals were perceived. In the beginning of the session, each participant performed two  
8 practice trials (the CC and the ADC conditions) in order to get acquainted with the task  
9 requirements.

10 <<<Figure 1 about here>>>  
11

12 Each of the four conditions consisted of four trials, resulting in a total of 16  
13 experimental trials. For each of the three experimental conditions with an adaptation period  
14 (the AC, the ADC and the AIC), the four trials consisted in the ambiguous squares stimulus  
15 being presented twice after the ‘downwards’ and twice after the ‘upwards’ unambiguous  
16 squares stimulus. The 16 trials were organized in two experimental blocks, separated by a 5  
17 min break. Each block contained eight randomly presented trials, with two trials from every  
18 experimental condition and one of each of the two unambiguous orientations: ‘upwards’ and  
19 ‘downwards’. The participants were asked to let the perceptual reversals occur naturally and  
20 were instructed not to intentionally manipulate their percepts. No feedback on performance  
21 was given.

## 22 **Statistical analyses**

23 There were two dependent measures used in the analyses: (1) the first responses  
24 regarding the percept of the orientation of the ambiguous squares stimulus (hereafter

1 Orientation First Percept) and (2) the average durations during the 30 sec test period,  
2 calculated separately for the ‘upwards’ and ‘downwards’ percepts (hereafter Perceptual  
3 Durations). The Orientation First Percept is a direct measure of the influence of the  
4 experimental manipulations on perception and the Perceptual Durations are metrics of the  
5 fluctuations of the perception. Concerning the analysis of Orientation First Percept, the initial  
6 interpretations were coded in terms of whether the ambiguous squares stimulus was  
7 perceived in the predicted (score = 1) or unpredicted (score = 0) orientation with respect to  
8 the expected effects from the experimental stimulus conditions (for further information about  
9 calculating the predicted scores see Intaité et al.), and an average score was used for statistical  
10 analyses (for further information on statistical analyses see Intaité et al., 2013).

11 A mixed ANOVA with one between-participant factor of GROUP (ASC and TA) and  
12 one within-participant factor of CONDITION (AC, CC, ADC, AIC) were conducted for the  
13 dependent variable Orientation First Percept. One-way ANOVAs (Bonferroni-Holm  
14 corrected, hereafter B-H) (Holm, 1979) were used as post-hoc tests to compare different  
15 conditions in case of a significant effect of CONDITION.

16 For the statistical analyses of the Perceptual Durations, successive presses of the same  
17 key during ambiguous squares stimulus observation (e.g. several subsequent indications of  
18 ‘upwards’ => ‘upwards’ or ‘downwards’ => ‘downwards’ percepts) were treated as errors  
19 and removed from the analyses. This happened on average ~1.97 times (SD = 3.04) for the  
20 ASC and ~ 1.41 times (SD = 1.12) for TA participants. Due to the limit in the ambiguous  
21 squares stimulus presentation duration (i.e., 30 s), a number of percepts were truncated as a  
22 result of the end of stimulus display and thus they were not included in the analyses.  
23 Kolmogorov-Smirnov tests revealed that Perceptual Durations values did not meet the

1 condition of normality. The distribution of the raw scores were leptokurtic and positively  
2 skewed, thus lognormal transformations were applied to these data (Howell, 2009).

3 A mixed ANOVA with one between-participant factor of GROUP (ASC and TA) and  
4 three within-participant factors of CONDITION (AC, CC, ADC, AIC), ADAPTING (or  
5 CONTEXT) STIMULUS (upwards and downwards) and PERCEPTUAL RESPONSE  
6 (upwards and downwards) were conducted on the data of Perceptual Durations (LOG  
7 transformed). After obtaining a significant effect of CONDITION and significant  
8 interactions with GROUP, and in order to correct for individual differences, we calculated  
9 the difference scores for each Perceptual Duration by subtracting the unpredicted Perceptual  
10 Durations from the predicted Perceptual Durations. After this, the predicted difference scores  
11 obtained for ‘upwards’ and ‘downwards’ perceptual orientations were averaged together  
12 creating ‘effect of manipulation’ variables for each condition. Subsequent one-way  
13 ANOVAs (B-H) were used as post-hoc tests to compare different ‘effect of manipulation’  
14 variables for each GROUP separately. In all cases of significant violations of sphericity,  
15 Huynh-Feldt corrections were applied.

#### 16 **Control experiment: Eye-movement recording and analysis**

17 In order to determine whether the participants were able to maintain central fixation  
18 during the experimental task we have performed a short control experiment. The stimuli and  
19 procedure were identical to the main experiment with the following exceptions: each of the  
20 conditions was presented only twice in order to reduce the total time of the experiment from  
21 90 min to 45 min.

22 During this control experiment, eye-tracking data (sample frequency 250 Hz) were  
23 recorded with an Eyelink 1000 (SR Research, Ottawa, Ontario, Canada). The device was

1 individually calibrated with a 9-point calibration routine at the beginning of the experiment.  
2 Sixteen ASC (eleven male, mean age = 45.44 years, SD = 8.41) and thirteen TA participants  
3 (five male, mean age = 41.08 years, SD = 7.29) took part in the control experiment. ASC and  
4 TA groups were matched on age, verbal, and performance IQ (*largest t* = 1.47). Eleven ASC  
5 participants (seven male) and seven TA participants (four male) took part in both the main  
6 and the control experiments, the latter of which was performed on another day and therefore  
7 would have not influenced behavioral performance during the main experiment. The data  
8 were prescreened for co-morbidity and medication in the same manner as the data of the main  
9 experiment.

10         Fixations were calculated based on the recorded gaze behavior. A fixation duration  
11 threshold of 150 ms was used. Fixations that had the same position and were separated by a  
12 blink were concatenated. To calculate dwell time (i.e., the percent of time spent fixating on  
13 the interest area), consecutive fixations were concatenated. For pupil data analysis, the pupil  
14 area values at 1 sec before each perceptual reversal were averaged together. Two rectangular  
15 areas of interest (AOI), one comprising the size of the ambiguous squares stimulus ( $1.51^\circ \times$   
16  $1.51^\circ$ ) (hereafter AOI-1) and one comprising the size of the context stimuli ( $5.82^\circ \times 5.82^\circ$ )  
17 (hereafter AOI-2) were defined. As the responses to the provided test stimuli depended on  
18 the participants' performance during the adaptation period, for the AOI-1 we analyzed the  
19 fixations' data both for the adaptation period (120 s before presenting the ambiguous squares  
20 stimulus) and for the test period (30 s of the ambiguous squares stimulus presentation time).  
21 For the AOI-2 analyses, only fixations data obtained during the test period were analyzed  
22 (Table 2). Kruskal-Wallis tests were performed to compare study groups regarding the  
23 average count of fixations, the dwell times both for AOI-1 and for AOI-2, and pupil area

1 values at 1 sec before the presses indicating perceptual reversals between all conditions and  
2 groups.

3 <<< Table 2 about here >>>

4

## Results

### Eye tracking

Results revealed no statistically significant group differences between the average count of fixations (largest  $\chi^2 = .52$ ), the dwell times neither for AOI-1 (largest  $\chi^2 = 1.61$ ) nor for AOI-2 (largest  $\chi^2 = .52$ ), and pupil area values at 1 sec before the presses indicating perceptual reversals (largest  $\chi^2 = 2.02$ ) across all conditions and across groups (see Table 2).

### Orientation First Percept

We performed a mixed ANOVA with one between-subjects factor of GROUP (ASC, TA), one within-subjects factor of CONDITION (AC, CC, AIC, ADC) and Orientation First Percept as dependent variable. The first reported percepts of the ambiguous squares were differentially influenced by experimental manipulations as indicated by a significant effect of CONDITION ( $F(3,153) = 15.71, p < .001, \eta p^2 = .24$ ). Effects of these experimental manipulations were comparable across GROUPS (GROUP effect, CONDITION  $\times$  GROUP interaction: largest  $F < 1$ ). First, we wanted to check whether adaptation or context had a stronger influence on the first ambiguous squares figure percept. Thus, we compared the adaptation and context conditions independently and this revealed that irrespective of GROUP ( $F < 1$ ) the manipulation of context was stronger than that of adaptation ( $F(1, 51) = 25.15, p < .001, \eta p^2 = .33$ ) (Fig. 2A). Further, we solely compared those conditions that had the same predicted orientations of responses (Intaitè et al., 2013): AC with the ADC (adaptation-matching first interpretation of ambiguous squares stimulus) and the CC with the AIC (context-matching first interpretation of the ambiguous squares stimulus). Subsequent comparisons revealed that the predicted effect was stronger in response to the ADC than in response to the AC ( $F(1,51) = 38.56, p < .001, \eta p^2 = .43$ ), and the predicted effect was



1 ambiguous stimulus, we have conducted a mixed ANOVA with CONDITION (AC, CC,  
2 AIC, ADC), ADAPTING (or CONTEXT) STIMULUS, PERCEPTUAL RESPONSE  
3 (downwards, upwards) as within-subjects factors and GROUP (ASC, TA) as between-  
4 subjects factor and Perceptual Durations as dependent variable. The mixed ANOVA showed  
5 significant effects of CONDITION ( $F(3,153) = 7.69, p < .001, \eta^2 = .13$ ) and PERCEPTUAL  
6 RESPONSE ( $F(1,51) = 14.01, p < .001, \eta^2 = .22$ ) revealing that perceptual ‘downwards’  
7 responses were longer than ‘upwards’ responses, a finding termed the *perceptual orientation*  
8 *bias*, i.e. the perception of the ambiguous cube-like figure (e.g., Necker cube) is typically  
9 biased towards a ‘front-side-down’ interpretation. Significant PERCEPTUAL RESPONSE  
10  $\times$  GROUP ( $F(1,51) = 5.21, p < .05, \eta^2 = .09$ ), CONDITION  $\times$  PERCEPTUAL RESPONSE  
11 ( $F(3,153) = 3.33, p < .05, \eta^2 = .06$ ), CONDITION  $\times$  PERCEPTUAL RESPONSE  $\times$  GROUP  
12 ( $F(3,153) = 2.90, p < .05, \eta^2 = .05$ ) and CONDITION  $\times$  ADAPTING STIMULUS  $\times$   
13 PERCEPTUAL RESPONSE ( $F(3,153) = 49.07, p < .001, \eta^2 = .49$ ) interactions were  
14 obtained, whereas the effect of GROUP was not significant ( $F < 1$ ). In order to simplify our  
15 data, we have calculated difference scores between the predicted and unpredicted Perceptual  
16 Durations (the selection of predicted and unpredicted Perceptual Durations were based on  
17 the predicted and unpredicted Orientation First Percept scores). The difference scores were  
18 calculated separately for ‘upwards’ and ‘downwards’ orientations and then averaged together  
19 to obtain a single ‘effect of manipulation’ variable per condition. In order to further explore  
20 the significant GROUP interactions obtained in the mixed ANOVA, subsequent one-way  
21 ANOVAs were performed for each GROUP separately.

1 For both ASC and TA groups, there was no difference between the adaptation (the  
2 AC) and the context (the CC) conditions (both F-values < 1), meaning that both conditions  
3 were influencing the perception of both groups at a comparable strength.

4 <<<Figure 3 about here>>>  
5

6 **ASC.** The ‘effect of manipulation’ variables were different between experimental  
7 conditions (as shown by significant effect of CONDITION ( $F(3,90) = 16.29, p < .001, \eta p^2 =$   
8  $.35$ )). Subsequent one-way ANOVAs, comparing pairwise all the experimental conditions  
9 (B-H corrected), revealed that the ‘effect of manipulation’ was larger in response to the ADC  
10 condition compared to all other conditions ( $F(1,30) \geq 11.92, p < .003, \eta p^2 \geq .28$ ). That is, the  
11 experimental manipulation was strongest in the ASC group, when bottom-up and top-down  
12 manipulations were congruent (Fig. 3).

13 **TA.** The effect of manipulations was different between experimental conditions as  
14 shown by a significant effect of CONDITION ( $F(3,63) = 6.34, p < .003, \eta p^2 = .23$ ).  
15 Subsequent one-way ANOVAs, comparing pairwise all experimental conditions (B-H  
16 corrected), revealed that the ‘effect of manipulation’ was stronger in response to the condition  
17 where bottom-up and top-down manipulations were congruent (the ADC) compared to the  
18 values obtained in response to the context condition (the CC), or the condition where bottom-  
19 up and top-down manipulations were incongruent (the AIC) ( $F(1,21) \geq 10.61, p < .005, \eta p^2$   
20  $\geq .34$ ). Furthermore, the effect of context alone (the CC) was stronger than that of AIC  
21 ( $F(1,21) = 7.98, p < .02, \eta p^2 = .28$ ).

22 More results (<https://osf.io/nhrxk/>), as well as the spreadsheet of the summary  
23 data

1 (<https://mfr.osf.io/render?url=https://osf.io/4bcqz/?action=download%26mode=render>), and our experimental code (<https://osf.io/y73mn/>), are available online.

2 <<<Table 3 about here>>>

3  
4 Results containing dependent measures ‘Response to the First Reversal’ (standard  
5 measurement of effects of adaptation on ambiguous figure perception) and ‘Reversal Rate’  
6 (standard measurement of ambiguous figure perception) are included in the supplement  
7 results section online. Furthermore, Spearman correlation coefficients were used to examine  
8 the relationship between task performance and symptom severity indexed in AQ scores as  
9 well as systemizing tendency indexed in IPT scores (<https://osf.io/nhrxk/>).

10

## Discussion

1  
2 We aimed to examine perceptual processing in ASC and a matched TA group using  
3 a previously validated paradigm, which, for the first time allows the investigation of both the  
4 independent and the combined influence of bottom-up and top-down influences on  
5 perception of ambiguous figures in ASC. We replicated previous findings of adaptation  
6 (Intaitè et al., 2013; Long & Moran, 2007; Long et al., 2002) and context (Intaitè et al., 2013;  
7 Sundaeswara & Schrater, 2008) effects on the perception of ambiguous figures in TA and  
8 found no differences between the ASC and the TA groups. The current results confirmed  
9 both an adaptation effect in the adaptation condition (the AC) and a context effect in the  
10 context condition (the CC), the latter of which surpassed the adaptation effect for both groups.  
11 Furthermore, we obtained the perceptual orientation bias, which is a typical finding showing  
12 that the cube-like ambiguous figures are perceived in ‘front-side-down’ orientation for longer  
13 durations (Dobbins & Grossmann, 2010; Kersten & Yuille, 2003; Murata et al., 2003;  
14 Sundaeswara & Schrater, 2008; Toppino & Long, 2015; Troje & McAdam, 2010; Washburn  
15 et al., 1931). With respect to group differences, we found that the effect of manipulation on  
16 Perceptual Durations for the ASC group was strongest when bottom-up manipulation was  
17 further strengthened by the top-down manipulation (the ADC) compared to all other  
18 conditions. The results of the TA group revealed that the same manipulation (the ADC) was  
19 comparable to independent adaptation (the AC) condition, but stronger than independent  
20 context (the CC) or incongruous bottom-up and top-down manipulation (the AIC). To sum  
21 up, our results add to converging evidence of additivity of bottom-up and top-down processes  
22 operating in the human visual system (Intaitè et al., 2013; Long & Toppino, 2004; Toppino,

1 2003) and show that general characteristics of basic visuo-perceptual functioning appear  
2 intact in ASC (for an extended discussion please see: (<https://osf.io/nhrxk/>)).

3 A Bayesian account (Pellicano & Burr, 2012) proposes that the interpretation of  
4 sensory information is less biased by prior experiences in ASC. Adaptation is a form of  
5 experience-dependent plasticity in which the current sensory experience is modified by the  
6 stimuli presented only up to several seconds before. Certain forms of this adaptation are  
7 reduced in ASC (Turi et al., 2015; Turi, Karaminis, Pellicano, & Burr, 2016). However, the  
8 type of adaptation tested in this study is not reduced in ASC, as in this paradigm the  
9 presentation time of at least one and a half minutes is necessary to attain the effect (Long et  
10 al., 1992). A shorter stimulus presentation exhibits a priming effect, that is the subsequent  
11 ambiguous figure is typically perceived in the same interpretation as the preceding  
12 unambiguous figure (Bugelski & Alampay, 1961; Long et al., 1992).

13 Our results also add to the view of perception in ASC being characterized by relative  
14 autonomy from top-down or contextual information resulting in more veridical perception as  
15 lined out by the Enhanced Perceptual Functioning account (Mottron et al., 2006). However,  
16 our results indicate that altered top-down modulation is stimulus- and task-specific  
17 (Greenaway & Plaisted, 2005). Mitchell and colleagues (2010) suggest that some of the  
18 induced top-down effects are atypically engaged in autistic perception. Top-down effects  
19 might be attenuated in autistic cognition, especially when they are not task-relevant; TA  
20 individuals are compelled to engage top-down processing to a much greater degree.  
21 Perceptual ambiguity and its manipulations reveal that we actively interpret the available  
22 visual information, rather than passively view it. Our data suggest that top-down processing  
23 exhibited by the ASC participants might as well depend on the respective bottom-up

1 processing, hence in such cases where top-down manipulation was supported by bottom-up  
2 manipulation (i.e., the ADC), we observe intact top-down processing. However, if the top-  
3 down manipulation is not supported by bottom-up manipulation (i.e., the AIC), the ASC  
4 participants show reduced interpretation of the available visual information, thus their top-  
5 down processing might be merely stimulus or task-dependent.

6 In summary, this study examines whether top-down processing in ASC is a general  
7 or a task-specific deficit. Our findings indicate that both bottom-up and top-down processes  
8 can influence perception concurrently and independently for the TA participants. We have  
9 observed a perturbation in the dynamics of perceptual ambiguity in ASC individuals when  
10 the presented top-down manipulation was not supporting the bottom-up manipulation. Thus  
11 ASC participants show the atypical task-specific bottom-up and top-down interactions that  
12 imply minor abnormalities in their visual perception and attention, thus supporting the view  
13 of stimulus-specific top-down modulation in ASC.

#### 14 **Acknowledgements**

15 This study was supported by the fellowship (SFRH/BPD/77563/2011) to M.I., from  
16 Portuguese Foundation of Science and Technology. We thank Melanie Rohde, Susanne  
17 Holocher, Marius Jonas, and Judith Schmitz for their help with recruitment and testing  
18 participants, Mathis Jording for help with experimental setup, and Dr. Brian Richardson from  
19 SR Research for help with eye-tracking data management. We would like to thank the  
20 reviewers for their helpful comments and suggestions.

21

## References

- 1 Aster, M., Neubauer, A., & Horn, R. (2006). WIE. *Wechsler Intelligenztest Für*  
2 *Erwachsene (WIE). Deutschsprachige Bearbeitung Und Adaptation Des WAIS-III von*  
3 *David Wechsler. Harcourt Test Services: Frankfurt.*
- 4 Bagby, R. M., Parker, J. D. A., & Taylor, G. J. (1994). The twenty-item Toronto  
5 Alexithymia Scale-I. Item selection and cross-validation of the factor structure.  
6 *Journal of Psychosomatic Research, 38*, 23–32.  
7 [http://doi.org/http://dx.doi.org/10.1016/0022-3999\(94\)90005-1](http://doi.org/http://dx.doi.org/10.1016/0022-3999(94)90005-1)
- 8 Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The  
9 Autism-Spectrum Quotient (AQ): Evidence from Asperger Syndrome/High-  
10 Functioning Autism, Males and Females, Scientists and Mathematicians. *Journal of*  
11 *Autism and Developmental Disorders, 31*, 5–17.  
12 <http://doi.org/10.1023/A:1005653411471>
- 13 Baron-Cohen, S., Wheelwright, S. (2004). The Empathy Quotient: an investigation of  
14 adults with Asperger syndrome and high functioning autism and normal sex  
15 differences. *Journal of Autism and Developmental Disorders, 34*, 163–175.  
16 <http://doi.org/10.1023/B:JADD.0000022607.19833.00>
- 17 Baron-Cohen, S., Wheelwright, S., Hill, J., Raste, Y., & Plumb, I. (2001). The “Reading  
18 the Mind in the Eyes” Test revised version: A study with normal adults, and adults  
19 with Asperger syndrome or high-functioning autism. *Journal of Child Psychology and*  
20 *Psychiatry, 42*, 241–251. <http://doi.org/10.1111/1469-7610.00715>
- 21 Baron-Cohen, S., Wheelwright, S., Scahill, V., Lawson, J., & Spong, A. (2001). Are  
22 intuitive physics and intuitive psychology independent? A test with children with  
23 Asperger syndrome. *Journal of Developmental and Learning Disorders, 5*, 47–78.
- 24 Baron-Cohen, S., Richler, J., Bisarya, D., Gurunathan, N., & Wheelwright, S. (2003). The  
25 Systemizing Quotient: an investigation of adults with Asperger syndrome and high  
26 functioning autism, and normal sex differences. *Philosophical Transactions of the*  
27 *Royal Society B: Biological Sciences, 358*, 361–374.  
28 <http://doi.org/0.1098/rstb.2002.1206>
- 29 Bird, G., Catmur, C., Silani, G., Frith, C., & Frith, U. (2006). Attention does not modulate  
30

1 neural responses to social stimuli in autism spectrum disorders. *NeuroImage*, *31*,  
2 1614–1624. <http://doi.org/10.1016/j.neuroimage.2006.02.037>

3 Bugelski, B. R., & Alampay, D. A. (1961). The role of frequency in developing perceptual  
4 sets. *Canadian Journal of Psychology*, *15*, 205–211. <http://doi.org/10.1037/h0083443>

5 Cook, J., Barbalat, G., & Blakemore, S. J. (2012). Top-down modulation of the perception  
6 of other people in schizophrenia and autism. *Frontiers in Human Neuroscience*, *6*, 1–  
7 10. <http://doi.org/10.3389/fnhum.2012.00175>

8 Dakin, S., & Frith, U. (2005). Vagaries of visual perception in autism. *Neuron*, *48*, 497–  
9 507. <http://doi.org/10.1016/j.neuron.2005.10.018>

10 Dobbins, A. C., & Grossmann, J. K. (2010). Asymmetries in perception of 3D orientation.  
11 *PLoS ONE*, *5*, 1–5. <http://doi.org/10.1371/journal.pone.0009553>

12 Freitag, C. M., Konrad, C., Häberlen, M., Kleser, C., von Gontard, A., Reith, W., Krick,  
13 C. (2008). Perception of biological motion in autism spectrum disorders.  
14 *Neuropsychologia*, *46*, 1480–1494.  
15 <http://doi.org/10.1016/j.neuropsychologia.2007.12.025>

16 Frith, U. (2003). *Autism: Neural basis and treatment possibilities*. Hoboken, NJ, Wiley,  
17 1st edition.

18 Greenaway, R., & Plaisted, K. (2005). Top-Down modulation in Autistic Spectrum  
19 Modulation disorders Is stimulus-specific. *Psychological Science*, *16*, 987–994.  
20 <http://doi.org/10.1111/j.1467-9280.2005.01648.x>

21 Happé, F. G. E. (1996). Studying weak coherence at low levels: Children with autism do  
22 not succumb to visual illusions. A research note. *Psychology and Psychiatry*, *37*, 873–  
23 877. <http://doi.org/10.1111/j.1469-7610.1996.tb01483.x>

24 Happé, F. G. E., & Frith, U. (2006). The weak coherence account: Detail-focused  
25 cognitive style in autism spectrum disorders. *Journal of Autism and Developmental*  
26 *Disorders*, *36*, 5–25. <http://doi.org/10.1007/s10803-005-0039-0>

27 Hautzinger, M., Keller, F., & Kühner, C. (2006). BDI-II. *Beck Depressions-Inventar*  
28 *(BDI-II). Revision. Frankfurt/Main: Harcourt Test Services*.

29 Hill, E. L. (2004). Executive dysfunction in autism. *Trends in Cognitive Sciences*, *8*, 26–  
30 32. <http://doi.org/10.1016/j.tics.2003.11.003>

31 Holm, S. (1979). A Simple Sequentially Rejective Multiple Test Procedure. *Scandinavian*

1        *Journal of Statistics*, 6, 65–70. <http://doi.org/10.2307/4615733>

2        Howell, D. C. (2009). *Statistical methods for psychology* (7th ed.). Belmont, CA:  
3        Wadsworth Publishing.

4        Intaitė, M., Noreika, V., Šoliūnas, A., & Falter, C. M. (2013). Interaction of bottom-up  
5        and top-down processes in the perception of ambiguous figures. *Vision Research*, 89,  
6        24–31. <http://doi.org/10.1016/j.visres.2013.06.011>

7        Kersten, D., & Yuille, A. (2003). Bayesian models of object perception. *Current Opinion*  
8        *in Neurobiology*, 13, 150–158. [http://doi.org/10.1016/S0959-4388\(03\)00042-4](http://doi.org/10.1016/S0959-4388(03)00042-4)

9        Long, G. M., Stewart, J. A., & Glancey, D. E. (2002). Configural biases and reversible  
10        figures: Evidence of multilevel grouping effects. *American Journal of Psychology*,  
11        115, 581–607. <http://doi.org/10.2307/1423528>

12        Long, G. M., Toppino, T. C., & Mondin, G. W. (1992). Prime time: Fatigue and set effects  
13        in the perception of reversible figures. *Perception & Psychophysics*, 52, 609–616.  
14        <http://doi.org/10.3758/BF03211697>

15        Long, G., & Moran, C. J. (2007). How to keep a reversible figure from reversing: Teasing  
16        out top – down and bottom – up processes. *Perception*, 36, 431–445.  
17        <http://doi.org/10.1068/p5630>

18        Long, G., & Toppino, T. (2004). Enduring interest in perceptual ambiguity: Alternating  
19        views of reversible figures. *Psychological Bulletin*, 130, 748–768.  
20        <http://doi.org/10.1037/0033-2909.130.5.748>

21        Loth, E., Gómez, J. C., & Happé, F. (2010). When seeing depends on knowing: Adults  
22        with Autism Spectrum Conditions show diminished top-down processes in the visual  
23        perception of degraded faces but not degraded objects. *Neuropsychologia*, 48, 1227–  
24        1236. <http://doi.org/10.1016/j.neuropsychologia.2009.12.023>

25        Mitchell, P., Mottron, L., Soulières, I., & Ropar, D. (2010). Susceptibility to the Shepard  
26        illusion in participants with autism: Reduced top-down influences within perception?  
27        *Autism Research*, 3, 113–119. <http://doi.org/10.1002/aur.130>

28        Mottron, L., Dawson, M., Soulières, I., Hubert, B., Burack, J. (2006). Enhanced perceptual  
29        functioning in autism: An update, and eight principles of autistic perception. *Journal*  
30        *of Autism and Developmental Disorders*, 36, 27–43. [http://doi.org/10.1007/s10803-](http://doi.org/10.1007/s10803-005-0040-7)  
31        [005-0040-7](http://doi.org/10.1007/s10803-005-0040-7)

- 1 Murata, T., Matsui, N., Miyauchi, S., Kakita, Y., & Yanagida, T. (2003). Discrete  
2 stochastic process underlying perceptual rivalry. *Neuroreport*, *14*, 1347–1352.  
3 <http://doi.org/10.1097/01.wnr.0000077553.91466.41>
- 4 Muth, A., Hönekopp, J., & Falter, C. M. (2014). Visuo-Spatial Performance in Autism: A  
5 Meta-analysis. *Journal of Autism and Developmental Disorders*, *44*, 3245–3263.  
6 <http://doi.org/10.1007/s10803-014-2188-5>
- 7 Nelson, S. B., & Valakh, V. (2015). Excitatory / Inhibitory balance and circuit  
8 homeostasis in Autism Spectrum Disorders. *Neuron*, *87*, 684–698.  
9 <http://doi.org/10.1016/j.neuron.2015.07.033>
- 10 Pellicano, E., & Burr, D. (2012). When the world becomes “too real”: A Bayesian  
11 explanation of autistic perception. *Trends in Cognitive Sciences*, *16*, 504–510.  
12 <http://doi.org/10.1016/j.tics.2012.08.009>
- 13 Plaisted, K., O’Riordan, M., & Baron-Cohen, S. (1998). Enhanced visual search for a  
14 conjunctive target in autism: A research note. *Journal of Child Psychology and*  
15 *Psychiatry, and Allied Disciplines*, *39*, 777–783. [http://doi.org/10.1111/1469-](http://doi.org/10.1111/1469-7610.00376)  
16 [7610.00376](http://doi.org/10.1111/1469-7610.00376)
- 17 Ropar, D., Mitchell, P., & Ackroyd, K. (2003). Do children with autism find it difficult to  
18 offer alternative interpretations to ambiguous figures? *British Journal of*  
19 *Developmental Psychology*, *21*, 387–395.  
20 <http://doi.org/10.1348/026151003322277766>
- 21 Shah, A., & Frith, U. (1983). An islet of ability in autistic children: A research note.  
22 *Journal of Child Psychology and Psychiatry*, *24*, 613–620.  
23 <http://doi.org/10.1111/j.1469-7610.1983.tb00137.x>
- 24 Shah, A., & Frith, U. (1993). Why do autistic individuals show superior performance on  
25 the block design task? *Journal of Child Psychology and Psychiatry*, *34*, 1351–1364.  
26 <http://doi.org/10.1111/j.1469-7610.1993.tb02095.x>
- 27 Simmons, D. R., Robertson, A. E., McKay, L. S., Toal, E., McAleer, P., & Pollick, F. E.  
28 (2009). Vision in autism spectrum disorders. *Vision Research*, *49*, 2705–2739.  
29 <http://doi.org/10.1016/j.visres.2009.08.005>
- 30 Sobel, D. M., Capps, L. M., & Gopnik, A. (2005). Ambiguous figure perception and  
31 theory of mind understanding in children with autistic spectrum disorders. *British*

1        *Journal of Developmental Psychology*, 23, 159–174.  
2        <http://doi.org/10.1348/026151004X20694>

3        Sundareswara, R., & Schrater, P. R. (2008). Perceptual multistability predicted by search  
4        model for Bayesian decisions. *Journal of Vision*, 8, 1–19. <http://doi.org/10.1167/8.5.12>

5        Toppino, T. C. (2003). Reversible-figure perception: Mechanisms of intentional control.  
6        *Perception & Psychophysics*, 65, 1285–1295. <http://doi.org/10.3758/BF03194852>

7        Toppino, T. C., & Long, G. M. (2015). Time for a change: What dominance durations  
8        reveal about adaptation effects in the perception of a bi-stable reversible figure.  
9        *Attention, Perception, & Psychophysics*, 77, 867–882. <http://doi.org/10.3758/s13414->  
10        014-0809-x

11        Troje, N. F., & McAdam, M. (2010). The viewing-from-above bias and the silhouette  
12        illusion. *I-Perception*, 1, 143–148. <http://doi.org/10.1068/i0408>

13        Turi, M., Burr, D. C., Igliozzi, R., Aagten-murphy, D., Muratori, F., & Pellicano, E.  
14        (2015). Children with autism spectrum disorder show reduced adaptation to number.  
15        *Proceedings of the National Academy of Sciences of the United States of America*,  
16        112, 7868–7872. <http://doi.org/10.1073/pnas.1504099112>

17        Turi, M., Karaminis, T., Pellicano, E., & Burr, D. (2016). No rapid audiovisual  
18        recalibration in adults on the autism spectrum. *Scientific Reports*, 6, 1–7.  
19        <http://doi.org/10.1038/srep21756>

20        Von Grünau, M. W., Wiggin, S., & Reed, M. (1984). The local character of perspective  
21        organization. *Perception & Psychophysics*, 35, 319–324.  
22        <http://doi.org/10.3758/BF03206335>


23        Washburn, M., Mallay, H., & Naylor, A. (1931). The influence of the size of an outline  
24        cube on the fluctuations of its perspective. *American Journal of Psychology*, 43, 484–  
25        489. <http://doi.org/10.2307/1414619>

26        WHO. (1992). International Statistical Classification of Diseases and Related Health  
27        Problems, 10th revision, 10.


28  
29  
30  
31

1 **FIGURES**

2



Schematic representation of a single trial:




3

4 **Figure 1.** Schematic representation of all experimental conditions: (a) the adaptation  
 5 condition (AC), (b) the context condition (CC), (c) the adaptation different from context  
 6 condition (ADC), and (d) the adaptation identical to context condition (AIC). A schematic  
 7 representation of a single experimental trial is depicted in the lower part of the figure.

8

2



1


2 **Figure 2.** Median number of times that autistic spectrum condition (ASC) and typical adults  
3 (TA) first reported the “predicted” orientation of the ambiguous squares stimulus, in  
4 accordance with the orientation elicited by the respective experimental condition. Violin  
5 plots depict the shape of the distribution and region inside the violin contains all of the  
6 observed data. Values obtained in the wider parts of the violin are more probable than those  
7 in narrower parts. The median and interquartile ranges are displayed by overlaying a box  
8 plot. Error bars represent the lowest and the highest data points still within 1.5 interquartile  
9 ranges (IQR).

10

11

12

3



1

2 **Figure 3.** Median "effect of manipulation" variables. Violin plots depict the shape of the  
3 distribution and region inside the violin contains all of the observed data. Values  
4 obtained in the wider parts of the violin are more probable than those in narrower  
5 parts. The median and interquartile ranges are displayed by overlaying a box plot. Error  
6 bars represent the lowest and the highest data points still within 1.5 IQR.

7

8

9

10

11

12

13

1  
2  
3  
  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

**TABLES**

	ASC (N=31; 20 male)			TA (N=22; 12 male)		
	Mean	SD	Range	Mean	SD	Range
<b>Age</b>	43:6	8:6	20:2 – 55:9	39:3	8:7	22:2 – 53:0
<b>VIQ</b>	113	17	81 – 135	115	13	94 – 137
<b>PIQ</b>	108	18	67 – 141	107	14	73 – 132
<b>FIQ</b>	113	18	78 – 140	111	13	83 – 139
<b>AQ</b>	41	4	27 – 48	17	6	6 – 29
<b>IPT</b>	8	3	2 – 15	10	2	4 – 14

**Table 1:** Means, standard deviations, and ranges of age (years:months), verbal IQ (VIQ), performance IQ (PIQ), full IQ (FIQ), Autism-Spectrum Quotient (AQ), number of mistakes made in Intuitive Physics Test (IPT) of participants with autism spectrum condition (ASC) and typical adults (TA).

		ASC (N=12; 10 male)			TA (N=7; 4 male)		
		Fix	DT	PA	Fix	DT	PA
AOI-1	AP (SD)	70.8 (2.8)	83.6 (35.7)	627.2 (134.9)	77.8 (2.4)	94.6 (29.9)	548.6 (95.0)
	TP (SD)	71.2 (3.4)	21.1 (11.3)	629.5 (144.9)	74.5 (3.1)	23.2 (9.7)	558.4 (93.4)
AOI-2	AP (SD)						
	TP (SD)	98.8 (14.4)	28.2 (4.3)	623.2 (146.9)	98.5 (3.7)	28.7 (1.4)	560.1 (88.4)

**Table 2:** Means (SD) of eye tracking data for fixations (Fix.; %) dwell times (DT; s) and Pupil area (PA; arbitrary units) of participants with an autism spectrum condition (ASC) and typical adults (TA) for the two Areas-of-Interest: AOI-1 (size: 1.51° × 1.51°) and AOI-2 (size: 5.82° × 5.82°) during Adaptation (AP) and Test periods (AP).

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18

Dependent variable	Findings for experimental Conditions and Groups	
<b>Orientation First Percept (1stP)</b>	Main ANOVA: CONDITION ( $F(3,153) = 15.71, p < .001, \eta p^2 = .24$ )	
	<b>AC &lt; CC:</b> ( $F(1, 51) = 25.15, p < .001, \eta p^2 = .33$ ) <b>AC &lt; ADC:</b> ( $F(1,51) = 38.56, p < .001, \eta p^2 = .43$ ) <b>AIC &lt; CC:</b> ( $F(1,51) = 6.47, p < .05, \eta p^2 = .11$ )	
	Group interactions: all F-values < 1 Group effect: $F < 1$	
<b>Perceptual Durations</b>	Main ANOVA: CONDITION ( $F(3,153) = 7.69, p < .001, \eta p^2 = .13$ )	
	PERCEPTUAL RESPONSE ( $F(1,51) = 14.01, p < .001, \eta p^2 = .22$ )	
	Group interactions: PERCEPTUAL RESPONSE $\times$ GROUP ( $F(1,51) = 5.21, p < .05, \eta p^2 = .09$ ), CONDITION $\times$ PERCEPTUAL RESPONSE $\times$ GROUP ( $F(3,153) = 2.90, p < .05, \eta p^2 = .05$ ). All other GROUP interactions F-values < 1 Group effect: $F < 1$	
	<b>ASC (N = 31; 20 male)</b>	<b>TA (N = 22; 12 male)</b>
<b>'Effect of Manipulation'</b>	<b>AC <math>\approx</math> CC</b> ( $F < 1$ ) <b>ADC &gt; AC, CC, AIC:</b> ( $F(1,30) \geq 11.92, p < .003, \eta p^2 \geq .28$ )	<b>AC <math>\approx</math> CC</b> ( $F < 1$ ) <b>ADC &gt; CC and AIC:</b> ( $F(1,21) \geq 10.61, p < .005, \eta p^2 \geq .34$ ) <b>AIC &lt; CC:</b> ( $F(1,21) = 7.98, p < .02, \eta p^2 = .28$ )

1

2 **Table 3:** Overview of results across dependent variables and conditions (AC: Adaptation Condition,  
3 CC: Context Condition, ADC: Adaptation different from Context, AIC: Adaptation identical to Context)  
4 of participants with an autism spectrum condition (ASC) and typical adults (TA) for all dependent  
5 variables of the experiment.

6

7

8