

# Associations between Lifestyle Factors and Parkinson's Disease in an Urban Sri Lankan Clinic Study

ORIGINAL

Ruwani Wijeyekoon<sup>1</sup>,  
Vindika Suriyakumara<sup>2</sup>,  
Ranjanie Gamage<sup>3</sup>, Tharushi Fernando<sup>2</sup>,  
Amila Jayasuriya<sup>2</sup>, Dhanusha Amarasinghe<sup>2</sup>,  
Harsha Gunasekara<sup>4</sup>,  
Dharshana Sirisena<sup>7</sup>, Dhammika Amaratunga<sup>5</sup>,  
Chanaka Muthukuda<sup>6</sup>, Roger A Barker<sup>1,8</sup>,  
Caroline Williams-Gray<sup>1</sup>, Ranil De Silva<sup>2</sup>

- 1 John van Geest Centre for Brain Repair, Department of Clinical Neurosciences, University of Cambridge, UK.
- 2 Genetic, Diagnostic and Research Laboratory, Department of Anatomy, Faculty of Medical Sciences, University of Sri Jayewardenepura, Nugegoda, Sri Lanka.
- 3 Institute of Neurology, National Hospital of Sri Lanka, Colombo, Sri Lanka.
- 4 Sri Jayewardenepura General Hospital, Nugegoda, Sri Lanka.
- 5 Independent Statistical Consultant and Researcher, Colombo, Sri Lanka
- 6 University of Alabama at Birmingham, USA.
- 7 Neurology Unit, Teaching Hospital, Kurunegala, Sri Lanka.
- 8 MRC-Wellcome Trust Cambridge Stem Cell Institute.

## Abstract

**Background:** Associations between certain environmental and lifestyle factors and Parkinson's disease (PD) have been reported in several studies, but information on these factors and Parkinson's Disease (PD) in South Asia, is limited.

**Objective:** To determine associations between lifestyle factors and PD in an urban clinic-based study in Sri Lanka.

**Methods:** In this case-control study, demographic and lifestyle factor data (including diet, coffee/tea drinking, smoking, alcohol status) was collected from an unselected cohort of PD patients and age and gender-matched controls attending clinics in Greater Colombo, Sri Lanka. Associations between lifestyle factors and PD status were assessed using Logistic Regression analysis, while links with age of PD onset were explored with Kaplan Meier and Cox Regression survival analyses. Results with  $p < 0.05$  were considered to be statistically significant.

**Findings:** Of 229 patients with parkinsonism, 144 had Idiopathic PD using standard diagnostic criteria. Controls numbered 102. Coffee drinkers and smokers were significantly less likely to have PD (coffee,  $p < 0.001$ ; Odds Ratio (OR)=0.264; smoking,  $p = 0.043$ ; OR=0.394). Coffee drinkers were older at PD onset ( $p < 0.001$ ). Similar trends seen with tea drinking were not statistically significant.

## Contact information:

Ranil De Silva.

**Address:** Department of Anatomy, Faculty of Medical Sciences, University of Sri Jayewardenepura, Nugegoda, Sri Lanka.

**Tel:** +94 11 5662609.

✉ [ranil@sjp.ac.lk](mailto:ranil@sjp.ac.lk)

**Conclusions:** This is the first formal study of PD and these lifestyle factors in South Asia. It demonstrates an inverse association between coffee drinking, smoking and PD, and an association between coffee drinking and later age of PD onset. This is in line with other studies done worldwide, suggesting biological associations with global relevance.

#### Keywords

Parkinson's Disease; Lifestyle; Coffee; Smoking; Sri Lanka.

## Introduction

Parkinson's Disease (PD) has a multifactorial aetiology, with a complex interaction of genetic and environmental risk factors. Several environmental and lifestyle factors have been associated with PD and especially smoking, coffee and tea intake have been linked to reduced risk [1, 2]. Meta-analyses of global studies have confirmed a decreased relative risk for smoking (pooled Relative Risk (RR) 0.59) and coffee intake (pooled RR 0.75) [2]. Further dose dependent effects on delaying the age of PD onset have been seen with coffee intake, while tea intake has also been associated with decreased risk and delayed age of onset [3].

Few studies have investigated these associations in South Asian populations, which have different genetic and lifestyle exposure profiles compared to Western and East Asian populations. Studies on PD in Indian and other South Asian populations have also found that PD patients indigenous to this part of the world have some phenotypic differences compared to Caucasian populations. In particular, an increased proportion of patients have levodopa hypo-responsiveness, bradykinesia-dominant disease and early cognitive dysfunction. This difference in phenotype has been suggested to be linked to higher rates of diabetes and hypertension in the

populations along with dietary and genetic factors [4]. Considering these differences, few studies have investigated associations between PD and lifestyle factors in South Asia to ascertain if these are different compared to those established in Caucasian and other populations. Studies in India have found associations between increased PD risk and non-smoking status, well water drinking and pesticide use [5, 6] but none have investigated factors such as coffee and tea intake.

Sri Lanka is a multi-ethnic country with an agricultural background and one of the fastest ageing populations in South Asia. Neurodegenerative diseases such as PD and Alzheimer's disease (AD) are thus becoming an increasingly important healthcare issue, but few formal studies have been performed. A recent neuropathological study on neurodegenerative pathology in elderly brains found more PD related Lewy-body pathology in Sri Lankan subjects compared to Indian subjects, while AD related pathology was similar in both groups, potentially indicating variability in neurodegenerative disease pathology even among South Asian countries [7].

There is currently limited data available on PD and no data on lifestyle risk factors and PD, in Sri Lanka. Due to cultural reasons, the lifestyle factor profiles in South Asian countries can be very different

from those seen in Western countries. For example in Sri Lanka, smoking and alcohol intake have a strong male predominance. Furthermore, most Sri Lankans drink tea regularly, while coffee intake is much lower than in the West. Therefore, this study aimed to investigate, for the first time, the association between lifestyle factors and PD in a clinic based population in Colombo, Sri Lanka.

## Methods

This was a retrospective case-control study. Patients were recruited from the Neurology and Movement Disorders clinics in hospitals in Greater Colombo, Sri Lanka over a 4 year period. All patients who had been clinically diagnosed with Parkinson's disease (PD) or parkinsonism by a certified Neurologist were approached regarding the study and all consenting patients (~80%) were recruited. Retrospective detailed case note review of recruited patients was conducted in order to identify those who fulfilled UK Queen Square Brain Bank criteria for Idiopathic PD, and only those fulfilling these criteria were included in the final study. Controls consisted of patients without neurological disorders recruited from non-neurological clinics or relatives of PD patients, who had no neurological complaints. Control status was considered to be the absence of a diagnosis of PD up to the point of assessment.

The study was performed with appropriate approval from the University of Sri Jayewardenepura Ethics Committee (No. 427/09) and informed consent was obtained from all participants.

Demographic, lifestyle and medical history information was collected using a pro-forma data sheet. This included information on province of birth, ethnicity, education level, diet, smoking, alcohol, coffee and tea intake, pesticide exposure and presence of medical co-morbidities (ischaemic heart disease, diabetes and hypertension). Details of the patients' neurological history and examination were obtained from their medical records.

Variables were further categorised as appropriate prior to analysis. Due to the location of the clinics most participants were from the Western province. Therefore 'province of birth' was categorised into Western Province and non-Western Province. Education level was categorised as 'below A-Level' (i.e. left school before the age of 18) and 'A-Level and above'. Coffee drinking was categorised as "coffee drinkers" and "coffee non-drinkers", where "coffee non-drinkers" were considered to be those who drank <2 cups/month. As all participants drank tea, tea intake was divided into "standard intake" (<=3 cups/day) and "high intake" (>3 cups/day). Presence of pesticide exposure was considered to involve regular occupational spraying of pesticides. Other data was categorised as follows -: diet (vegetarian/non-vegetarian); smoking (ever-smokers/never-smokers); alcohol (ever-drinkers/never-drinkers); ischaemic heart disease, diabetes, hypertension (presence/absence).

Statistical analyses were performed using SPSS version 21. Between-group comparisons were performed on percentage numbers, using T tests and Chi-squared tests. Logistic Regression analysis was performed to determine the effect of the demographic and lifestyle factors on the risk of PD, with PD status as the dependent variable. Kaplan Meier survival analysis and Cox Regression analysis were performed to assess the effect of these factors on the age of onset of PD. Results with a p value <0.05 were considered to be statistically significant.

## Results

### Demographics

229 patients with parkinsonism were recruited. Of these, 144 fulfilled UK Queen Square Brain Bank Criteria for Idiopathic PD based on retrospective case note review. An age and gender matched group of 102 controls were also recruited.

No significant differences in age, gender, ethnicity and province of birth were seen between the patient and control groups (**Table 1**). Education level was higher amongst the PD group, which may be related to higher educated PD patients being more likely to attend clinic.

**Table 1.** Key demographics and summary of the measured lifestyle factors and co-morbidities.

Demographics	Patients	Controls	p value
	%	%	
Gender			
Male	59.7	54.9	0.260
Female	40.3	45.1	
Age at Assessment			
Average ± SD	61.8 ± 8.2	61.7 ± 7.9	0.950
Ethnicity			
Burgher	0.7	1.0	0.914
Muslim	7.1	4.9	
Sinhala	87.3	89.2	
Tamil	4.9	4.9	
Province of Birth			
Western Province	72.9	65.0	0.187
non Western Province	27.1	35.0	
Education level			
Below A Level	73.6	86.1	0.022*
A Level and above	26.4	13.9	
<b>Lifestyle Factors and Co-morbidities</b>			
Diet			
Vegetarian	11.3	9.8	0.701
Smoking status			
ever smokers	28.1	44.1	0.010*
Alcohol status			
ever drinkers	35.0	44.1	0.151
Coffee drinkers	34.8	67.6	<0.001**
Tea drinking			
high intake	14.8	24.5	0.060
Ischaemic Heart Disease	6.0	7.8	0.571
Diabetes	20.6	17.64	0.570
Hypertension	27.6	30.4	0.640
Pesticide exposure history	3.6	4.9	0.624
Total Number	144	102	

SD: Standard Deviation; \*: p<0.05; \*\*: p<0.01.

## Lifestyle Factors and PD

Simple group comparison analyses suggested that a significantly smaller percentage of PD patients were coffee drinkers or ex/current smokers (**Table 1**). Logistic regression analysis, with PD versus control status as the dependent variable and age, gender and above stated lifestyle factors/comorbidities as predictor variables, indicated a highly significant negative association between coffee drinking and PD (p<0.001, Odds Ratio(OR) 0.264; 95% Confidence Interval (CI) 0.14–0.499) (**Table 2**). There was also a significant negative association between 'ever smoking status' and PD (p=0.043, OR 0.394; 95% CI 0.159–0.972). The negative association between high tea intake and PD did not reach statistical significance (p=0.106, OR=0.523; 95% CI 0.238–1.148).

Considering the significant effect of gender, the data was also analysed separately for males and fe-

**Table 2.** Summary table displaying the results of the Logistic Regression analysis of factors contributing to the risk of PD.

Variable	B coefficient	p value	Odds Ratio-Exp (B)	95% Confidence Intervals for Exp(B)	
				Lower	Upper
Gender	-1.079	0.015	0.340*	0.142	0.813
Age At Assessment	-0.005	0.810	0.995	0.957	1.035
Diet	-0.212	0.671	0.809	0.304	2.151
Smoking	-0.932	0.043	0.394*	0.159	0.972
Alcohol	-0.619	0.194	0.539	0.212	1.370
Coffee	-1.330	0.000	0.264**	0.140	0.499
Tea	-0.649	0.106	0.523	0.238	1.148
Education	0.653	0.109	1.921	0.864	4.272
IHD	-1.084	0.143	0.338	0.079	1.443
Diabetes	-0.222	0.611	0.801	0.341	1.881
Hypertension	-0.263	0.472	0.769	0.375	1.576
Pesticide Exposure	-0.683	0.373	0.505	0.112	2.272
Constant	4.676	0.014	107.305		

\*: p<0.05; \*\*: p<0.01

males. The association between decreased PD and smoking was more significant in males ( $p=0.019$ , OR 0.313; 95% CI 0.118 – 0.826). Coffee drinking and decreased PD were more significantly associated in females ( $p<0.001$ , OR 0.096; 95% CI 0.031–0.299).

### Lifestyle Factors and PD Age of Onset

Survival analysis with gender stratification was performed to investigate the relationship between each of the lifestyle factors and age of onset of PD symptoms. Kaplan Meier survival analysis demonstrated that regular coffee drinking was associated with a later age of onset of PD (overall,  $p<0.001$ ; females,

$p<0.001$ ; males,  $p= 0.013$ ; Log Rank (Mantel-Cox), with the significance withstanding Bonferroni correction for multiple testing for the population as a whole and also for female participants (**Figure 1a**). In males, ever-smoking was associated with delayed PD onset ( $p=0.048$ ; Log Rank (Mantel-Cox) (**Figure 1b**).

Higher education level was associated with earlier PD onset ( $p=0.01$ ; Breslow (Generalised Wilcoxon), which may indicate that those with increased education were more aware of the significance of early PD symptoms. There was no significant effect with other variables.


Cox regression analysis confirmed the significant association between coffee intake and delayed PD onset following adjustment for age, gender and all measured variables ( $p<0.001$ ; Exp(B) 2.229) (**Table 3**).

**Table 3.** Summary table displaying the results of the Cox Regression analysis of age of onset of PD with all measured variables.

Variable	B coefficient	p value	Odds Ratio-Exp (B)	95% Confidence Intervals for Exp(B)	
				Lower	Upper
Gender	0.095	0.721	1.099	0.654	1.848
Diet	0.234	0.465	1.264	0.675	2.367
Smoking	0.455	0.146	1.576	0.854	2.909
Alcohol	-0.209	0.495	0.811	0.445	1.479
Coffee	0.802	0.000	2.229**	1.471	3.378
Tea	0.354	0.211	1.425	0.818	2.481
Education	-0.149	0.537	0.862	0.536	1.384
IHD	0.800	0.181	2.226	0.689	7.196
Diabetes	0.065	0.817	1.067	0.618	1.842
Hypertension	0.419	0.094	1.521	0.931	2.485
Pesticide Exposure	-0.074	0.902	0.928	0.286	3.015

\*:  $p<0.05$ ; \*\*:  $p<0.01$

**Figure 1:** Kaplan Meier curves of age of onset of PD in different patient groups.



a) Coffee drinkers versus non-drinkers; b) Male smokers versus non-smokers.

## Conclusions and Discussion

This is the first study evaluating associations between lifestyle factors and PD in Sri Lanka. It indicates a significant association between coffee drinking and a decreased prevalence (~70% less) and delayed age of onset of PD. It also demonstrates an association between smoking and non-PD status. These findings are in keeping with other studies from around the world [2] and support the contention that these factors are strongly associated with the development of PD regardless of genetic and ethnic background.

Tea intake has been associated with a decreased risk of PD [2]. The association between high tea intake and decreased PD status seen in this study was not statistically significant. This may be due to the relatively small sample size and the lack of non-tea drinking participants.

There are many possible theories to explain the biological basis of the effects seen with caffeine intake, smoking/nicotine and PD. In the past, these associations have been considered to be due to a lack of addictive behaviours in the premorbid personalities of patients likely to develop PD but subsequent research has shown dose dependent protective effects on risk and age of onset, with beneficial effects of caffeine in therapeutic trials, possibly through its action as an Adenosine A2 receptor antagonist [2, 8, 9]. Several studies have also demonstrated evidence for the neuroprotective effects of coffee and caffeine in other neurodegenerative diseases, including in the progression from Mild Cognitive Impairment to AD dementia [10]. The effect of caffeine on the gut microbiota has also been suggested to play a role [11].

Nicotine may also have some neuroprotective effects, acting via nicotinic acetylcholine receptors in neurones and glia [12]. However, PD patients have also been found to have less difficulty quitting smoking than those without PD, suggesting that decreased responsiveness to nicotine could be a prodromal feature of PD itself. This potentially

indicates a role for reverse causation in explaining the association between smoking and decreased PD - namely people are less likely to smoke regularly because they are more likely to develop PD [13].

This is the first study investigating these lifestyle factors and PD in Sri Lanka, but it has some limitations. Due to the recruitment strategy, most of the participants are from regions around the capital, Colombo, which may not be fully representative of the population of the entire country. Smoking and alcohol intake are generally very low among females in Sri Lanka for cultural reasons, which may attenuate the overall effects seen, although the separate analysis for males and females does take account of this.

The lack of detailed information on the lifestyle factors (e.g. age at start of regular coffee intake, number of years of intake and number of cups/day), is a further limitation. Thus, it is not possible to determine if the exposure started prior to the onset of PD symptoms (which would support a more causative role) or afterwards (which would simply indicate a correlation). Finally, the study relies on patient reported information and patients' interpretation of lifestyle factor associated questions which can sometimes be less reliable.

Nevertheless, the results from this unique Sri Lankan study are in keeping with more detailed prospective studies from around the world, suggesting that the findings are likely to be valid. Details on the phenotypes of the PD patients in this study were not available, but considering the previously observed phenotypic differences in South Asian PD patients [4], this study may provide some indication that the observed lifestyle factor associations have global, biological relevance to all phenotypes of Idiopathic PD. However, more detailed, larger studies will be required to confirm the observed associations between PD, coffee intake and smoking, as well as to further probe other potential associations (eg. with tea intake) in Sri Lanka. Investigation of

the biological mechanisms underlying these associations will be valuable in identifying potential therapeutic and preventative strategies.

## Acknowledgements

The authors thank all the participants in this study. The authors also thank Dr. R.P.P. Dinalee for her contribution to data collection.

## Financial support

University of Sri Jayewardenepura and The World Health Organisation, Sri Lanka.

NIHR Cambridge Biomedical Research Centre and MRC-Wellcome Trust Cambridge Stem Cell Institute, UK.

The authors declare that there are no conflicts of interest regarding the publication of this paper.

## Abbreviations

PD: Parkinson's Disease; OR: Odds Ratio; CI: Confidence Interval; SD: Standard Deviation

## References

1. de Lau LML, Breteler MMB. Epidemiology of Parkinson's disease. *Lancet Neurol* 2006; 5:525–35. doi:10.1016/S1474-4422(06)70471-9
2. Wirdefeldt K, Adami H-O, Cole P, et al. Epidemiology and etiology of Parkinson's disease: a review of the evidence. *Eur J Epidemiol* 2011; 26 Suppl 1:S1–58. doi:10.1007/s10654-011-9581-6
3. Tan EK, Tan C, Fook-Chong SMC, et al. Dose-dependent protective effect of coffee, tea, and smoking in Parkinson's disease: A study in ethnic Chinese. *J Neurol Sci* 2003; 216:163–7. doi: 10.1016/j.jns.2003.07.006
4. Hu MTM, Chaudhuri KR, Jarosz J, et al. An imaging study of parkinsonism among African-Caribbean and Indian London communities. *Mov Disord* 2002; 17:1321-8. doi: 10.1002/mds.10261
5. Behari M, Srivastava AK, Das RR, et al. Risk factors of Parkinson's disease in Indian patients. *J Neurol Sci* 2001; 190:49-55. <http://www.sciencedirect.com/science/article/pii/S0022510X01005780>
6. Das K, Ghosh M, Nag C, et al. Role of familial, environmental and occupational factors in the development of Parkinson's disease. *Neurodegener Dis* 2011; 8:345-51. doi:10.1159/000323797
7. Wijesinghe P, Shankar SK, Chickabasaviah YT, et al. Cytoskeletal Pathologies of Age-Related Diseases between Elderly Sri Lankan (Colombo) and Indian (Bangalore) Brain Samples. *Curr Alzheimer Res* 2016; 13:268-80.
8. Postuma RB, Lang AE, Munhoz RP, et al. Caffeine for treatment of Parkinson disease: a randomized controlled trial. *Neurology* 2012; 79:651-8. doi: 10.1212/WNL.0b013e318263570d
9. Schwarzschild MA, Xu K, Oztas E, et al. Neuroprotection by caffeine and more specific A2A receptor antagonists in animal models of Parkinson's disease. *Neurology* 2003; 61:555-61. doi: 10.1212/01.WNL.0000095214.53646.72
10. Cao C, Loewenstein D a., Lin X, et al. High blood caffeine levels in mci linked to lack of progression to dementia. *J Alzheimer's Dis* 2012; 30:559-72. doi: 10.3233/JAD-2012-111781
11. Derkinderen P, Shannon KM, Brundin P. Gut feelings about smoking and coffee in Parkinson's disease. *Mov Disord* 2014; 29:976-9. doi: 10.1002/mds.25882
12. Quik M, Perez X a., Bordia T. Nicotine as a potential neuroprotective agent for Parkinson's disease. *Mov Disord* 2012; 27:947-57. doi: 10.1002/mds.25028
13. Ritz B, Lee P-C, Lassen CF, et al. Parkinson disease and smoking revisited Ease of quitting is an early sign of the disease. *Neurology* 2014; 1396-402. doi: 10.1212/WNL.0000000000000879.

### Publish in International Archives of Medicine

International Archives of Medicine is an open access journal publishing articles encompassing all aspects of medical science and clinical practice. IAM is considered a megajournal with independent sections on all areas of medicine. IAM is a really international journal with authors and board members from all around the world. The journal is widely indexed and classified Q2 in category Medicine.