

Trends in Parasitology

Helminth microbiomes – a hidden treasure trove?

--Manuscript Draft--

Manuscript Number:	TREPAR-D-18-00201R1
Article Type:	Opinion
Keywords:	macrobiota; microbiota; helminth; nematode; host-parasite interactions
Corresponding Author:	Cinzia Cantacessi Cambridge, UNITED KINGDOM
First Author:	T.P. Jenkins
Order of Authors:	T.P. Jenkins P.J. Brindley R.B. Gasser Cinzia Cantacessi
Abstract:	<p>There is increasing attention on the complex interactions occurring between gastrointestinal parasitic helminths and the microbial flora (microbiota) inhabiting the host gut. However, little is known about the occurrence, structure and function of microbial populations residing within parasite organs and tissues, referred to hereafter as the 'parabiome'. In this article, we argue that an in depth understanding of the parasite-parabiome interplay may significantly enhance current understanding of parasite biology and physiology, and may lead to the discovery of entirely novel, anthelmintic-independent interventions against parasites and parasitic diseases.</p>



28 October 2018

Dear Doctor Kong,

REJOINER: MS. No. (TREPAR-D-18-00201)

Entitled 'Helminth Microbiomes - a Hidden Treasure Trove?' *Trends in Parasitology*.

We are very grateful for the reviewers' and editor's time and efforts in evaluating this manuscript. The points raised by the reviewers are very constructive and have contributed to enhance the manuscript. We have provided detailed responses to individual points in the following rejoinder:

Reviewer #1:

I am not sure that Figure 1 is absolutely necessary. Perhaps a figure depicting the selective acquisition of bacteria by *Trichuris*- and then how these bacteria and murine host bacteria then aid the parasite may be of greater interest?

RESPONSE: The authors thank the reviewer for this comment and have provided an additional figure depicting the proposed microbiome acquisition strategies of *Trichuris muris*, *Haemonchus contortus*, and *Brugia malayi*. The focus of this figure lies on acquisition rather than mechanisms of gut microbial benefits, since the latter remains highly speculative.

Reading this article made me think of "microbes" other than bacteria that may inhabit parasites- i.e. viruses and fungi- and how these affect parasite development/host responses. Given it is an opinion article, could the authors speculate this field as one that will emerge in future? I am aware of some fungi that negatively affect plant nematodes. With the new technologies for culturing and sequencing it may be pertinent to identify more than just bacterial species?

RESPONSE: The authors thank the reviewer for the comment and agree that other microbes could be of importance and should be investigated alongside bacteria. Following the reviewer's suggestion, an additional "outstanding question" has been added and discussed in the "Concluding Remarks" section (lines 251-252).

Reviewer #2:

My only comment/concern is with the term 'parabiome'. Although it is defined within the text, I don't know that it is essential. My original interpretation of the term was that it referred to the parasitic component of a host microbiome (similar to virome, mycobiome referring to viral and fungal components). When I did an informal poll of colleagues and trainees to see what they thought a 'parabiome' referred to, the responses were either as above, or that it could refer to the microbiome of mice that undergone parabiosis. Overall, I think it would be clearer to

continue to refer to the 'helminthic microbiome' as laid out in the title, and interspersed throughout the text.

RESPONSE: The authors thank the reviewer for this constructive comment and agree that the term 'parabiome' may raise some confusion. Therefore, following the reviewer's suggestion, we have replaced it with "helminth microbiome".

In conclusion, we consider that we have rigorously addressed each of the points raised by each of the reviewers. We agree with the points raised, and modified the text accordingly. These comments have led to an enhanced manuscript, which we consider to meet the standard for publication in *Trends in Parasitology*.

Yours sincerely,
Cinzia Cantacessi
On behalf of all authors.

Highlights

- In spite of a plethora of evidence supporting key roles of resident bacteria for parasite fitness and survival (i.e. in filarial nematodes and whipworms), little is known of microbial populations inhabiting organs and tissues of gastrointestinal nematodes of major socio-economic significance.
- To date, one of the few well-characterised examples is the symbiotic relationship between bacteria of the genus *Wolbachia* and filarial nematodes.
- Notably, our understanding of this symbiosis has already been successfully exploited for therapeutic purposes.
- The relentless advancement in sequencing techniques, together with the rise of novel microbiome editing tools provide unprecedented opportunities to investigate helminth parabiomes and exploit parasite-parabiome relationships for the development of novel strategies of parasite control.

1 Helminth microbiomes – a hidden treasure trove?

2
3 T.P. Jenkins,¹ P.J. Brindley,² R.B. Gasser,³ and C. Cantacessi¹

4 ¹ Department of Veterinary Medicine, University of Cambridge, Cambridge, CB3 0ES, UK

5 ² Department of Microbiology, Immunology & Tropical Medicine, and Research Center for
6 Neglected Diseases of Poverty, School of Medicine & Health Sciences, George Washington
7 University, Washington DC, 20037, USA

8 ³ Department of Veterinary Biosciences, Melbourne Veterinary School, Faculty of Veterinary
9 and Agricultural Sciences, The University of Melbourne, Parkville, Victoria 3010, Australia

10 * Correspondence: cc779@cam.ac.uk (C. Cantacessi)

11
12 **Keywords:** macrobiota, microbiota, helminth, nematode, host-parasite interactions

13 There is increasing attention on the complex interactions occurring between gastrointestinal
14 parasitic helminths and the microbial flora (microbiota) inhabiting the host gut. However, little
15 is known about the occurrence, structure and function of microbial populations residing within
16 parasite organs and tissues. In this article, we argue that an in depth understanding of the
17 interplay between parasites and their microbiomes may significantly enhance current
18 knowledge of parasite biology and physiology, and may lead to the discovery of entirely novel,
19 anthelmintic-independent interventions against parasites and parasitic diseases.

20 **Gastrointestinal helminths – worm guts within host guts**

21 The gastrointestinal (GI) tract of vertebrates is inhabited by 10–100 trillion microorganisms,
22 including bacteria, viruses, fungi, protists and archaea, which are collectively known as the
23 ‘gut **microbiota**’ (see Glossary) [1, 2]. This complex ecosystem exerts a number of highly
24 specialised functions that are essential to host physiology, including the absorption of nutrients,
25 synthesis of essential organic compounds, development of adaptive immunity and protection
26 against pathogens [3-6]. In particular, amongst such pathogens are metazoan parasites (called
27 ‘the **macrobiota**’, which includes parasitic nematodes such as roundworms, hookworms and
28 whipworms) that are traditionally considered detrimental to the vertebrate host, as they can
29 subtract nutrients, damage host tissues and release toxic waste products (reviewed by [7]).
30 These parasites live in close association with the vertebrate microbiota and, over the last few
31 years, evidence has started to emerge of the existence of complex **mutualistic** relationships
32 between these two players that might facilitate the long-term establishment of nematodes in
33 the gut of the host [8-11]. Similar to their vertebrate hosts, parasitic nematodes have complete,
34 tubular digestive systems responsible for nutrient uptake, processing and absorption; yet, for
35 most species of socio-economically important parasites, and GI nematodes in particular, little
36 is known about the occurrence, structure and function of populations of resident gut microbes
37 [12]. Nevertheless, critical evidence of the existence of essential **symbiotic** relationships
38 between parasites and bacteria is provided by filarial nematodes, e.g., *Onchocerca volvulus*
39 (causing river blindness) and *Wuchereria bancrofti* and *Brugia malayi* (causing lymphatic
40 filariasis) (reviewed by [13]), whose propagation and survival are dependent on a genus of
41 bacteria, i.e. *Wolbachia*, which has become the target of intense investigations aimed to
42 develop novel filaricidal compounds [14-18]. This evidence supports the hypothesis, strongly
43 corroborated by recent experimental findings [19, 20], that the digestive system as well as other
44 organs and tissues of GI nematodes may also harbour resident microbes with essential roles in

45 parasite physiology and survival. Fully characterizing and understanding the structure and
46 function of **helminth microbiomes**, and determining the role/s they play in key aspects of
47 parasite biology and host-parasite interactions, could not only have broad implications for
48 future studies of the origin of parasitism itself, but might also lead to the discovery of radically
49 new interventions against these worms.

50 Here, we (i) summarize current knowledge of the origin, structure and function of helminth
51 microbiomes; (ii) identify some key knowledge gaps for future research in this field; and (iii)
52 discuss the potential therapeutic advances that such knowledge could deliver, with a particular
53 focus on the development of novel, helminth microbiome-based strategies for the control of
54 infections by GI nematodes of major socio-economic significance.

55

56 **Strategies of acquisition and maintenance of helminth microbiomes**

57 Investigations of the interactions between parasites and their resident bacteria rely on a
58 thorough understanding of the dynamics of microbiome acquisition. Whilst for GI nematodes
59 knowledge in this area is relatively limited, several investigations have documented the fine
60 strategy *via* which bacteria of the genus *Wolbachia* are propagated through successive
61 generations of filarial parasites (reviewed by [21]) (cf. Figure 1). In the filarial nematode *B.*
62 *malayi*, these bacteria inhabit the lateral chords of both adult male and female worms, and the
63 reproductive system of the latter sex, where they colonise the ovaries, oocytes and early
64 embryos within the uteri [22]. Upon egg fertilisation, populations of *Wolbachia* segregate
65 asymmetrically in the developing embryo, which results in an uneven distribution of these
66 bacteria in the tissues of the resultant microfilariae [22]. In particular, the numbers of vertically
67 transmitted *Wolbachia* remain stable (~70 per embryo [17]) throughout development of the
68 new generation of filarial parasites into infective third-stage larvae (L3s) in the mosquito vector

69 (reviewed by [21]). Upon L3-invasion of a new, susceptible vertebrate host, the number of
70 *Wolbachia* bacteria rapidly increases in the hypodermal cord of developing worms, with a
71 further expansion occurring in the reproductive tissues of sexually mature females (reviewed
72 by [21]). Crucially, embryonic development is entirely dependent on *Wolbachia*, as treatment
73 with tetracycline antibiotics results in a marked reduction of viable microfilariae (reviewed by
74 [22]). To daeujte, the **obligate relationship** between filarial nematodes and *Wolbachia*
75 represents the only known example of a mutualistic association between parasitic nematodes
76 and bacteria.

77 Members of another group of helminth parasites, the digenean trematodes, are known to
78 harbour populations of *Neorickettsia* **endosymbionts** which share numerous genetic
79 similarities with *Wolbachia* (reviewed by [23]). *Neorickettsia* inhabit a range of environments
80 suitable for the development of the infective stages of digenean parasites and their intermediate
81 hosts (e.g., aquatic molluscs), thus lending credit to the hypothesis that a proportion of these
82 bacteria are horizontally transmitted (reviewed by [23, 24]). Nevertheless, in *Plagiorchis*
83 *elegans*, a common GI helminth of a range of fishes, birds and mammals (including humans)
84 [25], *Neorickettsia* is predominantly transmitted vertically across generations of parasites [26].
85 However, unlike *Wolbachia* in filarial nematodes, transmission of *Neorickettsia* in *P. elegans*
86 occurs also through the asexual stages of this parasite [27]. Furthermore, since transmission
87 rates of *Neorickettsia* from adult *P. elegans* to the offspring vary from 11% to 91% [27], it has
88 been suggested that the life cycle of this flatworm is not dependent on their neorickettsial
89 endosymbionts, but rather that *Neorickettsia* utilises *P. elegans* as a vehicle for transmission to
90 vertebrate hosts [28].

91 For GI nematodes, experimental evidence of microbiome acquisition strategies is available for
92 two species, namely *Haemonchus contortus*, an abomasal roundworm of small ruminants [19]
93 and *Trichuris muris*, a large intestinal whipworm of rodents [20] (cf. Figure 1). For the former,

94 a recent study [19] localised selected genera of bacteria (i.e. *Weissella* and *Leuconostoc*) to the
95 gut of adult worms and to the uterus of sexually mature females by fluorescence *in situ*
96 hybridization (FISH) and transmission electron microscopy; using DNA fingerprinting, the
97 same genera could be identified in eggs laid by these females and, following larval culture,
98 their L3 offspring [19]. Notably, these microorganisms could not be identified in the faecal
99 matter on which larval culture was performed, thus providing evidence of maternal
100 transmission of these bacteria [19]. Other bacterial genera (i.e. *Lactococcus* and *Streptococcus*)
101 could be identified in the distal uterus of sexually mature females of *H. contortus* [19].
102 Nevertheless, these bacteria were not detected in newly deposited eggs and developed L3s.
103 Since these bacteria occur in the rumen of the host [29], the authors hypothesized that female
104 worms acquired them by ingesting ruminal fluid [19]. Whilst the roles that species of
105 *Lactococcus* and *Streptococcus* might play in the fundamental biology of *H. contortus* is
106 presently unknown, these data suggest that this parasite might employ a ‘hybrid’ microbiome
107 acquisition strategy, with some ‘core’ endosymbionts (i.e. *Weissella* and *Leuconostoc*), which
108 may play essential roles in parasite fitness and survival, being vertically transmitted, and others
109 (i.e. *Lactococcus* and *Streptococcus*) being acquired from the host to underpin a certain level
110 of microbiome plasticity and capacity for environmental adaptation.

111 A clear strategy of microbiome acquisition from the mammalian host has been recently
112 demonstrated for *T. muris* [20] (cf. Figure 1). The essential role that the host microbiome plays
113 in the development and propagation of this parasite had already been demonstrated in a seminal
114 study [30], which showed that parasite egg hatching in the large intestine of the mouse was
115 dependent on the microbial flora within the host gut. Recently, a study by the same research
116 group [20] demonstrated that, following egg hatching, *T. muris* acquires populations of bacteria
117 that, together, form a ‘core’ nematode microbiome, which is markedly distinct from the
118 microbiome inhabiting the environment in which the worms reside [20]. Although the

119 *Trichuris* microbiome described predominantly comprised Firmicutes and Bacteroidetes,
120 similar to the host microbiome, it was also rich in Proteobacteria. Proteobacteria constituted
121 9% of the entire *T. muris* microbiome – a 31-fold and 13-fold increase in relative proportions
122 of this bacterial group compared with the microbiome of uninfected mice and *T. muris*-infected
123 mice, respectively [20]. This process of selective microbiome acquisition was demonstrated to
124 be independent from the initial host microbiome composition, and the administration of broad-
125 spectrum antibiotics to adult *T. muris ex vivo* resulted in a marked decrease in parasite fitness
126 and survival rates, thus providing cogent evidence for an essential role of the host-acquired
127 microbiome for the successful completion of the whipworm life cycle. Whether *T. muris*
128 acquires its own microbiome passively, or actively selects populations of bacteria with
129 functional properties which are able to facilitate its survival in the vertebrate host, remains to
130 be established (cf. [20]). Nevertheless, the observation that mono-colonisation of germ-free
131 mice with a single species of bacterium, i.e. *Bacteroides thetaiotaomicron*, resulted in
132 successful egg hatching and establishment of chronic *T. muris* infection [20] provides a unique
133 opportunity to design targeted experiments that can shed light on the precise mechanisms of
134 acquisition of the *T. muris* microbiome. In turn, this knowledge will form the necessary basis
135 to answer fundamental questions regarding helminth microbiome structure and function.

136

137 **Are helminth microbiomes key to parasite fitness?**

138 Studies of the structure and function of helminth microbiomes are in their infancy.
139 Nevertheless, over the past decades, evidence has emerged about the functional association
140 between the free-living nematode *Caenorhabditis elegans* and the bacteria inhabiting it [31,
141 32]. Indeed, *C. elegans* is known to host a species-rich bacterial community, dominated by
142 Proteobacteria, such as *Enterobacteriaceae* and members of the genera *Pseudomonas*,
143 *Stenotrophomonas*, *Ochrobactrum*, and *Sphingomonas* [31, 32]. Crucially, the relative

144 proportions of bacterial populations forming the *C. elegans* microbiome vary according to the
145 developmental stage of this nematode [32], thus suggesting that worm development relies on a
146 range of bacterial functions that differ over time. In support of this hypothesis, worms
147 experimentally colonised with a subset of bacterial isolates representing the *C. elegans* ‘core’
148 **microbiome** displayed increased fitness and survival rates, and were maintained under
149 stressful conditions of temperature and osmolarity, compared with worms colonised solely by
150 *Escherichia coli* [32]. In addition, compared with *E. coli*-colonised *C. elegans*, worms fed with
151 the soil bacterium *Comamonas* displayed accelerated development, which was attributed to the
152 ability of this bacterial group to up-regulate the expression of genes associated with the
153 nematode’s moulting program [33]. The *C. elegans* microbiome has also been demonstrated to
154 play important roles in worm defence against pathogens; indeed, *Pseudomonas* isolates
155 detected amongst the worm resident populations of bacteria produce anti-mycotic compounds
156 that prevent colonisation by fungal agents [32].

157 Evidence from investigations of *C. elegans*, employed as a model for nematode-microbiome
158 interactions, points to a likely functional role of the microbiomes of parasitic helminths for
159 worm physiology, development and survival. However, the parasite microbiome itself may
160 benefit from the protected and nutrient-rich environment that the worm host offers [34, 35].
161 For instance, the Gammaproteobacteria *Photorhabdus* and *Xenorhabdus*, that inhabit the gut
162 of the entomopathogenic nematodes *Heterorhabditis* and *Steinernema*, are released upon
163 infection of the insect host by the infective juveniles; following their release, these bacteria
164 actively replicate and kill the insect host, while converting the insect carcass into a source of
165 nutrients to support nematode growth and development [36, 37].

166 For parasitic nematodes of medical and veterinary importance, the mutualistic association
167 between filarial nematodes and *Wolbachia* offers a key example of the fundamental functions
168 that the helminth microbiome exerts in the biology of its worm host, and *vice versa*. Indeed,

169 besides its known role in the development and survival of filarial embryos (reviewed by [21]),
170 *Wolbachia* is essential for worm nutrition and metabolism. The bacterium synthesizes haem,
171 riboflavin (vitamin B₂), and flavin adenine dinucleotide, which the parasite host is unable to
172 synthesize and that have been inferred to play an important role in filarial reproduction and
173 development, as well as nucleotides, which are required during oogenesis and embryogenesis
174 [18, 38, 39]. In addition, members of the genus *Wolbachia* participate in pathways aimed at
175 preventing apoptosis of filarial reproductive, embryonic and somatic cells [17, 40], likely by
176 the direct targeting of the apoptotic signalling cascade [41-43]. Finally, in the filarial parasite
177 of cattle *Onchocerca ochengi*, *Wolbachia* has been demonstrated to play a key role in host
178 immune evasion, specifically by attracting host neutrophils and, thus, averting a potentially
179 lethal effector response by degranulating eosinophils [44].

180 Over the years, the fundamental roles that *Wolbachia* play in pathways linked to reproduction,
181 metabolism and immune defence of filarial nematodes have been the subject of intense
182 scrutiny, focusing on developing novel chemotherapeutics to disrupt this mutualistic
183 relationship; some have been successful [15, 17, 21, 45]. For instance, the administration of 4-
184 week courses of doxycycline (belonging to the tetracycline family of antibiotics) and
185 rifampicin have been deemed effective in reducing the transmission of *O. volvulus*
186 microfilariae to mosquito intermediate hosts and filarial embryogenesis, respectively
187 (reviewed by [13]). Nevertheless, the length of drug administration required to achieve
188 significant effects, along with the severe adverse reactions that tetracyclines can cause in
189 children and pregnant women (e.g. permanent dental staining, teratogenic effects and
190 potentially fatal hepatotoxicity [46]), limit the use of these antibiotics in mass drug
191 administration (MDA) programs in areas where filariases are endemic. Nevertheless, these
192 findings raise the question as to whether a deep exploration of the microbiomes of other
193 helminth parasites of major socio-economic significance could hold promise for the

194 identification of novel targets for the development of antibiotic-independent control strategies
195 against the diseases caused by these worms.

196

197 **A new generation of helminth microbiome-targeting chemotherapeutics?**

198 Globally, more than two billion people are at risk of infection by GI nematodes, mainly the
199 hookworms *Ancylostoma duodenale* and *Necator americanus*, the whipworm *T. trichiura*, the
200 roundworm *Ascaris lumbricoides*, and the threadworm, *Strongyloides stercoralis*.
201 Collectively, these nematodes are responsible for more than 5.5 million disability-adjusted life
202 years (DALYs) (reviewed by [47]). Moreover, GI nematodes inflict significant production
203 losses in livestock due to the extensive morbidity and mortality associated with a range of
204 diseases that they cause (reviewed by [48]). Complete reliance on anthelmintics for the control
205 of these parasites (*via* MDA or targeted strategic worming programmes in humans and
206 livestock) bears substantial risks, linked to the global threat of emerging anthelmintic
207 resistance, as already observed in several GI nematodes of veterinary importance (reviewed by
208 [49, 50]). Yet, the discovery of alternative strategies for parasite control should be built on a
209 thorough understanding of the fundamental biology of these pathogens, and of key mechanisms
210 of interactions with their vertebrate hosts. A deeper knowledge of the structure and function of
211 the microbiomes of parasitic helminths, and of mechanisms of microbiome acquisition and
212 transmission, could lead to unprecedented discoveries in parasite physiology, pathology and
213 reproduction, and thus, to the development of completely novel control tools. Nevertheless, for
214 such discoveries to be harnessed, fundamental information needs to be acquired. We propose
215 that, in the first instance, the microbiomes of representative species of GI nematodes of
216 considerable medical and veterinary significance (Figure 2) could be qualitatively
217 characterized using high-throughput sequencing of the bacterial 16S rRNA gene (Box 1). The
218 selection of specimens of a range of parasite species from different hosts and geographical

219 locations would assist the determination of species-specific ‘core’ parasite microbiomes.
220 Following the establishment of reference 16S rRNA databases for each key parasite species,
221 shotgun metagenomic sequencing of their microbiomes would provide important information
222 on the relative abundance of each ‘core’ microbial species, and clues about their functional
223 potential. The latter, coupled to investigations of the proteomes and metabolomes of the
224 microbial communities inhabiting these parasites, could lead to a better understanding of the
225 possible role/s that the microbiomes of parasitic helminths play in the biology and physiology
226 of individual worms.

227 Key information on the modes of transmission of helminth microbiomes could be acquired *via*
228 experimental infections of vertebrate hosts with selected GI nematodes, followed by qualitative
229 and quantitative comparative analyses of the host microbiomes and key parasite developmental
230 stages. Furthermore, for selected GI nematodes (e.g. *Nippostrongylus brasiliensis*),
231 experimental infections of germ-free or antibiotic-treated mice re-colonised with fluorescently
232 labelled bacteria might provide clues on host-parasite microbiome transfer using *in vivo*
233 imaging (cf. [51]). Similar techniques could be used to localise species or groups of bacteria
234 in parasite organs and tissues, thus providing additional clues on the functions of such
235 microorganisms in worm biology. Together, this information would form a basis for
236 experimentation, aimed at interfering with such functions that may potentially lead to the
237 discovery of entirely novel, antibiotic-independent strategies for parasite control (Figure 3), for
238 example, *via* cutting-edge **microbiome editing** techniques including CRISPR/Cas9,
239 engineered probiotics, and/or bactericidal bacteriophages (reviewed by [52]).

240

241 **Concluding Remarks**

242 In spite of substantial evidence that points towards key role(s) of microbial species inhabiting
243 parasitic helminths in the fundamental biology of these pathogens and host-parasite
244 interactions, e.g. in filarial nematodes and, more recently, whipworms, current knowledge of
245 the microbiomes of key parasites of major socio-economic significance, such as GI nematodes
246 of humans and livestock, is scarce and fragmented. Nonetheless, the relentless progress in
247 microbiome investigation and editing technologies (Box 1), and novel high-throughput
248 bioinformatics pipelines, provides us with unprecedented opportunities to thoroughly
249 characterize the structures and functions of such microbial populations. At the core lie
250 questions surrounding modes of helminth microbiome acquisition and propagation to
251 successive generations of parasites, the localisation of endosymbiont microorganisms in the
252 organs and tissues of parasites, the functions that helminth microbiomes (including bacteria,
253 viruses and fungi) play in parasite biology and physiology, and the effects that disrupting
254 parasite-microbiome interactions may exert on parasite propagation and survival (see
255 Outstanding Questions). In turn, the new knowledge can be expected to provide us with a
256 plethora of opportunities to exploit parasite-microbiome associations to our advantage, for
257 example, by applying cutting-edge microbiome-editing techniques as novel intervention
258 strategies against parasitic nematodes and the diseases that they cause.

259

260 **Acknowledgements**

261 T.P.J. is the grateful recipient of a PhD scholarship by the Biotechnology and Biological
262 Sciences Research Council (BBSRC) of the United Kingdom. Research in the C.C. laboratory
263 is funded by grants by the Isaac Newton Trust, the Isaac Newton Trust/Wellcome Trust/
264 University of Cambridge joint grant scheme and by the Royal Society (UK). Research in the
265 R.B.G. laboratory is supported predominantly by the Australian Research Council (ARC), the

- 266 National Health and Medical Research Council (NHMRC), Melbourne Water Corporation and
267 Yourgene Bioscience.

268 **References**

- 269 1 Rajilic-Stojanovic, M. and de Vos, W.M. (2014) The first 1000 cultured species of the
270 human gastrointestinal microbiota. *FEMS Microbiol Rev* 38, 996-1047
- 271 2 Turnbaugh, P.J., *et al.* (2007) The human microbiome project. *Nature* 449, 804-810
- 272 3 Brestoff, J.R. and Artis, D. (2013) Commensal bacteria at the interface of host metabolism
273 and the immune system. *Nat Immunol* 14, 676-684
- 274 4 Brown, E.M., *et al.* (2013) The role of the immune system in governing host-microbe
275 interactions in the intestine. *Nat Immunol* 14, 660-667
- 276 5 Hooper, L.V., *et al.* (2012) Interactions between the microbiota and the immune system.
277 *Science* 336, 1268-1273
- 278 6 Tremaroli, V. and Backhed, F. (2012) Functional interactions between the gut microbiota
279 and host metabolism. *Nature* 489, 242-249
- 280 7 McCarty, T.R., *et al.* (2014) Global progress towards eliminating gastrointestinal helminth
281 infections. *Curr Opin Gastroenterol* 30, 18-24
- 282 8 Bancroft, A.J., *et al.* (2012) Life on the edge: the balance between macrofauna, microflora
283 and host immunity. *Trends Parasitol* 28, 93-98
- 284 9 Glendinning, L., *et al.* (2014) The microbiota and helminths: sharing the same niche in the
285 human host. *Parasitology* 141, 1255-1271
- 286 10 Reynolds, L.A., *et al.* (2015) Cohabitation in the intestine: Interactions among helminth
287 parasites, bacterial microbiota, and host immunity. *J Immunol* 195, 4059-4066
- 288 11 Kreisinger, J., *et al.* (2015) Interactions between multiple helminths and the gut
289 microbiota in wild rodents. *Philos Trans R Soc Lond B Biol Sci* 370, 20140295
- 290 12 Dheilly, N.M., *et al.* (2017) Parasite microbiome project: Systematic investigation of
291 microbiome dynamics within and across parasite-host interactions. *Msystems* 2, e00050-
292 e000517
293

- 294 13 Bouchery, T., *et al.* (2013) The symbiotic role of *Wolbachia* in Onchocercidae and its
295 impact on filariasis. *Clin Microbiol Infect* 19, 131-140
- 296 14 Hoerauf, A., *et al.* (2003) Doxycycline as a novel strategy against bancroftian filariasis—
297 depletion of *Wolbachia* endosymbionts from *Wuchereria bancrofti* and stop of microfilaria
298 production. *Med Microbiol Immunol* 192, 211-216
- 299 15 Johnston, K.L., *et al.* (2017) Identification and prioritization of novel anti-*Wolbachia*
300 chemotypes from screening a 10,000-compound diversity library. *Sci Adv* 3, eaao1551
- 301 16 Johnston, K.L., *et al.* (2014) Repurposing of approved drugs from the human
302 pharmacopoeia to target *Wolbachia* endosymbionts of onchocerciasis and lymphatic
303 filariasis. *Int J Parasitol Drugs Drug Resist* 4, 278-286
- 304 17 Landmann, F., *et al.* (2011) Anti-filarial activity of antibiotic therapy is due to extensive
305 apoptosis after *Wolbachia* depletion from filarial nematodes. *PLoS Pathog* 7, e1002351
- 306 18 Slatko, B.E., *et al.* (2010) The *Wolbachia* endosymbiont as an anti-filarial nematode
307 target. *Symbiosis* 51, 55-65
- 308 19 Sinnathamby, G., *et al.* (2018) The bacterial community associated with the sheep
309 gastrointestinal nematode parasite *Haemonchus contortus*. *PLoS One* 13, e0192164
- 310 20 White, E.C., *et al.* (2018) Manipulation of host and parasite microbiotas: Survival
311 strategies during chronic nematode infection. *Sci Adv* 4, eaap7399
- 312 21 Slatko, B.E., *et al.* (2014) *Wolbachia* endosymbionts and human disease control. *Mol*
313 *Biochem Parasitol* 195, 88-95
- 314 22 Landmann, F., *et al.* (2010) Asymmetric *Wolbachia* segregation during early *Brugia*
315 *malayi* embryogenesis determines its distribution in adult host tissues. *PLoS Negl Trop Dis* 4,
316 e758
- 317 23 Tkach, V.V. and Greiman, S.E. (2016) Transmission biology, host associations,
318 distribution and molecular diagnostics of *Neorickettsia*. In *Rickettsiales: Biology, Molecular*
319 *Biology, Epidemiology, and Vaccine Development* (Thomas, S., ed), pp. 295-325, Springer
320 International Publishing

321 24 McNulty, S.N., *et al.* (2017) Genomes of *Fasciola hepatica* from the Americas reveal
322 colonization with *Neorickettsia* endobacteria related to the agents of potomac horse and
323 human sennetsu fevers. *PLoS Genet* 13, e1006537

324 25 Tkach, V.V., *et al.* (2008) Family Plagiorchiidae. In *Keys to the Trematoda* (Bray, R.A., *et*
325 *al.* eds), pp. 295–325, CABI

326 26 Vaughan, J.A., *et al.* (2012) Neorickettsial endosymbionts of the digenea: diversity,
327 transmission and distribution. *Adv Parasitol* 79, pp. 253-297

328 27 Greiman, S.E., *et al.* (2013) Transmission rates of the bacterial endosymbiont,
329 *Neorickettsia risticii*, during the asexual reproduction phase of its digenean host, *Plagiorchis*
330 *elegans*, within naturally infected lymnaeid snails. *Parasit Vectors* 6, 303

331 28 Greiman, S.E., *et al.* (2014) Large scale screening of digeneans for *Neorickettsia*
332 endosymbionts using real-time PCR reveals new *Neorickettsia* genotypes, host associations
333 and geographic records. *PLoS One* 9, e98453

334 29 De Barbieri, I., *et al.* (2015) Production attributes of Merino sheep genetically divergent
335 for wool growth are reflected in differing rumen microbiotas. *Livestock Science* 178, 119-129

336 30 Hayes, K.S., *et al.* (2010) Exploitation of the intestinal microflora by the parasitic
337 nematode *Trichuris muris*. *Science* 328, 1391-1394

338 31 Clark, L.C. and Hodgkin, J. (2014) Commensals, probiotics and pathogens in the
339 *Caenorhabditis elegans* model. *Cell Microbiol* 16, 27-38

340 32 Dirksen, P., *et al.* (2016) The native microbiome of the nematode *Caenorhabditis elegans*:
341 gateway to a new host-microbiome model. *BMC Biol* 14, 38

342 33 MacNeil, L.T., *et al.* (2013) Diet-induced developmental acceleration independent of TOR
343 and insulin in *C. elegans*. *Cell* 153, 240-252

344 34 Sansonetti, P.J. and Medzhitov, R. (2009) Learning tolerance while fighting ignorance.
345 *Cell* 138, 416-420

346 35 Zeng, Q., *et al.* (2017) Models of microbiome evolution incorporating host and microbial
347 selection. *Microbiome* 5, 127

348 36 Goodrich-Blair, H. and Clarke, D.J. (2007) Mutualism and pathogenesis in *Xenorhabdus*
349 and *Photorhabdus*: two roads to the same destination. *Mol Microbiol* 64, 260-268

350 37 Clarke, D.J. (2008) *Photorhabdus*: a model for the analysis of pathogenicity and
351 mutualism. *Cell Microbiol* 10, 2159-2167

352 38 Foster, J., *et al.* (2005) The *Wolbachia* genome of *Brugia malayi*: Endosymbiont evolution
353 within a human pathogenic nematode. *PLoS Biol* 3, 599-614

354 39 Taylor, M.J., *et al.* (2005) *Wolbachia* bacterial endosymbionts of filarial nematodes. *Adv*
355 *Parasitol* 60, 245-284

356 40 Foray, V., *et al.* (2018) *Wolbachia* control stem cell behavior and stimulate germline
357 proliferation in filarial nematodes. *Dev Cell* 45, 198-211 e193

358 41 Faherty, C.S. and Maurelli, A.T. (2008) Staying alive: bacterial inhibition of apoptosis
359 during infection. *Trends Microbiol* 16, 173-180

360 42 James, E.R. and Green, D.R. (2004) Manipulation of apoptosis in the host-parasite
361 interaction. *Trends Parasitol* 20, 280-287

362 43 Hacker, G. and Fischer, S.F. (2002) Bacterial anti-apoptotic activities. *FEMS Microbiol*
363 *Lett* 211, 1-6

364 44 Hansen, R.D.E., *et al.* (2011) A worm's best friend: recruitment of neutrophils by
365 *Wolbachia* confounds eosinophil degranulation against the filarial nematode *Onchocerca*
366 *ochengi*. *Proc Biol Sci* 278, 2293-2302

367 45 Turner, J.D., *et al.* (2017) Albendazole and antibiotics synergize to deliver short-course
368 anti-*Wolbachia* curative treatments in preclinical models of filariasis. *Proc Natl Acad Sci U S*
369 *A* 114, E9712-E9721

370 46 Aronson, J.K. (2015) Tetracyclines. In *Meyler's side effects of drugs: the international*
371 *encyclopedia of adverse drug reactions and interactions* (16th edn) (Aronson, J.K., ed), pp.
372 772-788, Elsevier

373 47 Jourdan, P.M., *et al.* (2018) Soil-transmitted helminth infections. *Lancet* 391, 252-265

374 48 Charlier, J., *et al.* (2014) Chasing helminths and their economic impact on farmed
375 ruminants. *Trends Parasitol* 30, 361-367

376 49 Kaplan, R.M. and Vidyashankar, A.N. (2012) An inconvenient truth: global warming and
377 anthelmintic resistance. *Vet Parasitol* 186, 70-78

378 50 Rose, H., *et al.* (2015) Widespread anthelmintic resistance in European farmed ruminants:
379 a systematic review. *Vet Rec* 176, 546

380 51 Geva-Zatorsky, N., *et al.* (2015) *In vivo* imaging and tracking of host-microbiota
381 interactions *via* metabolic labeling of gut anaerobic bacteria. *Nat Med* 21, 1091-1100

382 52 Sheth, R.U., *et al.* (2016) Manipulating bacterial communities by *in situ* microbiome
383 engineering. *Trends Genet* 32, 189-200

384 53 Franzosa, E.A., *et al.* (2015) Sequencing and beyond: integrating molecular 'omics' for
385 microbial community profiling. *Nat Rev Microbiol* 13, 360-372

386 54 Goodwin, S., *et al.* (2016) Coming of age: Ten years of next-generation sequencing
387 technologies. *Nat Rev Genet* 17, 333-351

388 55 Salter, S.J., *et al.* (2014) Reagent and laboratory contamination can critically impact
389 sequence-based microbiome analyses. *BMC Biol* 12, 87

390 56 Glassing, A., *et al.* (2016) Inherent bacterial DNA contamination of extraction and
391 sequencing reagents may affect interpretation of microbiota in low bacterial biomass
392 samples. *Gut Pathog* 8, 24

393 57 Howe, K.L., *et al.* (2017) WormBase ParaSite - a comprehensive resource for helminth
394 genomics. *Mol Biochem Parasitol* 215, 2-10

395 58 Roumpeka, D.D., *et al.* (2017) A review of bioinformatics tools for bio-prospecting from
396 metagenomic sequence data. *Front Genet* 8, 23

397 59 Tropini, C., *et al.* (2017) The gut microbiome: Connecting spatial organization to
398 function. *Cell Host Microbe* 21, 433-442

399 60 Parkinson, J., *et al.* (2004) A transcriptomic analysis of the phylum Nematoda. *Nat Genet*
400 36, 1259-1267

401 **Glossary**

402 **Core microbiome:** The group of microbes consistently found within a host microbiome, which
403 demonstrate a persistent association and provide a critical function within the habitat in which
404 they are detected.

405 **Endosymbionts:** Any organism that lives within the body or cells of another organism in a
406 symbiotic relationship with the host body or cell, often but not always to mutual benefit.

407 **Macrobiota:** Collective term used to describe the macroorganisms (e.g. helminths) that live in
408 a particular niche, on or in a living being, and are large enough to be seen with the naked eye.

409 **Microbiota:** Collective term used to describe the microorganisms that live in a particular niche,
410 on or in a living being, and exhibit a symbiotic relationship with the host.

411 **Microbiome editing:** Targeted manipulation of microbiota through the use of technology
412 and/or other microbiota.

413 **Mutualism:** A symbiosis which is beneficial to both organisms involved.

414 **Obligate relationship:** One or both of the symbionts entirely depend on each other for
415 survival.

416 **Helminth microbiome:** The collective microbiome of parasitic helminths.

417 **Symbiosis:** A relationship between two or more organisms that live closely together.

418 **Box 1: Sequencing and analysis of helminth microbiomes – a need for**
419 **standardisation**

420 The most commonly used techniques for microbiome profiling include high-throughput
421 sequencing of the bacterial 16S rRNA gene (often referred to as ‘16S rRNA sequencing’), and
422 whole genome sequencing of microbial communities (‘metagenome sequencing’) [53, 54].
423 Whilst these techniques are highly suitable for the study of vertebrate microbiomes, their
424 application to investigations of the microbiomes of helminth parasites, and GI nematodes in
425 particular, will require careful considerations surrounding sample processing, as well as data
426 analysis and interpretation. First, GI nematode specimens are usually directly harvested from
427 vertebrate hosts (e.g. experimentally infected animals), or cultured in host faecal matter, which
428 dramatically increases the risk of ‘contaminating’ microbial populations, for instance coating
429 the outer nematode cuticle, in the parasite DNA of interest. In order to overcome this potential
430 limitation, the cuticle of worms collected from infected hosts should be sterilized (e.g. using
431 sodium hypochlorite) prior to microbial DNA extraction. In addition, in order to generate
432 meaningful data on parasite microbiome structure and function, the determination of ‘core’
433 taxa, specific to a given parasite species and/or species group will be necessary. Importantly,
434 negative (‘no-DNA template’) samples must be processed alongside samples of interest, in
435 order to ensure that sequences generated from microorganisms contaminating laboratory
436 surfaces and equipment, glassware and plastic ware, as well as nucleic acid isolation and
437 amplification and sequencing reagents [55, 56] are correctly identified and subtracted from the
438 final sequence dataset(s). Whilst existing high throughput genome sequencing datasets
439 generated from a range of GI nematodes (particularly as part of whole genome sequencing
440 projects; [57]) might include useful sequence information on the occurrence of resident
441 microbial populations within parasite tissues (which, in such datasets, are traditionally
442 considered ‘contaminant sequences’ and therefore discarded), the absence of adequate controls

443 in these experiments largely prevents the retrospective use of these data in parasite
444 microbiome-sequencing studies. Once microbial sequence data have been obtained and
445 annotated [58], information on the localization of specific taxa of interest in parasite tissues
446 can be obtained using microscopy techniques, including fluorescence *in situ* hybridization (=
447 FISH), immunofluorescence, and transmission electron microscopy (TEM) (reviewed in [59]).
448 Whilst all of these techniques require highly specialized staff and are relatively costly and time
449 consuming to use, they might provide useful insights into means of helminth microbiome
450 acquisition and function.

451 **Figure 1. Proposed helminth microbiome acquisition strategies for *Brugia***
452 ***malayi*, *Trichuris muris*, and *Haemonchus contortus*.**

453 (A) *B. malayi* microfilariae acquire *Wolbachia* microorganisms via the female germ line, and
454 populations of resident microbes expand throughout larval development through to infective
455 third stage larvae in the mosquito intermediate host. In adult male and female *B. malayi*, the
456 bacteria localise to the lateral chords of both sexes and the female reproductive system, where
457 they colonise the ovaries, oocytes and early embryos within the uteri (B) Unembryonated *T.*
458 *muris* eggs are passed through murine faeces and embryos develop inside the eggs. Upon
459 ingestion by a murine host, the eggs hatch in the small intestine and release larvae that acquire
460 selected populations of bacteria from the gut of their rodent hosts. Thereafter, the larvae mature
461 and establish themselves as adult males and females in the colon. (C) Female *H. contortus*
462 transfer selected populations of bacteria to the offspring via the germline. Eggs shed in the
463 environment with the faeces of the ruminant host hatch and release first-stage larvae. Upon
464 ingestion of the latter by a new ruminant host, the developing worms acquire further
465 populations of bacteria from the rumen of the latter. Red arrows indicate helminth microbiome
466 acquisition events. Empty blue circles indicate the absence of a microbiome.

467

468 **Figure 2. Identification of key helminth taxa to be investigated as**
469 **representatives of their taxonomic clades.**

470 Asterisks (*) indicate taxa for which published data on parasite microbiome structure and/or
471 function(s) is available. The figure was adapted from Parkinson et al. [60].

472

473 **Figure 3. Plan of action for an efficient investigation and consequent**
474 **exploitation of helminth microbiomes.**

475 (A) 16S rRNA gene sequencing of helminth microbiomes to establish core microbiota present.
476 (B) Shotgun metagenomic sequencing (i), proteomic (ii), and metabolomic (iii) analyses of the
477 helminth microbiota to annotate functional roles to the core microbiota. (C) Implementation of
478 animal models in wild type and germ-free mice to assess helminth-microbiota acquisition and
479 transmission. (D) Localization of microbiota through fluorescent light- and electron
480 microscopy. (E) Identification and targeting of key microbiota to decrease parasite fitness
481 (based on the previously assessed parameters) through the implementation of microbiome
482 editing techniques.

Outstanding questions

- Which microbial taxa form the core microbiomes of different helminth species?
- How do helminths acquire their microbiomes?
- Where do endosymbionts localise within the parasite host?
- What are the functions of the microbiomes of parasitic helminths?
- What role/s do microbes other than bacteria (i.e. viruses and fungi), play in parasite biology and physiology?
- Can helminth-microbiome relationships be exploited for the development of new strategies for parasite control?



28 October 2018

Dear Doctor Kong,

REJOINER: MS. No. (TREPAR-D-18-00201)

Entitled 'Helminth Microbiomes - a Hidden Treasure Trove?' *Trends in Parasitology*.

We are very grateful for the reviewers' and editor's time and efforts in evaluating this manuscript. The points raised by the reviewers are very constructive and have contributed to enhance the manuscript. We have provided detailed responses to individual points in the following rejoinder:

Reviewer #1:

I am not sure that Figure 1 is absolutely necessary. Perhaps a figure depicting the selective acquisition of bacteria by *Trichuris*- and then how these bacteria and murine host bacteria then aid the parasite may be of greater interest?

RESPONSE: The authors thank the reviewer for this comment and have provided an additional figure depicting the proposed microbiome acquisition strategies of *Trichuris muris*, *Haemonchus contortus*, and *Brugia malayi*. The focus of this figure lies on acquisition rather than mechanisms of gut microbial benefits, since the latter remains highly speculative.

Reading this article made me think of "microbes" other than bacteria that may inhabit parasites- i.e. viruses and fungi- and how these affect parasite development/host responses. Given it is an opinion article, could the authors speculate this field as one that will emerge in future? I am aware of some fungi that negatively affect plant nematodes. With the new technologies for culturing and sequencing it may be pertinent to identify more than just bacterial species?

RESPONSE: The authors thank the reviewer for the comment and agree that other microbes could be of importance and should be investigated alongside bacteria. Following the reviewer's suggestion, an additional "outstanding question" has been added and discussed in the "Concluding Remarks" section (lines 251-252).

Reviewer #2:


My only comment/concern is with the term 'parabiome'. Although it is defined within the text, I don't know that it is essential. My original interpretation of the term was that it referred to the parasitic component of a host microbiome (similar to virome, mycobiome referring to viral and fungal components). When I did an informal poll of colleagues and trainees to see what they thought a 'parabiome' referred to, the responses were either as above, or that it could refer to the microbiome of mice that undergone parabiosis. Overall, I think it would be clearer to

continue to refer to the 'helminthic microbiome' as laid out in the title, and interspersed throughout the text.

RESPONSE: The authors thank the reviewer for this constructive comment and agree that the term 'parabiome' may raise some confusion. Therefore, following the reviewer's suggestion, we have replaced it with "helminth microbiome".



In conclusion, we consider that we have rigorously addressed each of the points raised by each of the reviewers. We agree with the points raised, and modified the text accordingly. These comments have led to an enhanced manuscript, which we consider to meet the standard for publication in *Trends in Parasitology*.

Yours sincerely,
Cinzia Cantacessi
On behalf of all authors.



[Click here to access/download](#)

Author Supplementary Material
TREPAR-D-18-00201.R1_tracked.docx



Your Postscript file "Figure1.eps" cannot be opened and processed.
Please see the common list of problems, and suggested resolutions below.

Reason: The submitted postscript could not be processed. Please check
the file and resubmit.

Other Common Problems When Creating a PDF from a Postscript file

You will need to convert your Postscript file to another format or fix
the current Postscript file, then re-submit it.

Your Postscript file "Figure2.eps" cannot be opened and processed.
Please see the common list of problems, and suggested resolutions below.

Reason: The submitted postscript could not be processed. Please check
the file and resubmit.

Other Common Problems When Creating a PDF from a Postscript file

You will need to convert your Postscript file to another format or fix
the current Postscript file, then re-submit it.

Your Postscript file "Figure3.eps" cannot be opened and processed.
Please see the common list of problems, and suggested resolutions below.

Reason: The submitted postscript could not be processed. Please check
the file and resubmit.

Other Common Problems When Creating a PDF from a Postscript file

You will need to convert your Postscript file to another format or fix
the current Postscript file, then re-submit it.