

1 Unusual Activity of a *Chlamydomonas* TET/JBP Family Enzyme

2 L. Aravind,^{*,†} Shankar Balasubramanian,^{*,‡,§} and Anjana Rao^{*,§,||}

3 [†]National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, Maryland
4 20894, United States

5 [‡]Department of Chemistry and Cancer Research UK Cambridge Institute, University of Cambridge, Cambridge CB2 0RE, U.K.

6 [§]La Jolla Institute for Immunology and Sanford Consortium for Regenerative Medicine, La Jolla, California 92037, United States

7 ^{||}Department of Pharmacology and Moores Cancer Center, University of California, San Diego, California 92093, United States

8 **T**he Great Oxygenation Event, which occurred on earth
9 around 2.45 billion years ago, opened up the niche for
10 enzymes that could directly oxidize substrates using molecular
11 oxygen (O₂), including the 2-oxoglutarate (2OG) and Fe(II)-
12 dependent dioxygenases. 2OGFe-dioxygenases possess a
13 double-stranded b-helix fold, with an iron center chelated by
14 two histidines and an aspartate.¹ 2OGFe-dioxygenases catalyze
15 the incorporation of one atom of O₂ into 2-oxoglutarate,
16 oxidizing it to succinate, and the second atom into an organic
17 substrate. The AlkB family and related clades catalyze this
18 reaction on alkyl adducts on nitrogens of bases, whereas the
19 TET/JBP family catalyzes the oxidation of methyl groups
20 attached to carbons of bases.¹ As enzymes that generate
21 epigenetic marks by modifying DNA or RNA, members of
22 both the AlkB and TET/JBP families have been repeatedly
23 recruited to different eukaryotic lineages.

24 In eukaryotes, enzymes of the TET/JBP family were first
25 recognized in kinetoplastids (e.g., the human parasite
26 *Trypanosoma*), where they act on thymine in a pathway that
27 results in the synthesis of the hypermodified base J in DNA
28 (Figure 1A). In metazoans and fungi, however, TET/JBP
29 enzymes catalyze the serial oxidation of 5mC to 5-hydroxy-, 5-
30 formyl-, and 5-carboxylcytosine (5hmC, 5fC, and 5caC,
31 respectively) (Figure 1B, top). All three oxidized methyl-
32 cytosines are intermediates in DNA demethylation; in
33 mammalian cells, 5hmC marks highly expressed genes and
34 active enhancers.²

35 The chlorophyte alga *Chlamydomonas reinhardtii* possesses a
36 lineage-specific expansion of at least 12 genes predicted to
37 encode active TET/JBP enzymes (in genome assembly
38 Genbank GCA_000002595.3). A recent study of one of
39 these enzymes,³ termed CMD1, revealed an unexpected
40 feature. Unlike other characterized 2OGFe-dioxygenases,
41 *Escherichia coli*-expressed CMD1 uses ascorbate instead of
42 2OG as its essential co-substrate *in vitro* (Figure 1B, bottom).
43 Enzymes typically do not generate a mixture of stereoisomeric
44 products, but CMD1 produces a skewed (3:1) mixture of
45 stereoisomeric glyceryl modifications of 5mC as its major
46 products, with the glyceryl moiety directly linked to the -CH₂
47 group at position 5 of cytosine [5-glyceryl-methylcytosines
48 (5gmC)]. This has functional implications, because protein
49 readers of major groove modifications would typically show a
50 preference for a defined stereochemistry. On the basis of
51 biosynthetic isotope tracer experiments in *E. coli*, the authors
52 propose that the two stereoisomeric 5gmCs are produced
53 directly from the ascorbate co-substrate via its oxidation to
54 CO₂ and glyoxylate, along with transfer of the glyceryl moiety

to 5mC. 5hmC and 5caC are produced in minor amounts, 55
through a mechanism that apparently does not require 2OG 56
but needs ascorbate. 57

Ascorbate (>1 mM) is believed to help reduce Fe(III) at the 58
active site of 2OGFe-dioxygenases back to Fe(II), thereby 59
“resetting” the active site for catalysis. Addition of ascorbate 60
results in a dose-dependent improvement in catalysis by 61
mammalian TET1 and an increased amount of 5hmC in cells.⁴ 62
Thus, instead of the usual 2OG substrate, *C. reinhardtii* CMD1 63
appears to have evolved to use a cofactor normally involved in 64
restoration of enzyme activity as its essential co-substrate, to 65
produce both the glyceryl adducts and the minor oxidized 66
5mC products. 67

From the known structures of TET/JBP proteins, a total of 68
10 residues other than the three (H, D/E, and H) that form 69
the iron-chelating center can be inferred to be proximal to the 70
2OG. Of these, the most important is the nearly absolutely 71
conserved arginine, which forms a salt bridge with the distal 72
carboxylate of 2OG. This residue is also conserved in all *C.* 73
reinhardtii TET/JBP proteins and might form a salt bridge with 74
ionized ascorbic acid in CMD1. This basic mechanism of 75
ascorbate binding is likely to be conserved across the TET/JBP 76
superfamily. The remaining less conserved positions might play 77
a role in the increased selectivity of CMD1 for ascorbate over 78
2OG. In principle, certain family members could utilize 79
ascorbate analogues to catalyze distinct modifications. 80

The 5gmC base constitutes a very minor fraction (~0.25%) 81
of 5mC in genomic DNA of *C. reinhardtii*, ~1000 5gmC in the 82
entire *C. reinhardtii* genome. Nevertheless, like oxidized 5mC 83
generated by mammalian and fungal TET/JBP enzymes, 5gmC 84
functions to counter the repressive effects of DNA cytosine 85
methylation. Mutants lacking CMD1 were compromised for 86
nonphotochemical quenching, a photoprotective mechanism 87
required for growth under high-light conditions. This 88
phenotype was traced to a decreased level of expression and 89
an increased level of cytosine methylation at two genes 90
encoding LHCSR3 (light-harvesting complex stress-related 91
protein 3). *C. reinhardtii* possesses two additional genes that are 92
very closely related to CMD1 (>99% sequence identity at the 93
protein level over the entire alignable length), possibly 94
explaining why CMD1 mutants retain ~40% of the wild-type 95
levels of 5gmC. 96

How does 5gmC promote cytosine demethylation? Unlike 97
most other organisms of the Viridiplantae lineage, *C. reinhardtii* 98




Figure 1. (A) Pathways for the transformations from C to 5caC or from C to 5gmC. (B) JBP-mediated oxidation of T to 5hmU followed by glycosylation to form base J.

99 lacks a member of the Demeter family of DNA glycosylases,
100 the primary enzymes involved in cytosine demethylation in
101 plants. However, it possesses a homologue of the TDG family,
102 which has been shown to be involved in removing the 5fC and
103 5caC modifications from metazoan DNA. Hence, it is possible
104 that the large adduct generated by CMD1 is a target for the
105 cognate DNA glycosylase, which then helps demethylate
106 regulatory regions of genes involved in photoadaptation to
107 high-light conditions. This view is supported by the
108 observation that in the CMD1 mutant in which global 5gmC
109 levels drop by 40%, global 5mC levels are doubled. The same
110 doubling of global 5mC occurs when ascorbate biosynthesis is
111 removed. Because basal 5mC levels are 400-fold higher than
112 5gmC levels in wild-type cells, this phenomenon cannot be

113 explained simply by conversion of 5mC to 5gmC. Rather,
114 5gmC formation is likely a necessary intermediate on a high-
115 flux DNA demethylation pathway (whether replication-
116 dependent or glycosylase-mediated) that turns over 5mC.

117 Certain eukaryotic lineages, including fungi such as
118 *Coprinopsis cinereus*, show moderate to large lineage-specific
119 expansions and frequent associations with DNA transposons of
120 the KDZ (Kykuja–Dileera–Zisupton) family.⁵ These DNA
121 transposons have arguably played major roles in disseminating
122 the associated TET/JBP enzymes across their host genomes
123 and may utilize the enzymes to regulate both their own
124 expression and that of other self-ish elements in the genome.
125 CMD1 is related to these enzymes and was likely acquired by
126 the Chlamydomonadales from such a source, especially given

127 that some versions in these algae are still linked to the
128 transposons. Future studies will reveal whether the unusual
129 catalytic activity reported for CMD1 is more widely distributed
130 across the TET/JBP family.

131 ■ AUTHOR INFORMATION

132 Corresponding Authors

133 *E-mail: aravind@ncbi.nlm.nih.gov.

134 *E-mail: sb10031@cam.ac.uk.

135 *E-mail: arao@lji.org.

136 ORCID

137 Shankar Balasubramanian: 0000-0002-0281-5815

138 Anjana Rao: 0000-0002-1870-1775

139 Funding

140 L.A. is supported by the intramural funds of the National
141 Library of Medicine. Work in the Rao lab is supported by
142 National Institutes of Health (NIH) Grant R35 CA210043 to
143 A.R. Work in the Balasubramanian lab is supported by core
144 funding (C14303/A17197) and program funding (C9681/
145 A18618) from Cancer Research UK and by an Investigator
146 Award from the Wellcome Trust (209441/z/17/z).

147 Notes

148 The authors declare the following competing financial
149 interest(s): S.B. is a founder, and A.R. is a member of the
150 scientific advisory board, of Cambridge EpiGenetix. L.A.
151 declares no competing financial interests.

152 ■ ACKNOWLEDGMENTS

153 The authors thank Rahul Kohli for valuable discussions.

154 ■ REFERENCES

- 155 (1) Hausinger, R. P., and Schofield, C. J. (2015) *2-Oxoglutarate-*
156 *dependent dioxygenases*, Royal Society of Chemistry.
- 157 (2) Scott-Browne, J. P., Lio, C-W J, and Rao, A. (2017) TET
158 proteins in natural and induced differentiation. *Curr. Opin. Genet. Dev.*
159 *46*, 202–208.
- 160 (3) Xue, J. H., Chen, G. D., Hao, F., Chen, H., Fang, Z., Chen, F. F.,
161 Pang, B., Yang, Q. L., Wei, X., Fan, Q. Q., Xin, C., Zhao, J., Deng, X.,
162 Wang, B. A., Zhang, X. J., Chu, Y., Tang, H., Yin, H., Ma, W., Chen,
163 L., Ding, J., Weinhold, E., Kohli, R. M., Liu, W., Zhu, Z. J., Huang, K.,
164 Tang, H., and Xu, G. L. (2019) A Vitamin C-derived DNA
165 modification catalyzed by an algal TET homologue. *Nature* *569*,
166 581–585.
- 167 (4) Blaschke, K., Ebata, K. T., Karimi, M. M., Zepeda-Martínez, J. A.,
168 Goyal, P., Mahapatra, S., Tam, A., Laird, D. J., Hirst, M., Rao, A.,
169 Lorincz, M. C., and Ramalho-Santos, M. (2013) Vitamin C induces
170 Tet-dependent DNA demethylation and a blastocyst-like state in ES
171 cells. *Nature* *500*, 222–226.
- 172 (5) Iyer, L. M., Zhang, D., de Souza, R. F., Pukkila, P. J., Rao, A., and
173 Aravind, L. (2014) Lineage-specific expansions of TET/JBP genes
174 and a new class of DNA transposons shape fungal genomic and
175 epigenetic landscapes. *Proc. Natl. Acad. Sci. U. S. A.* *111*, 1676–1683.