

Imaging suicidal thoughts and behaviors: a comprehensive review of two decades of neuroimaging studies

Lianne Schmaal^{*1,2}, Anne-Laura van Harmelen^{*3}, Vasiliki Chatzi³, Elizabeth T.C. Lippard⁴, Yara J. Toenders^{1,2}, Lynnette A. Averill⁵, Carolyn M. Mazure⁶, Hilary P. Blumberg⁷

* shared first authorship

¹Orygen, The National Centre of Excellence in Youth Mental Health, Parkville, Australia

²Centre for Youth Mental Health, The University of Melbourne, Melbourne, Australia

³Psychiatry, University of Cambridge, Cambridge, UK

⁴Psychiatry, Dell Medical School, University of Texas, Austin, USA

⁵Psychiatry, Yale School of Medicine, New Haven, USA and Department of Veterans Affairs National Center for PTSD, Clinical Neurosciences Division, West Haven, USA

⁶Psychiatry and Women's Health Research at Yale, Yale School of Medicine, New Haven, USA

⁷Psychiatry, Radiology and Biomedical Imaging, and the Child Study Center, Yale School of Medicine, New Haven, USA

Corresponding author:

Name: Hilary Blumberg, M.D.

Affiliation: Yale School of Medicine

Address: Department of Psychiatry, 60 Temple Street, Suite 6B, New Haven, CT
06510

Telephone: 203-785-6180

Email: hilary.blumberg@yale.edu

Word count: Text: 9386 words, 3 Figures, 3 Tables

ABSTRACT

Identifying brain alterations that contribute to suicidal thoughts and behaviors (STBs) are important to developing more targeted and effective strategies to prevent suicide. In the last decade, and especially in the last 5 years, there has been exponential growth in the number of neuroimaging studies reporting structural and functional brain circuitry correlates of STBs. Within this narrative review, we conducted a comprehensive review of neuroimaging studies of STBs published to date and summarize the progress achieved on elucidating neurobiological substrates of STBs, with a focus on converging findings across studies. We review neuroimaging evidence across differing mental disorders for structural, functional and molecular alterations in association with STBs, that converges particularly in regions of brain systems that subserve emotion and impulse regulation including the ventral prefrontal cortex (VPFC) and dorsal PFC (DPFC), insula and their mesial temporal, striatal and posterior connection sites, as well as in the connections between these brain areas. The reviewed literature suggests that impairments in medial and lateral VPFC regions and their connections may be important in the excessive negative and blunted positive internal states that can stimulate suicidal ideation, and that impairments in a DPFC and inferior frontal gyrus (IFG) system may be important in suicide attempt behaviors. A combination of VPFC and DPFC system disturbances may lead to very high risk circumstances in which suicidal ideation is converted to lethal actions via decreased top-down inhibition of behavior and/or maladaptive, inflexible decision-making and planning. The dorsal anterior cingulate cortex and insula may play important roles in switching between these VPFC and DPFC systems, which may contribute to the transition from suicide thoughts to behaviors. Future neuroimaging research of larger sample sizes, including global efforts, longitudinal designs, careful consideration of developmental stages, and sex and gender, will facilitate more effectively targeted preventions and interventions to reduce loss of life to suicide.

Keywords

Attempted Suicide; Suicidal Ideation; Suicide; Neuroimaging; Functional Neuroimaging, Magnetic Resonance Imaging

1 **1. INTRODUCTION**

2 Around one million people die by suicide annually¹. Globally, suicide is the tenth leading
3 cause of death for all ages and the second leading cause of death among young people aged
4 15-29 years¹. In 2013, it was estimated that 9.3 million adults 18 years and older in the United
5 States had suicidal thoughts and 1.3 million attempted suicide². Additionally, a 2011 report
6 estimated that 13% of adolescents planned a suicide attempt in the previous year and 8%
7 attempted suicide³. Unfortunately, suicide death rates have continued to rise. For example,
8 since 1999, rates in the U.S. have increased 30%⁴. Predictive value of currently identified
9 non-biological risk factors for suicide is limited⁵, and a reliable biological risk marker has yet to
10 be identified. In order to prevent suicide more effectively, there is an urgent need to better
11 understand the mechanisms that confer increased risk for suicidal thoughts and behaviors
12 (STBs), and to identify biological markers of risk to generate more targeted successful
13 prevention strategies and monitor responses to them. In the last decade, and especially in the
14 last 5 years, there has been exponential growth in the number of neuroimaging studies
15 reporting structural and functional brain circuitry correlates of STBs (Figure 1). In the last 10
16 years, a number of excellent reviews on aspects of this research have emerged⁶⁻¹⁰. Here we
17 review research across structural, functional and neurochemical neuroimaging modalities,
18 providing a narrative review of 130 neuroimaging studies with a focus on the most researched
19 brain circuitries and findings that converge across studies.

20
21 **2. METHODS**

22 A search was performed in PubMed for original research articles published before March 12,
23 2018. The following terms were used: "MRI", "SPECT", "PET", "magnetic resonance
24 imaging", "positron emission tomography", "single photon emission computed tomography",
25 "DTI", "diffusion tensor imaging", "diffusion weighted imaging", "neuroimaging", "fMRI",
26 "functional magnetic resonance imaging", "spectroscopy" (separated by OR) in combination
27 with the terms "suicide", "suicidal", "suicidality" (separated by OR). We selected articles that:
28 (i) were published in a peer-reviewed journal in English and (ii) included groups with suicidal
29 ideation (SI) and/or history of suicide attempt(s) (SA). Of note, non-suicidal self-injury (NSSI)
30 was not included in this review, and studies that did not differentiate NSSI from suicidal
31 behaviors were excluded, because these can be differentiated on the basis of intention,
32 frequency, and lethality and may have partly distinct underlying mechanisms¹¹.

33
34 **3. RESULTS**

35 We identified 131 unique articles meeting review criteria (Tables 1-3). The populations
36 studied reflect reports that the majority of people with STBs have a diagnosable mental
37 illness. Major depressive disorder (MDD) or bipolar disorder (BD) account for over half of
38 suicide deaths¹². After mood disorders and borderline personality disorder (BPD), the
39 prevalence of suicide deaths is highest among people with substance use disorders and
40 schizophrenia (SZ), followed by post-traumatic stress disorder (PTSD) and anxiety

41 disorders¹². The mental disorders researched varied by study, with each study typically
42 including a single disorder, and the majority of studies conducted in individuals with mood
43 disorders. Most studies compared people with a mental disorder and a history of suicide
44 attempts (suicide attempters, SAs) to people with a mental disorder and/or healthy controls
45 (HCs) without a history of attempt. Fewer studies focused on SI. Most studies employed a
46 cross-sectional design and a single structural or functional imaging modality. The majority
47 were conducted with adults; only a small proportion with adolescents (Figure 1). A subset of
48 studies provided preliminary findings of associations between neuroimaging measures and
49 key risk factors for suicide, e.g. medical lethality of prior attempts, emotion dysregulation,
50 anhedonia, impulsiveness, and reduced cognitive control (for reviews see¹³⁻¹⁵).

51

52 Despite modest sample sizes of studies, and the heterogeneity of their clinical samples and
53 neuroimaging acquisition and analysis methods, converging evidence is emerging to support
54 roles for specific brain regions/circuitries in STBs. These are particularly in cortico-
55 striatolimbic systems that subserve emotion and impulse regulation and include prefrontal,
56 cingulate and insula cortices, amygdala, hippocampus, thalamus and striatum regions
57 (Tables 1-3). Within the prefrontal cortex (PFC), studies vary widely in the selection of regions
58 studied, with regions of interest (ROIs) often including overlapping regions that encompass
59 ventral and dorsal, medial as well as lateral, PFC. Thus, while we identify the importance for
60 future study of specific PFC subregions, given their differing connectivity, cellular and
61 molecular features and functions, we discuss the PFC grouped broadly into the ventral
62 prefrontal cortex (VPFC; divided into medial and lateral portions), which has the highest
63 concentration of reported findings, the dorsal prefrontal cortex (DPFC; divided into lateral and
64 medial portions), and the anterior cingulate cortex (ACC). We also discuss findings in the
65 insula, and mesial temporal (hippocampus, amygdala), subcortical (basal ganglia, thalamus)
66 and posterior regions (posterior cingulate cortex, lateral temporal lobes and cerebellum). See
67 Figure 2 for definitions of the brain regions. Since the VPFC, DPFC, ACC, insula, mesial
68 temporal, basal ganglia, thalamus and posterior regions have been studied most frequently in
69 relation to STBs and because most converging evidence exist for the involvement of these
70 regions, we specifically focus our review on findings within these brain areas, as well as in the
71 connections between them. However, additional studies and positive and negative findings
72 not discussed below can be found in Tables 1-3.

73

74 We focus our discussion below on findings that are primarily derived from comparisons of
75 individuals with a mental disorder and STBs versus individuals with a mental disorder without
76 STBs (diagnostic controls, DCs), rather than comparisons with HCs, unless otherwise
77 specified. Findings based on a comparison between DCs and individuals with the same
78 mental disorder plus STBs are more commonly reported in the literature and are more likely
79 to reflect specific effects of STBs in that disorder, whereas comparisons with HCs may
80 include more general effects of having a mental disorder. Below we first detail structural,

81 functional and neurochemical findings within regions and end each section on a region with a
82 summary. We then follow with a section devoted to studies of connectivity among regions
83 within major implicated brain systems.

84

85 **Ventral Prefrontal Cortex (VPFC)**

86 The *lateral* VPFC (VLPFC) refers to the inferior lateral areas of the frontal cortex
87 encompassing lateral orbitofrontal cortex (OFC, brodmann area (BA)47, lateral BA11), inferior
88 frontal gyrus (IFG, BAs 44, 45), and lateral aspects of the rostral PFC (RLPFC, lateral BA10)
89 (Figure 2). The lateral VPFC plays a key role in cognitive control, including response
90 inhibition, and is activated when behavioral responses are modulated in response to the
91 emotional or motivational context^{16,17}. The *medial* VPFC (VMPFC) refers to the medial OFC
92 (medial BA11) and medial aspects of the rostral PFC (RMPFC, medial BA10) (Figure 2). The
93 medial VPFC has a well-established role in self-reflection¹⁸, appraisal of internally-generated
94 emotions (both positive and negative)^{16,19}, appraisal of past and imagined future events and
95 reward processing^{20,21}. Structural, functional and neurochemical alterations in these regions
96 have been associated with maladaptive strategies for regulation of negative affect (e.g.,
97 rumination), negative self-referential thinking^{22,23}, and diminished positive affect (e.g.,
98 anhedonia)²⁴. Evidence is mounting that emotion dysregulation has a central role in the
99 generation of STBs. This includes elevations in negative, and blunting in positive, subjective
100 emotions, self-referential thoughts, and responses to valenced stimuli²⁵. These alterations are
101 thought to contribute to key clinical risk symptoms for STBs including depression, anxiety,
102 rumination, guilt, reduced self-esteem, helplessness, anhedonia and hopelessness^{22-24,26-31}.

103

104 *VLPFC*

105 Structural MRI studies have consistently shown lower gray matter volumes of the VLPFC in
106 adult SAs, including SAs with MDD³²⁻³⁴, BD^{35,36} and schizoaffective disorder (SZA)³⁷.
107 Findings of lateral OFC volume decreases, extending to medial OFC, in adolescent and
108 young adult SAs with BD³⁶, suggest these may be early differences. Lower VLPFC thickness,
109 but not volume, is one of the rare findings related to SI in adults with MDD³⁸, suggesting the
110 VLPFC may be involved not only in suicide behavior but also in the ideation that may
111 generate it.

112

113 Gray matter volume decreases in VLPFC were also associated with high lethality of prior
114 suicide attempts in MDD²⁴ and BPD^{36,37}. This converges with longstanding findings from both
115 postmortem studies demonstrating VPFC (ROIs including both VLPFC and VMPFC)
116 differences in people who died by suicide³⁹, and positron emission tomography (PET) studies
117 of high lethality or high intent SAs, showing VPFC (VLPFC and VMPFC) alterations in
118 serotonin (5-HT) synthesis, transporters and 5-HT1a receptors^{40,41} (Table 2). Lower 5-HT1a
119 binding in the OFC was associated with interim SI during a 2-year follow-up in adults with
120 MDD and past attempts⁴². Although conflicting findings exist⁴³⁻⁴⁶, these data provide some

121 consistent findings of a location and a potential mechanism, i.e. lateral extending to medial
122 VPFPC serotonergic dysfunction, as a potential biomarker of risk for high lethality STBs. There
123 have been few molecular imaging studies of other neurotransmitter systems and
124 neurochemicals implicated in STBs.

125

126 Functional MRI (fMRI) studies performed while subjects conducted specific behavioral tasks
127 provide evidence that STBs are associated with VLPFC functional abnormalities in response
128 to emotional and other hedonically-valenced stimuli. Increased activation of the IFG and
129 lateral and medial OFC while viewing angry (but not happy, sad or neutral) faces was
130 reported in adult SAs with MDD^{47,48}. Higher IFG activation in response to angry faces was
131 also associated with poorer attempt planning and higher impulsivity in adult SAs with MDD⁴⁹.
132 Furthermore, young adult SAs with BPD displayed higher lateral OFC activation while
133 instructed to experience and regulate negative autobiographical memories⁵⁰. Increased
134 activation in a region of interest that included both the medial and lateral OFC was also seen
135 in response to winning a reward⁴⁷ in adult SAs with MDD.

136

137 The IFG also plays a critical in cognitive control and response inhibition⁵¹. During
138 performance of a continuous performance task, higher IFG, RLPFC and lateral OFC
139 responses were associated with both attempts and SI in adults with mood disorders with
140 psychotic features, in the absence of task performance differences⁵². Higher lateral OFC was
141 also reported during error trials in a response inhibition task in veterans with SI⁵³. In contrast,
142 a second study in adult SAs with SZ using the same continuous performance task showed
143 reduced activation in a cluster encompassing the RLPFC and IFG, extending to the VMPFC
144 and ventral ACC, that was associated with SI but did not further distinguish between ideators
145 with and without a history of attempt⁵⁴.

146

147 *VMPFC*

148 In addition to the structural and PET study findings reported for the VLPFC above that
149 extended to the VMPFC or that were based on an ROI including both VLPFC and VPFPC,
150 lower medial VPFPC cortical thickness was also associated with greater motor impulsivity in
151 adolescent SAs with MDD⁵⁵. As cortical thickness and surface area contributions to volume
152 are thought to be genetically independent⁵⁶ and result from different neurobiological
153 processes⁵⁷, it is important to examine these separately in studies of STBs. However, the
154 majority of studies have either not examined thickness and surface area separately from
155 volume or only thickness but without examination of surface area in SAs with MDD^{32,58}.
156 Cortical thickness is thought to be influenced by the number and the size of cells within a
157 column, packing density, as well as by the number of connections and the extent of their
158 myelination, while cortical surface area is driven by the number ontogenetic columns that run
159 perpendicular to the surface of brain⁵⁹.

160

161 Functionally, in addition to higher activation in the lateral and medial OFC in response to
162 angry faces and to winning a reward in adult SAs with MDD⁴⁷ as reported above, a recent
163 study using machine learning to investigate adolescent SAs (with and without SI) showed that
164 the medial VMPFC was among the most discriminating regions, within a multivariate pattern
165 of fMRI brain activation in response to actively thinking about life- and death-related concepts,
166 for distinguishing between adolescent suicidal ideators with and without a history of attempt,
167 although in a very small sample size⁶⁰.

168

169 *Summary*

170 Structural neuroimaging studies have consistently shown that alterations in the VLPFC and
171 VMPFC are implicated in SA across a range of mental disorders and age ranges. Reduced
172 VLPFC volumes were also associated with lethality of attempts, potentially mediated by
173 serotonergic dysfunction, although findings of serotonergic dysregulation remain inconsistent.
174 The involvement of structural and functional VLPFC and VMPFC alterations in SI remains
175 understudied. Viewing and regulating negative emotions and motivationally-valenced stimuli
176 has been linked to increased activation of the lateral and medial OFC in adult (including
177 young adults) SAs with MDD and BPD, and associated with poorer attempt planning and
178 higher impulsivity. Finally, higher VLPFC activity, including in the IFG, RLPFC and lateral
179 OFC, during cognitive control and response inhibition in relation to SA and SI in adults with
180 mood disorders has been reported across a number of studies. These increased activations
181 may, in the absence of task performance differences, reflect a need for greater engagement
182 of these regions for reaching similar performance in these individuals with STBs. In contrast,
183 adult SAs with SZ showed reduced activation in these regions during cognitive control, which
184 may suggest that there are some differences in the neural signatures of STBs between mood
185 and psychotic disorders, which will be an important direction for future study.

186

187 **Dorsal Prefrontal Cortex (DPFC)**

188 The DPFC can be broadly divided into dorsolateral (DLPFC) and dorsomedial PFC (DMPFC).
189 The DMPFC and the DLPFC together support top-down control of emotions and behaviors,¹⁷
190 cognitive flexibility and complex decision-making⁶¹. Deficits in these processes are thought to
191 have an important role in STBs, particularly in the transition from SI to behavior, as the
192 threshold to acting is lowered by decreased top-down behavioral inhibition, and diminished
193 flexibility in generating alternate and more adaptive behavioral choices^{15,62}. Neuroimaging
194 evidence suggests that the DMPFC (medial portions of BA 8 and 9) is robustly recruited
195 during tasks that require mental state inference^{63,64}. The DMPFC is further involved in
196 tracking decision conflict and reinforcement history⁶⁵, as well as in emotion regulation⁶⁶. The
197 DLPFC (BA46, lateral BA9) is involved in the conscious active control of planned behavior
198 and cognition, as well as working memory⁶⁷. Access of the DLPFC to memory processing in
199 hippocampal regions is shared by the rostralateral PFC (lateral frontal pole BA10), which has
200 been implicated in meta-cognitive awareness^{68,69}.

201

202 *DLPFC*

203 Although the amount of evidence to date has been less than in VPFC, accumulating findings
204 also support a role for the DLPFC in STBs. Structural MRI studies support lower volume in
205 DLPFC in adult SAs across MDD^{32,33,70} and BD^{33,35}. Studies showed lower DLPFC thickness
206 in adult SAs with MDD⁵⁸ and SZ⁷¹, but have not examined cortical surface area. In addition,
207 lower DLPFC volume was associated with attempt lethality in mood and psychotic
208 disorders^{32,37}. Higher baseline 5-HT1a receptor binding potential in the DLPFC was also
209 associated with higher lethality of future attempts and SI during a 2-year follow-up⁴².

210

211 Functionally, decreased lateral (and medial) DPFC when ten adults with self-reported
212 depression listened to their own narrative of their attempt was reported in a study in which
213 imaging was conducted close to the time of the attempts (one to four weeks prior⁷²). A study
214 of adolescents with a history of SI showing lower right DLPFC activation during passive
215 viewing of negative emotional scenes, suggests that DLPFC decreases during processing of
216 negative emotional stimuli might be present early in the course of SI⁷³. Higher right DLPFC
217 engagement was observed during regulation of responses to negative emotional scenes in
218 the same adolescents, suggesting the direction of DLPFC differences depends on the specific
219 task requirements (passive viewing versus regulating)⁷³. Another study in adolescent SAs
220 with MDD suggests that the direction may also relate to the specific emotion, as passive
221 viewing of angry faces, but not happy faces, elicited higher DLPFC responses⁷⁴, perhaps due
222 to the high sensitivity to criticism and social rejection previously reported in individuals with
223 STBs^{75,76}.

224

225 The DLPFC's critical role in decision-making in the context of evaluating the motivational
226 value of choices⁷⁷ may be especially relevant to STBs. Blunted DLPFC activation was
227 observed when evaluating risky versus safe options in adult SAs with MDD⁴⁷ and when
228 evaluating lower immediate rewards versus larger delayed rewards in older adults with MDD
229 and well-planned suicide attempts⁷⁸. This is in line with a behavioral study of older adult SAs
230 with MDD showing lower levels of delay discounting (or impulsive decision-making) and better
231 planning in suicide attempts⁷⁹. Increased DLPFC activation has also been observed in adults
232 with STBs across a range of "cold" cognitive control tasks, especially those requiring inhibition
233 of automatic response tendencies. These included paradigms such as continuous
234 performance, stop signal, go-no go and stroop tasks studied in adult SAs with MDD or BD
235 with psychotic features⁵², SZ⁸⁰, and ideators with SZ⁸¹ or PTSD and MDD in veterans⁵³.
236 Elevated activation in DLPFC was observed in suicidal ideators with past attempts compared
237 to ideators without attempts while performing a continuous performance task⁵². Single photon
238 emission tomography (SPECT) and PET studies showing lower resting regional cerebral
239 blood flow (rCBF) and glucose metabolic rates (rCMRglu) in the DLPFC in adult SAs with

240 mood disorders^{82,83}. Moreover, lower DLPFC rCMRglu was associated with higher lethality of
241 attempt⁸⁴ and with SI with a plan versus ideation without a specific plan⁸⁵ in adults with MDD.

242

243 *DMPFC*

244 Structural MRI studies also support lower volume in DMPFC in adult SAs with MDD^{32,70} and
245 BD^{33,35}. Although less studied than DLPFC in functional neuroimaging, lower DMPFC was
246 reported in the study in which adults with depression listened to their own narrative of their
247 recent attempt, which was especially pronounced during mental pain aspects of the
248 narrative⁷². Furthermore, decreased activation was also found in DMPFC during viewing of
249 angry faces in adult SAs with MDD⁴⁸.

250

251 *Summary*

252 Structural alterations in both the DLPFC and DMPFC have been consistently observed in
253 adults across mental disorders and structural alterations in the DLPFC have been associated
254 with attempt lethality. This latter finding, together with findings of serotonergic dysfunction in
255 the DLPFC being associated with higher lethality of future attempts during a 2-year follow-up,
256 implicates DLPFC structure and serotonergic system functioning in STB risk. With regard to
257 DLPFC and DMPFC functioning, there is convergence in showing differences related to STBs
258 during processing of negative emotional stimuli, although the direction of effects (activation
259 increases versus decreases) in the DLPFC differed across studies, with contributions to the
260 differences unclear as the studies differed across multiple variables. Elevated activation while
261 performing cognitive control tasks (in the absence of performance differences) together with
262 lower resting regional cerebral blood flow and glucose metabolic rates in the DLPFC could
263 suggest that increased DPFC may be recruited for reaching similar task performance perhaps
264 due to lower baseline levels of activation in the DPFC regions. Findings from one study
265 suggests that functional DLPFC alterations during cognitive control can discriminate between
266 suicidal ideators with past attempts and ideators without attempts. Moreover, blunted DLPFC
267 activation when evaluating the value of different decision options may also represent a risk
268 marker for SA, and especially for well-planned attempts. Well-planned versus impulsive
269 suicide attempts have been suggested to be different phenotypes⁸⁶. From papers reviewed,
270 there was a greater concentration of findings in the VPFC in impulsive SAs and in DPFC in
271 planful SAs. Therefore, we speculate that the relative ventral versus dorsal localization of the
272 PFC abnormalities may contribute to the differing phenotypes.

273

274 **Anterior cingulate cortex (ACC)**

275 The ventral ACC consists of BA25 and ventral BA32 sub- and pre-genual to the corpus
276 callosum and plays a critical role in valuation and control of autonomic viscerosensory
277 signals, the modulation of physiological responses to stress and the appraisal of internal
278 feelings¹⁶. The dorsal ACC (dorsal BA24 and 32) plays an important role in the appraisal of

279 actions (and adaptively adjust behaviour as a consequence) and reward-based decision-
280 making¹⁶.
281
282 Structural MRI studies support lower volume in both ventral and dorsal ACC in adult SAs with
283 MDD^{34,58} and BD³⁵, and that was related to a higher number of attempts in adolescents with
284 BPD and MDD⁸⁷ and to higher lethality of attempts in adults with BPD⁸⁸ and psychotic BD³⁷.
285 Lower dorsal ACC (dACC) rCMRglu was associated with higher lethality of attempt⁸⁴. In
286 addition, lower 5-HT_{1a} binding in the ACC was associated with interim SI during a 2-year
287 follow-up in adults with MDD and past attempts⁴². A few studies that investigated
288 neurotransmitter systems other than 5-HT implicated the ACC in relation to SI. For example,
289 a positive association was shown between SI and ACC monoamine oxidase-A (MAO-A)
290 density in adults with BPD⁸⁹. A relation between SI and increased ACC neuroinflammation (as
291 assessed by translocator protein (TSPO) availability) was reported in adults with MDD⁹⁰.
292 Furthermore, dACC GABA concentrations were lower in adult female SA+SI compared to
293 clinical controls without SA or SI, however, this effect was no longer significant after
294 correcting for age⁹¹.
295
296 Viewing of angry faces elicited higher dACC responses in adolescent SAs with MDD⁷⁴,
297 perhaps due to the high sensitivity to criticism and social rejection previously reported in
298 individuals with STBs^{75,76}. In contrast, decreased activation was found in ACC (ROI capturing
299 both ventral and dorsal ACC) during viewing of sad faces in adult SA with MDD⁴⁷. With regard
300 to positive stimuli, blunted ventral ACC responses were found during the anticipation of
301 reward in adult⁹², including elderly⁹³, SAs with MDD. Elevated responses in the ventral ACC
302 have also been reported in relation to positive stimuli. For example, higher activation in
303 ventral ACC was seen in response to happy facial expressions⁴⁸ and in response to actual
304 winning⁴⁷ (in contrast to blunted response during reward anticipation) in adult SAs with MDD.
305
306 *Summary*
307 The ACC has mostly been studied in relation to emotional processing. Although various
308 studies have observed dorsal and ventral ACC activation alterations in adolescents and
309 adults with MDD and SA, the direction of alterations seem to be complex and dependent on
310 task condition and stimulus type (positive versus negative). One could perhaps interpret the
311 findings of increased dorsal and ventral activation in response to angry faces and to positive
312 stimuli, together with blunted ventral ACC activation during reward anticipation, as negative
313 biases, as they may reflect reduced reward anticipation (anticipation phase) versus increased
314 activation in response to negative stimuli and in relation to positive prediction errors in
315 response to positive stimuli (outcome phase). The finding of a positive relation between SI
316 and ACC neuroinflammation, together with findings of increased inflammatory markers in the
317 ACC in postmortem studies of people who died by suicide⁴⁸ and in blood and cerebrospinal

318 fluid in people with ideation and a history of violent or high intent attempts^{49,50}, suggests that
319 neuroinflammation in the ACC may constitute a promising target for future studies of STBs.

320

321 **Insula**

322 The insular cortex is a key hub in emotional processing with connectivity to PFC, particularly
323 VPFC, as well as mesial temporal structures⁹⁴. The insula plays an important role in
324 interoceptive awareness for positive and negative internal states⁹⁵, including emotional and
325 other types of pain, and understanding and sharing of other people's emotional states^{96,97}.

326 Only in more recent studies has insula structure and function and related-behavior been
327 investigated for its role in STBs. For example, on a behavioral level, interoceptive deficits
328 have been reported among SAs compared to individuals who only thought about or planned
329 suicide among general psychiatric outpatient adults⁹⁸ and predicted SI severity at 6-month
330 follow-up in community adolescents⁹⁹.

331

332 Smaller insula volume has been reported in adult SAs with BPD¹⁰⁰, in a combined group with
333 SZ/SZA/psychotic BD³⁷ and elderly with MDD⁷⁰. Lower insula thickness was observed in
334 adults in relation to suicide attempts in SZ⁷¹ and SI in MDD³⁸. Smaller insula volume was
335 associated with higher attempt lethality and lower impulsivity in BPD^{88,100}. In contrast, larger
336 insula volumes were reported in relation to attempt lethality in adults with BD¹⁰¹. It is possible
337 that the type of insula differences relate to specific characteristics of the high lethality
338 attempters, since larger insula volumes were also found in association with higher lifetime
339 history of aggression in BPD⁸⁸. Some findings in the PFC noted above in MDD extended to
340 the insula, including of associations between baseline 5-HT1a binding potential with SI and
341 lethality of future attempts within a 2-year follow-up period⁴² and of increased
342 neuroinflammation (TPSO availability)⁹⁰.

343

344 SPECT research showed higher insula rCBF in adult SAs with MDD¹⁰² at rest and higher
345 insula fMRI activation was found in adults with MDD or BD with psychotic features during a
346 cognitive control task with insula activity related to higher intensity of SI⁵². Higher insula fMRI
347 activation was also associated with lower subjective value of gain and loss in adult SAs with
348 MDD⁹². Lower activation in the posterior insula during social exclusion was found in adult
349 SA's with MDD or BD, which was suggested to indicate a higher tolerance to pain via
350 repeated exposure to painful and provocative experiences in subjects vulnerable for
351 suicide¹⁰³.

352

353 *Summary*

354 Smaller insula volume has been associated with SAs and lower impulsivity in adults across
355 various mental disorders, whilst, both smaller and larger insula volumes have been
356 associated with higher attempt lethality. fMRI studies found higher insular activation related to
357 during reward processing and cognitive control in adult SA with MDD, while lower insula

358 activation was associated with a higher tolerance to social pain in adult SAs with MDD or BD.
359 Thus, there is preliminary evidence for an involvement of insular structural and functional
360 alterations in SI and SAs. However, since very few studies have focussed on the insula and
361 both decreases and increases in these alterations have been reported, more research is
362 needed to elucidate the role of the insula in STBs. Interestingly, immune challenges activate
363 interoceptive brain pathways (including the insula), triggering alterations in mood and
364 cognition, motivation, and neurovegetative processes¹⁰⁴. Together with preliminary evidence
365 of increased neuroinflammation in the insula related to SI, this suggests that the insula may
366 be an important region for future studies of neuroinflammation and STBs.

367

368 **Amygdala and Hippocampus**

369 Due to their roles in processing of emotion, emotional memory and the stress response^{105–108},
370 the mesial temporal amygdala, hippocampus and entorhinal cortex (BA28, within the adjacent
371 parahippocampal gyrus) are also thought to be involved in STBs. However, findings have
372 been inconsistent. Larger amygdala volumes were reported in adult SAs with MDD¹⁰⁹ and
373 SZ¹¹⁰, but more studies have not detected significant amygdala findings^{33,38,100,101,111–113}.
374 Smaller hippocampus volumes were reported in adult SAs with MDD¹¹⁴ and adolescent and
375 young adult SAs with BD³⁶, and one study reported a smaller parahippocampal gyrus¹¹⁵.
376 However, more studies have not detected associations of hippocampus or parahippocampus
377 volume with STBs^{33–35,38,101,109–112}. While difficulties detecting differences in these small mesial
378 temporal structures may relate to imaging methods, it may be that mesial temporal alterations
379 are only apparent in specific subgroups of people with STBs. For example, high lethality
380 attempts were associated with smaller volumes of the hippocampus and parahippocampal
381 gyrus in adult SAs with BPD^{88,100}. In addition, amygdala and hippocampus volumes were
382 negatively associated with impulsivity in patients with low lethality attempts⁸⁸, and amygdala
383 volume positively associated with self-aggression in SAs with SZ¹¹⁰.

384

385 PET studies have shown preliminary evidence for a role of serotonergic alterations in the
386 amygdala and hippocampus in STBs. Increased hippocampus 5-HT_{2a} receptor binding and
387 5-HT release was observed in adult SAs with BPD¹¹⁶ and in adults with MDD and high
388 lethality attempts⁴⁰ respectively, compared to HCs. Recently, baseline 5-HT_{1a} receptor
389 binding in the amygdala, hippocampus and parahippocampal gyrus was associated with
390 higher SI during a 2-year follow-up in adults with MDD⁴².

391

392 Few fMRI studies have focused on mesial temporal ROIs. An activation study focusing on the
393 amygdala found no association between SI and amygdala functioning during emotion
394 processing in 10 ideators¹¹⁷. Increases were observed during autobiographical recall of
395 mental pain experienced during an ideator's own attempt in right parahippocampal gyrus
396 versus suicide action in left hippocampus⁷². In contrast, parahippocampal gyrus activation
397 was blunted in adult SAs with MDD choosing between a smaller immediate reward versus a

398 larger but delayed reward, especially when the two rewards were more than 1 year apart⁷⁸.
399 Given the role of the parahippocampal gyrus in prospection¹¹⁸, its blunted response to
400 prospects with longer versus shorter delays may represent a neural substrate of impaired
401 prospection in SAs¹¹⁹⁻¹²¹, potentially undermining the deterrents and alternative solutions
402 during a suicidal crisis.

403

404 *Summary*

405 Although structural alterations in the amygdala and the hippocampus have been consistently
406 implicated in mental disorders¹²²⁻¹²⁴, the majority of studies reviewed do not report structural
407 alterations in these regions in relation to STBs. These mixed findings could perhaps be
408 explained if additional involvement of the amygdala and hippocampus in STBs beyond their
409 role in mental disorders is subtle with small effects only apparent in studies with very large
410 sample sizes. This is consistent with a post-hoc power analysis based on observed effect
411 sizes in the largest study on subcortical volumes in STBs to date¹¹¹. Alternatively, mesial
412 temporal structural alterations may only become apparent in specific subgroups of people
413 with STBs. Preliminary evidence suggests serotonergic alterations in the amygdala and
414 hippocampus linked to SA as well as SI across mental disorders, and altered functioning in
415 these regions in relation to increased autobiographical recall of mental pain, blunted
416 immediate reward processing and impaired prospection in patients with SA, although
417 molecular and functional studies focussing on these regions are still scarce.

418

419 **Striatum and Thalamus**

420 The ventral striatum includes the nucleus accumbens and ventral parts of the putamen and
421 caudate¹²⁵ and is a core region of the reward network¹²⁶. Dorsal striatum, including dorsal
422 caudate and putamen, functions include initiating action, inhibitory control, and stimulus-
423 response learning¹²⁷. The striatum projects to the frontal lobe via the thalamus¹²⁵, which is
424 also involved in sensory processing¹²⁸.

425

426 Lower caudate and putamen volumes have been reported in adult SAs with MDD in
427 comparison to MDD non-attempters^{34,129} and HCs¹³⁰, and putamen volumes were negatively
428 associated with impulsivity¹²⁹. Lower putamen binding of the serotonin transporter (5-HTT)
429 was also reported in adult SAs with MDD compared to HCs¹³¹, and was negatively associated
430 with impulsivity¹³². However, striatal 5-HT binding was positively associated with SI in adult
431 SAs with MDD¹³³. With regard to the thalamus, higher volumes were reported in veterans with
432 traumatic brain injury and suicide attempts¹³⁴ and higher 5-HT synthesis reported in adult SAs
433 with a mix of psychiatric diagnoses⁴⁰. However, other studies, including the largest study to
434 date of individuals with STBS (N=451), have not detected associations of striatum and
435 thalamus volume with STBs^{33,38,111}.

436

437 Few fMRI studies have examined striatum and thalamus regions. Positive correlations were
438 observed between intensity of past SI and dorsal striatum responses during cognitive control
439 in adults with MDD or BD with psychotic features⁵². Lower putamen activation in adults with
440 BD during a motor task was associated with higher SI¹³⁵. Higher thalamus activation was
441 observed when viewing knives (versus landscapes)¹³⁶ and higher thalamus activation during
442 response inhibition (go-no-go task) was associated with higher levels of mental pain and
443 suicide intent¹³⁷ in adults with SAs and MDD.

444

445 *Summary*

446 Mixed findings were reported for the involvement of structural alterations in the striatum and
447 thalamus in relation to STBs, with the largest sample to date showing no associations in
448 people with MDD. Increased dorsal striatum responses were found during cognitive control in
449 the absence of performance differences in individuals with SI, suggesting greater
450 engagement of this region to reach similar levels of cognitive control. Higher thalamus
451 activations were reported during emotion processing and inhibition and associated with SI in
452 adult SA with MDD. Structural and 5-HTT alterations in the dorsal striatum specifically linked
453 to impulsivity in adult MDD with SA converge with findings of functional alterations in the
454 dorsal striatum in relation to diminished cognitive and affective control associated with SI. Of
455 note, alterations in the ventral striatum have been proposed to underlie reduced reward
456 anticipation and anhedonia in individuals with STBs¹³⁸. No studies reported ventral striatal
457 *activity*, however, ventral striatal *connectivity* findings during reward processing and under
458 rest are discussed in the “Structural and Functional Connectivity” section below.

459

460 **Posterior Structures**

461 The temporal association cortices are involved in perceptual processing of faces and other
462 complex object features^{139,140}, auditory information and language¹⁴¹. Consistent with this,
463 structural MRI findings in lateral temporal cortex were observed in adult SAs with
464 SZ/SZA/psychotic BD and other disorders, such as mood disorders, in which psychotic
465 misperceptions can be observed. Lower middle and superior temporal gyrus volume was
466 found in adult SAs with primary psychotic disorders^{37,142} and BPD¹⁰⁰, and lower thickness of
467 middle and superior temporal gyri were observed in adult SAs with SZ⁷¹. Lower middle and
468 superior temporal volumes were also associated with high lethality attempts in adults SAs
469 with BPD^{88,100}. Serotonin system studies have yielded various results including lower 5-HTT
470 temporal binding associated with higher impulsivity in adult SAs with different mental
471 disorders¹³², and higher baseline 5-HT1a temporal lobe receptor binding in adults SAs with
472 MDD⁴² associated with higher levels of SI at 2-year follow-up. Functional MRI studies also
473 suggest a role of the lateral temporal lobe in emotion processing in STBs. Specifically,
474 adolescent SAs with MDD showed enhanced right middle temporal gyrus activation during
475 passive viewing of angry, happy and neutral facial expressions⁷⁴ and during recall and re-
476 imagination of suicidal episodes in adult SAs with MDD⁷². SI was associated with increased

477 superior temporal activation during error processing in veterans with traumatic brain injury⁵³.
478 In contrast, lower perfusion in these temporal regions during rest (measured by rCBF) was
479 reported in adults with MDD and SA^{82,143}.

480

481 A few studies implicate other posterior brain regions including posterior cingulate cortex
482 (PCC) and cerebellum in STBs (Tables 1-3). The PCC is implicated in psychological
483 processes that may be linked to STBs, including controlling the vividness of negative mental
484 imagery¹⁴⁴ and enhancing self-referential processing¹⁴⁵. Lower PCC gray matter volume was
485 found in adult SAs with MDD¹⁴⁶. Decreased PCC activation was observed during cognitive
486 control in adult SAs with psychotic mood disorders⁵² and during self-referential processing in
487 adolescent ideators with MDD¹⁴⁷, although adult SAs with depressive disorders, compared to
488 HCs, showed increased PCC response when viewing knives¹³⁶.

489

490 The cerebellum is increasingly recognized for its involvement in emotional processes^{148,149}.
491 Lower volumes of the cerebellum were reported in adult and adolescent SAs with MDD or
492 BD^{36,70,146,150}. Functionally, while adult SAs with MDD showed increased cerebellum activation
493 during recall and re-imagination of their own suicidal episode⁷² and while viewing angry faces,
494 they showed decreased activation while viewing happy faces⁴⁸. Decreased activation was
495 also observed while passively viewing negative emotional pictures in adolescents with a
496 history of SI⁷³. Finally, ketamine-induced reductions in SI were associated with increases in
497 rCMRglu including in cerebellum¹⁵¹.

498

499 *Summary*

500 Lower middle and superior temporal gyri volumes have been reported in 6 studies across a
501 range of mental disorders, and related to high lethality attempts and higher impulsivity.
502 Serotonergic alterations in these regions have not been extensively investigated and
503 directions of reported effects are mixed. Increased activation in middle and superior temporal
504 gyri have also been reported in adults and adolescents with SA and MDD, especially in
505 relation to emotion processing. Preliminary evidence suggests a role for the PCC and
506 cerebellum in STBs, especially in relation to self-referential, and (autobiographical) emotion
507 processing, but studies investigating these regions remain scarce. Of interest, one study
508 suggests that ketamine-induced changes in SI are associated with ketamine-induced
509 increased in cerebellum rCMRglu. This medication-related finding is a potential lead in
510 understanding brain mechanisms that may be helpful targets for suicide prevention
511 interventions, but requires replication.

512

513 **Structural and Functional Connectivity**

514 Disturbances in the structure and function *within* brain regions can result in alterations in brain
515 networks, including the ability of brain regions to coordinate their activity in a system. System
516 dysfunction can also result from abnormalities in the connections between regions.

517 Increasingly, abnormalities in the structural and functional connections *between* brain regions
518 within larger-scale brain networks have been reported in studies of STBs.

519

520 Connections of the medial VPFC with other cortical midline structures (PCC, precuneus) and
521 temporal and parietal regions are implicated in the brain's default mode network and play an
522 important role in self-referential processes, social cognition, autobiographical memory and
523 prospective imaging¹¹⁸. Lower resting functional connectivity was reported in the default mode
524 network in adolescents with MDD and SI¹⁵². These findings are in line with findings of lower
525 resting functional connectivity of rostral ACC with medial OFC, precuneus and temporal pole
526 in adults with MDD and SI¹⁵³ and within the precuneus in young adult SAs with MDD¹⁵⁴.

527 Structurally, lower fractional anisotropy (FA; thought to reflect the structural integrity of white
528 matter and the neuronal connections it contains¹⁵⁵) in the medial VPFC was reported in adult
529 SAs with BD, which was associated with higher motor impulsivity¹⁵⁶. Lower FA was also
530 reported in the ventral cingulum (connecting posterior and temporal default mode network
531 regions) in adults with MDD and SI³⁸. Lower FA of the corpus callosum genu, that provides
532 connections of interhemispheric anterior default mode network regions, was associated with a
533 higher number of SAs in BD, MDD and BPD^{157,158}.

534

535 The limbic network includes the amygdala, medial and lateral OFC, medial temporal regions,
536 thalamus and basal ganglia and is involved in emotional and autonomic processes¹⁵⁹. Using
537 task fMRI, lower amygdala-medial VPFC/rostral PFC connectivity was found in adolescent
538 and young adult BD SAs while viewing happy and neutral facial expression, associated with
539 higher lethality of attempts and current SI³⁶. During rest, greater amygdala connectivity with
540 lateral OFC, insula and middle temporal gyrus was found in adult SAs with MDD, with greater
541 amygdala-parahippocampus connectivity associated with SI¹⁶⁰. These functional connectivity
542 alterations are in line with lower FA in the uncinate fasciculus, that provides major amygdala
543 connections, in adolescents SAs with BD³⁶.

544

545 The medial OFC together with ventral striatum and the ventral tegmental area form core hubs
546 of a reward network, with additional limbic regions, DLPFC and dACC forming a wider reward
547 network subserving reward-related memory and evaluation¹²⁶. An fMRI study investigating
548 connectivity during reward processing showed a positive correlation between SI and
549 connectivity of the left ventral striatum with dACC, DMPFC and DLPFC during loss trials in
550 adults with MDD¹⁶¹. Using resting state fMRI, Kim et al.¹⁶² found reduced connectivity in
551 circuitry resembling this reward network, including the OFC, striatum and thalamus, in adults
552 with MDD and recent (past month) SI. This is in line with findings of lower structural
553 connectivity between VPFC/OFC and striatal regions in adults with MDD and SI (33% also
554 had prior SA)¹⁶³, and between the ACC and OFC (as measured by graph theory) in adult SAs
555 with MDD¹⁶⁴. Lower FA was also reported in the anterior limb of the internal capsule

556 (connecting striatal and thalamic regions with the PFC) in adults with MDD and
557 attempts^{165,166}.

558

559 The left and right DLPFC and DMPFC together with parietal regions comprise a network key
560 in cognitive control of thought, emotion and behavior (executive control network¹⁶⁷). Lower
561 executive control network coherence during rest has been associated with both lifetime SI
562 and past attempts in adolescents with MDD¹⁵², in line with findings of lower DLPFC resting
563 state connectivity in young adult SAs with MDD associated with higher impulsivity¹⁶⁸ and
564 reduced white matter integrity (FA) in the DMPFC in adult SAs with MDD¹⁶⁹.

565

566 Alterations in dACC connectivity have been linked to STBs in the context of conflict-
567 monitoring, with different patterns of dACC connectivity associated with SI versus SA in
568 adults with recent-onset SZ¹⁷⁰. That is, the presence of lifetime SI was *positively* associated
569 with magnitude of functional connectivity of dACC with the precuneus, a core hub of the
570 default mode network. This may suggest a reduced capacity of the dACC to 'switch off'
571 default mode network activity associated with more internally focused attention, when
572 activation of externally focused cognitive processing is required¹⁷⁰. In contrast, history of
573 suicide attempts was *negatively* associated with dACC connectivity with DPFC (BA9, 8,
574 lateral BA10), lateral VPFC (BA45), PCC, parietal regions (BA7,40) and superior and middle
575 temporal gyri (BA22, 39, 40)¹⁷⁰. These findings may suggest that SI and SA have divergent
576 bases in dACC connectivity with default mode network versus lateral PFC circuitries,
577 respectively, in the context of monitoring conflict in adult SAs with recent-onset SZ. This is
578 supported by findings of abnormal conflict-related dACC connectivity with VLPFC, OFC,
579 insula and striatum associated with SI intensity, but altered dACC connectivity with DLPFC
580 and frontal motor regions associated with past suicide attempts in adults with MDD or BD with
581 psychotic features¹⁷¹. In addition, decreased functional connectivity of the dACC with bilateral
582 insula while viewing angry faces was also reported in adolescent SAs with MDD⁷⁴.

583 Connectivity between the dACC and insula plays an important role in detecting salient internal
584 and external stimuli to guide behavior¹⁷² and has been implicated in the anticipation of
585 aversive experiences, especially in depressed individuals¹⁷³. Reduced connectivity between
586 the dACC and bilateral insula may indicate inefficient strategies to process the salience of,
587 and select contextually appropriate behavioral responses to, negative emotional stimuli.

588

589 *Summary*

590 Emerging evidence suggests that resting state functional connectivity in the default mode
591 network, and white matter tracts connecting regions within this network, play a role in both SI
592 and SA across mental disorders. Functional and structural connectivity alterations in the
593 affective network have also been associated with SI and SA, as well as lethality of attempts,
594 both during emotion processing and during rest. Connectivity abnormalities in the reward
595 network have mostly been examined in adults with MDD, and both structural and functional

596 connectivity changes in regions of this network have been associated with SI and SA in this
597 group. Connectivity changes within and between regions implied in the cognitive control
598 network has been less extensively studied in relation to STBs, and the few studies conducted
599 suggest a role of lower resting state functional connectivity within this network in STBs in
600 adolescents and young adults with MDD. In addition, functional connectivity of dACC is
601 implicated in STBs, but with divergent connectivity patterns related to SI versus SA.

602
603

604 **4. DISCUSSION**

605 While the literature primarily includes cross-sectional studies with small sample sizes,
606 differing clinical populations and a wide range of imaging methods, there is emerging
607 convergence in the brain regions implicated in STBs. Taken together with the recent
608 increased momentum in studies on STBs (Figure 1), it is very hopeful that significant
609 advances in our understanding of brain mechanisms contributing to STBs are on the horizon.
610 A critical frontier is to identify markers for elevated risk, especially short term risk in the
611 transition from SI to attempt. While the majority of studies to date were on SAs, some studies
612 specifically investigated associations with SI, and brain regions found to be associated with
613 STBs have known roles in processes thought to contribute to STBs. Below we briefly
614 summarize the most convergent findings from the literature reviewed above, and propose
615 directions for future neuroimaging studies on the neurobiology of STB.

616

617 **A Tentative Brain Model of STBs**

618 Figure 3 summarizes convergent findings emerging from the reviewed literature on brain
619 alterations associated with STBs. Abnormalities in an extended VPFC system, including
620 regions of the default mode, affective and reward networks such as the ventral and rostral
621 ACC, insula, medial and lateral OFC, mesial temporal regions, ventral striatum and posterior
622 structures (lateral temporal, PCC, precuneus, cerebellum), and in the connections among
623 these regions, may be important in the excessive negative and blunted positive internal states
624 that can stimulate SI. This is in line with the well-established role of this extended VPFC
625 system in functions implicated in SI including appraisal of internally-generated emotions, self-
626 referential processing, recall of emotional episodic memories, imagining future positive and
627 negative events, valuation of rewards, and integrating environmental stimuli to modulate
628 subjective emotional states^{16,17,67,118,126,148}. A more lateral and dorsal system, including
629 DMPFC, DLPFC and dACC, and together with the IFG, may facilitate suicide behaviors
630 through their role in cognitive control of thought, emotion and behavior as well as cognitive
631 flexibility, complex decision-making (e.g. valuation of different decision options) and
632 planning^{16,17}. A combination of VPFC and DPFC/IFG system disturbances may lead to very
633 high risk circumstances in which SI may convert to lethal actions via decreased top-down
634 inhibition of behavior or maladaptive, and inflexible decision-making and planning.

635

636 Alterations in the connections between these systems may contribute to the transition from
637 suicidal thoughts to behaviors. For example, the dACC has connections to both the
638 “emotional” ventral limbic system and the “cognitive” dorsal prefrontal system¹⁷⁴. Consistent
639 with this, findings suggest differential connectivity of the dACC in relation to SI versus a
640 history of attempt, with dACC connectivity with VPFC, insula and striatal regions primarily
641 associated with SI and dACC connectivity with DPFC regions associated with suicide
642 attempts. The insula is also implicated in both SI and attempts and may perhaps be important
643 for the transition from SI to attempt. Although the involvement of the insula in STBs has had
644 little direct research focus, its critical involvement in interoceptive processing, detecting
645 salient internal and external stimuli, experiencing emotion and self-awareness^{175–177} suggests
646 an important role of the insula in SI. Additionally, the insula is implicated in disconnection from
647 bodily experiences, which in turn may lower the threshold to engaging in behaviors that harm
648 the body (in line with the acquired capability theory¹⁷⁸) thus suggesting a role of the insula in
649 suicide behaviors. In line with important roles of dACC and insula circuitry (as part of the
650 ‘salience network’) in mediating or switching between the extended medial VPFC (default
651 mode, affective, reward) systems and the DPFC/IFG (executive control) system^{179–181}, the
652 dACC and insula may represent integral hubs that facilitate the transition from SI to attempt.
653 However, this suggestion of the dACC and insula’s ability of mediating dynamic interactions
654 between the medial VPFC and DPFC/IFG systems and through these interactions playing a
655 role in the transition from SI to attempt remains highly speculative and will need to be
656 confirmed in future, preferably longitudinal, studies on the neurobiology of STB.

657

658 **Future directions**

659 The literature reviewed underscores the need for future studies to include larger sample sizes
660 and careful attention to developmental stages. In addition, studies employing longitudinal
661 designs are critically needed to identify risk markers for future suicide attempts in order to
662 develop improved preventive strategies. Recent preliminary evidence, from a rare longitudinal
663 study of adolescents and young adults with mood disorders over about 3 years, showed that
664 those with future attempts had lower VPFC volume and decreased FA in VPFC and DPFC
665 connections¹⁸², suggesting that these may be potential predictors for STBs already present in
666 adolescence. Longitudinal study over short time intervals are largely absent from the literature
667 and are critically-needed to assess proximal suicide risk.

668

669 Furthermore, studies focusing on identifying brain alterations that predict or fluctuate with
670 changes in STBs following treatment could provide urgently needed biomarkers for response
671 of STBs to existing interventions and could help develop novel treatments specifically
672 targeting these biomarkers. Preliminary evidence has implicated brain circuitry that may
673 mediate the reduction of suicide risk by treatments such as lithium and ketamine^{183–185}. For
674 example, ketamine-induced reductions in SI were associated with increases in baseline
675 rCMRglu in a cluster including the cerebellum and occipital cortex¹⁵¹. In addition, in 57 adults

676 with BD with and without a history of attempt, lowest DLPFC, OFC, ACC, superior temporal
677 cortex, parietal and occipital cortex volumes were observed in SAs off-lithium, followed by
678 SAs and non-attempters on lithium, with the largest volumes in people with non-attempting
679 BD patients on lithium³⁵. However, the study of other pharmacological and non-
680 pharmacological treatments that can directly (e.g. deep brain stimulation) or indirectly target
681 the involved circuitry is needed.

682

683 Although adolescents have been less studied than adults, some findings have been similar.
684 For instance, alterations in structure of the VMPFC, VLPFC, DLPFC, DMPFC, ACC, lateral
685 temporal regions and parahippocampal gyrus were observed in both adolescents and adults
686 with SA. Furthermore, lower connectivity in regions related to the default mode network
687 during rest have been consistently reported across adolescents, young adults and adults with
688 SI and/or SA. Greater DLPFC activity in response to angry faces was also observed in both
689 adolescents and adults with STBs. The highly limited number of studies in adolescents and a
690 lack of different life stages in single studies prevents drawing conclusions about the overlap in
691 functional brain alterations across different stages of life and brain maturation. Thus, there is
692 some converging evidence across adolescent and adult samples, however, not all adult
693 findings have been observed in adolescents (see Tables 1-3 for a complete overview). This
694 may at least in part be due to the continued maturation of involved brain systems so that not
695 all features may be expressed until adulthood^{186,187}.

696

697 Most studies only included a single disorder, impeding conclusions around shared versus
698 unique neural substrates of STBs across different mental disorders. Different studies
699 employed different in- and exclusion criteria, imaging methods and STB assessments.
700 Nonetheless, gray matter alterations in the VLPFC, DLPFC, ACC, and insula have been
701 consistently reported across the diagnostic categories reviewed (i.e., MDD, BPD, BD, SZ),
702 while lateral temporal alterations were more uniquely observed in psychotic disorders and
703 BPD. Reduced white matter integrity in ventral PFC regions was reported in relation to STBs
704 in both BD and MDD and lower FA in the corpus callosum was associated with a higher
705 number of attempts across BD, MDD and BPD. Functional brain alterations in relation to
706 emotion processing and regulation was investigated only in MDD, BD and BPD, with
707 consistent findings of increased VLPFC activation in response to negative emotional stimuli
708 across MDD and BPD, while the only study of adolescents with BD focused on amygdala-
709 PFC connectivity³⁶. Higher DLPFC activation during cognitive control tasks, in the absence of
710 task performance differences, was also consistently reported across MDD, BD, SZ and
711 PTSD. In contrast, higher versus lower IFG activation during cognitive control was observed
712 in mood disorders versus SZ, respectively. Alterations in dACC connectivity with default
713 mode, affective and reward network related regions in relation to SI, and alterations in dACC
714 connectivity with DLPFC in relation to SA during cognitive control, has been observed across
715 MDD, BD and SZ. Other functional domains such as reward processing, decision making,

716 social exclusion and self-referential processing, as well as resting state fMRI, have only been
717 investigated in a single disorder, i.e. depression. In one of the few studies to assess SAs
718 across MDD and BD (published subsequent to our literature search), VPFC gray matter
719 volume reductions and uncinate fasciculus FA reductions were common to both disorders,
720 suggesting that they may be important in risk for attempts across mood disorders¹⁸⁸.
721 Preliminary findings of that study also indicated there may be differences in involved regions
722 in SAs between the disorders, with greater uncinate involvement in attempters in BD and
723 dorsal frontal white matter in MDD. These findings suggest both transdiagnostic and unique
724 gray and white matter targets for suicide prevention across mental disorders.

725

726 The examination of sex differences in STBs is a major gap in the neuroimaging literature and
727 a critical area for study. There are well-established sex-dependent features known in STBs¹⁸⁹,
728 such as a higher rate of attempts in females and a higher lethality of attempts in males¹⁹⁰⁻¹⁹².
729 Moreover, increased death rates by suicide between 1999 and 2017 as reported by the CDC
730 showed the rate of increase was substantially higher for females than males (53% and 26%
731 respectively). Female suicide rate increases were particularly high in youth/early adulthood
732 (ages 10-24) and middle age (45-64 years), when females have their highest risk, while the
733 highest risk for males is age 75 and older¹⁹³. Of the studies in this review, 11 (8.4%) had
734 exclusively female and 15 (11.5%) exclusively male samples. Authors of six studies (4.6%)
735 commented on this homogeneity as a limitation, and 14 (10.7%) provided a rationale for
736 single sex studies. These included the need to avoid potential confounds of previously
737 reported sex-based effects e.g., previous reports of sex differences in cortical responses with
738 similar fMRI tasks (n=5) and corpus callosum structure (n=2); high male representation in the
739 group under investigation, such as military veterans (n=3); female-skewed prevalence of BPD
740 (n=1); males and females imaged on different scanners (n=1); small number of male SAs
741 (n=1); and an attempt to replicate previous work (n=1). Ninety studies (68.7%) included sex
742 as a covariate, controlling for potential differences. A small percentage of studies (13.8%)
743 included sex as a variable to assess potential interactions with STBs. Five studies (3.8%)
744 reported sex-related findings, though no relationship with STBs^{37,147,194-197} and one study
745 (0.8%) reported more females than males studied were SAs but reported no neuroimaging-
746 related findings¹⁹⁸. Chase and colleagues¹⁹⁹, in a study that controlled for sex, noted that one
747 participant identified as transmale. Careful consideration is warranted in how sex and gender
748 are evaluated and categorized for analyses, including the importance of allowing subjects to
749 identify by gender and this self-identification of gender is considered in all studies. This is
750 particularly important because transgender and sexual-minority individuals are at increased
751 risk for STBs and death by suicide^{200,201}. Collectively, these STB neuroimaging research data
752 highlight the urgent need for future work on sex and gender.

753

754 Integration of neuroimaging research across differential mechanistic levels including genetic,
755 molecular, social, and environmental risk factors will be crucial to the elucidation of a holistic

756 STB pathophysiology and mechanisms associated with vulnerability and resilience and thus
757 tailored intervention development. Sophisticated analytic methods, such as machine learning
758 techniques⁶⁰, can be utilized to allow imaging risk markers to be identified at the level of the
759 individual instead of at a group level, a key ingredient for clinically viable biomarkers²⁰² and
760 precision medicine. In addition, the use of high resolution ultra-high field strength MRI
761 methods and more specific functional neuroimaging tasks may further enhance ability to
762 parse roles of specific brain regions, such as PFC subregions²⁰³. Molecular imaging has
763 produced important leads, such as in serotonergic and inflammation mechanisms (Table 2),
764 consistent with post-mortem, genetic and peripheral biomarker studies implicating an
765 important role of the serotonergic system and inflammatory mechanisms in STBs^{204,205}.
766 Imaging of molecular mechanisms other than serotonin remains scarce, with for example only
767 four magnetic resonance spectroscopy (¹H-MRS) studies examining glutamatergic and
768 Gamma-Aminobutyric Acid (GABA)-ergic mechanisms that have been published to date
769 (Table 2). Given post-mortem findings of altered glutamatergic and GABA-ergic gene
770 expression and receptor availability in suicide victims mainly in the prefrontal cortex, ACC and
771 hippocampus^{206–208}, future ¹H-MRS should clarify the role of these neurotransmitters in STBs
772 in vivo. In addition, a generation of new methods to identify other currently implicated and
773 novel molecular mechanisms is needed. For example, a link between oxytocin and SI was
774 recently suggested²⁰⁹.

775

776 Future studies should also further investigate the more elusive subjective aspects of suicide
777 risk, such as SI, as well as implicated psychological constructs such as hopelessness,
778 rumination and anhedonia^{210–212}, which may be relevant across diagnoses. Similarly, social
779 experiences such as childhood maltreatment and peer bullying form a strong prelude to STB
780 in later life^{209,213,214}, and impact on the neural structures implicated in STB (e.g. DMPFC
781 structure and function^{215,216}). Therefore, adverse experiences should be taken into account in
782 future studies on the neurobiology of STB.

783

784 Links identified between neuroimaging measures and behaviors outside of the scanner could
785 facilitate the development of less invasive and more easily and widely disseminated risk
786 detection methods. Furthermore, investigations of larger samples can be facilitated by
787 international collaborations that pool existing data across many different samples, such as the
788 MQ HOPES (<https://www.mqmentalhealth.org/research/profiles/overcome-and-predict-the-emergence-of-suicide>) and ENIGMA STB (<http://enigma.ini.usc.edu/ongoing/enigma-stb/>)
789 consortia. These initiatives represent cost-effective ways to substantially increase statistical
790 power, which could provide more robust and reliable findings²¹⁷, and ability to study age,
791 gender and sex effects, as well as unique and shared mechanisms associated with STBs
792 across different mental disorders. Furthermore, large combined datasets will provide a unique
793 opportunity to identify and test reproducibility of different pathways to suicide that may differ
794 across many parameters including a range from the impulsive to the highly planned. Such
795

796 efforts will need to take into account the variety of assessment methods used for STBs across
797 studies. One way to address with this important challenge is through the examination of the
798 presence versus absence of suicidal behavior (attempts), and/or suicidal ideation (with or
799 without intent and/or a plan) across assessment types (see for example Renteria et al.¹¹¹).
800 Another approach could involve standardization of scores across different instruments by
801 developing a common metric²¹⁸. Standardization of STB assessment across future studies
802 would significantly facilitate the sharing of data, and thereby advancing our understanding of
803 brain-based STB vulnerability.

804

805 **Conclusions**

806 More than two decades of neuroimaging studies on STBs suggests a transdiagnostic model
807 for STBs in which an extended VPFC system may be important in the excessive negative and
808 blunted positive internal states that can stimulate SI and a DPFC/IFG system that may
809 facilitate suicide attempt behaviors. Interactions between these systems are likely important in
810 the transition from ideation to attempt, perhaps mediated by dACC and insula regions, but
811 require further investigation. With the exponential growth of research on STBs, including the
812 initiation of large global efforts, it is hopeful that suicide prevention will soon be more
813 effectively targeted, reducing the tragic loss of life to suicide.

Acknowledgments

This work was supported by the MQ Brighter Futures Award MQBFC/2 (ALvH, HB, LS), by the National Institute of Mental Health of the National Institutes of Health under Award Number R01MH117601 (LS), RC1MH088366 (HPB), R01MH113230 (HPB), R61MH111929 (HPB), T32MH014276 (ETCL), and T32DA022975 (ETCL). LS is supported by a NHMRC Career Development Fellowship (1140764). ALVH is supported by a Royal Society Dorothy Hodgkin Fellowship (DH15017). LAA is supported by the American Foundation for Suicide Prevention, Brain and Behavior Foundation/NARSAD, Robert E. Leet and Clara M. Guthrie Patterson Trust, and Department of Veterans Affairs. HPB is supported by the American Foundation for Suicide Prevention and For the Love of Travis Foundation.

Conflicts of interest

The authors declare no competing financial interests. HPB received an honorarium for a talk at Aetna.

REFERENCES

- 1 World Health Organization. Public health action for the prevention of suicide: A framework. Geneva, Switzerland, 2012.
- 2 Substance Abuse and Mental Health Services Administration, Results from the 2013 National Survey on Drug Use and Health: Mental Health Findings, NSDUH Series H-49, HHS Publication No. (SMA) 14-4887. Rockville, MD, 2014.
- 3 Eaton D, Kann L, Kinchen S, Shanklin S, Flint K, Hawkins J *et al.* Youth risk behavior surveillance—United States, 2011. *Morbidity and Mortality Weekly Report. Surveillance Summaries. Surveill Summ* 2012; **61**: 1–162.
- 4 Curtin SC, Warner M, Hedegaard H. NCHS Data Brief No. 241: Increase in Suicide in the United States, 1999 – 2014 Key findings. 2016. <http://www.cdc.gov/nchs/products/databriefs/db241.htm>.
- 5 Franklin JC, Ribeiro JD, Fox KR, Bentley KH, Kleiman EM, Huang X *et al.* Risk factors for suicidal thoughts and behaviors: A meta-analysis of 50 years of research. *Psychol Bull* 2017; **143**: 187–232.
- 6 Balcioglu YH, Kose S. Neural substrates of suicide and suicidal behaviour: from a neuroimaging perspective. *Psychiatry Clin Psychopharmacol* 2018; **28**: 314–328.
- 7 Bani-Fatemi A, Tasmim S, Graff-Guerrero A, Gerretsen P, Strauss J, Kolla N *et al.* Structural and functional alterations of the suicidal brain: An updated review of neuroimaging studies. *Psychiatry Res - Neuroimaging* 2018; **278**: 77–91.
- 8 Domínguez-Baleón C, Gutiérrez-Mondragón LF, Campos-González AI, Rentería ME. Neuroimaging studies of suicidal behavior and non-suicidal self-injury in psychiatric patients: A systematic review. *Front Psychiatry* 2018; **9**. doi:10.3389/fpsy.2018.00500.
- 9 Jollant F, Lawrence NL, Olié E, Guillaume S, Courtet P. The suicidal mind and brain: A review of neuropsychological and neuroimaging studies. *World J Biol Psychiatry* 2011; **12**: 319–339.
- 10 van Heeringen K, Bijttebier S, Desmyter S, Vervaeke M, Baeken C. Is there a neuroanatomical basis of the vulnerability to suicidal behavior? A coordinate-based meta-analysis of structural and functional MRI studies. *Front Hum Neurosci* 2014; **8**: 1–8.
- 11 Hamza CA, Stewart SL, Willoughby T. Examining the link between nonsuicidal self-injury and suicidal behavior: A review of the literature and an integrated model. *Clin Psychol Rev* 2012; **32**: 482–495.
- 12 Chesney E, Goodwin GM, Fazel S. Risks of all-cause and suicide mortality in mental disorders: A meta-review. *World Psychiatry* 2014; **13**: 153–160.
- 13 Turecki G, Brent DA. Suicide and suicidal behaviour. *Lancet* 2016; **387**: 1227–1239.
- 14 Hawton K, van Heeringen K. Suicide. *Lancet* 2009; **373**: 1372–1381.
- 15 Dombrovski AY, Hallquist MN. The decision neuroscience perspective on suicidal behavior: Evidence and hypotheses. *Curr Opin Psychiatry* 2017; **30**: 7–14.

- 16 Dixon ML, Thiruchselvam R, Todd R, Christoff K. Emotion and the prefrontal cortex: An integrative review. *Psychol Bull* 2017; **143**: 1033–1081.
- 17 Mitchell DGV. The nexus between decision making and emotion regulation: A review of convergent neurocognitive substrates. *Behav Brain Res* 2011; **217**: 215–231.
- 18 Murray RJ, Schaer M, Debbané M. Degrees of separation: A quantitative neuroimaging meta-analysis investigating self-specificity and shared neural activation between self- and other-reflection. *Neurosci Biobehav Rev* 2012; **36**: 1043–1059.
- 19 Bechara A, Damasio AR. The somatic marker hypothesis: A neural theory of economic decision. *Games Econ Behav* 2005; **52**: 336–372.
- 20 D'Argembeau A, Van der Linden M. Predicting the phenomenology of episodic future thoughts. *Conscious Cogn* 2012; **21**: 1198–1206.
- 21 Levy DJ, Glimcher PW. The root of all value: a neural common currency for choice. *Curr Opin Neurobiol* 2012; **22**: 1027–1038.
- 22 Urry HL, Reekum CM van, Johnstone T, Kalin NH, Thurow ME, Schaefer HS *et al*. Amygdala and Ventromedial Prefrontal Cortex Are Inversely Coupled during Regulation of Negative Affect and Predict the Diurnal Pattern of Cortisol Secretion among Older Adults. *J Neurosci* 2006; **26**: 4415–4425.
- 23 Mitchell JP, Banaji MR, Macrae CN. The link between social cognition and self-referential thought in the medial prefrontal cortex. *J Cogn Neurosci* 2005; **17**: 1306–1315.
- 24 Drevets WC. Neuroimaging and neuropathological studies of depression: Implications for the cognitive-emotional features of mood disorders. *Curr Opin Neurobiol* 2001; **11**: 240–249.
- 25 Jollant F, Lawrence NL, Olié E, Guillaume S, Courtet P. The suicidal mind and brain: A review of neuropsychological and neuroimaging studies. *World J Biol Psychiatry* 2011; **12**: 319–339.
- 26 Bratman GN, Hamilton JP, Hahn KS, Daily GC, Gross JJ. Nature experience reduces rumination and subgenual prefrontal cortex activation. *Proc Natl Acad Sci* 2015; **112**: 8567–8572.
- 27 Bertossi E, Ciaramelli E. Ventromedial prefrontal damage reduces mind-wandering and biases its temporal focus. *Soc Cogn Affect Neurosci* 2016; **11**: 1783–1791.
- 28 Burke TA, Connolly SL, Hamilton JL, Stange JP, Lyn Y, Alloy LB. Cognitive Risk and Protective Factors for Suicidal Ideation: A Two Year Longitudinal Study in Adolescence. *J Abnorm Child Psychol* 2016; **44**: 1145–1160.
- 29 Ducasse D, Loas G, Dassa D, Gramaglia C, Zeppegno P, Guillaume S *et al*. Anhedonia is associated with suicidal ideation independently of depression: A meta-analysis. *Depress Anxiety* 2018; **35**: 382–392.
- 30 Miranda R, Valderrama J, Tsypes A, Gadol E, Gallagher M. Cognitive inflexibility and suicidal ideation: Mediating role of brooding and hopelessness. *Psychiatry Res* 2013; **210**: 174–181.

- 31 Smith JM, Alloy LB, Abramson LY. Cognitive Vulnerability to Depression, Rumination, Hopelessness, and Suicidal Ideation: Multiple Pathways to Self-Injurious Thinking. *Suicide Life-Threatening Behav* 2006; **36**: 443–454.
- 32 Ding Y, Lawrence N, Olié E, Cyprien F, Le Bars E, Bonafé A *et al*. Prefrontal cortex markers of suicidal vulnerability in mood disorders: A model-based structural neuroimaging study with a translational perspective. *Transl Psychiatry* 2015; **5**: e516.
- 33 Gosnell SN, Velasquez KM, Molfese DL, Molfese PJ, Madan A, Fowler JC *et al*. Prefrontal cortex, temporal cortex, and hippocampus volume are affected in suicidal psychiatric patients. *Psychiatry Res - Neuroimaging* 2016; **256**: 50–56.
- 34 Wagner G, Koch K, Schachtzabel C, Schultz CC, Sauer H, Schlösser RG. Structural brain alterations in patients with major depressive disorder and high risk for suicide: Evidence for a distinct neurobiological entity? *Neuroimage* 2011; **54**: 1607–1614.
- 35 Benedetti F, Radaelli D, Poletti S, Locatelli C, Falini A, Colombo C *et al*. Opposite effects of suicidality and lithium on gray matter volumes in bipolar depression. *J Affect Disord* 2011; **135**: 139–147.
- 36 Johnston JAY, Wang F, Liu JJ, Blond BN, Wallace A, Liu JJ *et al*. Multimodal neuroimaging of frontolimbic structure and function associated with suicide attempts in adolescents and young adults with bipolar disorder. *Am J Psychiatry* 2017; **174**: 667–675.
- 37 Giakoumatos CI, Mathew IT. Are Structural Brain Abnormalities Associated With Suicidal Behaviour in Patients With Psychotic Disorder? 2014; **47**: 1389–1395.
- 38 Taylor WD, Boyd B, McQuoid DR, Kudra K, Saleh A, MacFall JR. Widespread white matter but focal gray matter alterations in depressed individuals with thoughts of death. *Prog Neuro-Psychopharmacology Biol Psychiatry* 2015; **62**: 22–28.
- 39 Arango V, Underwood MD, Mann JJ. Serotonin brain circuits involved in major depression and suicide. *Prog Brain Res* 2002; **136**: 443–453.
- 40 Leyton M, Paquette V, Gravel P, Rosa-Neto P, Weston F, Diksic M *et al*. α -[11C]methyl-L-tryptophan trapping in the orbital and ventral medial prefrontal cortex of suicide attempters. *Eur Neuropsychopharmacol* 2006; **16**: 220–223.
- 41 Sullivan GM, Oquendo MA, Milak M, Miller JM, Burke A, Ogden RT *et al*. Positron emission tomography quantification of serotonin1A receptor binding in suicide attempters with major depressive disorder. *JAMA Psychiatry* 2015; **72**: 169–178.
- 42 Oquendo MA, Galfalvy H, Sullivan GM, Miller JM, Milak MM, Elizabeth Sublette M *et al*. Positron emission tomographic imaging of the serotonergic system and prediction of risk and lethality of future suicidal behavior. *JAMA Psychiatry* 2016; **73**: 1048–1055.
- 43 Parsey R V., Oquendo MA, Ogden RT, Olvet DM, Simpson N, Huang YY *et al*. Altered serotonin 1A binding in major depression: A [carbonyl-C-11] WAY100635 positron emission tomography study. *Biol Psychiatry* 2006; **59**: 106–113.
- 44 Miller JM, Everett BA, Oquendo MA, Ogden RT, Mann JJ, Parsey R V. Positron Emission Tomography Quantification of Serotonin Transporter Binding in Medication-

- Free Bipolar Disorder. *Synapse* 2016; **70**: 24–32.
- 45 Mann JJ, Metts A V., Ogden RT, Mathis CA, Rubin-Falcone H, Gong Z *et al.* Quantification of 5-HT1A and 5-HT2A receptor Binding in Depressed Suicide Attempters and Non-Attempters. *Arch Suicide Res* 2018; **1118**: 1–12.
- 46 Cannon DM, Ichise M, Fromm SJ, Nugent AC, Rollis D, Gandhi SK *et al.* Serotonin Transporter Binding in Bipolar Disorder Assessed using [11C]DASB and Positron Emission Tomography. *Biol Psychiatry* 2006; **60**: 207–217.
- 47 Olié E, Ding Y, Le Bars E, de Champfleury NM, Mura T, Bonafé A *et al.* Processing of decision-making and social threat in patients with history of suicidal attempt: A neuroimaging replication study. *Psychiatry Res - Neuroimaging* 2015; **234**: 369–377.
- 48 Jollant F, Lawrence NS, Giampietro V, Brammer MJ, Fullana MA, Drapier D *et al.* Orbitofrontal cortex response to angry faces in men with histories of suicide attempts. *Am J Psychiatry* 2008; **165**: 740–748.
- 49 Vanyukov PM, Szanto K, Siegle GJ, Hallquist MN, Reynolds CF, Aizenstein HJ *et al.* Impulsive traits and unplanned suicide attempts predict exaggerated prefrontal response to angry faces in the elderly. *Am J Geriatr Psychiatry* 2015; **23**: 829–839.
- 50 Silvers JA, Hubbard AD, Chaudhury S, Biggs E, Shu J, Grunebaum MF *et al.* Suicide attempters with Borderline Personality Disorder show differential orbitofrontal and parietal recruitment when reflecting on aversive memories. *J Psychiatr Res* 2016; : 71–78.
- 51 Zhang R, Geng X, Lee TMC. Large-scale functional neural network correlates of response inhibition: an fMRI meta-analysis. *Brain Struct Funct* 2017; **222**: 3973–3990.
- 52 Minzenberg MJ, Lesh TA, Niendam TA, Yoon JH, Cheng Y, Rhoades RN *et al.* Control-related frontal-striatal function is associated with past suicidal ideation and behavior in patients with recent-onset psychotic major mood disorders. *J Affect Disord* 2015; **188**: 202–209.
- 53 Matthews S, Spadoni A, Knox K, Strigo I, Simmons A. Combat-exposed war veterans at risk for suicide show hyperactivation of prefrontal cortex and anterior cingulate during error processing. *Psychosom Med* 2012; **74**: 471–475.
- 54 Minzenberg MJ, Lesh TA, Niendam TA, Yoon JH, Rhoades RN, Carter CS. Frontal cortex control dysfunction related to long-term suicide risk in recent-onset schizophrenia. *Schizophr Res* 2014; **157**: 19–25.
- 55 Fradkin Y, Khadka S, Bessette KL, Stevens MC. The relationship of impulsivity and cortical thickness in depressed and non-depressed adolescents. *Brain Imaging Behav* 2017; **11**: 1515–1525.
- 56 Winkler AM, Kochunov P, Blangero J, Almasy L, Zilles K, Fox PT *et al.* Cortical thickness or grey matter volume? The importance of selecting the phenotype for imaging genetics studies. *Neuroimage* 2010; **53**: 1135–1146.
- 57 Rakic P. A small step for the cell, a giant leap for mankind: a hypothesis of neocortical expansion during evolution. *Trends Neurosci* 1995; **18**: 383–388.

- 58 Wagner G, Schultz CC, Koch K, Schachtzabel C, Sauer H, Schlösser RG. Prefrontal cortical thickness in depressed patients with high-risk for suicidal behavior. *J Psychiatr Res* 2012; **46**: 1449–1455.
- 59 Rakic P. Specification of Cerebral Cortical Areas. *Science* 1988; **241**: 170–176.
- 60 Just MA, Pan L, Cherkassky VL, McMakin D, Cha C, Nock MK *et al.* Machine learning of neural representations of suicide and emotion concepts identifies suicidal youth. *Nat Hum Behav* 2017; **1**: 911–919.
- 61 Fellows LK, Farah MJ. Different underlying impairments in decision-making following ventromedial and dorsolateral frontal lobe damage in humans. *Cereb Cortex* 2005; **15**: 58–63.
- 62 Richard-Devantoy S, Berlim MT, Jollant F. A meta-analysis of neuropsychological markers of vulnerability to suicidal behavior in mood disorders. *Psychol Med* 2014; **44**: 1663–1673.
- 63 Andrews-Hanna JR, Smallwood J, Spreng RN. The default network and self-generated thought: Component processes, dynamic control, and clinical relevance. *Ann N Y Acad Sci* 2014; **1316**: 29–52.
- 64 Wagner DD, Haxby J V., Heatherton TF. The representation of self and person knowledge in the medial prefrontal cortex. *Wiley Interdiscip Rev Cogn Sci* 2012; **3**: 451–470.
- 65 Miller EK, Cohen JD. An integrative theory of prefrontal cortex function. *Annu Rev Neurosci* 2001; **24**: 167–202.
- 66 Rive MM, Van Rooijen G, Veltman DJ, Mary ML, Schene AH, Ruhé HG. Neural correlates of dysfunctional emotion regulation in major depressive disorder. A systematic review of neuroimaging studies. *Neurosci Biobehav Rev* 2013; **37**: 2529–2553.
- 67 Petrides M. Lateral prefrontal cortex: Architectonic and functional organization. *Philos Trans R Soc B Biol Sci* 2005; **360**: 781–795.
- 68 Baird B, Smallwood J, Gorgolewski KJ, Margulies DS. Medial and Lateral Networks in Anterior Prefrontal Cortex Support Metacognitive Ability for Memory and Perception. *J Neurosci* 2013; **33**: 16657–16665.
- 69 De Martino B, Fleming SM, Garrett N, Dolan RJ. Confidence in value-based choice. *Nat Neurosci* 2013; **16**: 105–110.
- 70 Hwang JP, Lee TW, Tsai SJ, Chen TJ, Yang CH, Lirng JF *et al.* Cortical and subcortical abnormalities in late-onset depression with history of suicide attempts investigated with MRI and voxel-based morphometry. *J Geriatr Psychiatry Neurol* 2010; **23**: 171–184.
- 71 Besteher B, Wagner G, Koch K, Schachtzabel C, Reichenbach JR, Schlösser R *et al.* Pronounced prefronto-temporal cortical thinning in schizophrenia: Neuroanatomical correlate of suicidal behavior? *Schizophr Res* 2016; **176**: 151–157.
- 72 Reisch T, Seifritz E, Esposito F, Wiest R, Valach L, Michel K. An fMRI study on mental

- pain and suicidal behavior. *J Affect Disord* 2010; **126**: 321–325.
- 73 Miller AB, McLaughlin KA, Busso DS, Brueck S, Peverill M, Sheridan MA. Neural Correlates of Emotion Regulation and Adolescent Suicidal Ideation. *Biol Psychiatry Cogn Neurosci Neuroimaging* 2018; **3**: 125–132.
- 74 Pan LA, Hassel S, Segreti AM, Nau SA, Brent DA, Phillips ML. Differential patterns of activity and functional connectivity in emotion processing neural circuitry to angry and happy faces in adolescents with and without suicide attempt. *Psychol Med* 2013; **43**: 2129–2142.
- 75 Hagan CR, Joiner TE. The Indirect Effect of Perceived Criticism on Suicide Ideation and Attempts. *Arch Suicide Res* 2017; **21**: 438–454.
- 76 Hames JL, Rogers ML, Silva C, Ribeiro JD, Teale NE, Joiner TE. A Social Exclusion Manipulation Interacts with Acquired Capability for Suicide to Predict Self-Aggressive Behaviors. *Arch Suicide Res* 2018; **22**: 32–45.
- 77 Tsutsui KI, Grabenhorst F, Kobayashi S, Schultz W. A dynamic code for economic object valuation in prefrontal cortex neurons. *Nat Commun* 2016; **7**. doi:10.1038/ncomms12554.
- 78 Vanyukov PM, Szanto K, Hallquist MN, Siegle GJ, Reynolds CF, Forman SD *et al*. Paralimbic and lateral prefrontal encoding of reward value during intertemporal choice in attempted suicide. *Psychol Med* 2016; **46**: 381–391.
- 79 Dombrovski AY, Szanto K, Siegle GJ, Wallace ML, Forman SD, Sahakian B *et al*. Lethal forethought: Delayed reward discounting differentiates high- and low-lethality suicide attempts in old age. *Biol Psychiatry* 2011; **70**: 138–144.
- 80 Minzenberg MJ, Lesh T, Niendam T, Yoon JH, Cheng Y, Rhoades RN *et al*. Frontal motor cortex activity during reactive control is associated with past suicidal behavior in recent-onset schizophrenia. *Crisis* 2015; **36**: 363–370.
- 81 Lee KH, Pluck G, Lekka N, Horton A, Wilkinson ID, Woodruff PWR. Self-harm in schizophrenia is associated with dorsolateral prefrontal and posterior cingulate activity. *Prog Neuro-Psychopharmacology Biol Psychiatry* 2015; **61**: 18–23.
- 82 Willeumier K, Taylor D V., Amen DG. Decreased cerebral blood flow in the limbic and prefrontal cortex using SPECT imaging in a cohort of completed suicides. *Transl Psychiatry* 2011; **1**: e28-8.
- 83 Sublette ME, Milak MS, Galfalvy HC, Oquendo MA, Malone KM, Mann JJ. Regional Brain Glucose Uptake Distinguishes Suicide Attempters from Non-Attempters in Major Depression. *Arch Suicide Res* 2013; **17**: 434–447.
- 84 Oquendo MA, Placidi GPA, Malone KM, Campbell C, Keilp J, Brodsky B *et al*. Positron emission tomography of regional brain metabolic responses to a serotonergic challenge and lethality of suicide attempts in major depression. *Arch Gen Psychiatry* 2003; **60**: 14–22.
- 85 van Heeringen K, Wu GR, Vervaeke M, Vanderhasselt MA, Baeken C. Decreased resting state metabolic activity in frontopolar and parietal brain regions is associated

- with suicide plans in depressed individuals. *J Psychiatr Res* 2017; **84**: 243–248.
- 86 Oquendo MA. Impulsive versus planned suicide attempts: Different phenotypes? *J Clin Psychiatry* 2015; **76**: 293–294.
- 87 Goodman M, Hazlett EA, Avedon JB, Siever DR, Chu KW, New AS. Anterior cingulate volume reduction in adolescents with borderline personality disorder and co-morbid major depression. *J Psychiatr Res* 2011; **45**: 803–807.
- 88 Soloff P, White R, Diwadkar VA. Impulsivity, aggression and brain structure in high and low lethality suicide attempters with borderline personality disorder. *Psychiatry Res - Neuroimaging* 2014; **222**: 131–139.
- 89 Kolla NJ, Chiucciariello L, Wilson AA, Houle S, Links P, Bagby RM *et al.* Elevated Monoamine Oxidase-A Distribution Volume in Borderline Personality Disorder Is Associated with Severity Across Mood Symptoms, Suicidality, and Cognition. *Biol Psychiatry* 2016; **79**: 117–126.
- 90 Holmes SE, Hinz R, Conen S, Gregory CJ, Matthews JC, Anton-Rodriguez JM *et al.* Elevated Translocator Protein in Anterior Cingulate in Major Depression and a Role for Inflammation in Suicidal Thinking: A Positron Emission Tomography Study. *Biol Psychiatry* 2018; **83**: 61–69.
- 91 Prescott A, Sheth C, Legarreta M, Renshaw PF, McGlade E, Yurgelun-Todd D. Altered Cortical Gamma-Amino Butyric Acid in Female Veterans With Suicidal Behavior: Sex Differences and Clinical Correlates. *Chronic Stress* 2018; **2**.
doi:10.1177/2470547018768771.
- 92 Baek K, Kwon J, Chae JH, Chung YA, Kralik JD, Min JA *et al.* Heightened aversion to risk and loss in depressed patients with a suicide attempt history. *Sci Rep* 2017; **7**: 11228.
- 93 Dombrovski AY, Szanto K, Clark L, Reynolds CF, Siegle GJ. Reward signals, attempted suicide, and impulsivity in late-life depression. *JAMA Psychiatry* 2013; **70**: 1020–1030.
- 94 Uddin LQ, Kinnison J, Pessoa K, Anderson ML. Beyond the Tripartite Cognition–Emotion–Interoception Model of the Human Insular Cortex. *J Cogn Neurosci* 2014; **26**: 16–27.
- 95 Craig AD. Interoception: The sense of the physiological condition of the body. *Curr Opin Neurobiol* 2003; **13**: 500–505.
- 96 Singer T, Seymour B, Doherty JPO, Stephan KE, Dolan RJ, Frith CD. Understanding the Emergence of Neuropsychiatric Disorders with Network[1]. 2009; **439**: 466–469.
- 97 Singer T, Seymour B, O’Doherty J, Kaube H, Dolan RJ, Frith CD. Empathy for Pain Involves the Affective but not Sensory Components of Pain. *Science (80-)* 2004; **303**: 1157–1162.
- 98 Forrest LN, Smith AR, White RD, Joiner TE. (Dis)connected: An examination of interoception in individuals with suicidality. *J Abnorm Psychol* 2015; **124**: 754–763.
- 99 Brausch AM, Woods SE. Emotion Regulation Deficits and Nonsuicidal Self-Injury

- Prospectively Predict Suicide Ideation in Adolescents. *Suicide Life-Threatening Behav* 2018. doi:10.1111/sltb.12478.
- 100 Soloff PH, Pruitt P, Sharma M, Radwan J, White R, Diwadkar VA. Structural brain abnormalities and suicidal behavior in borderline personality disorder. *J Psychiatr Res* 2012; **46**: 516–525.
- 101 Duarte DGG, Neves M de CL, Albuquerque MR, Turecki G, Ding Y, de Souza-Duran FL *et al.* Structural brain abnormalities in patients with type I bipolar disorder and suicidal behavior. *Psychiatry Res - Neuroimaging* 2017; **265**: 9–17.
- 102 Amen DG, Prunella JR, Fallon JH, Amen B, Hanks C. A Comparative Analysis of Completed Suicide Using High Resolution Brain SPECT Imaging. *J Neuropsychiatry Clin Neurosci* 2009; **21**: 430–439.
- 103 Olié E, Jollant F, Deverdun J, De Champfleury NM, Cyprien F, Le Bars E *et al.* The experience of social exclusion in women with a history of suicidal acts: A neuroimaging study. *Sci Rep* 2017; **7**: 1–8.
- 104 Savitz J, Harrison NA. Interoception and Inflammation in Psychiatric Disorders. *Biol Psychiatry Cogn Neurosci Neuroimaging* 2018; **3**: 514–524.
- 105 Canteras NS, Swanson LW. Projections of the ventral subiculum to the amygdala, septum, and hypothalamus: A PHAL anterograde tract-tracing study in the rat. *J Comp Neurol* 1992; **324**: 180–194.
- 106 Zola-Morgan. Effects of lesions of perirhinal cortex or parahippocampal cortex on memory in monkeys. *Soc Neurosci Abstr* 1994; **9**: 4355–4370.
- 107 Patel D, Anilkumar S, Chattarji S, Buwalda B. Repeated social stress leads to contrasting patterns of structural plasticity in the amygdala and hippocampus. *Behav Brain Res* 2018; **347**: 314–324.
- 108 Doré BP, Rodrik O, Boccagno C, Hubbard A, Weber J, Stanley B *et al.* Negative Autobiographical Memory in Depression Reflects Elevated Amygdala-Hippocampal Reactivity and Hippocampally Associated Emotion Regulation. *Biol Psychiatry Cogn Neurosci Neuroimaging* 2018; **3**: 358–366.
- 109 Monkul ES, Hatch JP, Nicoletti MA, Spence S, Brambilla P, Lacerda ALT *et al.* Fronto- limbic brain structures in suicidal and non-suicidal female patients with major depressive disorder. *Mol Psychiatry* 2007; **12**: 360–366.
- 110 Spoletini I, Piras F, Fagioli S, Rubino IA, Martinotti G, Siracusano A *et al.* Suicidal attempts and increased right amygdala volume in schizophrenia. *Schizophr Res* 2011; **125**: 30–40.
- 111 Rentería ME, Schmaal L, Hibar DP, Couvy-Duchesne B, Strike LT, Mills NT *et al.* Subcortical brain structure and suicidal behaviour in major depressive disorder: A meta-analysis from the ENIGMA-MDD working group. *Transl Psychiatry* 2017; **7**: e1116.
- 112 Gifuni AJ, Ding Y, Olié E, Lawrence N, Cyprien F, Le Bars E *et al.* Subcortical nuclei volumes in suicidal behavior: nucleus accumbens may modulate the lethality of acts.

- Brain Imaging Behav* 2016; **10**: 96–104.
- 113 Lijffijt M, Rourke ED, Swann AC, Zunta-Soares GB, Soares JC. Illness-course modulates suicidality-related prefrontal gray matter reduction in women with bipolar disorder. *Acta Psychiatr Scand* 2014; **130**: 374–387.
- 114 Colle R, Chupin M, Cury C, Vandendrie C, Gressier F, Hardy P *et al.* Depressed suicide attempters have smaller hippocampus than depressed patients without suicide attempts. *J Psychiatr Res* 2015; **61**: 13–18.
- 115 Pan LA, Ramos L, Segreti AM, Brent DA, Phillips ML. Right superior temporal gyrus volume in adolescents with a history of suicide attempt. *Br J Psychiatry* 2015; **206**: 339–340.
- 116 Soloff PH, Price JC, Meltzer CC, Fabio A, Frank GK, Kaye WH. 5HT2A Receptor Binding is Increased in Borderline Personality Disorder. *Biol Psychiatry* 2007; **62**: 580–587.
- 117 Marchand WR, Lee JN, Garn C, Thatcher J, Gale P, Kreitschitz S *et al.* Aberrant emotional processing in posterior cortical midline structures in bipolar II depression. *Prog Neuro-Psychopharmacology Biol Psychiatry* 2011; **35**: 1729–1737.
- 118 Spreng RN, Mar RA, Kim ASN. The common neural basis of autobiographical memory, prospection, navigation, theory of mind, and the default mode: A quantitative meta-analysis. *J Cogn Neurosci* 2009; **21**: 489–510.
- 119 Schacter DL, Addis DR, Buckner RL. Remembering the past to imagine the future: the prospective brain. *Nat Rev Neurosci* 2007; **8**: 657.
- 120 Peters J, Büchel C. The neural mechanisms of inter-temporal decision-making: Understanding variability. *Trends Cogn Sci* 2011; **15**: 227–239.
- 121 Peters J, Büchel C. Episodic Future Thinking Reduces Reward Delay Discounting through an Enhancement of Prefrontal-Mediotemporal Interactions. *Neuron* 2010; **66**: 138–148.
- 122 van Erp TGM, Hibar DP, Rasmussen JM, Glahn DC, Pearlson GD, Andreassen O a *et al.* Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Mol Psychiatry* 2015; : 1–7.
- 123 Schmaal L, Veltman DJ, Van Erp TGM, Smann PG, Frodl T, Jahanshad N *et al.* Subcortical brain alterations in major depressive disorder: Findings from the ENIGMA Major Depressive Disorder working group. *Mol Psychiatry* 2016; **21**: 806–812.
- 124 Hibar DP, Westlye LT, Van Erp TGM, Rasmussen J, Leonardo CD, Faskowitz J *et al.* Subcortical volumetric abnormalities in bipolar disorder. *Mol Psychiatry* 2016; **21**: 1710–1716.
- 125 Cummings JL. Frontal-Subcortical Circuits and Human Behavior. *Arch Neurol* 1993; **50**: 873–880.
- 126 Bracht T, Linden D, Keedwell P. A review of white matter microstructure alterations of pathways of the reward circuit in depression. *J Affect Disord* 2015; **187**: 45–53.
- 127 Featherstone RE, McDonald RJ. Dorsal striatum and stimulus-response learning:

- Lesions of the dorsolateral, but not dorsomedial, striatum impair acquisition of a stimulus-response-based instrumental discrimination task, while sparing conditioned place preference learning. *Neuroscience* 2004; **124**: 23–31.
- 128 Alitto, H. J., & Usrey WM (2003). Corticothalamic feedback and sensory processing. *Curr Opin Neurobiol* 2003; **13**: 440–445.
- 129 Dombrovski AY, Siegle GJ, Szanto K, Clark L, Reynolds CF, Aizenstein H. The temptation of suicide: Striatal gray matter, discounting of delayed rewards, and suicide attempts in late-life depression. *Psychol Med* 2012; **42**: 1203–1215.
- 130 Vang FJ, Ryding E, Träskman-Bendz L, van Westen D, Lindström MB. Size of basal ganglia in suicide attempters, and its association with temperament and serotonin transporter density. *Psychiatry Res - Neuroimaging* 2010; **183**: 177–179.
- 131 Nye JA, Purselle D, Plisson C, Voll RJ, Stehouwer JS, Votaw JR *et al*. Decreased brainstem and putamen ser binding potential in depressed suicide attempters using [¹¹C]-zient pet imaging. *Depress Anxiety* 2013; **30**: 902–907.
- 132 Ryding E, Ahnlide JA, Lindström M, Rosén I, Träskman-Bendz L. Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. *Psychiatry Res - Neuroimaging* 2006; **148**: 195–203.
- 133 Yeh YW, Ho PS, Chen CY, Kuo SC, Liang CS, Yen CH *et al*. Suicidal ideation modulates the reduction in serotonin transporter availability in male military conscripts with major depression: A 4-[¹⁸F]-ADAM PET study. *World J Biol Psychiatry* 2015; **16**: 502–512.
- 134 Lopez-Larson M, King JB, McGlade E, Bueler E, Stoeckel A, Epstein DJ *et al*. Enlarged thalamic volumes and increased fractional anisotropy in the thalamic radiations in veterans with suicide behaviors. *Front Psychiatry* 2013; **4**: 1–13.
- 135 Marchand WR, Lee JN, Garn C, Thatcher J, Gale P, Kreitschitz S *et al*. Striatal and cortical midline activation and connectivity associated with suicidal ideation and depression in bipolar II disorder. *J Affect Disord* 2011; **133**: 638–645.
- 136 Kim YJ, Park HJ, Jahng GH, Lee SM, Kang WS, Kim SK *et al*. A pilot study of differential brain activation to suicidal means and DNA methylation of CACNA1C gene in suicidal attempt patients. *Psychiatry Res* 2017; **255**: 42–48.
- 137 Richard-Devantoy S, Olié E, Guillaume S, Courtet P. Decision-making in unipolar or bipolar suicide attempters. *J Affect Disord* 2016; **190**: 128–136.
- 138 Marchand WR, Lee JN, Johnson S, Thatcher J, Gale P, Wood N *et al*. Striatal and cortical midline circuits in major depression: Implications for suicide and symptom expression. *Prog Neuro-Psychopharmacology Biol Psychiatry* 2012; **36**: 290–299.
- 139 Booth MCA, Rolls ET. View-invariant representations of familiar objects by neurons in the inferior temporal visual cortex. *Cereb Cortex* 1998; **8**: 510–523.
- 140 Haxby J V, Hoffman EA, Gobbini MI. The distributed human neural system for face perception. *Trends Cogn Sci* 2000; **4**.

- 141 Howard M a, Volkov IO, Mirsky R, Garell PC, Noh MD, Granner M *et al.* Auditory cortex on the human posterior superior temporal gyrus. *J Comp Neurol* 2000; **416**: 79–92.
- 142 Aguilar EJ, García-Martí G, Martí-Bonmatí L, Lull JJ, Moratal D, Escartí MJ *et al.* Left orbitofrontal and superior temporal gyrus structural changes associated to suicidal behavior in patients with schizophrenia. *Prog Neuro-Psychopharmacology Biol Psychiatry* 2008; **32**: 1673–1676.
- 143 Audenaert K, Goethals I, Van laere K, Lahorte P, Brans B, Versijpt J *et al.* Spect neuropsychological activation procedure with the verbal fluency test in attempted suicide patients. *Nucl Med Commun* 2002; **23**: 907–916.
- 144 Motoyama H, Hishitani S. The brain mechanism that reduces the vividness of negative imagery. *Conscious Cogn* 2016; **39**: 59–69.
- 145 Feng C, Yan X, Huang W, Han S, Ma Y. Neural representations of the multidimensional self in the cortical midline structures. *Neuroimage* 2018; **183**: 291–299.
- 146 Peng H, Wu K, Li J, Qi H, Guo S, Chi M *et al.* Increased suicide attempts in young depressed patients with abnormal temporal-parietal-limbic gray matter volume. *J Affect Disord* 2014; **165**: 69–73.
- 147 Quevedo K, Ng R, Scott H, Martin J, Smyda G, Keener M *et al.* The Neurobiology of Self-Face Recognition in Depressed Adolescents with Low or High Suicidality. *J Abnorm Psychol* 2016; **125**: 1185–1200.
- 148 Adamaszek M, D'Agata F, Ferrucci R, Habas C, Keulen S, Kirkby KC *et al.* Consensus Paper: Cerebellum and Emotion. *Cerebellum* 2017; **16**: 552–576.
- 149 Womer FY, Wang F, Chepenik LG, Kalmar JH, Spencer L, Edmiston E *et al.* Sexually dimorphic features of vermis morphology in bipolar disorder. *Bipolar Disord* 2009; **11**: 753–758.
- 150 Lee YJ, Kim S, Gwak AR, Kim SJ, Kang SG, Na KS *et al.* Decreased regional gray matter volume in suicide attempters compared to suicide non-attempters with major depressive disorders. *Compr Psychiatry* 2016; **67**: 59–65.
- 151 Ballard ED, Lally N, Nugent AC, Furey ML, Luckenbaugh DA, Zarate CA. Neural correlates of suicidal ideation and its reduction in depression. *Int J Neuropsychopharmacol* 2015; **18**: 1–6.
- 152 Ordaz SJ, Goyer MS, Ho TC, Singh MK, Gotlib IH. Network basis of suicidal ideation in depressed adolescents. *J Affect Disord* 2018; **226**: 92–99.
- 153 Du L, Zeng J, Liu H, Tang D, Meng H, Li Y *et al.* Fronto-limbic disconnection in depressed patients with suicidal ideation: A resting-state functional connectivity study. *J Affect Disord* 2017; **215**: 213–217.
- 154 Zhang S, Chen J mei, Kuang L, Cao J, Zhang H, Ai M *et al.* Association between abnormal default mode network activity and suicidality in depressed adolescents. *BMC Psychiatry* 2016; **16**: 1–10.

- 155 Beaulieu C. The basis of anisotropic water diffusion in the nervous system - A technical review. *NMR Biomed* 2002; **15**: 435–455.
- 156 Mahon K, Burdick KE, Wu J, Ardekani BA, Szeszko PR. Relationship between suicidality and impulsivity in bipolar I disorder: A diffusion tensor imaging study. *Bipolar Disord* 2012; **14**: 80–89.
- 157 Cyprien F, de Champfleury NM, Deverdun J, Olié E, Le Bars E, Bonafé A *et al.* Corpus callosum integrity is affected by mood disorders and also by the suicide attempt history: A diffusion tensor imaging study. *J Affect Disord* 2016; **206**: 115–124.
- 158 Lischke A, Domin M, Freyberger HJ, Grabe HJ, Mentel R, Bernheim D *et al.* Structural Alterations in the Corpus Callosum Are Associated with Suicidal Behavior in Women with Borderline Personality Disorder. *Front Hum Neurosci* 2017; **11**: 1–10.
- 159 Laird, P. Mickle Fox, Simon B. Eickhoff, Jessica A. Turner, Kimberly L. Ray, D. Reese McKay, David C. Glahn, Christian F. Beckmann, Stephen M. Smith and PTF. Behavioral Interpretations of Intrinsic Connectivity Networks. *J Cogn Neurosci* 2011; **23**: 4022–4037.
- 160 Kang SG, Na KS, Choi JW, Kim JH, Son YD, Lee YJ. Resting-state functional connectivity of the amygdala in suicide attempters with major depressive disorder. *Prog Neuro-Psychopharmacology Biol Psychiatry* 2017; **77**: 222–227.
- 161 Quevedo K, Ng R, Scott H, Kodavaganti S, Smyda G, Diwadkar V *et al.* Ventral Striatum Functional Connectivity during Rewards and Losses and Symptomatology in Depressed Patients. *Biol Psychol* 2017; **123**: 62–73.
- 162 Kim K, Kim SW, Myung W, Han CE, Fava M, Mischoulon D *et al.* Reduced orbitofrontal-thalamic functional connectivity related to suicidal ideation in patients with major depressive disorder. *Sci Rep* 2017; **7**: 15772.
- 163 Myung W, Han CE, Fava M, Mischoulon D, Papakostas GI, Heo JY *et al.* Reduced frontal-subcortical white matter connectivity in association with suicidal ideation in major depressive disorder. *Transl Psychiatry* 2016; **6**: e835–8.
- 164 Bijttebier S, Caeyenberghs K, van den Ameele H, Achten E, Rujescu D, Titeca K *et al.* The Vulnerability to Suicidal Behavior is Associated with Reduced Connectivity Strength. *Front Hum Neurosci* 2015; **9**. doi:10.3389/fnhum.2015.00632.
- 165 Jia Z, Huang X-Q, Wu Q-Z, Zhang T-J, Lui S, Zhang J *et al.* High-field magnetic resonance imaging of suicidality in patients with major depressive disorder. *Am J Psychiatry* 2010; **167**: 1381–1390.
- 166 Jia Z, Wang Y, Huang X, Kuang W, Wu Q, Lui S *et al.* Impaired frontothalamic circuitry in suicidal patients with depression revealed by diffusion tensor imaging at 3.0 T. *J Psychiatry Neurosci* 2014; **39**: 170–177.
- 167 Bressler SL, Menon V. Large-scale brain networks in cognition: emerging methods and principles. *Trends Cogn Sci* 2010; **14**: 277–290.
- 168 Cao J, Chen X, Chen J, Ai M, Gan Y, Wang W *et al.* Resting-state functional MRI of abnormal baseline brain activity in young depressed patients with and without suicidal

- behavior. *J Affect Disord* 2016; **205**: 252–263.
- 169 Olvet DM, Peruzzo D, Thapa-Chhetry B, Sublette ME, Sullivan GM, Oquendo MA *et al.* A diffusion tensor imaging study of suicide attempters. *J Psychiatr Res* 2014; **51**: 60–67.
- 170 Minzenberg M, Lesh T, Niendam T, Yoon J, Cheng Y, Rhoades R *et al.* Conflict-related anterior cingulate functional connectivity is associated with past suicidal ideation and behavior in recent-onset schizophrenia. *J Psychiatr Res* 2015; **65**: 95–101.
- 171 Minzenberg MJ, Lesh T, Niendam T, Yoon JH, Cheng Y, Rhoades R *et al.* Conflict-related anterior cingulate functional connectivity is associated with past suicidal ideation and behavior in recent-onset Psychotic Major Mood Disorders. *J Neuropsychiatry Clin Neurosci* 2016; **28**: 95–101.
- 172 Janes AC, Farmer S, Peechatka AL, Frederick BDB, Lukas SE. Insula-dorsal anterior cingulate cortex coupling is associated with enhanced brain reactivity to smoking cues. *Neuropsychopharmacology* 2015; **40**: 1561–1568.
- 173 Giesecke T, Gracely RH, Williams DA, Geisser ME, Petzke FW, Clauw DJ. The relationship between depression, clinical pain, and experimental pain in a chronic pain cohort. *Arthritis Rheum* 2005; **52**: 1577–1584.
- 174 Stevens F, Hurley R, Taber K. Anterior Cingulate Cortex: Unique Role in Cognition and Emotion. *J neuropsychiatry Clin Neurosci* 2011; **23**: 121–125.
- 175 Gasquoine PG. Contributions of the insula to cognition and emotion. *Neuropsychol Rev* 2014; **24**: 77–87.
- 176 Menon V, Uddin LQ. Saliency, switching, attention and control: a network model of insula function. *Brain Struct Funct* 2010; **214**: 655–667.
- 177 Nieuwenhuys R. *The insular cortex. A review.* 1st ed. Elsevier B.V., 2012 doi:10.1016/B978-0-444-53860-4.00007-6.
- 178 Deshpande G, Baxi M, Witte T, Robinson JL. A neural basis for the acquired capability for suicide. *Front Psychiatry* 2016; **7**: 1–19.
- 179 Goulden N, Khusnulina A, Davis NJ, Bracewell RM, Bokde AL, McNulty JP *et al.* The salience network is responsible for switching between the default mode network and the central executive network: Replication from DCM. *Neuroimage* 2014; **99**: 180–190.
- 180 Zhou Y, Friston KJ, Zeidman P, Chen J, Li S, Razi A. The Hierarchical Organization of the Default, Dorsal Attention and Salience Networks in Adolescents and Young Adults. *Cereb Cortex* 2018; **28**: 726–737.
- 181 Sridharan D, Levitin D, Menon V. A critical role for the right fronto-insular cortex in switching between central-executive and default-mode networks. *Proc Natl Acad Sci* 2008; **105**: 12569–12574.
- 182 Cox Lippard ET, Johnston JAY, Spencer L, Quatrano S, Fan S, Sankar A *et al.* Preliminary examination of gray and white matter structure and longitudinal structural changes in frontal systems associated with future suicide attempts in adolescents and

- young adults with mood disorders. *J Affect Disord* 2018; **in press**.
- 183 Wilkinson ST, Sanacora G. Ketamine: a Potential Rapid-Acting Antisuicidal Agent? *Depress Anxiety* 2016; **33**: 711–717.
- 184 Cipriani A, Hawton K, Stockton S, Geddes JR. Lithium in the prevention of suicide in mood disorders: updated systematic review and meta-analysis. *BMJ* 2013; **346**: f3646.
- 185 Al Jurdi RK, Swann A, Mathew SJ. Psychopharmacological Agents and Suicide Risk Reduction: Ketamine and Other Approaches. *Curr Psychiatry Rep* 2015; **17**. doi:10.1007/s11920-015-0614-9.
- 186 Kundu P, Benson BE, Rosen D, Frangou S, Leibenluft E, Luh W-M *et al*. The integration of functional brain activity from adolescence to adulthood. *J Neurosci* 2018; **38**: 1864–17.
- 187 Kochunov P, Glahn DC, Lancaster J, Thompson PM, Kochunov V, Rogers B *et al*. Fractional anisotropy of cerebral white matter and thickness of cortical gray matter across the lifespan. *Neuroimage* 2011; **58**: 41–49.
- 188 Fan S, Lippard E, Sankar A, Wallace A, Johnston J, Wang F *et al*. Grey and white matter differences in adolescents and young adults with prior suicide attempts across bipolar and major depressive disorders. *J Affect Disord* 2018; **in press**.
- 189 Fox KR, Millner AJ, Mukerji CE, Nock MK. Examining the role of sex in self-injurious thoughts and behaviors. *Clin Psychol Rev* 2017; : 0–1.
- 190 Olfson M, Blanco C, Wall M, Liu SM, Saha TD, Pickering RP *et al*. National Trends in Suicide Attempts Among Adults in the United States. *JAMA psychiatry* 2017; **74**: 1095–1103.
- 191 Choi NG, DiNitto DM, Marti CN, Kaplan MS, Conwell Y. Suicide Means among Decedents Aged 50+ Years, 2005–2014: Trends and Associations with Sociodemographic and Precipitating Factors. *Am J Geriatr Psychiatry* 2017; **25**: 1404–1414.
- 192 Kaplan MS, McFarland BH, Huguet N. Characteristics of adult male and female firearm suicide decedents: Findings from the National Violent Death Reporting System. *Inj Prev* 2009; **15**: 322–327.
- 193 Hedegaard H, Curtin S, Warner M. Suicide Mortality in the United States, 1999–2017. *NCHS Data Brief* 2018; **330**: 1–8.
- 194 Nery-Fernandes F, Rocha M V., Jackowski A, Ladeia G, Guimarães JL, Quarantini LC *et al*. Reduced posterior corpus callosum area in suicidal and non-suicidal patients with bipolar disorder. *J Affect Disord* 2012; **142**: 150–155.
- 195 Gabbay V, Bradley KA, Mao X, Ostrover R, Kang G, Shungu DC. Anterior cingulate cortex γ -aminobutyric acid deficits in youth with depression. *Transl Psychiatry* 2017; **7**. doi:10.1038/tp.2017.187.
- 196 Soloff PH, Meltzer CC, Becker C, Greer PJ, Kelly TM, Constantine D. Impulsivity and prefrontal hypometabolism in borderline personality disorder. *Psychiatry Res* 2003; **123**: 153–163.

- 197 Soloff PH, Chiappetta L, Mason NS, Becker C, Price JC. Effects of serotonin-2A receptor binding and gender on personality traits and suicidal behavior in borderline personality disorder. *Psychiatry Res* 2014; **222**: 140–148.
- 198 Ehrlich S, Noam GG, Lyoo IK, Kwon BJ, Clark MA, Renshaw PF. White matter hyperintensities and their associations with suicidality in psychiatrically hospitalized children and adolescents. *J Am Acad Child Adolesc Psychiatry* 2004; **43**: 770–776.
- 199 Chase HW, Segreti AM, Keller TA, Cherkassky VL, Just MA, Pan LA *et al*. Alterations of functional connectivity and intrinsic activity within the cingulate cortex of suicidal ideators. *J Affect Disord* 2017; **212**: 78–85.
- 200 di Giacomo E, Krausz M, Colmegna F, Aspesi F, Clerici M. Estimating the risk of attempted suicide among sexual minority youths: a systematic review and meta-analysis. *JAMA Pediatr* 2018; **172**: 1145–1152.
- 201 Mueller S, De Cuyper G, T'Sjoen G. Transgender research in the 21st century: a selective critical review from a neurocognitive perspective. *Am J Psychiatry* 2017; **174**: 1155–1162.
- 202 Woo CW, Chang LJ, Lindquist MA, Wager TD. Building better biomarkers: Brain models in translational neuroimaging. *Nat Neurosci* 2017; **20**: 365–377.
- 203 Duyn JH. The future of ultra-high field MRI and fMRI for study of the human brain. *Neuroimage* 2012; **62**: 1241–1248.
- 204 Courtet P, Giner L, Seneque M, Guillaume S, Olie E, Ducasse D. Neuroinflammation in suicide: Toward a comprehensive model. *World J Biol Psychiatry* 2016; **17**: 564–586.
- 205 Oquendo MA, Sullivan GM, Sudol K, Baca-Garcia E, Stanley BH, Sublette ME *et al*. Toward a biosignature for suicide. *Am J Psychiatry* 2014; **171**: 1259–1277.
- 206 Sequeira A, Mamdani F, Ernst C, Vawter MP, Bunney WE, Lebel V *et al*. Global brain gene expression analysis links Glutamatergic and GABAergic alterations to suicide and major depression. *PLoS One* 2009; **4**: 21–23.
- 207 Zhao J, Verwer RWH, Gao SF, Qi XR, Lucassen PJ, Kessels HW *et al*. Prefrontal alterations in GABAergic and glutamatergic gene expression in relation to depression and suicide. *J Psychiatr Res* 2018; **102**: 261–274.
- 208 Poulter MO. Altered organization of GABAA receptor mRNA expression in the depressed suicide brain. *Front Mol Neurosci* 2010; **3**: 1–10.
- 209 Lebowitz E, Blumberg H, Silverman W. Negative peer social interactions and oxytocin levels linked to suicidal ideation in anxious youth. *J Affect Disord* 2018; **in press**.
- 210 Miranda R, Nolen-Hoeksema S. Brooding and reflection: Rumination predicts suicidal ideation at 1-year follow-up in a community sample. *Behav Res Ther* 2007; **45**: 3088–3095.
- 211 Williams JMG, Van Der Does AJW, Barnhofer T, Crane C, Segal ZS. Cognitive reactivity, suicidal ideation and future fluency: Preliminary investigation of a differential activation theory of hopelessness/suicidality. *Cognit Ther Res* 2008; **32**: 83–104.

- 212 Auerbach RP, Millner AJ, Stewart JG, Esposito EC. Identifying differences between depressed adolescent suicide ideators and attempters. *J Affect Disord* 2015; **186**: 127–133.
- 213 Peyre H, Hoertel N, Stordeur C, Lebeau G, Blanco C, McMahon K *et al.* Contributing Factors and Mental Health Outcomes of First Suicide Attempt During Childhood and Adolescence. *J Clin Psychiatry* 2015; **78**: e622.
- 214 Björkenstam C, Kosidou K, Björkenstam E. Childhood adversity and risk of suicide: cohort study of 548 721 adolescents and young adults in Sweden. *BMJ* 2017; **357**: j1334.
- 215 van Harmelen A-L, van Tol M-J, van der Wee NJA, Veltman DJ, Aleman A, Spinhoven P *et al.* Reduced Medial Prefrontal Cortex Volume in Adults Reporting Childhood Emotional Maltreatment. *Biol Psychiatry* 2010; **68**: 832–838.
- 216 Van Harmelen AL, Hauber K, Moor BG, Spinhoven P, Boon AE, Crone EA *et al.* Childhood emotional maltreatment severity is associated with dorsal medial prefrontal cortex responsivity to social exclusion in young adults. *PLoS One* 2014; **9**: e85107.
- 217 Button K, Ioannidis J, Mokrysz C, Nosek B, Flint J, Robinson E *et al.* Power failure: why small sample size undermines the reliability of neuroscience. *Nat Rev Neurosci* 2013; **14**: 365–375.
- 218 Fischer HF, Rose M. Scoring Depression on a Common Metric: A Comparison of EAP Estimation, Plausible Value Imputation, and Full Bayesian IRT Modeling. *Multivariate Behav Res* 2018; **54**: 85–99.
- 219 Rüsç N, Spoletini I, Wilke M, Martinotti G, Bria P, Trequattrini A *et al.* Inferior frontal white matter volume and suicidality in schizophrenia. *Psychiatry Res - Neuroimaging* 2008; **164**: 206–214.
- 220 Matsuo K, Nielsen N, Nicoletti MA, Hatch JP, Monkul ES, Watanabe Y *et al.* Anterior genu corpus callosum and impulsivity in suicidal patients with bipolar disorder. *Neurosci Lett* 2010; **469**: 75–80.
- 221 Gifuni AJ, Olié E, Ding Y, Cyprien F, le Bars E, Bonafé A *et al.* Corpus callosum volumes in bipolar disorders and suicidal vulnerability. *Psychiatry Res - Neuroimaging* 2017; **262**: 47–54.
- 222 Harenski CL, Brook M, Kosson DS, Bustillo JR, Harenski KA, Caldwell MF *et al.* Socio-neuro risk factors for suicidal behavior in criminal offenders with psychotic disorders. *Soc Cogn Affect Neurosci* 2017; **12**: 70–80.
- 223 Baldaçara L, Nery-Fernandes F, Rocha M, Quarantini LC, Rocha GGL, Guimarães JL *et al.* Is cerebellar volume related to bipolar disorder? *J Affect Disord* 2011; **135**: 305–309.
- 224 Benedetti F, Riccaboni R, Poletti S, Radaelli D, Locatelli C, Lorenzi C *et al.* The serotonin transporter genotype modulates the relationship between early stress and adult suicidality in bipolar disorder. *Bipolar Disord* 2014; **16**: 857–866.
- 225 Cyprien F, Courtet P, Malafosse A, Maller J, Meslin C, Bonafé A *et al.* Suicidal

- behavior is associated with reduced corpus callosum area. *Biol Psychiatry* 2011; **70**: 320–326.
- 226 Kim B, Oh J, Kim MK, Lee S, Tae WS, Kim CM *et al*. White matter alterations are associated with suicide attempt in patients with panic disorder. *J Affect Disord* 2015; **175**: 139–146.
- 227 Thomas LA, Bellis MD De. Pituitary Volumes in Pediatric Maltreatment-Related Posttraumatic Stress Disorder. *Biol Psychiatry* 2004; **55**: 725–758.
- 228 Caplan R, Siddarth P, Levitt J, Gurbani S, Shields WD, Sankar R. Suicidality and brain volumes in pediatric epilepsy. *Epilepsy Behav* 2010; **18**: 286–290.
- 229 Ehrlich S, Noam GG, Lyoo IK, Kwon BJ, Clark MA, Renshaw PF. Subanalysis of the location of white matter hyperintensities and their association with suicidality in children and youth. *Ann N Y Acad Sci* 2003; **1008**: 265–268.
- 230 Ehrlich S, Breeze JL, Hesdorffer DC, Noam GG, Hong X, Alban RL *et al*. White matter hyperintensities and their association with suicidality in depressed young adults. *J Affect Disord* 2005; **86**: 281–287.
- 231 Pompili M, Ehrlich S, De Pisa E, Mann JJ, Innamorati M, Cittadini A *et al*. White matter hyperintensities and their associations with suicidality in patients with major affective disorders. *Eur Arch Psychiatry Clin Neurosci* 2007; **257**: 494–499.
- 232 Pompili M, Innamorati M, Mann JJ, Oquendo MA, Lester D, Del Casale A *et al*. Periventricular white matter hyperintensities as predictors of suicide attempts in bipolar disorders and unipolar depression. *Prog Neuro-Psychopharmacology Biol Psychiatry* 2008; **32**: 1501–1507.
- 233 Ahearn EP, Jamison KR, Steffens DC, Cassidy F, Provenzale JM, Lehman A *et al*. MRI correlates of suicide attempt history in unipolar depression. *Biol Psychiatry* 2001; **50**: 266–270.
- 234 Sachs-Ericsson N, Hames JL, Joiner TE, Corsentino E, Rushing NC, Palmer E *et al*. Differences between suicide attempters and nonattempters in depressed older patients: Depression severity, white-matter lesions, and cognitive functioning. *Am J Geriatr Psychiatry* 2014; **22**: 75–85.
- 235 Lee SJ, Kim B, Oh D, Kim MK, Kim KH, Bang SY *et al*. White matter alterations associated with suicide in patients with schizophrenia or schizophreniform disorder. *Psychiatry Res - Neuroimaging* 2016; **248**: 23–29.
- 236 Audenaert K, Van Laere K, Dumont F, Slegers G, Mertens J, Van Heeringen C *et al*. Decreased frontal serotonin 5-HT₂receptor binding index in deliberate self-harm patients. *Eur J Nucl Med* 2001; **28**: 175–182.
- 237 Van Heeringen C, Audenaert K, Van Laere K, Dumont F, Slegers G, Mertens J *et al*. Prefrontal 5-HT₂receptor binding index, hopelessness and personality characteristics in attempted suicide. *J Affect Disord* 2003; **74**: 149–158.
- 238 Fountoulakis K, Lacovides A, Fotiou F, Nimatoudis J, Bascialla F, Ioannidou C *et al*. Neurobiological and psychological correlates of suicidal attempts and thoughts of

- death in patients with major depression. *Neuropsychobiology* 2004; **49**: 42–52.
- 239 Henningsson S, Borg J, Lundberg J, Bah J, Lindström M, Ryding E *et al.* Genetic Variation in Brain-Derived Neurotrophic Factor Is Associated with Serotonin Transporter but Not Serotonin-1A Receptor Availability in Men. *Biol Psychiatry* 2009; **66**: 477–485.
- 240 Bah J, Lindström M, Westberg L, Mannerås L, Ryding E, Henningsson S *et al.* Serotonin transporter gene polymorphisms: Effect on serotonin transporter availability in the brain of suicide attempters. *Psychiatry Res - Neuroimaging* 2008; **162**: 221–229.
- 241 Lindström MB, Ryding E, Bosson P, Ahnlide JA, Rosén I, Träskman-Bendz L. Impulsivity related to brain serotonin transporter binding capacity in suicide attempters. *Eur Neuropsychopharmacol* 2004; **14**: 295–300.
- 242 Yeh YW, Ho PS, Chen CY, Kuo SC, Liang CS, Ma KH *et al.* Incongruent reduction of serotonin transporter associated with suicide attempts in patients with major depressive disorder: A positron emission tomography study with 4-[18F]-ADAM. *Int J Neuropsychopharmacol* 2015; **18**: 1–9.
- 243 Miller JM, Hesselgrave N, Ogden RT, Sullivan GM, Oquendo MA, Mann JJ *et al.* Positron Emission Tomography Quantification of Serotonin Transporter in Suicide Attempters with Major Depressive Disorder. *Biol Psychiatry* 2013; **74**: 287–295.
- 244 Jollant F, Near J, Turecki G, Richard-Devantoy S. Spectroscopy markers of suicidal risk and mental pain in depressed patients. *Prog Neuro-Psychopharmacology Biol Psychiatry* 2017; **73**: 64–71.
- 245 Rocha MV, Nery-Fernandes F, Guimarães JL, De Castro Quarantini L, De Oliveira IR, Ladeia-Rocha GG *et al.* Normal metabolic levels in prefrontal cortex in euthymic bipolar I patients with and without suicide attempts. *Neural Plast* 2015; : 1–9.
- 246 Cao J, mei Chen J, Kuang L, Ai M, dong Fang W, Gan Y *et al.* Abnormal regional homogeneity in young adult suicide attempters with no diagnosable psychiatric disorder: A resting state functional magnetic imaging study. *Psychiatry Res - Neuroimaging* 2015; **231**: 95–102.
- 247 Pan LA, Batezati-Alves SC, Almeida JRC, Segreti A, Akkal D, Hassel S *et al.* Dissociable patterns of neural activity during response inhibition in depressed adolescents with and without suicidal behavior. *J Am Acad Child Adolesc Psychiatry* 2011; **50**: 602–611.
- 248 Pan L, Segreti A, Almeida J, Jollant F, Lawrence N, Brenta D *et al.* Preserved hippocampal function during learning in the context of risk in adolescent suicide attempt. *Psychiatry Res Neuroimaging* 2013; **211**: 112–118.
- 249 Jollant F, Lawrence NS, Olie E, O'Daly O, Malafosse A, Courtet P *et al.* Decreased activation of lateral orbitofrontal cortex during risky choices under uncertainty is associated with disadvantageous decision-making and suicidal behavior. *Neuroimage* 2010; **51**: 1275–1281.
- 250 Cullen KR, Westlund MK, Klimes-Dougan B, Mueller BA, Houry A, Eberly LE *et al.*

Abnormal amygdala resting-state functional connectivity in adolescent depression. *JAMA Psychiatry* 2014; **71**: 1138–1147.

- 251 Zhang H, Wei X, Tao H, Mwansisya TE, Pu W, He Z *et al.* Opposite Effective Connectivity in the Posterior Cingulate and Medial Prefrontal Cortex between First-Episode Schizophrenic Patients with Suicide Risk and Healthy Controls. *PLoS One* 2013; **8**: 1–8.
- 252 Marchand WR, Lee JN, Johnson S, Gale P, Thatcher J. Differences in functional connectivity in major depression versus bipolar II depression. *J Affect Disord* 2013; **150**: 527–532.
- 253 Tsujii N, Mikawa W, Tsujimoto E, Adachi T, Niwa A, Ono H *et al.* Reduced left precentral regional responses in patients with major depressive disorder and history of suicide attempts. *PLoS One* 2017; **12**: e0175249.
- 254 Pu S, Nakagome K, Yamada T, Yokoyama K, Matsumura H, Yamada S *et al.* Suicidal ideation is associated with reduced prefrontal activation during a verbal fluency task in patients with major depressive disorder. *J Affect Disord* 2015; **181**: 9–17.

FIGURE LEGENDS

Figure 1: Number of neuroimaging studies on suicidal thoughts and behaviors published in the last two decades.


The figure was based on the studies included in this review, calculated separately for studies including adolescents and studies only including adults and divided into separate 4-year time bins for publication date.




Figure 2: Overview of brain regions and structural connections included in this review

(A) Brain regions that have been most reported in neuroimaging studies investigating structural, functional and molecular brain alterations associated with suicidal thoughts and behaviors, with a subset of regions grouped more broadly into ventral prefrontal cortex, dorsal prefrontal cortex, insula, mesial temporal, subcortical, and posterior regions. (B) White matter tracts implicated in suicidal thoughts and behaviors reported in diffusion tensor imaging studies. DMPFC, dorsomedial prefrontal cortex; dACC, dorsal anterior cingulate cortex; RMPFC, rostromedial prefrontal cortex; mOFC, medial orbitofrontal cortex; vACC, ventral anterior cingulate cortex; PCC, posterior cingulate cortex; Thal, thalamus; VS, ventral striatum; Hippo, hippocampus; Amyg, amygdala; DLPFC, dorsolateral prefrontal cortex; RLPFC, rostral prefrontal cortex; IFG, inferior frontal gyrus; IOFC, lateral orbitofrontal cortex; Put, putamen; Caud, caudate.


Figure 3: A tentative brain circuitry model of suicidal thoughts and behaviors.

Medial VPFC (ventral ACC, OFC, RPFC), insula, amygdala, hippocampus, lateral temporal regions, posterior midline structures (posterior cingulate cortex and precuneus), dACC, ventral striatum, thalamus and cerebellum contribute to the generation of suicidal ideation through their roles in excessive negative and blunted positive internal states, negative self-referencing, impairments in future thinking and rumination. DPFC (DLPFC and DMPFC), IFG and dACC alterations further exacerbate suicidal thoughts and facilitate suicide behaviors due to their involvement in diminished cognitive control of thought, emotion and behavior and impairments in cognitive flexibility and valuation of different decision options. Alterations in bottom-up and top-down connections between these extended medial VPFC and DPFC/IFG systems may contribute to the transition from suicidal thoughts to behaviors. The dACC and insula may mediate this transition. Dashed lines indicate speculative associations that need further confirmation by future structural and functional connectivity studies. DMPFC, dorsomedial prefrontal cortex; dACC, dorsal anterior cingulate cortex; RPFC, rostral prefrontal cortex; OFC, orbitofrontal cortex; vACC, ventral anterior cingulate cortex; PCC, posterior cingulate cortex; Thal, thalamus; VS, ventral striatum; Hippo, hippocampus; Amyg, amygdala; DLPFC, dorsolateral prefrontal cortex; IFG, inferior frontal gyrus; Put, putamen; Caud, caudate.





- Striatum & Thalamus ● Dorsal Prefrontal Cortex (DPFC) ● Posterior regions ● Mesial temporal
- Ventral Prefrontal Cortex (VPFC) ● Anterior Cingulate Cortex (ACC)



● Diminished cognitive control of emotion and behavior, cognitive inflexibility, impaired valuation of different decision options



Mediating dynamic interactions between the extended VPFC and DPFC/IFG systems



● Enhanced negative and blunted positive internal states, negative self-referencing, impairments in future thinking, rumination

Table 1. Findings from Structural Imaging Studies of Suicidal Thoughts and Behaviors

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
STRUCTURAL MAGNETIC RESONANCE IMAGING STUDIES OF GRAY AND WHITE MATTER							
<i>Suicide attempt studies</i>							
Goodman et al 2011 ⁸⁷	BPD + MDD	13 SA	13 HC	20 (77)	SA: 15.8 (1.1), HC: 16.2 (0.8)	ROI volume: OFC, ACC, DMPFC, DLPFC	Number of SA associated with ↓ ACC (BA24) volume (combined WM and GM), and ↑ WM (not GM) in posterior cingulate (BA 23), however, similar association with BPD symptom severity for both findings.
Fradkin et al 2017 ⁵⁵	MDD	29 SA	29 HC	46 (79)	SA: 17.6 (1.82), HC:16.9 (1.72)	WB cortical thickness and surface area	In HC: motor impulsivity associated with ↑ RMPFC/RLPFC thickness. In SA: motor impulsivity associated with ↓ RMPFC/RLPFC thickness. In SA: non-planning impulsivity associated with ↑ paracentral lobule thickness. No findings medicated vs unmedicated.
Cao et al 2016 ¹⁶⁸	DD	35 SA	18 DC, 47 HC	66 (66)	SA: 20.63 (3.65), DC: 21.39 (3.05), HC: 20.53 (1.84)	WB VBM GMV and WMV (SPM)	SA + SI vs DC + SI: No differences between groups
Johnston et al 2017 ³⁶	BD	26 SA	42 DC, 45 HC	43 (63)	SA: 20.5(3.0), DC:20.6 (3.2), HC:20.8(3.3)	WB, VBM GMV (SPM)	SA +SI vs DC: ↓ GMV in right medial/lateral OFC (BA11/47), hippocampus, bilateral cerebellum
Pan et al 2015 ¹¹⁵	MDD	28 SA	31 DC, 41 HC	ND	SA: 16.0 (1.27), DC: 16.06 (1.47), HC: 14.48 (1.84)	WB GMV, WMV and cortical thickness	SA + SI vs DC: ↓ caudal middle frontal gyrus (BA8) volume, temporal pole (BA38) thickness, parahippocampal gyrus (BA34) volume
Peng et al 2014 ¹⁴⁶	MDD	20 SA	18 DC, 28 HC	38 (58)	SA: 27.75 (7.21), DC: 31.06 (7.39), HC: 28.61 (5.45)	WB VBM GMV (SPM)	SA vs DC (SI not reported): ↓ left PCC, which correlated negatively with dysfunctional attitudes.
Gosnell et al 2016 ³³	MDD, BD, AA, ANX	20 SA	20 DC, 20 HC	28 (47)	SA: 28.9 (9.98), DC: 29.25 (11.1), HC: 28.9 (10.0)	ROI volume: thalamus, insula, basal ganglia, hippocampus, amygdala, corpus callosum, cortical lobes.	SA vs DC: ↓ right precentral gyrus, right IFG, right caudal middle frontal gyrus (DMPFC), left precentral lobule and total temporal cortex. No association between SI (regardless of history of SA) and any of the ROIs
Soloff et al 2012 ¹⁰⁰	BPD	44 SA	24 DC, 52 HC	76 (63)	SA: 29.6 (8.0), DC: 25.9 (5.7), HC: 25.9 (7.2)	ROI VBM GMV (SPM), ROIs: IFG, OFC, ACC, middle and superior temporal cortex, insula, hippocampus, parahippocampus, fusiform gyrus, lingual gyrus and amygdala	SA vs DC: ↓ insula and larger lingual and middle and superior temporal gyri. In SA: HL associated with ↓ lateral OFC, middle and superior temporal gyri, insula, fusiform gyrus, lingual gyrus and parahippocampal gyrus
Monkul et al 2007 ¹⁰⁹	MDD	7 SA	10 DC, 17 HC	34 (100)	SA: 31.4 (13.9), DC: 36.5 (7.5), HC: 31.3 (8.3)	ROI manual shape tracing, ROIs: OFC, ACC, PCC, amygdala and hippocampus. DLPFC, subgenual PFC, thalamus, temporal lobe, caudate and lateral ventricles in exploratory analyses	SA vs DC: ↑ right amygdala

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Besteher et al 2016 ⁷¹	SCZ	14 SA	23 DC, 50 HC	46 (53)	SA: 34.4 (12.1), DC: 28.8 (9.7), HC: 29.5 (7.9)	WB cortical thickness and mean curvature	SA vs DC: SA ↓ in right DLPFC and superior and middle temporal gyri, temporopolar cortex, and insula
Giakoumatos et al 2014 ³⁷	SCZ, SZA or BD-P	148 SA (97 HL, 51 LL)	341 DC, 262 HC	387 (52)	SA-HL: 35.6 (11.7), SA-LL: 36.9 (12.2), DC: 35.9 (13.3), HC: 38.1 (12.5)	WB GMV, cortical surface area and thickness	SA vs DC: ↓ GMV in bilateral superior and middle frontal gyri (DLPFC), and inferior and superior temporal gyri, left superior parietal and supramarginal cortex, and right insula and thalamus. High (vs. low) lethality: ↓ GMV in left lingual area and right cuneus across all attempters; left dorsal ACC (BA32), left inferior parietal cortex, left inferior temporal gyrus, and right middle temporal gyrus in BD-P; left lingual gyrus, bilateral pericalcarine, right cuneus and right lateral occipital cortex in SZ; left middle frontal gyrus (DLPFC) in SZA.
Rüsch et al 2008 ²¹⁹	SCZ	10 SA	45 DC, 55 HC	42 (38)	SA: 30.3 (6.5): DC 37.3 (11.6)	WB VBM GMV and WMV (SPM)	SA vs DC: ↑ WMV in bilateral posterior lateral OFC and IFG. No GMV differences
Matsuo et al 2010 ²²⁰	BD	10 SA	10 DC, 27 HC	47 (100)	SA: 36.2 (10.1), DC:44.2 (12.5), HC: 36.9 (13.8)	ROI volume, manual shape tracing of CC genu, anterior body, posterior body, isthmus and splenium	SA vs DC: no significant differences. In SA: impulsivity associated with ↓ anterior corpus callosum genu
Lijffijt et al 2014 ¹¹³	BD	51 SA	42 DC, 45 HC	138 (100)	SA: 36.6 (10.7), DC: 41.1 (11.3)	ROI volume, ROIs: superior frontal gyrus, rostral and caudal middle frontal gyrus, frontal pole, IFG, medial and lateral OFC, rostral and caudal ACC	SA vs DC: no differences. Lower PFC GMV only in SA with previous hospitalization
Soloff et al 2014 ⁸⁸	BPD	51 SA (16 HL, 35 LL)		41 (80)	SA-HL: 36.1 (9.2): SA-LL: 27.4 (5.9).	ROI volume VBM of GMV (SPM), ROIs: OFC, ACC, middle and superior temp gyrus, insula, (para-)hippocampus, lingual gyrus, amygdala	HL vs LL SA: ↓ GMV in bilateral middle and superior temporal gyri, left lingual gyrus, bilateral lateral OFC, right insula, bilateral fusiform gyrus, right parahippocampus, left ventral and dorsal ACC and left hippocampus. In HL SA: aggression associated with ↑ ventral and dorsal ACC, lateral OFC, middle and superior temporal gyri and right insula, and impulsivity associated with ↑ right middle and superior temporal gyri and ↓ insula. In LL SA: impulsivity associated with ↓ right middle and superior temporal gyri, bilateral insula, bilateral lingual gyrus, ventral ACC, fusiform gyrus, lateral OFC, hippocampus and amygdala
Aguilar et al 2008 ¹⁴²	SCZ	13 SA	24 DC	0 (0)	SA: 37.1 (11.0), DC: 42.7 (10.2)	WB VBM GMV (SPM)	SA vs DC: ↓ medial OFC and superior temporal gyrus
Ding et al 2015 ³²	Past MDD/BD	67 SA	82 DC, 82 HC	98 (42)	SA: 39.2 (10.6), DC: 39.4 (9.7), HC: 37.8 (8.1)	WB and ROI VBM GMV (SPM) and cortical thickness and surface area, ROIs: OFC, VLPFC, VMPFC (including ACC), and DLPFC	SA vs DC: ↓ OFC (BA47, lateral part of BA11) in exploratory WB analysis. In SA: lethality last SA associated with ↓ right DPFC BA8/9/46), OFC, left VLPFC (BA44/45). Number SA associated with ↓ right DPFC and left OFC. No associations with SI.

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Gifuni et al 2017 ²²¹	Past MDD/BD	61 SA	75 DC, 73 HC	120 (47)	SA: 38.3 (10.7), DC: 38.4 (9.1), HC: 39.2 (7.0)	ROI volume SBM, ROI: corpus callosum	SA vs DC: no differences. No correlation corpus callosum volume and SI, age at first SA and number of SA
Gifuni et al 2016 ¹¹²	Past DD/BD	73 SA	89 DC, 91 HC	120 (47)	SA: 39.2 (10.6), DC: 39.4 (9.5), HC: 38.3 (8.2)	ROI volume, ROIs: amygdala, hippocampus, caudate, globus pallidus, putamen, nucleus accumbens, ventral diencephalon and thalamus	SA vs DC: no differences. In SA: lethality SA associated with ↓ left and right nucleus accumbens
Harenski et al 2017 ²²²	SCZ, SZA, BD-P, or MDD-P	18 SA	18 DC, 59 HC, 26 HC	0 (0)	SA: 38.9 (11.73), DC: 40.2 (10.23), HC: 32.5 (11.16), CHC: 33.0 (9.49)	WB and ROI VBM GMV (SPM), ROIs: posterior superior temporal cortex, temporal poles and medial PFC	SA vs DC: ↓ left and right temporal pole
Nery-Fernandes et al 2012 ¹⁹⁴	BD	19 SA	21 DC, 22 HC	41 (66)	SA: 39.8 (11.4), DC: 42.0 (8.6), HC: 37.7 (13.5)	ROI volume VBM GMV (SPM), ROI: corpus callosum	SA vs DC: no differences
Vang et al 2010 ¹³⁰	MDD, AD	7 SA	6 HC	ND	SA: 40 (11.83), matched HC, no details	ROI volume, ROIs: subcortical structures	SA vs HC: ↓ globus pallidus and caudate, and correlated with 5-HTT binding. In SA: non-impulsive temperament associated with ↓ globus pallidus GMV
Baldaçara et al 2011 ²²³	BD	20 SA	20 DC, 22 HC	41 (66)	SA: 39.94 (11.2), DC: 41.9 (8.9), HC: 37.7 (13.6)	ROI volume VBM GMV and WMV (SPM), ROIs: cerebellum	SA vs DC: no differences total brain volume or cerebellar volume
Wagner et al 2011 ³⁴	MDD	10 with SA and/or first-degree relative with SA	15 DC, 30 HC	50 (83)	SA: 41.0 (12.5): DC: 34.1 (10.5), HC: 35.1 (10.4).	WB VBM GMV(SPM)	SA vs DC: ↓ rostral ACC (BA24) and right caudate
Wagner et al 2012 ⁵⁸	MDD	10 with SA and/or first-degree relative with SA	15 DC, 30 HC	50 (83)	SA: 41.0 (12.5), DC: 34.1 (10.5), HC: 35.1 (10.4)	WB cortical thickness	SA vs DC: ↓ VLPFC (BA47), DLPFC (BA46) and dorsal ACC (BA32). Patients with own versus relative with SA: no differences in dorsal ACC, VLPFC, DLPFC.
Duarte et al 2017 ¹⁰¹	BD	20 SA	19 DC, 20 HC	34 (57)	SA: 41.10 (12.64): DC: 42.26 (11.70), HC: 37.40 (10.20)	WB and ROI volume VBM GMV (SPM), ROIs: OFC, DLPFC (including IFG), ACC, amygdala, hippocampus, thalamus and insula	SA vs DC: ↑ right rostral ACC (BA24). In SA: HL SA was associated with ↑ insula, LL SA was associated with ↓ OFC (BA47)

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Benedetti et al 2014 ²²⁴	BD	32 SA	104 DC	93 (68)	SA (5-HTTLPR l/l): 41.4 (10.7), SA (5-HTTLPR *s): 46.4 (12.8), DC (5-HTTLPR l/l): 48.5 (10.4), DC (5-HTTLPR *s): 46.8 (12.6)	WB GMV (SPM)	SA vs DC: no differences.
Lee et al 2016 ¹⁵⁰	MDD	19 SA	19 DC, 20 HC	41 (73)	SA: 42.0 (10.8), DC: 41.1 (15.2)	ROI volume GMV (SPM), ROIs not specified	SA vs DC: ↓ right cerebellum and left angular gyrus
Spoletini et al 2011 ¹¹⁰	SCZ	14 SA	36 DC, 50 HC	35 (39)	SA: 42.9 (11.3), DC: 39.8 (11.4): HC: 40.0 (16.6)	ROI volume GMV (FSL), ROIs: lateral ventricles, thalamus, caudate, putamen, pallidum, hippocampus, amygdala, nucleus accumbens	SA vs DC: ↑ right amygdala
Benedetti et al 2011 ³⁵	BD	19 SA (with/without Lithium)	38 DC (with/without Lithium)	38 (67)	SA-L-: 43.6 (10.4), SA-L+: 45.6 (11.3), DC-L-: 45.9 (10.5), DC-L+: 46.2 (13.3)	WB VBM GMV (SPM)	SA vs DC: ↓ DPFC (BA6/8/9), RLPFC (BA10), OFC (BA11/47), dorsal ACC (BA32), parietal and occipital cortex and ↑ in bilateral superior temporal gyrus. SA with lithium vs without: ↑ DPFC (BA6/8), OFC (BA11/47), ACC (BA24/32), parietal and occipital cortex and ↓ in bilateral superior temporal gyrus
Colle et al 2015 ¹¹⁴	MDD	24 SA	39 DC	39 (62)	SA: 44.2 (11.9), DC: 47.7 (12.6)	ROI volume GMV (SACHA, automatic segmentation), ROI: hippocampus.	SA vs DC: ↓ hippocampus. No difference between SA in last month versus SA >1 month ago
Dombrovski et al 2012 ¹²⁹	MDD	13 SA	20 DC, 19 HC	30 (58)	SA: 66.0 (6.4), DC: 67.7 (7.0), HC: 70.5 (7.5)	ROI voxel count basal ganglia (caudate, putamen, pallidum)	SA vs DC: ↓ putamen, associative and ventral striatum voxel count. In SA: delay discounting associated with ↓ putamen voxel count
Cyprien et al 2011 ²²⁵	MDD, ANX, BD	21 SA or SI	234 DC, 180 HC	222 (51)	SA: 72.2 (4.3), DC: 71.0 (3.8), HC: 71.0 (3.8)	ROI volume manual shape tracing, ROI: corpus callosum	SA vs DC: ↓ posterior third of corpus callosum
Hwang et al 2010 ⁷⁰	MDD	27 SA	43 DC, 26 HC	0 (0)	SA: 79.1 (5.6), DC: 79.6 (5.1), HC: 79.5 (4.3)	WB VBM GMV and WMV (SPM)	SA vs DC: ↓ GMV in DPFC (BA6/8/9/46), precentral gyrus, postcentral gyrus, superior parietal lobe, inferior parietal lobe, cuneus, superior temporal gyrus, insula, cerebellum, midbrain, ↓ WMV in DPFC (BA6/8/9/46), precentral and postcentral gyrus, inferior parietal lobe, precuneus, occipital lobe, external capsule, cerebellum
Lopez-Larson et al 2013 ¹³⁴	TBI	19 SA	40 DC, 15 HC	0 (0)	18-55	ROI volume, ROI: thalamus	SA vs DC: ↑ right thalamus
Jia et al 2010 ¹⁶⁵	MDD	16 SA	36 DC, 52 HC	55 (53)	SA: 34.2 (13.7), DC: 34.7 (12.5), HC: 37.1 (16.0)	WB VBM GMV and WMV (SPM)	SA vs DC: no differences

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Kim et al 2015 ²²⁶	PD	12 SA	25 DC	23 (64)	16-60	WB VBM GMV and WMV (SPM)	SA vs DC: no differences
Rentería et al 2017 ¹¹¹	MDD	153 SA or SI+plan, 298 SI-plan	650 DC, 1996 HC	ND	SA+SI: 21.10-53.8 (across 7 samples), DC: 22.9-54.8, HC: 22.9-55.4.	ROI volume, ROIs: nucleus accumbens, amygdala, caudate, hippocampus, pallidum, putamen, thalamus, ICV	SA+SI with plan vs DC: no differences. SI vs DC: no differences

<i>Suicidal ideation studies</i>							
Thomas et al 2004 ²²⁷	PTSD + Childhood maltreatment	47 SI, 17 SA	14 DC, 121 HC	89 (49)	DC: 11.71 (2.6); HC: 11.74 (2.5)	ROI volume manual shape tracing (IMAGE), ROI: pituitary	SI (with SA) vs DC: ↑ pituitary
Taylor et al 2015 ³⁸	MDD	21 SI, including 10 with past SA	53 DC, 91 HC	108 (65)	SI: 33.5 (9.1), DC:37.5 (8.9), HC: 29.9 (9.1)	WB and ROI volume GMV and cortical thickness, ROIs: OFC, cingulate cortex, insula, amygdala, parahippocampus, thalamus, basal ganglia	SI+SA vs DC: ↓ cortical thickness of left insula, left caudal middle frontal gyrus (DLPFC), left superior parietal cortex, left superior temporal gyrus. No GMV differences
Caplan et al 2010 ²²⁸	Epilepsy & MDD, ANX, ADHD	11 SI (No past SA)	40 DC	28 (55)	SI: 11.04 (2.06), DC:9.43 (2.07)	ROI volume manual shape tracing, ROIs: middle frontal gyrus, superior frontal gyrus, OFC, temporal lobe	SI vs DC: ↓ right orbital frontal gyrus WMV and ↑ left temporal lobe GMV

MAGNETIC RESONANCE IMAGING STUDIES OF WHITE MATTER HYPERINTENSITIES

<i>Suicide attempt studies</i>							
Ehrlich et al 2003 ²²⁹	MDD, BD, PsD, conduct/ADHD	43 SA	110 DC	39 (26)	Entire sample: 14.6 (3.4). No further details provided.	WMH (Modified version of Coffey scale)	SA vs DC: ↑ DWMHs in parietal lobes. All SA subjects had lesions in right posterior parietal lobe
Ehrlich et al 2004 ¹⁹⁸	MDD, BD, PsD, conduct/ADHD	43 SA	110 DC	41 (27)	Entire sample: 14.6 (3.4). No further details provided.	WMH (Modified version of Coffey scale)	↑ WMH associated with past SA, driven by PVH. SI not associated with WMH
Ehrlich et al 2005 ²³⁰	MDD	62 SA	40 DC	68 (67)	Entire sample: 26.7 (5.5). No further details provided.	WMH (Modified version of Fazekas scale)	SA vs DC: ↑ PVH, not DWMH. No association with SI
Pompili et al 2007 ²³¹	MDD, BD	29 SA	26 DC	41 (63)	SA: 42.2 (13.5), DC: 44.6 (14.0)	WMH (Modified version of Fazekas scale)	SA vs DC: ↑ WMH. SI was not associated with WMH

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Pompili et al 2008 ²³²	MDD, BD	44 SA	55 DC	57 (58)	SA: 45.57 (16.10), DC: 47.327(14.54)	WMH (Modified version of Fazekas scale)	SA vs DC: ↑ PVH, no difference DWMH
Ahearn et al 2001 ²³³	MDD	20 SA	20 DC	17 (85)	SA: 66.0 (5.8), DC: 66.4 (5.7)	WMH (Coffey and Boyko scales)	SA vs DC: ↑ subcortical GM hyperintensities, and trend towards more PVH
Sachs-Ericsson et al 2014 ²³⁴	MDD	23 SA	223 DC	149 (67)	SA: 66.74 (6.6), DC: 69.8 (7.5)	WMH (Duke Neuropsychiatric Imaging Research Laboratory modified version of MrX software)	SA vs DC: ↑ WM lesions in the left hemisphere. ↑ increase over time in bilateral WMH in SA, which was predicted by the number of depressive episodes.

DIFFUSION TENSOR IMAGING STUDIES							
<i>Suicide attempt studies</i>							
Johnston et al 2017 ³⁶	BD	26 SA	42 DC, 45 HC	43 (63)	14-25	WB FA maps (SPM)	SA vs DC: ↓ FA in left uncinate fasciculus, right uncinate fasciculus and right cerebellum
Lischke et al 2017 ¹⁵⁸	BPD	13 SA	8 DC, 20 HC	41 (100)	18-45	Tractography based FA and MD with seeds in genu, splenium and body of corpus callosum	SA vs DC: no differences. Number of attempts associated with ↓ FA and MD in splenium and FA in the genu
Lee et al 2016 ²³⁵	SCZ	15 SA	41 DC	41 (73)	18-60	WB FA, MD, AD and RD maps (FSL)	SA vs DC: ↑ FA in left corona radiata (anterior, superior, posterior), superior longitudinal fasciculus, posterior limb and retrolenticular part of internal capsule, external capsule, posterior thalamic radiation, sagittal stratum (including inferior longitudinal fasciculus and inferior fronto-occipital fasciculus), ↓ AD in retrolenticular part of internal capsule, posterior thalamic radiation and sagittal stratum. No differences in MD and RD.
Kim et al 2015 ²²⁶	PD	12 SA	25 DC	23 (64)	16-60	FA, MD, AD and RD maps (FSL), Tracts: corona radiata, inferior longitudinal fasciculus, inferiorfronto-occipital fasciculus, superior longitudinal fasciculus, posterior thalamic radiation, internal capsule, splenium of corpus callosum	SA vs DC: ↑ FA in posterior and superior corona radiata, sagittal stratum (including inferior longitudinal fasciculus and inferior fronto-occipital fasciculus), superior longitudinal fasciculus, posterior thalamic radiation, retrolenticular part of internal capsule and splenium of the corpus callosum. No differences in MD, RD and AD. In SA: positive correlation between suicidal ideation and FA of the right retrolenticular part of internal capsule and right and left posterior thalamic radiation. In DC: positive correlation between suicidal ideation and FA of splenium, right retrolenticular part of internal capsule and left posterior thalamic radiation
Mahon et al 2012 ¹⁵⁶	BD	14 SA	15 DC, 15 HC	18 (41)	SA: 33.3 (14.1), DC: 36.5 (12.3), HC: 33.7 (12.6)	WB FA maps	SA vs DC: ↓ FA in white matter tract in medial VPFC. In SA: medial VPFC FA negatively correlated with motor impulsivity

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Olvet et al 2014 ¹⁶⁹	MDD	13 SA	39 DC, 46 HC	52 (53)	18-65	ROI FA and ADC maps (FSL), ROIs: mOFC, DMPFC, rACC, dACC	SA vs DC: ↓ FA in DMPFC. No difference in AD
Jia et al 2010 ¹⁶⁵	MDD	16 SA	36 DC, 52 HC	55 (53)	SA: 34.2 (13.7), DC: 34.7 (12.5), HC: 37.1 (16.0)	WB and ROI analysis of FA, MD and RD maps (DTIstudio), ROIs: bilateral lentiform nucleus, bilateral hippocampus, and bilateral thalamus	SA vs DC: ↓ FA and AD in the left anterior limb of internal capsule and ↓ FA and ↑ RD in right lentiform nucleus
Jia et al 2014 ¹⁶⁶	MDD	23 SA	40 DC, 46 HC	59 (54)	SA: 36.3 (14.5), DC: 34.0 (14.5), HC: 33.3 (11.4)	Tractography based FA with seed in left anterior limb of the internal capsule	SA vs DC: ↓ percentage of projecting fibers connecting the anterior limb of the internal capsule to the left OFC and left thalamus
Lopez-Larson et al 2013 ¹³⁴	TBI	19 SA	40 DC, 15 HC	0 (0)	18-55	ROI FA maps (FSL), ROI: anterior thalamic radiation	SA vs DC: ↑ FA in bilateral anterior thalamic radiation. Positive correlation impulsivity and FA in right anterior thalamic radiation
Cyprien et al 2016 ¹⁵⁷	BD, MDD	45 SA	46 DC, 30 HC	121 (100)	18-50	ROI FA, MD, RD, AD maps (FSL), ROIs: genu, body and splenium of corpus callosum	SA vs HC: no differences that survived multiple comparison correction. Number of attempts associated with ↓ FA in genu, body and splenium of corpus callosum. FA in splenium negatively correlated with suicidal intent.
Bijttebier et al 2015 ¹⁶⁴	past MDD	13 SA	15 DC, 17 HC	32 (72)	18-65	Tractography combined with network-based statistics	SA vs DC: ↓ structural connectivity in network including medial VPFC, temporal gyrus, precuneus, cuneus, parietal cortex, amygdala, hippocampus, occipital regions. Decreased connectivity between left olfactory cortex and left ACC
<i>Suicidal ideation studies</i>							
Myung et al 2016 ¹⁶³	MDD	24 SI	25 DC, 31 HC	52 (65)	18-62	Tractography combined with network-based statistics and graph analysis	SA vs DC: ↓ structural connectivity in left hemisphere network including striatal regions, frontal regions (DLPFC, IFG, lateral OFC and RMPFC), lateral occipital and superior parietal regions. Betweenness centrality of left rostral middle frontal gyrus (DLPFC) positively correlated with suicidal ideation. Participation coefficient of left rostral middle frontal gyrus (DLPFC) positively correlated with impulsivity
Taylor et al 2015 ³⁸	MDD	21 SI, including 10 with past SA	53 DC, 91 HC	108 (65)	20-50	ROI FA and MD maps (FSL), ROIs: internal capsule, thalamic radiation, cingulum bundle, corpus callosum, uncinate fasciculus	SA vs DC: ↑ RD and ↓ FA in the corona radiata, the hippocampal region of the cingulum and the anterior thalamic radiation

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
---------------	-----------------	--------------------	-------------------------	---------------	-----	---------	------------

Symbols & Abbreviations: *Percentages are rounded to the nearest whole number; **Results are reported for SA or SI in comparison with diagnostic controls. If no diagnostic controls were included in the study, results based on SA or SI compared to healthy controls are reported; **AA:** alcohol abuse; **ACC:** anterior cingulate cortex; **AD:** adjustment disorder; **ADHD:** attention deficit hyperactivity disorder; **ANX:** anxiety disorder; **BA:** Broadman's Area; **BD:** bipolar disorder; **BD-P:** bipolar disorder with psychotic symptoms; **BPD:** borderline personality disorder; **DC:** diagnostic controls; **DD:** depressive disorder; **DLPFC:** dorsolateral prefrontal cortex; **DMPFC:** dorsomedial prefrontal cortex; **DWMH:** deep white matter hyperintensities; **FA:** fractional anisotropy; **GM:** grey matter; **GMV:** grey matter volume; **HC:** healthy controls; **HL:** high lethality; **IFG:** inferior frontal gyrus; **LL:** low lethality; **MD:** mood disorder; **MDD:** major depressive disorder; **MDD-P:** major depressive disorder with psychotic features; **ND:** not detailed; **OFC:** orbitofrontal cortex; **PCC:** posterior cingulate cortex; **PD:** personality disorder; **PFC:** prefrontal cortex; **PsD:** psychosis; **PVH:** periventricular hyperintensities; **RLPFC:** rostralateral prefrontal cortex; **RMPFC:** rostromedial prefrontal cortex; **ROI:** region of interest; **SA:** suicide attempt; **SBM:** surface based morphometry; **SCZ:** schizophrenia; **SI:** suicidal ideation; **SUD:** substance use disorder; **SZA:** schizoaffective disorder; **SPM:** Statistical Parametric Mapping toolbox; **TBI:** traumatic brain injury; **VBM:** voxel based morphometry; **VLPFC:** ventrolateral prefrontal cortex; **VMPFC:** ventromedial prefrontal cortex; **WB:** whole brain; **WM:** white matter; **WMH:** white matter hyperintensities; **WMV:** white matter volume; **5-HTTLPR l/l:** serotonin transporter long/long genotype; **5-HTTLPR*s:** serotonin transporter s allele carriers genotype.

Table 2. Findings from Molecular Imaging Studies of Suicidal Thoughts and Behaviors

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
SINGLE PHOTON EMISSION TOMOGRAPHY STUDIES							
<i>Suicide attempt studies</i>							
Audenaert et al 2001 ²³⁶	MDD, AD, PsD	9 SA	12 HC	8 (38)	19-48	123I-5-I-R91150 for 5-HT2a receptors in PFC	SA vs HC: ↓ binding potential of 5-HT2a receptors in PFC
Audenaert et al 2002 ¹⁴³	MDD	20 SA	20 HC	24 (60)	19-50	99mTc-Ethyl Cystine Dimer rCBF SPECT during letter and category fluency	SA vs HC: ↓ perfusion in IFG, ACC, temporal gyrus, hypothalamus during verbal fluency task
van Heeringen et al 2003 ²³⁷	MDD, AD, PsD	9 SA	13 HC	ND	19-47	123I-5-I-R91150 for 5-HT2a receptors in PFC	SA vs HC: ↓ binding potential of 5-HT2a receptors in PFC
Amen et al 2009 ¹⁰²	MDD	12 SA	12 DC, 12 HC	3 (8)	19-64	99mTc HMPAO SPECT to assess rCBF	SA vs DC: ↓ rCBF in subgenual ACC, ↑ rCBF in right insula, dorsal ACC
Willeumier et al 2011 ⁸²	MD	21 SA	36 DC, 27 HC	5 (24)	15-66	99mTc HMPAO SPECT to assess rCBF	SA vs DC: ↓ rCBF in frontal, temporal and parietal regions
Fountoulakis et al 2004 ²³⁸	MDD	13 SA, 10 SI	33 DC	ND	21-60	99mTc HMPAO SPECT to assess rCBF	SA vs DC: no differences. SI vs no-SI: no differences
Henningsson et al 2009 ²³⁹	MDD, PD	9 SA	9 HC	ND	23-67	123I-β-CIT for 5-HTT binding potential, assessment of Val66Met polymorphisms	Within SA: carriers of the Val/Val genotype of Val66Met had ↑ 5HTT binding potential in the parietal cortex and in the occipital lobes
Bah et al 2008 ²⁴⁰	MDD, AD, PD	9 SA	9 HC	0 (0)	23-67	123I-β-CIT for 5-HTT binding potential, assessment of SLC6A4 polymorphisms	SA vs HC: no differences. In SA: presence of S-allele of 5-HTTLPR genotype associated with lower 5-HTT binding potential in frontal, parietal and occipital cortex
Lindström et al 2004 ²⁴¹	MDD, AD, PD	12 SA	12 HC	4 (17)	23-67	123I-β-CIT methods to separate 5-HTT and DAT uptake	SA vs HC: no differences in 5-HTT or DAT binding. In SA: impulsivity associated with ↓ whole brain 5-HTT binding
Ryding et al 2006 ¹³²	MDD, AD, ANX, PD	12 SA	12 HC	4 (17)	23-67	123I-β-CIT methods to separate 5-HTT and DAT uptake	SA vs HC: no differences in 5-HTT or DAT binding. In SA: impulsivity associated with ↓ 5-HTT binding potential in inferior/orbital frontal cortex, temporal regions, midbrain, thalamus, basal ganglia
Vang et al. 2010 ¹³⁰	MDD, AD	7 SA	6 HC	3 (23)	SA: 40 (11.83), matched HC, no details	123I-β-CIT methods to separate 5-HTT and DAT uptake	SA vs HC: not reported. In SA: significant negative correlation between 5HTT binding and globus pallidus volume
<i>Suicidal ideation studies</i>							
Fountoulakis et al 2004 ²³⁸	MDD	13 SA, 10 SI	33 DC	ND	21-60	99mTc HMPAO SPECT to assess rCBF	SI vs DC: no differences. SI vs no-SI: no differences

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
POSITRON EMISSION TOMOGRAPHY STUDIES							
<i>Suicide attempt studies</i>							
Soloff et al 2003 ¹⁹⁶	BPD	13 SA	9 HC	22 (100)	18-49	[18F]FDG PET during rest	SA vs HC: ↓ rCMRglu in bilateral medial OFC
Yeh et al 2015 ¹³³	MDD	5 SA	5 DC, 10 HC	0 (0)	20-25	4-[18F]-ADAM for SERT availability	SA vs DC: ↑ SERT binding potential in the midbrain, thalamus, striatum and PFC. Suicidal ideation associated with ↑ SERT binding potential in the same 4 regions
Soloff et al 2014 ¹⁹⁷	BPD	21 SA	12 DC, 27 HC	32 (53)	BPD: 27.5 (7.2), HC: 28.8 (8.2)	[18F]altanserin for 5-HT2a receptor binding potential	SA vs DC: ↑ binding potential of 5-HT2a receptors in the occipital cortex in females only
Soloff et al 2007 ¹¹⁶	BPD	12 SA	2 DC, 11 HC	25 (100)	19-46	[18F]altanserin for 5-HT2a receptor binding potential	SA vs HC: ↑ binding potential of 5-HT2a receptors in the hippocampus, medial temporal cortex and occipital cortex. No associations with number of attempts
Cannon et al 2006 ⁴⁶	BD	8 SA	10 DC, 37 HC	36 (65)	BD: 30 (9), HC: 32 (9)	[11C]DASB for 5-HTT binding potential	SA vs DC: ↓ 5-HTT binding in the midbrain and ↑ in the rostral ACC
Oquendo et al 2003 ⁸⁴	MDD	25 SA (9 LL, 16 HL)		15 (60)	LL: 30.4 (8.7), HL: 42.9 (10.4)	[18F]FDG PET, fenfluramine vs. placebo challenge	HL vs LL SA: ↓ rCMRglu in ACC (BA24/32), IFG (BA44) and DPFC (BA6/8/9), more pronounced after fenfluramine challenge. Lower VMPFC rCMRglu associated with lower impulsivity, higher suicidal intent and higher lethality
Sullivan et al 2015 ⁴¹	MDD	29 SA	62 DC	59 (65)	18-65	[11C]WAY-100635 for 5-HT1a receptor binding potential	SA vs DC: no differences in 5-HT1a receptor binding. High vs Low lethality SA: ↑ 5-HT1a receptor binding potential in the raphe nuclei. Positive association 5-HT1a receptor binding potential in the raphe nuclei and suicidal intent. Positive association 5-HT1a receptor binding potential in the raphe nuclei and PFC and suicidal ideation
Miller et al 2016 ⁴⁴	BD, MDD	11 SA	6 DC, 31 HC	29 (60)	21 - 61	[11C]DASB for 5-HTT binding potential	SA vs DC: no differences in 5-HTT binding
Yeh et al 2015 ²⁴²	MDD	8 SA	9 DC, 17 HC	18 (53)	20-65	4-[18F]-ADAM for SERT availability	SA vs DC: no differences in SERT binding in individual regions, but ↑ PFC/midbrain SERT binding ratio. Suicide intent positively associated with PFC/midbrain SERT binding ratio
Leyton et al 2006 ⁴⁰	MD, PD, SUD (all HL)	10 SA	16 HC	7 (28)	SA: 37.7 (6.4), HC: 35.5 (12.0)	Alpha-11C-methyl-L-tryptophan trapping as index of 5-HT synthesis	SA vs HC: ↓ 5-HT synthesis in lateral and medial OFC extending into VMPFC, ↑ 5-HT synthesis in thalamus, paracentral lobule, occipital cortex and hippocampus. Negative correlation suicide intent and serotonin synthesis in lateral OFC and VMPFC
Parsey et al 2006 ⁴³	MDD	9 SA	16 DC, 43 HC	49 (69)	MDD: 38.0 (13.4), HC: 38.8 (15.9)	[11C]McN 5652 for 5-HTT binding potential	SA vs DC: no differences in SERT binding

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Miller et al 2013 ²⁴³	MDD	15 SA	36 DC, 32 HC	41 (49)	SA: 38.5 (11.5), DC: 41.0 (10.5), HC: 32.6 (11.3)	[11C]DASB for 5-HTT binding potential	SA vs DC: ↓ 5-HTT binding potential in midbrain
Nye et al 2013 ¹³¹	MDD	11 SA	10 HC	8 (38)	SA: 38.5 (13.6), HC: 21.3 (2.4)	[11C]ZIENT for SERT binding potential	SA vs HC: ↓ 5-HTT in the midbrain/pons and putamen
Mann et al 2018 ⁴⁵	MDD	8 SA	8 DC, 8 HC	ND	21-53	[11C]WAY-100635 for 5-HT1a receptor binding potential, [18F]altanserin for 5-HT2a receptor binding potential	SA vs DC: no difference in 5-HT1a and 5-HT2a receptor binding
Oquendo et al 2016 ⁴²	MDD	51 past SA, 15 future SA	49 DC	61 (61)	18-65	[11C]WAY-100635 for 5-HT1a receptor binding potential, [11C]DASB for 5-HTT binding potential	Future attempt versus no future attempt: no differences in 5-HTT binding. Higher lethality of future attempts associated with ↑ 5-HT1a receptor binding potential in insula, DPFC, ACC and raphe nuclei. SI at follow up associated with ↑ 5-HT1a receptor binding in raphe nuclei, amygdala, hippocampus, parahippocampul gyrus, temporal lobe, ACC, DPFC, medial PFC, OFC, insula, occipital lobe, parietal lobe
Sublette et al 2013 ⁸³	MDD, BD	13 SA	16 DC	19 (66)	SA: 36.0 (11.5), DC: 42.2 (13.0)	[18F]FDG PET, fenfluramine vs. placebo challenge	SA vs DC: ↓ rCMRglu in right DLPFC, more pronounced after fenfluramine challenge, and ↑ rCMRglu in VMPFC, not more pronounced after fenfluramine. SI negatively correlated with rCMRglu in DLPFC
<i>Suicidal ideation studies</i>							
Holmes et al 2018 ⁹⁰	MDD	9 SI	5 DC, 13 HC	13 (48)	MDD: 31 (12), HC: 33 (11)	[11C](R)-PK11195 for TSPO availability index of neuroinflammation)	SI vs no-SI: ↑ TSPO availability in ACC and insula
Kolla et al 2016 ⁸⁹	BPD		28 DC, 14 HC	56 (100)	18-51	[11C]Harmine for MAO-A VT (index of MAO-A density)	Positive association MAO-A VT in PFC and ACC with SI, but also with depressive symptom scores
Van Heeringen et al 2017 ⁸⁵	MDD	17 SI + plan, 11 SI	12 DC, 20 HC	38 (63)	SI+plans: 46.1 (10.9), SI: 42.6 (11.6), DC: 51.2 (6.5), HC: 43.8 (13.1)	[18F]FDG PET during rest	SI+plans vs SI: ↓ rCMRglu in RLPFC and inferior parietal lobe
Ballard et al 2015 ¹⁵¹	MDD	12 SI	8 DC	6 (30)	MDD: 48 (12)	[18F]FDG PET during rest, ketamine challenge	Baseline SI associated with ↑ rCMRglu in infralimbic cortex. Ketamine induced reductions in SI associated with reductions in rCMRglu in infralimbic cortex and increases in rCMRglu in cluster including lingual gyrus, occipital cortex and cerebellum

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
MAGNETIC RESONANCE SPECTROSCOPY STUDIES							
<i>Suicide attempt studies</i>							
Jollant et al 2017 ²⁴⁴	past MDD	15 SA	10 DC, 33 HC	35 (60)	15 -55	Proton MRS; Metabolites: glutamate, glutamine, N-acetylaspartate, myo-inositol, aspartate, glutathione, GABA, N-acetylaspartylglutamate, total choline. ROI: right dorsal PFC	SA vs DC: no significant differences. Choline levels positively correlated with current suicidal ideation
Prescot et al 2018 ⁹¹	history or current MDD, PTSD and/or SUD	57 with SA or SI	24 DC	16 (20)	SA+SI: 37.2 (9.1), DC: 36.2 (9.7)	Proton Magnetic Resonance Spectroscopy. Metabolites: GABA, N-acetylaspartylglutamate, glutamine, glutamate, creatine. ROI: dorsal ACC	SA vs DC: no significant differences. In females only, ↓ GABA in SA+SI compared to DC but no longer significant after correcting for age differences
Rocha et al 2015 ²⁴⁵	past BD	19 SA	21 DC, 22 HC	41 (66)	SA: 39.8 (11.4), DC: 42.0 (8.6), HCL 37.7 (13.5)	Proton MRS; Metabolites: N-acetylaspartate, choline, creatine, myo-inositol. ROI: medial OFC	SA vs DC: no significant differences
<i>Suicidal ideation studies</i>							
Gabbay et al 2017 ¹⁹⁵	MDD		44 DC, 36 HC	46 (50)	12-21	Proton MRS; Metabolites: GABA and GLX. ROI: rostral ACC	No correlation between GABA and GLX levels in rostral ACC and SI

Symbols & Abbreviations: *Percentages are rounded to the nearest whole number; **Results are reported for SA or SI in comparison with diagnostic controls. If no diagnostic controls were included in the study, results based on SA or SI compared to healthy controls are reported; **ACC:** anterior cingulate cortex; **AD:** adjustment disorder; **ANX:** anxiety disorder; **BD:** bipolar disorder; **BPD:** borderline personality disorder; **DAT:** dopamine transporter; **DC:** diagnostic controls; **DLPFC:** dorsolateral prefrontal cortex; **DPFC:** dorsolateral prefrontal cortex; **FDG:** fludeoxyglucose; **GABA:** gamma-aminobutyric acid; **GLX:** glutamate + glutamine; **HC:** healthy controls; **HL:** high lethality; **IFG:** inferior frontal gyrus; **LL:** low lethality; **MAO-A:** monoamine oxidase A; **MDD:** major depressive disorder; **MRS:** magnetic resonance spectroscopy; **ND:** not detailed; **OFC:** orbitofrontal cortex; **PET:** positron emission tomography; **PD:** panic disorder; **PFC:** prefrontal cortex; **PsD:** psychotic disorder; **rCBF:** regional cerebral blood flow; **rCMRglu:** regional cerebral metabolic rate for glucose; **RLPFC:** rostrolateral prefrontal cortex; **SA:** suicide attempt; **SERT:** serotonin transporter; **SI:** suicidal ideation; **SP:** social phobia; **SPECT:** single photon emission computed tomography; **SUD:** substance use disorder; **TSPO:** translocator protein, **VMPFC:** ventromedial prefrontal cortex; **5-HT:** serotonin, **5-HTT:** serotonin transporter

Table 3. Findings from Functional Imaging Studies of Suicidal Thoughts and Behaviors

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
FUNCTIONAL MAGNETIC RESONANCE IMAGING STUDIES							
<i>Suicide attempt studies</i>							
<i>Resting State fMRI</i>							
Cao et al 2016 ¹⁶⁸	DD	35 SA	18 DC, 47 HC	66 (66)	SA: 20.63 (3.65), DC: 21.39 (3.05), HC: 20.53 (1.84)	WB fractional zALFF	SA vs DC: ↑ zALFF in right superior temporal gyrus, left middle temporal gyrus, left middle occipital gyrus, left angular gyrus, ↓ zALFF in left RLPFC. In SA: negative correlation impulsivity and zALFF in left RLPFC
Cao et al 2015 ²⁴⁶	No DX	19 SA	20 HC	22 (56)	SA: 19.8 (1.6), HC: 20.3 (1.7)	WB ReHo	SA vs HC: ↓ ReHo in left fusiform gyrus, lateral OFC (BA47), hippocampus, right angular gyrus, bilateral parahippocampal gyrus, DLPFC (BA46) and cerebellum, ↑ ReHo in right inferior parietal lobe, left precuneus, right medial OFC (BA11)
Zhang et al 2016 ¹⁵⁴	DD	35 SA	18 DC, 47 HC	66 (66)	SA: 20.63 (3.65), DC: 21.26 (3.02), HC: 20.48 (1.86)	ICA of DMN	SA vs DC: ↓ FC in the right precuneus, ↑ FC in the left lingual gyrus and left cerebellum
Kang et al 2017 ¹⁶⁰	MDD	19 SA	19 DC	20 (53)	SA: 42.0 (10.8), DC: 41.1 (15.2)	Amygdala seed-based FC	SA vs DC: ↑ FC left amygdala with right insula and left OFC (BA11), ↑ FC right amygdala with left middle temporal gyrus. In SA: positive correlation between SI and right amygdala FC with right parahippocampal gyrus
<i>Cognitive control</i>							
Pan et al 2011 ²⁴⁷	MDD	15 SA	15 DC, 14 HC	25 (57)	12-17	Go-no-go response inhibition, WB	SA vs DC: ↓ dorsal ACC and insula activation during response inhibition, driven by greater activity in DC compared to both SA and HC (i.e., no evidence for abnormal response inhibition circuitry in SA)
Richard-Devantoy et al 2016 ¹³⁷	MDD	25 SA	22 DC, 27 HC	47 (61)	18-55	Go-no-go response inhibition, WB	SA vs DC: no significant differences. Suicidal intent was positively associated with thalamus activity during response inhibition
Minzenberg et al 2014 ⁵⁴	SCZ	8 SA+SI, 10 SI only	17 DC	10 (26)	18-50	Continuous performance, ROI frontal cortex	SA+SI vs SI: ↓ goal-related left dorsal premotor cortex (BA6) activity. SI vs no-SI: ↓ goal-related ventral ACC, VMPFC, VLPFC, DLPFC (BA9), RMPFC (BA10), RLPFC (BA10) extending to DMPFC (BA8), dorsal ACC (BA24/32). Intensity of ideation negatively correlated with goal-related activity in DMPFC (BA6/8), dorsal ACC (BA32), DLPFC (BA9), RMPFC (BA10), RLPFC (BA10), IFG (BA44/45)

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Minzenberg et al 2015 ⁵²	MD-P	8 SA+SI, 8 SI only	14 DC	13 (43)	18-50	Continuous performance, WB	SA+SI vs SI: ↓ activity in PCC, cuneus and precuneus and ↑ activity right OFC (BA47), right RLPFC (BA10), premotor cortex (BA6), DLPFC (BA9/46), insula during goal-representation. Positive correlation between intensity of SI and goal-related IFG (BA45), lateral and medial OFC (BA11/47), insula and dorsal striatum activity
Minzenberg et al 2015 ¹⁷⁰	SCZ	8 SA+SI, 7 SI only	17 DC	6 (19)	18-50	Continuous performance, dorsal ACC seed-based FC	SA+SI vs SI: ↓ dorsal ACC FC with RMPFC and RLPFC (BA10), DMPFC (BA8), DLPFC (BA9), dorsal ACC (BA32), IFG (BA45), superior temporal gyrus, middle temporal gyrus, precuneus, and PCC during conflict monitoring. SI vs no-SI: ↑ dorsal ACC-precuneus FC during conflict monitoring. Intensity of ideation was positively correlated with dorsal ACC FC with paracentral lobe, precuneus, left caudate, right putamen, right lateral globus pallidus and left thalamus.
Minzenberg et al 2016 ¹⁷¹	MD-P	8 SA+SI, 8 SI only	14 DC	13 (43)	18-50	Continuous performance, dorsal ACC seed-based FC	SA+SI vs SI: ↑ dorsal ACC FC with left DLPFC (BA9), frontal motor areas (BA4/6), inferior temporal gyrus, middle temporal gyrus, dorsal ACC (BA24/32) during conflict monitoring. SI vs no-SI: ↑ dorsal ACC FC with left DLPFC (BA9), DMPFC (BA8), premotor cortex (BA6), superior parietal cortex, inferior parietal cortex, superior temporal gyrus, middle temporal gyrus. Intensity of ideation positively correlated with dorsal ACC FC with bilateral premotor cortex (BA6), inferior and superior parietal cortex, middle and inferior temporal gyri, middle occipital gyrus and occipital regions, negatively correlated with IFG OFC (BA11/47), insula, putamen, globus pallidum, premotor area (BA6), somatosensory cortex (BA5/7) during conflict monitoring.
Minzenberg et al 2015 ⁸⁰	SCZ	8 SA (all past SI)	9 DC (all past SI)	—	18-50	Stroop, ROI frontal cortex	SA vs DC: ↑ left DPFC (BA6/8) during cognitive control.
Vanyukov et al 2016 ⁷⁸	MDD	13 SA	13 DC, 22 HC	30 (63)	46-90	Delay discounting, WB	SA vs DC: ↓ left DLPFC (BA9) activation with increasing value of smaller immediate reward, with a larger decrease in people with better planned SA. Longer versus shorter delay of delayed reward associated with ↓ left parahippocampal gyrus and middle occipital gyrus activation during trials
Decision making and reward processing							
Pan et al 2013 ²⁴⁸	MDD	15 SA	14 DC, 13 HC	23 (55)	12-17	Iowa gambling, WB	SA vs DC: ↓ right thalamus activation during risky choices. No association between activity in thalamus and lethality of attempt and severity of SI

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Olié et al 2015 ⁴⁷	MDD	15 SA	23 DC, 35 HC	0 (0)	18-60	Iowa gambling, ROI OFC, VLPFC, MPFC, ACC, DPFC	SA vs DC: ↓ activation in left DPFC (BA8/9/46) during risky choices, ↑ activation in bilateral DPFC, right OFC (BA11/47), right dorsal and ventral ACC (BA24/32) during winning
Baek et al 2017 ⁹²	past MDD	10 SA	12 DC, 22 HC	24 (55)	18-44	Risk and loss aversion, ROI striatum, OFC, VMPFC, ventral ACC, midbrain	SA vs DC: ↓ subgenual ACC (BA25) activity in response to potential gain. In SA: insula activity correlated negatively with the subjective value of probabilistic gain and loss
Jollant et al 2010 ²⁴⁹	MDD	13 SA	12 DC, 15 HC	0 (0)	22-59	Iowa gambling, ROI OFC, ACC, occipital cortex, precuneus/angular gyrus, thalamus, cerebellum, caudate, cuneus, superior frontal gyrus, parietal cortex	SA vs DC: ↓ lateral OFC (BA47) and occipital cortex activation during risky choices. No differences in activation during gain vs loss trials. No associations with SI
Dombrowski et al 2013 ⁹³	MDD	15 SA	18 DC, 20 HC	31 (59)	60+	Probabilistic reversal learning, WB	SA vs DC: ↓ activity in ventral ACC (BA24/25/32) during expected reward. Poor planning of attempt associated with ↓ activity of paralimbic network (ventral ACC/VMPFC, PCC, precuneus) during expected reward
Memory							
Reisch et al 2010 ⁷²	SR-depression	8 SA		8 (100)	SA: 38.5 (13.1)	Recall of a mental pain, suicide action and neutral conditions using autobiographical scripts of a recent episode of SA, WB	Recall of own suicidal episodes (mental pain and suicide action) vs neutral condition: ↓ activation in left DLPFC (BA46), right RLPFC (BA10), left DMPFC (BA6), ↑ right parahippocampal gyrus, right cuneus, left middle temporal gyrus and cerebellum. Recall of suicide action vs mental pain: ↑ activation in the left DMPFC (BA6), right dorsal ACC (BA32) and left hippocampus
Silvers et al 2016 ⁵⁰	BPD	46 SA	14 DC	60 (100)	SA: 30.0 (9.8), DC: 26.7 (5.0)	Reappraisal of negative autobiographical memories, ROI lateral OFC	SA vs DC: ↑ lateral OFC activation during both reappraisal and immersion, and ↓ precuneus and cuneus activation during reappraisal of memories. In SA: regulation success associated with greater cuneus and precuneus activity
Emotion processing							
Pan et al 2013 ⁷⁴	MDD	14 SA	15 DC, 15 HC	25 (57)	12-17	Viewing of angry, happy and neutral faces, WB activity and right dorsal ACC seed-based FC	SA vs DC: ↑ activity in right dorsal ACC, bilateral primary sensory cortex, left DLPFC (BA9), right middle temporal gyrus, ↓ activity in insula and ↓ FC between dorsal ACC and bilateral insula in response to angry faces. In DC: suicidal ideation was negatively correlated with left DLPFC activation during angry faces
Johnston et al 2017 ³⁶	BD	26 SA	42 DC, 45 HC	43 (63)	14-25	Viewing of happy, neutral and fearful faces, amygdala seed-based FC	SA vs DC: ↓ FC amygdala with left OFC (BA11/47), RPF (BA10), ventral ACC (BA32) in response to happy and neutral faces. SA lethality associated with ↓ FC amygdala with left ventral PFC in response to happy, neutral and fearful faces. In SA: SI was negatively correlated with FC between amygdala and RPF (BA10)

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Olié et al 2015 ⁴⁷	MDD	15 SA	23 DC, 35 HC	0 (0)	18-60	Viewing of angry, happy, sad and neutral faces, ROI OFC, VLPFC, MPFC, ACC, DPFC	SA vs DC: ↑ activation in left OFC and VLPFC in response to angry faces, and ↓ activation in right ACC in response to sad faces
Kim et al 2017 ¹³⁶	DD	14 SA	22 HC	26 (72)	20-47	Viewing of angry, happy, sad and neutral faces, and pictures of suicidal means, WB	SA vs HC: ↑ activation in the left DLPFC, IFG, thalamus and PCC when viewing knives versus natural landscapes. No differences in activity while viewing emotional faces
Jollant et al 2008 ⁴⁸	MDD	13 SA	14 DC, 16 HC	0 (0)	SA: 40.3 (11.3), DC: 43.9 (10.6), HC: 32.4 (9.8)	Viewing of angry, happy and neutral faces, WB	SA vs DC: ↑ activation in in right lateral OFC (BA47) and ↓ activation in right DMPFC (BA6) in response to intense angry faces, ↑ activation in right rostral ACC (BA32) and ↓ activation in right cerebellum to mild happy faces, ↑ activation in right cerebellum in response to mild angry faces
Vanyukov et al 2015 ⁴⁹	MDD	18 SA	13 DC, 18 HC	26 (53)	60+	Angry and fearful faces versus shape matching, WB	SA vs DC: no differences. In SA, higher activation of IFG (BA44/45) while matching angry faces was associated with poorer planning of attempt
Self-referential processing							
Quevedo et al 2016 ¹⁴⁷	DD	43 HS	39 LS, 37 HC	72 (61)	HS: 14.9 (1.6), LS: 14.9 (1.8), HC: 14.5 (1.5)	Emotional Self-Other Morph-Query (ESOM-Q), WB	HS vs LS: ↓ activity in RMPFC (BA10) and in a cluster including parahippocampus, hippocampus, and amygdala during happy self faces versus happy other faces. When controlling for depression severity: HS vs LS: ↓ PCC/precuneus and rostral ACC/BA10 during self faces.
Social exclusion							
Olié et al 2017 ¹⁰³	past MD	36 SA	41 DC, 28 HC	105 (100)	19-54	Social exclusion (Cyberball), WB	SA vs DC: ↓ activation in left supramarginal gyrus and posterior insula during social exclusion
Suicidal ideation studies							
Resting State fMRI							
Cullen et al 2014 ²⁵⁰	MDD		41 DC, 29 HC	54 (77)	MDD: 15.7 (2), HC: 16.0 (2)	Amygdala seed-based FC	No significant correlation between amygdala FC and suicidal ideation
Ordaz et al 2018 ¹⁵²	MDD	40 SI (33% past SA)		30 (75)	14-17	ICA of DMN, ECN, and SN	↓ Network coherence of left ECN, anterior DMN and SN associated with ↑ lifetime SI, only association with left ECN remained after controlling for depressive and anxiety symptom severity. Left ECN also associated with past SA at a trend-level
Chase et al 2017 ¹⁹⁹	MDD	34 SI (18 past SA)	40 HC	50 (68)	18-35	PCC seed-based FC	SI vs HC: ↑ dorsal PCC FC with left middle temporal gyrus. SA vs no-SA: ↑ dorsal PCC FC with left IFG
Du et al 2017 ¹⁵³	MDD	28 SI	20 DC, 30 HC	55 (71)	SI: 32.5 (9.9), DC: 37.1 (10.6), 35.7 (10.2)	Rostral ACC seed-based FC	SI vs DC: ↓ FC right rostral ACC to medial OFC and right middle temporal gyrus, finding for right middle temporal gyrus remained when corrected for depressive symptom severity

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Kim et al 2017 ¹⁶²	MDD	23 SI (7 past SA)	23 DC, 36 HC	68 (83)	SA: 47, SI: 57, DC: 52.7, HC: 56.5	Graph theory and network-based analysis	SI vs DC: ↓ FC in a network including left lateral OFC, left and right caudate, right putamen, left middle temporal gyrus, left and right thalamus and left postcentral gyrus. Node strength, clustering coefficient, regional efficiency of lateral OFC, left and right thalamus negatively correlated with SI
Cognitive control							
Matthews et al 2012 ⁵³	MDD, PTSD, TBI	13 SI	13 DC	0 (0)	SI: 29.5 (4.7), DC: 27.1 (3.6)	Stop signal response inhibition, WB	SI vs DC: ↑ left dorsal ACC (BA24), RLPFC (BA10), DLPFC (BA9/46), supramarginal gyrus, OFC (BA11), DMPFC (BA6) and superior temporal gyrus during error processing
Lee et al 2015 ⁸¹	SCZ		28 DC, 17 HC	8 (18)	SH: 43.6 (11.3), DC: 38.9 (7.3), HC: 37.9 (12.9)	Go-no-go response inhibition, WB	Positive correlation right DLPFC (BA9) activity during response inhibition and current SI in self-harm but no association in no-self-harm group
Zhang et al 2013 ²⁵¹	SCZ	14 SI	19 DC, 15 HC	24 (50)	18-45	N-back, dynamic causal modelling with PCC and MPFC seeds	High vs Low risk: ↓ no difference in activation of and connectivity between VMPFC and PCC. VMPFC activity was positively related with suicide risk
Decision making and reward processing							
Quevedo et al 2017 ¹⁶¹	MDD		38 DC, 30 HC	44 (65)	DC: 30.7 (7.7), HC: 32.0 (6.1)	Card guessing, ventral striatum seed-based FC	Positive correlation between SI and left ventral striatum FC with DMPFC, DLPFC and dorsal ACC during loss trials.
Motor control							
Marchand et al 2012 ¹³⁸	MDD	5 SA	17 DC	0 (0)	22-45	Motor activation, putamen seed-based FC	Positive correlation between SI and left putamen FC with left DMPFC and right putamen, and right putamen FC with left putamen. Left putamen FC with DMPFC also associated with depressive symptom severity
Marchand et al 2013 ²⁵²	MDD, BD	22 SI	18 DC	ND	21-45	Motor activation, PCC seed-based FC	Positive correlation between SI and PCC FC with left DLPFC, DMPFC and IFGs in MDD but not BD. PCC FC with IFG and DLPFC also associated with depressive symptom severity
Marchand et al 2011 ¹³⁵	BD-II	10 SI	6 DC, 19 HC	0 (0)	BD-II: 32.9 (7.5), HC: 33.7 (12.5)	Motor activation, ROI putamen	Negative correlations between a history of SI and activation in left putamen
Emotion processing							
Marchand et al 2011 ¹¹⁷	BD	10 past SI (3 past SA)	6 DC, 19 HC	0 (0)	21-60	Viewing of happy, fearful and neutral faces, ROI amygdala and subgenual ACC	No correlation between brain activity during the task and history of SI

Authors, year	Mental disorder	SA group/ SI group	Groups w/o SA and/or SI	Female n (%)*	Age	Methods	Findings**
Just et al 2018 ⁶⁰	No DX	17 SI (9 past SA)	17 HC	26 (77)	SI: 22.9 (3.6), HC: 22.1 (2.8)	Neurosemantic analyses of concepts related to suicide, positive and negative affect, machine learning on voxels with stable semantic tuning curves	SI could be discriminated from HC with 91% accuracy. Most discriminating regions were the left VMPFC, left DMPFC extending to dorsal ACC, right middle temporal gyrus, left inferior parietal cortex, and left IFG. SI+SA group could be discriminated from SI without SA with 94% accuracy, with most discriminating regions including the left VMPFC, left DMPFC extending to dorsal ACC, right middle temporal gyrus.
Emotion regulation							
Miller et al 2018 ⁷³	No DX	14 SI (4 past SA)	32 without SI	29 (63)	13-20	Emotion regulation, WB	SI vs no-SI: ↓ activity in thalamus, IFG/DLPFC (BA44/9), temporoparietal junction and cerebellum and ↑ activity in temporal pole during passive viewing of negative pictures, ↑ activity in DLPFC (BA9) during regulation of negative emotional pictures
NEAR-INFRARED SPECTROSCOPY STUDIES							
Suicide attempt studies							
Tsujii et al 2017 ²⁵³	MDD	30 SA	38 DC, 40 HC	69 (64)	SA: 37.6 (10.0), DC: 38.8 (9.7), HC: 38.2 (10.2)	Verbal fluency	SA vs DC: ↓ verbal fluency task induced changes in mean oxy-Hb in left precentral gyrus
Suicidal ideation studies							
Pu et al 2015 ²⁵⁴	MDD	31 SI	36 DC, 67 HC	76 (57)	SI: 57.3 (15.7), DC: 58.7 (16.5), HC: 58.1 (17.8)	Verbal fluency	SI vs DC: ↓ verbal Fluency task induced changes in oxy-Hb in right DLPFC, lateral OFC and right RLPFC

Symbols & Abbreviations: *Percentages are rounded to the nearest whole number; **Results are reported for SA or SI in comparison with diagnostic controls. If no diagnostic controls were included in the study, results based on SA or SI compared to healthy controls are reported; **ACC:** anterior cingulate cortex; **BA:** Broadman's Area; **BD:** bipolar disorder; **BD-II:** bipolar II disorder; **BPD:** borderline personality disorder; **DC:** diagnostic controls; **DD:** depressive disorder; **DLPFC:** dorsolateral prefrontal cortex; **DMN:** default mode network; **DMPFC:** dorsomedial prefrontal cortex; **ECN:** executive control network; **FC:** functional connectivity; **HC:** healthy controls; **HS:** high suicidality; **ICA:** independent component analysis; **IFG:** inferior frontal gyrus; **LS:** low suicidality; **MDD:** major depressive disorder; **MD-P:** psychotic mood disorder, **MPFC:** medial prefrontal cortex; **ND:** not detailed; **OFC:** orbitofrontal cortex; **oxy-HB:** oxygen-hemoglobin; **PCC:** posterior cingulate cortex; **PFC:** prefrontal cortex; **PTSD:** posttraumatic stress disorder; **ReHo:** regional homogeneity; **RLPFC:** rostrolateral prefrontal cortex; **RMPFC:** rostromedial prefrontal cortex; **SA:** suicide attempt; **SCZ:** schizophrenia; **SI:** suicidal ideation; **SN:** salience network; **SR-depression:** self-reported depression; **ROI:** region of interest; **TBI:** traumatic brain injury; **VLPCF:** ventrolateral prefrontal cortex, **VMPFC:** ventromedial prefrontal cortex; **WB:** whole brain; **zALFF:** z score amplitude of low frequency fluctuations