

Signatures of adaptation to a monocot host in the plant-parasitic cyst nematode *Heterodera sacchari*

Somnath S. Pokhare<sup>1,2</sup>, Peter Thorpe<sup>3,4</sup>, Pete Hedley<sup>3</sup>, Jennifer Morris<sup>3</sup>, Samer S. Habash<sup>1</sup>, Abdelnaser Elashry<sup>1#\*</sup>, Sebastian Eves-van den Akker<sup>5</sup>, Florian M.W. Grundler<sup>1</sup> & John T. Jones<sup>3,6,\*</sup>

<sup>1</sup>*Institute of Crop Science and Resource Conservation, Department of Molecular Phytomedicine, University of Bonn, 53115 Bonn, Germany.*

<sup>2</sup>Crop Protection Division, ICAR-National Rice Research Institute, Cuttack 753006, Odisha, India.

<sup>3</sup>*The James Hutton Institute, Invergowrie, Dundee, DD2 5DA, UK*

<sup>4</sup>*School of Medicine, University of St Andrews, North Haugh, St Andrews, KY16 9TZ*

<sup>5</sup>*Department of Plant Sciences, University of Cambridge, Cambridge, CB2 3EA,*

<sup>6</sup>*School of Biology, University of St Andrews, North Haugh, St Andrews, KY16 9TZ*

\*Corresponding author: john.jones@hutton.ac.uk

Running title: Parasitism by *Heterodera sacchari*

Key words: CLE peptides, Effectors, *Heterodera sacchari*, transcriptomics

#Current address: Strube Research GmbH & Co., Neue Str. 11, 38838 Schlanstedt, Germany

## Summary

Interactions between plant-parasitic nematodes and their hosts are mediated by effectors; secreted proteins that manipulate the plant to the benefit of the pathogen. To understand the role of effectors in host adaptation in nematodes we analysed the transcriptome of *Heterodera sacchari*, a cyst nematode parasite of rice and sugarcane. A multi-gene phylogenetic analysis showed that *H. sacchari* and the cereal cyst nematode *Heterodera avenae* share a common evolutionary origin and that they evolved to parasitise monocot plants from a common dicot-parasitic ancestor. We compared the effector repertoires of *H. sacchari* with those of the dicot parasites *H. glycines* and *Globodera rostochiensis* to understand the consequences of this transition. While, in general, effector repertoires are similar between the species, comparing effectors and non-effectors of *H. sacchari* and *G. rostochiensis* shows that effectors have accumulated more mutations than non-effectors. Although most effectors show conserved spatiotemporal expression profiles and likely function, some *H. sacchari* effectors are adapted to monocots. This is exemplified by the plant-peptide hormone mimics, the CLAVATA3/EMBRYO SURROUNDING REGION-like (CLE) effectors. Peptide hormones encoded by *H. sacchari* CLE effectors are more similar to those from rice than those from other plants, or those from other plant-parasitic nematodes. We experimentally validate the functional significance of these observations by demonstrating that CLE peptides encoded by *H. sacchari* induce a short root phenotype in rice, whereas those from a related dicot parasite do not. These data provide a functional example of effector evolution that co-occurred with the transition from dicot-parasitic to a monocot-parasitic lifestyle.

## Introduction

Plant-parasitic nematodes cause damage to world agriculture valued at approximately \$80 billion each year (Nicol *et al.*, 2011; Jones *et al.*, 2013). However, many growers (particularly in developing nations) are unaware of the damage caused by plant-parasitic nematodes because they are small, soil-dwelling and cause non-specific symptoms. The true extent of the damage caused by plant-parasitic nematodes is therefore likely to be considerably higher than this. The cyst forming nematodes (*Heterodera* and *Globodera* spp.) are one of the most damaging plant-parasitic nematode groups. These sedentary endoparasitic pathogens are obligate biotrophs and induce the formation of a large multinucleate, metabolically active syncytium in the roots of their hosts (reviewed in Perry, Moens & Jones, 2018). The syncytium is formed by progressive fusion of cells from an initial syncytial cell selected by the nematode after invasion of the host (Sobczak & Golinowski, 2011). Syncytium formation is associated with profound changes in host gene expression and modulation of the cell cycle (*e.g.* Siddique & Grundler, 2018).

The interactions between plants and their pathogens, including cyst nematodes, are mediated by effectors; secreted proteins that manipulate the host to the benefit of the pathogen. Effectors from cyst nematodes are primarily produced in the dorsal and subventral pharyngeal gland cells and are secreted into the host via the stylet. The availability of genome and/or transcriptome resources from a range of cyst nematodes (*e.g.* Gao *et al.*, 2003; Cotton *et al.*, 2014; Kumar *et al.*, 2014; Zheng *et al.*, 2015; Eves van den Akker *et al.*, 2016) has facilitated identification of effectors from these species. Strategies for identifying effectors from genome and transcriptome resources include identifying secreted proteins that are upregulated at parasitic stages of the nematode (*e.g.* Thorpe *et al.*, 2014; Espada *et al.*, 2016) or direct sequencing of mRNA extracted from aspirated gland cells (Maier *et al.*, 2013) followed in both cases by *in situ* hybridisation to confirm expression in the pharyngeal gland cells. More recently, it has been shown that promoters associated with genes expressed in the gland cells can be used to identify comprehensive lists of effectors from diverse PPN species (Eves-van den Akker & Birch, 2016; Espada *et al.*, 2018). As a result of these studies, effectors have been identified from a wide range of PPN with subsequent functional studies showing that they have roles in various stages of the plant-nematode interaction including metabolism of the plant cell wall to facilitate invasion and migration, suppression of host defences and initiation of the syncytium (reviewed in Jones & Mitchum, 2018).

The plant cell wall is the first significant barrier that any invading pathogen, including plant-parasitic nematodes, will need to overcome to infect a plant and plant-parasitic nematodes are well equipped with proteins that allow them to metabolise the plant cell wall. The first effector identified from any plant-parasitic nematode was a beta 1,4 endoglucanase (cellulase) from the potato cyst nematode *Globodera rostochiensis* (Smant *et al.*, 1998) and a range of cell wall degrading and modifying proteins

have subsequently been identified as cyst nematode effectors including pectate lyase (Popeijus *et al.*, 2000), GHF43 Arabinase (Cotton *et al.*, 2014), GH53 Arabinogalactan endo- 1,4 beta galactosidase (Vanholme *et al.*, 2009), expansins (Qin *et al.*, 2004) and proteins encoding carbohydrate binding domains (Hewezi *et al.*, 2008). All of these genes, as well as others encoding chorismate mutase (Jones *et al.*, 2003) and proteins potentially involved in vitamin biosynthesis (Craig *et al.*, 2008), have been acquired by horizontal gene transfer from bacteria (reviewed in Kikuchi *et al.*, 2017). Cyst nematode effectors have also been identified that suppress host defence responses, most notably several members of the SPRYSEC family of effectors (Postma *et al.*, 2012; Mei *et al.*, 2015) and a modified ubiquitin extension protein (Chronis *et al.*, 2013). The details of how cyst nematodes induce the formation of their syncytium in the roots of their hosts are less clear, although effectors that are likely to be important in this process have been characterised. Another novel effector (19C07) has been identified from *Heterodera glycines* and *Heterodera schachtii* that interacts with the LAX3 auxin influx transporter (Lee *et al.*, 2011). In addition, all cyst nematodes studied to date produce effectors that include variable numbers of C-terminal repeats encoding peptides similar to CLAVATA3/EMBRYO SURROUNDING REGION-like (CLE) peptides. Functional studies have shown that the nematode peptides can complement mutant *Arabidopsis* lacking these peptides (Wang *et al.*, 2005). The CLE proteins are modified by plant cell machinery in a manner similar to that of the endogenous proteins and the CLE peptides themselves subsequently interact with the CLAVATA2 receptor protein, which is required for nematode parasitism (Replogle *et al.*, 2011; Replogle *et al.*, 2013). In plants, CLE peptides regulate cell differentiation and thus contribute to the control of meristem maintenance in shoots, roots and vascular tissues. The ability to produce endogenous CLE peptides is likely to be a key factor in the ability to induce a feeding structure in plants.

Many plant species are parasitised by cyst nematodes, including monocots and dicots. However, each individual species of cyst nematode tends to have relatively narrow host range, with some exceptions. Although little is known about the molecular determinants of host range in plant parasitic nematodes, effectors have been shown to have a central role in this process in other pathogens (reviewed by Stam *et al.*, 2014). A view has emerged “non-host resistance” of closely related species is most likely due to recognition of effectors by a resistance gene, while failure to infect a more distantly related species is most likely due to the incompatibility of effectors with their cognate targets (Schulze-Lefert & Panstruga, 2011). For example, comparisons of the genomes of *Phytophthora infestans* and a closely related species *Phytophthora mirabilis*, which infects *Mirabilis jalapa* revealed that 82 of 345 genes which showed signs of positive selection could encode effector sequences (Raffaele *et al.*, 2010). Subsequent work on orthologous effectors that encode protease inhibitor from the two species showed that the protease inhibitor effectors from each of these species interact specifically with

protease targets from their respective host plants (Dong *et al.*, 2014). Host-pathogen co-evolution is therefore reflected in adaptations of effectors for function in the host.

*Heterodera sacchari* is an increasingly economically important pathogen of several monocot species including rice and sugarcane. Crop losses due to *H. sacchari* can exceed 40%, with more severe damage reported under rain fed upland conditions (Kyndt *et al.*, 2014). Compared to other characterised cyst nematodes, the biology of *H. sacchari* is unusual in that it is restricted to monocots and reproduces mainly by mitotic parthenogenesis (CABI, 2014). Here we sequenced and assembled the transcriptome of *H. sacchari* and use these data to reconstruct the evolutionary history of this and related species. These data suggest that *H. sacchari* and a related monocot parasite *H. avenae* evolved to parasitise monocot plants secondarily, from a dicot-parasitic ancestor. To explore the genetic changes associated with the evolutionary transition from dicot to monocot parasite, we identified homologues of previously identified effectors in *H. sacchari*. We show that in general the effectors have diversified in sequence more than non-effectors when compared to their most similar homologue in a dicot-parasite, and that while some effectors show conserved expression profiles and likely function, specific aspects of the effector repertoire of *H. sacchari* appear to be adapted to monocots.

## Results and Discussion

### The transcriptome of *H. sacchari*

A total of 17,086,132 paired end reads were obtained from the cDNA extracted from second stage juveniles (J2) and parasitic stage nematodes 15 days after infection. These sequence reads have been submitted to SRA (accession number PRJEB28025). The sequence reads were pooled and assembled into a single reference transcriptome of 44,230 transcripts after filtering with Transrate and removing contaminants. Assemblies are available at <https://zenodo.org/deposit/1324265>. CEGMA analysis showed that 88% of the core eukaryotic genes were present as full-length transcripts, with a further 6% represented by partial length transcripts. BUSCO analysis using the metazoan data set suggests that the assembly contains 76% complete BUSCO sequences and a further 6.0% fragmented sequences. Seventeen percent of the BUSCO genes were not identified. Comparisons with published gene models from cyst nematode genome sequences, showed that 34,203 (77.3%) and 30,482 (68.9%) of the *H. sacchari* transcripts matched sequences in *G. rostochiensis* and *G. pallida* respectively (evalue  $1e^{-10}$ ). Taken together, these data suggest that the assembly produced represents a large proportion of *H. sacchari* transcriptome of these life stages.

### *H. sacchari* evolved to parasitise monocots from a dicot-parasitic ancestor.

We used a subset of 96 core eukaryotic genes conserved in *H. sacchari* and 18 related species to reconstruct a multi-gene phylogeny (Figure 1). This phylogeny robustly positions *H. sacchari* and the

related cereal cyst nematode *H. avenae* in a mono-phyletic sub-clade of monocot parasites, nested within a clade of related dicot-parasites of the genera *Heterodera* and *Globodera*. The most parsimonious explanation for this is that 1) *H. sacchari* and *H. avenae* share a common, monocot-parasitic, ancestor, and 2) this last common monocot-parasitic ancestor evolved to parasitise a monocot host secondarily, from a dicot-parasitic ancestor. This provides the comparative framework to explore the genes conserved, and the genes diverged, during adaptation to monocot parasitism by nematodes.

### **Genes encoding cell wall modifying enzymes acquired via horizontal gene transfer in the *H. sacchari* transcriptome**

The plant cell wall is the first significant barrier to an invading pathogen, and while largely similar between dicots and monocot, there are noticeable differences in composition. Plant-parasitic nematodes in general are well equipped with proteins that allow them to modify and degrade specific components of the plant cell wall. Many of these genes were acquired via horizontal gene transfer from bacteria (Danchin *et al.*, 2010).

The transcriptome of *H. sacchari* contains representatives of most previously described cases of horizontal gene transfer in related plant-parasitic nematodes, including a wide range of cell wall degrading enzymes (Table 1) and several other sequences putatively acquired by horizontal gene transfer (e.g. the GH32 invertases and chorismate mutase (Danchin *et al.*, 2016; Jones *et al.*, 2003)). Analysis of the expression profiles of one of the *H. sacchari* GHF5 cellulases and the chorismate mutase showed that, as in other cyst nematodes, expression was restricted to the subventral pharyngeal gland cells in J2s and while the cellulase was upregulated at J2 the chorismate mutase was expressed throughout the life cycle (Figure 2). These sequences may therefore play a similar role in the biology of *H. sacchari* and other cyst nematodes.

Notably however, both the *H. sacchari* and *H. avenae* transcriptomes lack sequences similar to GH53 Arabinogalactan endo- 1,4 beta-galactosidase, in spite of the fact that all other cyst nematodes analysed to date, which parasitise dicots, have such proteins. While the absence of evidence in transcriptome datasets is not necessarily evidence of absence in the genome, it nevertheless reflects the host range of these species. Cell walls of commelinoid monocots (which include the main hosts of *H. sacchari* and *H. avenae*) have a different composition compared to those of dicots and contain relatively low amounts of pectic polysaccharides, including the substrate of the GH53 enzymes (Vogel, 2008). Without genomic resources we cannot conclude whether the conspicuous absence in the transcriptomes of the monocot parasites is because these genes have been lost entirely or that they are not expressed under these conditions.

### An overview of *H. sacchari* effector-like sequences

Effectors modulate plant process to promote disease, and are often finely tuned to their host. To determine whether the effector repertoire of *H. sacchari* reflects its secondary adaptation to a monocot host, we first identified and characterised effectors in the *H. sacchari* transcriptome by building on a detailed genome wide analysis of cyst nematode effectors performed for *G. rostochiensis* (Eves-van den Akker *et al.*, 2016a). Using *G. rostochiensis* effectors as a starting point for comparative purposes, 185 of the 295 identified a similar sequence in the *H. sacchari* transcriptome (at  $e$  value  $<1e^{-10}$ ). We compared the similarity of *H. sacchari* effector-like sequences and non-effector-like sequences to their most similar homologue in the *G. rostochiensis* genome. This showed that, on average, putative effectors of *H. sacchari* are more different to their closest homologue in *G. rostochiensis* than non-effectors are to their corresponding closest homologue (Figure 3). Taken together, these data suggest that while most effectors are apparently conserved, they have nevertheless accumulated more mutations than non-effectors since these species diverged. Further analysis of the differences between the variation in effectors and non-effectors was undertaken. 21% of the mutations were classed as nonsynonymous for the putative effectors, versus 22% for other transcripts, however, non-synonymous SNPs in the putative effectors were more often predicted to have a significant effect on the coding sequence compared to other transcripts: 0.93% vs 0.82% were classed as “high impact” for putative effectors and other transcripts respectively. All of the following high impact categories were more common in the putative effectors compared to other transcripts: gain of premature start codon, in frame deletions, frameshift variants, start loss, stop gain. Moreover, the rate of variants (i.e. length of transcript/ number of variants) was significantly higher for the putative effectors versus non-effectors (mean=1754 vs mean=1243 for putative effectors vs other transcripts respectively, Mann Whitney U test:  $1.98e^{-41}$ ).

Consistent with this observation, analysis of effectors from a range of pathogens, including PPN, has shown that they are under strong diversifying selection pressure. For example, whole genome resequencing of five pathotypes of *G. rostochiensis* showed that effectors contain significantly more variants and more nonsynonymous variants per gene than do randomly selected non-effector genes (Eves-van den Akker *et al.*, 2016a). On a more detailed scale, the SPRYSEC effector (RBP1) from *G. pallida*, which is recognised by the *Gpa2* resistance gene of potato, has been subjected to positive selection at several different residues, including the residue that determines recognition or evasion by *Gpa2* (Sacco *et al.*, 2009).

The SPRY domain family is greatly expanded in cyst nematodes with 295 sequences predicted in *G. pallida* of which approximately 30 have a signal peptide (and thus encode SPRYSEC effectors) (Cotton *et al.*, 2014), and 71 sequences in *G. rostochiensis* of which 17 may be SPRYSEC effectors (Eves-van den Akker *et al.*, 2016a). While we find a similar proportion of SPRYs to SPRYSECs in *H. sacchari* (80

SPRY domain encoding transcripts and 6 SPRYSECs), with similar spatial expression in the dorsal pharyngeal gland cell (Figure 4), the temporal expression is unusual. At least two *H. sacchari* SPRYSECs are upregulated in parasitic stage nematodes (Figure 4) and this contrasts with the *G. pallida* SPRYSECs which tend to be upregulated in J2s (Mei *et al.*, 2015). Several SPRYSECs have been shown to suppress host defence signalling (Postma *et al.*, 2012; Mei *et al.*, 2015; Mei *et al.*, 2018). What role the *H. sacchari* SPRYSECs play in infection is yet to be determined.

Other notable peculiarities in the effector repertoire of *H. sacchari* are the HYP effectors. The HYP effectors, first identified in *G. pallida* and *G. rostochiensis*, are secreted from the gland cells surrounding the main anterior sense organs and show unprecedented variability between individuals (Eves-van den Akker *et al.*, 2014). The HYP effectors are strongly upregulated in parasitic stages of *G. pallida* and can be subdivided into three subfamilies based on the presence and type of subfamily-specific tandem repeats. The transcriptome of *H. sacchari* contains two full length HYP-like transcripts (primarily represented in the 15 days post infection library) with unusual characteristics. A phylogenetic analysis of the *H. sacchari* HYP-like sequences with all other known full-length HYP effectors (n = 65) shows that the *H. sacchari* HYP-like sequences group in a separate subfamily (named Subfamily HYP-0, Figure 5). Comparison of the *H. sacchari* predicted sequences with those from *G. pallida* shows that these sequences have both highly conserved regions that flank the variable domain but they are the first to encode a major structural variant outside the hypervariable domain, and the region corresponding to the “hypervariable domain” contains a novel sequence that has a very limited repeat structure (sequence features summarised in Figure 5).

### **Specialisation of *H. sacchari* CLE-like effectors to parasitism of a monocot host**

One class of *H. sacchari* effectors shows signs of adaptation to a monocot host. The CLAVATA3/EMBRYO SURROUNDING REGION-like (CLE) effectors mimic plant peptide hormones and have been characterised in a number of plant-parasitic nematodes. The transcriptome of *H. sacchari* contained several partial transcripts that encode proteins with similarity to CLE-like effectors from other plant-parasitic nematodes. Using a novel approach based on identifying reads that map to partial transcripts and carrying out a local assembly, we were able to computationally assemble six unique transcripts to recapitulate the full length open reading frame, resulting in 5 unique polypeptide sequences from methionine to stop. As for other cyst nematodes, the *H. sacchari* sequences each encoded a signal peptide at the N-terminus followed by an N-terminal domain (Figure S2), which in other cyst nematodes enables translocation into the apoplast after the protein is secreted into the plant cell (Wang *et al.*, 2010a). All six of these *H. sacchari* transcripts encode a single canonical CLE domain at their C terminus. Another transcript (DN37996\_c0\_g2\_i1) encodes a tandemly repeated motif with no clear homology to canonical CLE domains (despite the similarity of the rest of the protein

sequence to CLE-effectors of other cyst nematodes). The six *H. sacchari* CLE-effector-like sequences can be divided into two groups based at least in part on their signal peptide and CLE domains. The CLE domains within each group are identical in protein and nucleic acid sequence (Figure 6A).

Given that CLE peptides vary between plant species in general, and monocots and dicots in particular, we hypothesised that the CLE domains of CLE-like effectors in *H. sacchari* may have specialised prior to/concurrent with the transition from dicot- to monocot-parasite. To test this hypothesis, we analysed a database of 391 CLE peptides from 20 plant species collated from Zhang *et al.* (2014) and Oelkers *et al.* (2008). We created an all by all matrix of similarity between plant CLE peptides and *H. sacchari* CLE peptides based on a normalized BLOSUM62 score. We then used this matrix to generate a CLE similarity network (Figure 6B), highlighting the host (rice *Oryza sativa*) and the nematode (*H. sacchari*) CLEs. Three of the *H. sacchari* CLEs form part of a well-connected cluster containing 14 other CLEs, 8 of which are from rice. On closer inspection, we found that three of these *H. sacchari* CLEs (Hsac\_DN37996\_c0\_g4\_i1, Hsac\_DN35920\_c0\_g3\_i1, and Hsac\_DN35920\_c0\_g2\_i1) were sequence-identical mimics of eight rice CLEs (OS\_GEN\_Os01g55080\_1, OSEST\_NP890021\_\_1, OSEST\_TC269510\_\_1, OS\_TA\_AK108976\_\_1, OS\_TA\_CA758496\_\_1 OS\_GEN\_Os01g48230\_1, OS\_GEN\_Os01g48260\_1, OSEST\_TC271220\_\_1) in the terminal 13 amino acids of its CLE domain. The 6 non-rice plant CLEs within this cluster are from *Arabidopsis* and *Populus*, but none have direct connections with *H. sacchari* CLEs. Three other *H. sacchari* CLEs (Hsac\_DN37996\_c0\_g1\_i1, Hsac\_DN49341\_c0\_g1\_i1, Hsac\_DN35920\_c0\_g1\_i1) form a second small cluster with one other rice CLE (OS\_GEN\_Os05g48730\_1). *H. sacchari* CLEs are the only nematode CLEs with a connection to a rice CLE and the only connections between *H. sacchari* CLEs and plant CLEs are with rice. Interestingly, there are several other sequence divergent CLE clusters in rice with no corresponding *H. sacchari* CLE, suggesting selective mimicry of a subset of this family. It is known that CLE family members in dicots have diverse roles (*e.g.* Mitchum *et al.*, 2008) and mimicry of a subset of rice CLEs by *H. sacchari* may reflect the need to target the function of a subset of the full rice CLE complement.

The functional significance of the similarity between the *H. sacchari* CLE peptides and those from rice was experimentally investigated by taking advantage of the short root phenotype observed from overexpression or exogenous application of CLEs (*e.g.* Fiers *et al.*, 2004 and Chen *et al.*, 2015). We synthesised a synthetic version of the *H. sacchari* CLE peptide that had the highest connectivity with rice CLEs in the network (sequence identical to the rice CLEs in the 13 terminal amino acids). In order to confirm that this is indeed a true *H. sacchari* sequence we cloned the gene encoding this peptide from *H. sacchari* gDNA (Figure S3). We analysed the effect of the peptide when applied exogenously to rice seedlings on root growth when compared to a randomised version of this CLE, or a CLE from

the dicot-parasitic *H. glycines* (Wang *et al.*, 2005). This analysis showed that the peptide from *H. sacchari* induced a short root phenotype in rice whereas peptides from *H. glycines* and a shuffled peptide used as a control had no effect (n=24 per condition,  $p < 0.001$ , Tukey's HSD, Figure 7).

## Conclusions

We use whole transcriptome sequencing to show that *H. sacchari* and the related *H. avenae* evolved to parasitise a monocot host from a last common dicot-parasitic ancestor. We mine these data to identify and characterise the cell wall degrading enzyme and effector complement of *H. sacchari*. Finally, we show that while *H. sacchari* has a similar effector arsenal to the related cyst nematodes that parasitise dicots, the CLE effectors provide a paradigm of functional adaptation for parasitism of a monocot host. These data build a foundation on which to explore novel effectors involved in monocot-parasitism.

## Materials and Methods

### Biological material

*Heterodera sacchari* was cultured on rice cv Nipponbare as described in Pokhare *et al.* (2019). Briefly, plants were grown in a potting mixture of sand, field soil and organic matter (70:29:1) and were infected with second stage juveniles. After 12 weeks, watering of the plants was stopped and the plants were allowed to dry for 2 weeks. Cysts were collected by Cobb's decanting and sieving method using standard protocols (Cobb, 1918). The cysts were surface sterilised and placed in 3 mM ZnCl<sub>2</sub> to initiate hatching. The resulting J2s were collected every 5 days and either frozen in liquid nitrogen or used immediately for infection of new plants grown on pluronic gel (Pokhare *et al.*, 2019). Parasitic stage female nematodes were collected by hand 15 days after infection and frozen in liquid nitrogen until use. The original population of *H. sacchari* was provided by Dr D. Coyne (International Institute for Tropical Agriculture). The population was originally derived from rice in Nigeria and was maintained at The International Institute for Tropical Agriculture on susceptible *Oryza sativa*. Although the original population was derived from a field population of cysts, this nematode reproduces by mitotic parthenogenesis meaning that genetic diversity within the population is likely to be low, facilitating much of the bioinformatic analysis.

### Transcriptome sequencing

RNA was extracted from second stage juvenile and parasitic stage female nematodes using a Nucleospin RNA XS kit (Macherey Nagel) following the manufacturer's instructions. The quantity and integrity of RNA were assessed using a Bioanalyser. Library preparation for RNAseq was performed

using the TruSeq RNA Library Prep Kit v2 (Illumina) as recommended by the manufacturer (Illumina; Protocol # 15026495, revision F). Separate libraries were generated from the juvenile (1 µg) and female (500 ng) total RNA samples, using single-end TruSeq Index Adapters AR002 (CGATGT) and AR004 (TGACCA), respectively. Each library was quality checked using a Bioanalyzer 2100 (Agilent) and quantified using a Qubit fluorometer (Thermo Fisher). Equal molarities of each library were combined and run at 12 pM on a MiSeq (Illumina) generating paired-end 2x 250 bp reads, as recommended. A fastq file was generated for each sample using MiSeq Control Software (version 2.6) for downstream QC and analysis.

### Transcriptome assembly, quality control, and annotation

Scripts used to analyse the data are available at: [https://github.com/peterthorpe5/Hsac\\_transcriptome](https://github.com/peterthorpe5/Hsac_transcriptome). Assemblies are available at <https://zenodo.org/deposit/1324265>. Raw reads are available under primary accession PRJEB28025 and secondary accession ERP110186.

The 17725370 read pairs (5,636,376 from juveniles and 11,329,919 from females with 759075 undetermined) were first quality control checked using FastQC (Andrews, 2010), then quality trimmed using Trimmomatic version 0.32 (Q15) (Bolger *et al.*, 2014). The resulting 17086132 read pairs were assembled using Trinity version 2.1.1 (kmer length 25) (Haas *et al.*, 2013). The resulting assembly was subjected to quality control filtering using Transrate version 1.0.1 (Smith-Unna *et al.*, 2016). Low quality/scoring transcripts were removed (based on read mapping to the assembly). Coding sequences were predicted using TransDecoder (Using DIAMOND BLASTP versus Swiss prot and HMM search versus Pfam A domain guides). The predicted coding sequences were DIAMOND BLASTP version 0.7.9 (Buchfink *et al.*, 2014) searched against GenBank NR database (May 2017). The results were interrogated for their alien index (AI) score using [https://github.com/peterthorpe5/public\\_scripts/tree/master/Lateral\\_gene\\_transfer\\_prediction\\_tool](https://github.com/peterthorpe5/public_scripts/tree/master/Lateral_gene_transfer_prediction_tool), which predicts putative horizontal gene transfer events. Any sequence with an AI greater than 25 and that had a BLAST hit identity greater than 70% to a non-metazoan was flagged as putative contamination. Putative contaminant sequences were removed, and the corresponding transcripts were removed from the assembly. BUSCO version 1.1b (Simão *et al.*, 2015) and CEGMA version 2.4 (Parra *et al.*, 2007) were used to quantify the completeness of the assembly. The resulting coding sequences were annotated using Trinotate (Grabherr *et al.*, 2011), HMMER (Finn *et al.*, 2011), Pfam (Yang & Smith 2013), SignalP 4.0 (Petersen *et al.*, 2011), TMHMM (Krogh *et al.*, 2001), BLAST (Altschul *et al.*, 1990), gene ontology (Ashburner *et al.*, 2000), eggNOG V3.0 (Powell *et al.*, 2012) and RNAmmer (Lagesen *et al.*, 2007). SPRYSECs were identified using HMMER, and Phobius (Käll *et al.*, 2007) for those

containing a secretion signal. CAZyme analysis was performed by the CAYZme analysis toolkit using both BLAST and PFAM analysis on the online server (Park *et al.*, 2010). BLAST2GO was used to identify and analyse GO terms (Conesa *et al.*, 2005). The *H. sacchari* transcripts were compared to all genes predicted in the *G. pallida* and *G. rostochiensis* genomes with matches identified using an e value cut off of  $10^{-10}$ . Similarly, candidate *H. sacchari* effectors were identified by comparisons with effectors from other cyst nematode species (Gao *et al.*, 2003; Thorpe *et al.*, 2014; Eves-van den Akker *et al.*, 2016a) using an e value cut off of  $10^{-10}$ .

### Phylogenetics

Ninety six CEGMA genes conserved in the genome and or transcriptome resources of 18 plant-parasitic nematodes species were used for phylogenetic reconstruction. The species used were *H. avenae* (Kumar *et al.*, 2014), *H. glycines* (Masonbrink *et al.*, 2019), *H. schachtii* (S. Eves-van den Akker, pers. comm.), *G. pallida* (Cotton *et al.*, 2014), *G. rostochiensis* (Eves-van den Akker *et al.*, 2016a), *G. ellingtonae* (Phillips *et al.*, 2017), *R. reniformis* (Eves-van den Akker *et al.*, 2016b), *R. similis* (Jacob *et al.*, 2008), *M. arenaria* (Blanc-Mathieu *et al.*, 2017), *M. javanica* (Blanc-Mathieu *et al.*, 2017), *M. incognita* (Blanc-Mathieu *et al.*, 2017), *M. hapla* (Opperman *et al.*, 2008), *P. coffeae* (Burke *et al.*, 2015), *N. aberrans* (Eves-van den Akker *et al.*, 2014), *B. xylophilus* (Kikuchi *et al.*, 2011), *C. elegans* (*C. elegans* sequencing consortium, 2018), *C. briggsae* (Hillier *et al.*, 2007) and *L. elongatus* (Danchin *et al.*, 2017). The protein sequences of CEGMA genes were aligned and refined individually using MUSCLE (Edgar, 2004). Individual alignments were concatenated and submitted to the IQtree online server with associated partition file. Model selection was carried out on each partition and a concatenated multi-gene phylogeny generated using the ultra-fast mode and 1000 bootstraps (Trifinopoulos *et al.*, 2016).

### CLE effector identification and network analysis

Several partial transcripts encoding proteins with similarity to CLE-effectors from various plant-parasitic nematodes were identified in the transcriptome assembly. Partial transcripts were computationally extended using an iterative approach of mapping and overlap assembly using the wrapper script provided with MITObim (Hahn *et al.*, 2013). Only the deduced amino acid sequences of full length CLEs were used for further analyses. The gene encoding the CLE peptide used for functional analyses was cloned and sequenced in order to ensure the validity of the computational approach. To compare *H. sacchari* CLE domains to those of plants a database of CLE peptides was collated from Zhang *et al.* (2014), and Oelkers *et al.* (2008). All CLE domains were combined into a fasta file and filtered for those with missing information. Custom python script 1 was used to construct

an all vs all similarity matrix based on the BLOSUM62 scores between amino acids. Custom python script 2 was used to parse the matrix and normalise self-normalise similarity scores. Custom python script 3 was used to convert the matrix to gefx, which was loaded into Gephi to visualise the network (Bastian *et al.*, 2009). Custom python scripts are available under github repository <https://github.com/sebastianevda/>.

### **Cloning and characterisation of candidate effector sequences**

The complete ORFs of selected genes were amplified from cDNA of pre-parasitic second stage juveniles, or complete genes from gDNA extracted from cysts. PCR products were purified using the Qiagen PCR Purification kit and cloned into the pGEMT Easy or pCR8/TOPO/GW vectors following the manufactures guidelines.

### **Analysis of expression profiles of candidate effectors**

The spatial expression patterns of candidate effectors from *H. sacchari* were investigated using *in situ* hybridisation of digoxigenin labelled probes to juvenile nematodes as previously described (Thorpe *et al.*, 2014). Negative controls for *in situ* hybridisations were carried out using sense probes and gave no signal (Figure S1). The expression of candidate effectors across the *H. sacchari* life cycle (eggs, pre-parasitic juveniles, 15 & 25 days post infective females) was analysed using quantitative PCR (qRT-PCR) with gene specific primers. Around 3000 eggs and pre-parasitic juveniles were collected for each replicate. To collect eggs, cysts were crushed under a binocular microscope and eggs were transferred into an Eppendorf tube in sterile water. Juveniles were harvested every 5 days as described above and flash frozen in liquid nitrogen. For the other life stages (15 & 25 days post inoculation), approximately 150 parasitic stage nematodes were collected from rice grown in pluronic gel as described above. Nematodes were stored at -80° C and used subsequently for RNA extraction. Total RNA was extracted as above and were processed using Agilent 2100 bioanalyzer system to check the quality and quantity of extracted RNA. The RNA samples with RNA integrity number (RIN) of more than 8 were used for cDNA synthesis using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems) and oligo-dT primer. The cDNA was tested for expression changes during the nematode life stages using the Stepone Plus Real-Time PCR System (Applied Biosystems) with cycling parameters of 95° C for 15 s and 60° C for 30 s (40 cycles) for amplification. Each sample well contained 10 µl of Fast SYBR Green qPCR Master Mix (Invitrogen), 9 µl of the gene specific primer mixture with a final concentration 1 µM for each primer and 1 µl of cDNA. The data was analysed using the Stepone Plus Real-Time PCR software to create Ct values and relative expression was calculated following Pfaffl (2001). Elongation Factor 1 alpha (EF1 alpha) was used as internal control for all experiments. Three biological replicates

from each stage and three technical replicates for each biological replicate were used for qPCR studies. Details of the primers used in this study are provided in Supplementary Table 1.

### Functional analysis of *H. sacchari* CLE sequences

To analyse the *in vivo* function of CLE peptides, we developed a protocol for exogenous application to rice seedlings, similar to that described for *A. thaliana* (Wang *et al.*, 2010). The terminal 13 amino acid CLE domain of nematode CLE-like sequences was synthesised with hydroxy prolines in positions 5 and 8 for *H. sacchari* (H-Lys-Arg-Leu-Ser-Hyp-Gly-Gly-Hyp-Asp-Pro-Gln-His-His-OH), *H. glycines* (H-Lys-Arg-Leu-Ser-Hyp-Ser-Gly-Hyp-Asp-Pro-His-His-His-OH). As a control, a shuffled version of the *H. sacchari* CLE sequence was also synthesised (H-Asp-His-Ser-Hyp-Gly-Gln-His-Arg-Hyp-Lys-Pro-Gly-Leu-OH). Seeds of rice (cv Nipponbare) were surface sterilised and allowed to germinate on sterile, wet filter paper. After 7 days, plants of similar size were transferred to plates containing ½ MS supplemented with 10 µM of the relevant peptide or an equivalent volume of sterile distilled water and left for 10 days at 25 °C in 16h light/8h dark. After this time plants were removed from the plates and the roots were washed to remove any adhering medium. The length of the longest root was measured for 24 biological replicates of each condition.

### Data availability statement

All sequence data, including assemblies, raw sequence reads and scripts used for processing, are publicly available as described in the materials and methods section. All other materials described in this MS are available upon request.

### Acknowledgements

Somnath Pokhare was supported by a fellowship provided by the Indian Council of Agricultural Research (ICAR), Government of India. This work was also supported by Royal Society International Exchanges award number IE150670 and by the Rural and Environmental Science and Analytical Services Division of the Scottish Government. SEvdA is supported by Biotechnology and Biological Sciences Research Council grants BB/R011311/1 and BB/S006397/1. This work benefited from interactions funded through COST Action FA1208. The authors thank Dr Danny Coyne (IITA) for providing the starter culture of the *H. sacchari* population used for this work. Bioinformatics and Computational Biology analyses were supported by the University of St Andrews Bioinformatics Unit which is funded by a Wellcome Trust ISSF award [grant 105621/Z/14/Z].

### Author contributions

SP, PT and SH carried out the majority of the experimental work and contributed to the writing and revision of the manuscript. PH and JM undertook and assisted in analysis of the sequencing data. JJ, FMW, SEvdA and AE designed the study, contributed to data analysis and interpretation of the results. JJ, PT and SEvdA led the writing of the manuscript with contributions from FMA and AE.

### Conflicts of interest

The authors have no conflicts of interest to declare.

## References

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. (1990). Basic local alignment search tool. *Journal of Molecular Biology* 215, 403-410.
- Andrews S. (2010). FastQC: a quality control tool for high throughput sequence data. Available online at: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>
- Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T. & Harris, M.A. (2000). Gene Ontology: tool for the unification of biology. *Nature Genetics* 25, 25-20.
- Bastian M., Heymann S. & Jacomy M. (2009). Gephi: an open source software for exploring and manipulating networks. International AAAI Conference on Weblogs and Social Media.
- Blanc-Mathieu, R., Perfus-Barbeoch, L., Aury, J. M., Da Rocha, M., Gouzy, J., Sallet, E., et al. (2017). Hybridization and polyploidy enable genomic plasticity without sex in the most devastating plant-parasitic nematodes. *PLoS genetics* 13(6), 1-36.
- Bolger, A.M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30, 2114-2120.
- Buchfink, B., Xie, C. & Huson, D.H. (2014). Fast and sensitive protein alignment using DIAMOND. *Nature Methods* 12, 59-60.
- Burke, M., Scholl, E.H., Bird, D. McK., Schaff, J.E., Colman, S.D., Crowell, R., Diener, S., Gordon, O., Graham, S., Wang, X., Windham, E., Wright, G.M. & Opperman, C.H. (2015). The plant parasite *Pratylenchus coffeae* carries a minimal nematode genome. *Nematology* 17, 621-637.
- C. elegans* sequencing consortium (1998). Genome sequence of the nematode *C. elegans*: A platform for investigating biology. *Science* 282, 2012-2018.
- CABI (2014). *Heterodera sacchari*. 139, 43–45.

- Chen, S., Lang, P., Chronis, D., Zhang, S., De Jong, W.S., Mitchum, M.G. & Wang, X. (2015). In planta processing and glycosylation of a nematode CLAVATA3/ENDOSPERM SURROUNDING REGION-like effector and its interaction with a host CLAVATA2-like receptor to promote parasitism. *Plant Physiology* 167, 262-272.
- Chronis, D., Chen, S., Lu, S., Hewezi, T., Carpenter, S.C.D., Loria, R., Baum, T.J. & Wang, X. (2013). A ubiquitin carboxyl extension protein secreted from a plant-parasitic nematode *Globodera rostochiensis* is cleaved *in planta* to promote plant parasitism. *The Plant Journal* 74, 185-196.
- Cotton, J.A., Lilley, C.J., Jones, L.M. *et al.* (2014). The genome and life-stage specific transcriptomes of *Globodera pallida* elucidate key aspects of plant parasitism by a cyst nematode. *Genome Biology* 15, R43. DOI: 10.1186/gb-2014-15-3-r43.
- Cobb, N.A. (1918). Estimating the nema population of soil, with special reference to the sugar-beet and root-gall nemas, *Heterodera schachtii* Schmidt and *Heterodera radicola* (Greef) Müller and with a description of *Tylencholaimus aequalis* n. sp. Washington, DC, USA, National Government Printing Office.
- Conesa, A., Götz, S., García-Gómez, J.M., Terol, J., Talón, M. & Robles, M. (2005). Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21, 3674-3676.
- Craig, J.P., Bekal, S., Hudson, M., Domier, L., Niblack, T. & Lambert, K.N. (2008). Analysis of a horizontally transferred pathway involved in vitamin B6 biosynthesis from the soybean cyst nematode *Heterodera glycines*. *Mol. Biol. Evol.* 25, 2085-2098.
- Danchin E.G.J., Rosso, M.N., Vieira, P., de Almeida-Engler, J., Coutinho, P.M., Henrissat, B. and Abad, P. (2010). Multiple lateral gene transfers and duplications have promoted plant parasitism ability in nematodes. *Proceedings of the National Academy of Sciences* 107 (41), 17651–17656.
- Danchin, E.G.J., Guzeeva, E., Mantelin, S., Berepiki, A. & Jones, J.T. (2016). Horizontal gene transfer from bacteria has enabled the plant-parasitic nematode *Globodera pallida* to feed on host-derived sucrose. *Molecular Biology and Evolution* 33, 1571-1579.
- Danchin, E.G.J., Perfus-Barbeoch, L., Rancurel, C., Thorpe, P., Da Rocha, M., Bajew, S., Neilson, R., Sokolova, E., Da Silva, C., Guy, J., Labadie, K., Esmenjoud, D., Helder, J., Jones, J. T. & Eves-van den Akker, S. (2017). The Transcriptomes of *Xiphinema index* and *Longidorus elongatus* suggest independent acquisition of some plant parasitism genes by horizontal gene transfer in early-branching nematodes. *Genes* 8(10): 287. DOI: 10.3390/genes8100287.
- Dong, A., Stam, R., Cano, L.M., Song, J., Sklenar, J., Yoshida, K., Bozkurt, T.O., Oliva, R., Liu, Z., Tian, M., Win, J., Banfield, M.J., Jones, A.M.E., van der Hoorn, R.A.L. & Kamoun, S. (2014). Effector specialization in a lineage of the Irish Potato Famine Pathogen. *Science* 343, 552-555.

- Edgar, R.C. (2004). MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* 5:113.
- Espada, M., Silva, A.C., Eves-van den Akker, S., Cock, P.J.A., Mota, M. & Jones, J.T. (2016). Identification and characterization of parasitism genes from the pine wilt nematode *Bursaphelenchus xylophilus* reveals a multi-layered detoxification strategy. *Molecular Plant Pathology* 17, 286-295.
- Espada, M., Eves-van den Akker, S., Maier, T., Vijaypalani, P., Baum, T.J., Mota, M. & Jones, J.T. (2018). STATAWAARS: A promoter motif associated with the tissue-specific transcriptome of the major effector-producing glands of the plant-parasitic nematode *Bursaphelenchus xylophilus*. *BMC Genomics* 19: 553.
- Eves-van den Akker, S. & Birch, P.R.J (2016). Opening the effector protein toolbox for plant–parasitic cyst nematode interactions. *Molecular Plant* 9, 1451-1453.
- Eves-van den Akker, S., Laetsch, D.R., Thorpe, P. *et al.* (2016a). The genome of the yellow potato cyst nematode, *Globodera rostochiensis*, reveals insights into the bases of parasitism and virulence. *Genome Biology* 17:124. DOI: 10.1186/s13059-016-0985-1.
- Eves-van den Akker, S., Lilley, C.J., Yusup, H.B., Jones, J.T. & Urwin, P.E. (2016b). Functional C-terminally encoded plant peptide (CEP) hormone domains evolved *de novo* in the plant parasite *Rotylenchulus reniformis*. *Molecular Plant pathology* 17(8), 1265–1275. DOI: 10.1111/mp.12402.
- Eves-van den Akker, S., Lilley, C.J., Jones, J.T. & Urwin, P.E. (2014). Identification and characterisation of a hyper variable apoplastic effector gene family of *G. pallida*. *PLoS Pathogens* 10(9): e1004391.
- Eves-van den Akker, S., Lilley, C.J., Danchin, E.G.J., Rancurel, C., Cock, P.J.A., Urwin, P.E. & Jones, J.T. (2014). The transcriptome of *Nacobbus aberrans* reveals insights into the evolution of sedentary endoparasitism in plant-parasitic nematodes. *Genome Biology and Evolution* 6 (9), 2181-2194.
- Fiers, M.A., Hause, G., Boutilier, K.A., Casamitjana-Martinez, E., Weijers, D., Offringa, R., van der Geest, L., Lookeren Campagne, M. van & Liu, C.M. (2004). Mis-expression of the CLV3/ESR-like gene CLE19 in *Arabidopsis* leads to a consumption of root meristem. *Gene*, 327, 37– 49.
- Finn, R.D., Clements, J. & Eddy, S.R. (2011). HMMER web server: interactive sequence similarity searching. *Nucleic Acids Research* 39, W29-W37.
- Gao, B., Allen, R., Maier, T., Davis, E.L., Baum, T.J. & Hussey, R.S. (2003). The parasitome of the phytonematode *Heterodera glycines*. *Molecular Plant-Microbe Interactions* 16, 720–726. <http://dx.doi.org/10.1094/MPMI.2003.16.8.720>.
- Gheysen, G. & Mitchum, M.G. (2011). How nematodes manipulate plant development pathways for infection. *Current Opinion in Plant Biology* 14, 1-7.

- Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R. & Zeng, Q. (2011). Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology* 29, 644-652.
- Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J., Couger, M.B., Eccles, D., Li, B., Lieber, M. & MacManes, M.D. (2013). *De novo* transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature Protocols* 8, 1494.
- Hahn, C., Bachmann, L., & Chevreur, B. (2013). Reconstructing mitochondrial genomes directly from genomic next-generation sequencing reads—a baiting and iterative mapping approach. *Nucleic acids research* 41(13), e129-e129.
- Hewezi, T., Howe, P., Maier, T., Hussey, R.S., Mitchum, M.G., Davis, E.L. & Baum, T.J. (2008). Cellulose binding protein from the parasitic nematode *Heterodera schachtii* interacts with *Arabidopsis* pectin methylesterase: cooperative cell wall modification during parasitism. *Plant Cell* 20, 3080-3093.
- Jacob, J., Mitreva, M., Vanholme, B. & Gheysen, G. (2008). Exploring the transcriptome of the burrowing nematode *Radopholus similis*. *Molecular Genetics and Genomics* 280, 1-17. <https://doi.org/10.1007/s00438-008-0340-7>.
- Hillier, L.W., Miller, R.D., Baird, S.E., Chinwalla, A., Fulton, L.A., Koboldt, D.C., & Waterston, R.H. (2007). Comparison of *C. elegans* and *C. briggsae* genome sequences reveals extensive conservation of chromosome organization and synteny. *PLoS biology* 5(7), e167.
- Jones, J.T., Furlanetto, C., Bakker, E., Banks, B., Blok, V.C., Chen, Q. & Prior, A. (2003). Characterisation of a chorismate mutase from the potato cyst nematode *Globodera pallida*. *Molecular Plant Pathology* 4, 43-50.
- Jones, J.T., Haegeman, A., Danchin, E.G.J., Gaur, H.S., Helder, J., Jones, M.G.K., Kikuchi, T., Manzanilla-López, R., Palomares-Rius, J.E., Wesemael, W.M.L. & Perry, R.N. (2013). Top 10 plant-parasitic nematodes in molecular plant pathology. *Molecular Plant Pathology* 14, 946–961. doi:10.1111/mpp.12057.
- Jones, J.T & Mitchum, M.G (2018). Biology of effectors. In Perry, R.N., Moens, M & Jones, J.T. (Eds). Cyst Nematodes. Wallingford, CABI, pp. 74-88.
- Käll, L., Krogh, A. & Sonnhammer, E.L. (2007). Advantages of combined transmembrane topology and signal peptide prediction—the Phobius web server. *Nucleic Acids Research* 35 (suppl\_2), W429-W432.
- Kikuchi, T., Cotton, J.A., Dalzell, J.J., Hasegawa, K., Kanzaki, N., McVeigh, P., et al. (2011). Genomic insights into the origin of parasitism in the emerging plant pathogen *Bursaphelenchus xylophilus*. *PLoS Pathogens* 7(9), e1002219.
- Kikuchi, T., Eves-van den Akker, S. & Jones, J.T. (2017). Genome evolution of plant-parasitic nematodes. *Annual Review of Phytopathology* 55, 333-354.

- Krogh, A., Larsson, B., Von Heijne, G. & Sonnhammer, E.L. (2001). Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *Journal of Molecular Biology* 305, 567-580.
- Kumar, M., Gantasala, N.P., Roychowdhury, T., Thakur, P.K., Banakar, P., Shukla, R.N., *et al.* (2014) De Novo transcriptome sequencing and Analysis of the Cereal Cyst Nematode, *Heterodera avenae*. *PLoS ONE* 9, e96311.
- Kyndt, T., Fernandez, D. & Gheysen G. (2014). Plant-parasitic nematode infections in rice: molecular and cellular insights. *Annual Review of Phytopathology* 52, 135-153.
- Lagesen, K., Hallin, P., Rødland, E.A., Stærfeldt, H.H., Rognes, T. & Ussery, D.W. (2007). RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Research*, 35, 3100-3108.
- Lee, C., Chronis, D., Kenning, C., Peret, B., Hewezi, T., Davis, E.L., Baum, T.J., Hussey, R.S., Bennet, M. & Mitchum, M.G. (2011). The novel cyst nematode effector protein 19C07 interacts with the *Arabidopsis* auxin influx transporter LAX3 to control feeding site development. *Plant Physiol.* 155, 866-880.
- Maier, T.R., Hewezi, T., Peng, J.Q. & Baum, T.J. (2013). Isolation of whole esophageal gland cells from plant-parasitic nematodes for transcriptome analyses and effector identification. *Mol. Plant-Microbe Interact.* 26, 31-35.
- Mei, Y., Thorpe, P., Guzha, A., Haegeman, A., Blok, V.C., MacKenzie, K., Gheysen, G., Jones, J.T & Mantelin, S. (2015). Only a small subset of the SPRY domain gene family in *Globodera pallida* is likely to encode effectors, two of which suppress host defences induced by the potato resistance gene Gpa2. *Nematology* 17, 409-424.
- Masonbrink, R., Maier, T. R., Muppirala, U., Seetharam, A. S., Lord, E., Juvale, P. S., ... & Mimee, B. (2019). The genome of the soybean cyst nematode (*Heterodera glycines*) reveals complex patterns of duplications involved in the evolution of parasitism genes. *BMC Genomics*, 20(1), 119
- Mei, Y., Wright, K.M., Haegeman, A., Bauters, L., Diaz-Granados, A., Goverse, A., Gheysen, G., Jones, J.T. & Mantelin, S. (2018). The *Globodera pallida* SPRYSEC effector GpSPRY-414-2 that suppresses plant defenses targets a regulatory component of the dynamic microtubule network. *Front. Plant Sci.* 9:1019.
- Mitchum, M.G., Wang, X. & Davis, E.L. (2008). Diverse and conserved roles of CLE peptides. *Current Opinion in Plant Biology* 11, 75-81.
- Nicol, J.M., Turner, S.J., Coyne, D.L., den Nijs, L., Hockland, S. & Maafi, Z.T. (2011). Current nematode threats to world agriculture. In: *Genomics and Molecular Genetics of Plant-Nematode Interactions* In Jones, J.T., Gheysen, G. and Fenoll, C. (Eds.) pp 21-44. Heidelberg, Germany: Springer. DOI 10.1007/978-94-007-0434-3\_1.

- Oelkers, K., Goffard, N., Weiller, G.F., Gresshoff, P.M., Mathesius, U. & Frickey, T. (2008). Bioinformatic analysis of the CLE signaling peptide family. *BMC Plant Biology* 8(1), 1.
- Opperman, C.H., Bird, D.M., Williamson, V.M., Rokhsar, D.S., Burke, M., Cohn, J., Cromer, J., Diener, S., et al. (2008). Sequence and genetic map of *Meloidogyne hapla*: A compact nematode genome for plant parasitism. *Proc. Natl Acad. Sci. USA* 105 (39), 14802–14807. <https://doi.org/10.1073/pnas.0805946105>.
- Park, B.H., Karpinets, T.V., Syed, M.H., Leuze, M.R. & Uberbacher, E.C. (2010). CAZymes Analysis Toolkit (CAT): web service for searching and analyzing carbohydrate-active enzymes in a newly sequenced organism using CAZy database. *Glycobiology* 20, 1574-1584.
- Parra, G., Bradnam, K. & Korf, I. (2007). CEGMA: a pipeline to accurately annotate core genes in eukaryotic genomes. *Bioinformatics* 23, 1061-1067.
- Perry, R.N., Moens, M & Jones, J.T. (2018). Cyst Nematodes. Wallingford, CABI.
- Pokhare, S.S., Habash, S.S., Jones, J.T., Elashry, A. And Grundler, F.M.W. (2019). *In vitro* life cycle of *Heterodera sacchari* on Pluronic gel. *Nematology* 21, 573-579.
- Popeijus, H., Olvermars, H., Jones, J.T., Blok, V.C., Goverse, A., Helder, J., Schots, A., Bakker J. & Smant, G. (2000). Degradation of plant cell walls by a nematode. *Nature* 406,36-37.
- Petersen, T.N., Brunak, S., von Heijne, G. & Nielsen, H. (2011). SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nature Methods* 8, 785.
- Phillips, W. S., Howe, D. K., Brown, A. M. V., Eves-van den Akker, S., Dettwyler, L., Denver, D. R. and Zasada, I. A. (2017). The Draft Genome of *Globodera ellingtonae*. *Journal of Nematology* 49(2), 127–128.
- Postma, W.J., Slootweg, E.J., Rehman, S., Finkers-Tomczak, A., Tytgat, T.O.G., van Gelderen, K., Lozano-Torres, J.L., Roosien, J., Pomp, R., van Schaik, C., Bakker, J., Goverse, A. & Smant, G. (2012). The effector SPRYSEC-19 of *Globodera rostochiensis* suppresses CC-NB-LRR mediated disease resistance in plants. *Plant Physiol.* 16, 944-954.
- Powell, S., Szklarczyk, D., Trachana, K., Roth, A., Kuhn, M., Muller, J., Arnold, R., Rattei, T., Letunic, I., Doerks, T. & Jensen, L.J. (2012). eggNOG v3. 0: orthologous groups covering 1133 organisms at 41 different taxonomic ranges. *Nucleic Acids Research*, 40(D1), D284-D289.
- Qin, L., Kudla, U., Roze, E., Goverse, A., Popeijus, H., Nieuwland, J., Overmars, H., Jones, J.T., Schots, A., Smant, G., Bakker, J. & Helder, H. (2004). Identification of a functional expansin, a non-enzymatic cell wall-loosening agent, from the plant parasitic nematode *Globodera rostochiensis*. *Nature* 427, 30.
- Raffaele, S., Farrer, R.A., Cano, L.M., Studholme, D.J., MacLean, D., Thines, M., Jiang, R.H., Zody, M.C., Kunjeti, S.G., Donofrio, N.M., Meyers, B.C., Nusbaum, C. & Kamoun, S. (2010). Genome evolution following host jumps in the Irish potato famine pathogen lineage. *Science* 330, 1540–1543.

Replogle, A., Wang, J., Bleckmann, A., Hussey, R.S., Baum, T.J., Shinichiro, S., Davis, E.L., Wang, X., Simon, R. & Mitchum, M.G. (2011). Nematode CLE signaling in *Arabidopsis* requires CLAVATA2 and CORYNE. *The Plant Journal* 65, 430-440.

Replogle, A., Wang, J., Paolillo, V., Smeda, J., Kinoshita, A., Durbak, A., Tax, F.E., Wang, X., Sawa, S. & Mitchum, M.G. (2013). Synergistic interaction of CLAVATA1, CLAVATA2, and RECEPTOR-LIKE PROTEIN KINASE 2 in cyst nematode parasitism of *Arabidopsis*. *Molecular Plant-Microbe Interactions* 26, 87-96.

Sacco, M.A., Koropacka, K., Grenier, E., Jaubert, M.J., Blanchard, A., Goverse, A., Smant, G. & Moffett, P. (2009). The cyst nematode SPRYSEC protein RBP-1 elicits Gpa2- and RanGAP2-dependent plant cell death. *PLoS Pathogens* 5, e1000564.

Schulze-Lefert P. & Panstruga R. (2011). A molecular evolutionary concept connecting nonhost resistance, pathogen host range, and pathogen speciation. *Trends Plant Sci.* 16, 117–125  
10.1016/j.tplants.2011.01.001.

Siddique S. & Grundler F.M.W. (2018) Parasitic nematodes manipulate plant development to establish feeding sites. *Current Opinion in Microbiology* 46, 102-108.

<https://doi.org/10.1016/j.mib.2018.09.004>

Smant, G., Stokkermans, J.P., Yan, Y., de Boer, J.M., Baum, T.J., Wang, X., Hussey, R.S., Gommers, F.J., Henrissat, B., Davis, E.L., Helder, J., Schots, A. & Bakker, J. (1998). Endogenous cellulases in animals: isolation of beta-1, 4-endoglucanase genes from two species of plant-parasitic cyst nematodes. *Proc. Natl Acad. Sci. USA* 95,4906-4911.

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V. & Zdobnov, E.M. (2015). BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31, 3210-3212.

Sobczak, M. and Golinowski, W. (2011). Cyst nematodes and syncytia. In: Jones, J.T., Gheysen, G. and Fenoll, C. (Eds). *Genomics and Molecular Genetics of Plant-Nematode Interactions*. Springer Academic Publishers, Dordrecht, The Netherlands, pp. 61-82. DOI 10.1007/978-94-007-0434-3\_4.

Stam, R., Mantelin, S., McLellan, H. & Thilliez, G. (2014). The role of effectors in nonhost resistance to filamentous plant pathogens. *Frontiers in Plant Science* 5, 582.  
<http://doi.org/10.3389/fpls.2014.00582>

Smith-Unna, R., Bournnell, C., Patro, R., Hibberd, J.M. & Kelly, S. (2016). TransRate: reference-free quality assessment of de novo transcriptome assemblies. *Genome Research* 26, 1134-1144.

Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A., & Minh, B.Q. (2016). W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Research*, 44 (Web Server issue), W232–W235. <http://doi.org/10.1093/nar/gkw256>.

- Thorpe, P., Mantelin, S., Cock, P.J.A. *et al.* (2014). Characterisation of the full effector complement of the potato cyst nematode *Globodera pallida*. *BMC Genomics* 15, 923. <https://doi.org/10.1186/1471-2164-15-923>.
- Vanholme, B., Haegeman, A., Jacob, J., Cannoot, B. & Gheysen, G. (2009). Arabinogalactan endo- 1,4-beta- galactosidase: a putative plant cell wall-degrading enzyme of plant parasitic nematodes. *Nematology* 11, 739-747.
- Vogel, J. (2008). Unique aspects of the grass cell wall. *Current Opinion in Plant Biology* 11, 301-307.
- Wang, X., Mitchum, M.G., Gao, B., Li, C., Diab, H., Baum, T.J., Hussey, R.S. & Davis, E.L. (2005). A parasitism gene from a plant-parasitic nematode with function similar to CLAVATA3/ESR (CLE) of *Arabidopsis thaliana*. *Molecular Plant Pathology* 6, 187-191.
- Wang, J., Joshi, S., Korkein, D. & Mitchum, M.G. (2010). Variable domain I of nematode CLEs directs post-translational targeting of CLE peptides to the extracellular space. *Plant Signaling & Behavior* 5(12), 1633-1635, DOI: 10.4161/psb.5.12.13774.
- Wang, J., Lee, C., Replogle, A., Joshi, S., Korkein, D., Hussey, R., Baum, T.J., Davis, E.L., Wang, X. And Mitchum, M.G. (2010a). Dual roles for the variable domain in protein trafficking and host-specific recognition of *Heterodera glycines* CLE effector proteins. *New Phytologist* 187, 1003–1017.
- Yang, Y. & Smith, S.A. (2013). Optimizing *de novo* assembly of short-read RNA-seq data for phylogenomics. *BMC Genomics* 14, 328.
- Zhang, Y., Yang, S., Song, Y., & Wang, J. (2014). Genome-wide characterization, expression and functional analysis of CLV3/ESR gene family in tomato. *BMC genomics* 15(1), 827.
- Zheng, M., Long, H., Zhao, Y., Lin, L., Xu, D., Zhang, H., *et al.* (2015) RNA-Seq based identification of candidate parasitism genes of cereal cyst nematode (*Heterodera avenae*) during incompatible infection to *Aegilops variabilis*. *PLoS ONE* 10: e0141095. <https://doi.org/10.1371/journal.pone.0141095>.

**Figure Legends:**

**Figure 1:** Phylogenetic analysis of 96 CEGMA genes conserved in 19 species of nematode shows that *H. sacchari* and *H. avenae* share a common monocot parasitic ancestor that most likely evolved from a parasite of dicots (indicated by the arrow). The scale bar indicates substitutions per base. Node labels indicate support values for 1000 bootstraps.

**Figure 2:** Analysis of spatio-temporal expression patterns of cellulase (left panels) and chorismate mutase (right panels) identified in the *H. sacchari* transcriptome. *In situ* hybridisation analysis showed that both genes are expressed in the subventral gland cells of second stage juveniles (J2s) of *H. sacchari* (purple staining; upper panels). Analysis using qPCR shows that the cellulase is expressed specifically at the J2 life stage while the chorismate mutase is expressed at all life stages tested (lower panels). Error bars represent standard errors across the biological replicates.

**Figure 3:** Histograms showing distributions of percentage identity between *H. sacchari* effectors (left panel) and non-effectors (right panel) with their homologues in *G. rostochiensis*. This analysis suggests that effector-like transcripts have accumulated more mutations than other transcripts since the divergence of these two species.

**Figure 4:** Analysis of spatio-temporal expression patterns of two putative SPRYSEC-encoding transcripts of *H. sacchari*. *In situ* hybridisation analysis showed that both genes are expressed in the dorsal gland cell of *H. sacchari* (purple staining, arrows; upper panels). Analysis by qPCR shows that both sequences are upregulated in later parasitic stage nematodes (lower panels). Error bars represent standard errors across the biological replicates.

**Figure 5:** Overview of the HYP gene family in cyst nematodes. Left, a schematic representation of the *H. sacchari* HYP-like sequences shows the position of a novel structural variant outside the hyper-variable domain. The faded region of the hyper-variable domain of HYP-1 and -3 represents the region of variable sequence. Right, a midpoint re-rooted phylogenetic construction of HYP sequences from *G. pallida* and *H. sacchari* position those of *H. sacchari* as a distinct sub-family (named here sub-family-0).

**Figure 6:** CLE-like effectors of *H. sacchari*. A) A midpoint re-rooted phylogenetic construction of the six complete CLE-like sequences in the *H. sacchari* transcriptome. The CLE domain containing effectors

form two groups, each of which has a distinct CLE domain (red box). B) A similarity network of CLE domains from plants and *H. sacchari*. CLEs from selected species are highlighted: Rice in green; *H. sacchari* in orange; *H. glycines* in purple; *G. rostochiensis* in red; *H. avenae* in dark blue; *R. reniformis* in light blue. Nodes in the network are scaled by connectivity. *H. sacchari* are the only nematode CLEs with connections to rice CLEs.

**Figure 7:** Functional adaptation of *H. sacchari* CLE effectors to monocot hosts. Exogenous application of synthetic *H. sacchari* CLE peptides causes a short root phenotype in rice when compared to a scrambled version of this peptide, a CLE peptide from the related dicot parasitic *H. glycines*, or no peptide control (n=24 per condition, p <0.001, Tukey's HSD).

**Figure S1:** Negative control *in situ* hybridisation reactions for chorismate mutase (left panel) and DN36333 (right panel). No staining is seen in these nematodes.

**Figure S2:** Alignment of CLE-effector-like sequences assembled from the *H. sacchari* transcriptome. Highlighted the position of the predicted signal peptide (green), the unknown repeat of DN37996\_c0\_g2\_i1 (not a CLE encoding peptide, purple), and the canonical CLE domain of the other six sequences (red). Arrow indicates polymorphism in CLE domain.

**Figure S3:** Cloned genomic CLE from *H. sacchari*. A) Alignment of the genomic clone and predicted transcript assembled from the transcriptome for the *H. sacchari* CLE-like gene Hsac\_DN37996F. Primers used to clone the gene are highlighted with grey bars. CLE domain (spanning an intron) is highlighted in purple. B) Mapping the RNAseq reads back to the genomic clone supports the deduced intron:exon structure, and reveals two potential splice variants (a read through intron in exon 4). The shorter splice variant results in a frame shift and consequently no CLE-like domain. The longer splice variant encodes a canonical CLE domain.

### Table legends

**Table 1:** *H. sacchari* sequences similar to putative cell wall degrading and modifying proteins

**Table S1:** Primers used in this study.

Substrate	Cellulose		Xylan	Arabinan	Pectin		Arabinogalactan		
Family	GH5	GH45	GH30	GH43	GH28	PL3	GH53	Expansins	CBM2 domains
<i>H. sacchari</i>	8	0	0	0	0	1	0	2	2
<i>H. avenae</i>	16	0	0	0	0	2	0	2	2
<i>H. schachtii</i>	9	0	0	1	0	9	3	4	2
<i>G. rostochiensis</i>	11	0	0	1	0	3	1	7	7
<i>G pallida</i>	16	0	0	1	0	7	2	9	6
<i>M. incognita</i>	21	0	6	2	1	30	0	20	9
<i>Nacobbus aberrans</i>	2	0	3	1	2	8	3	2	2
<i>Bursaphelenchus xylophilus</i>	0	11	0	0	0	15	0	8	0

**Table 1:** *H. sacchari* sequences similar to putative cell wall degrading and modifying proteins

<b>Primers for in situ hybridisation</b>	
Hsac_cellulase F (Hsac_DN30579)	CCTTGGCACTCCGATGTATT
Hsac_cellulase R (Hsac_DN30579)	ATCAGCTCGTCCAGCATTTT
Hsac_Chor Mutase F (Hsac_DN28871)	GCTGCTTTAACTTCCGCAAC
Hsac_Chor Mutase R (Hsac_DN28871)	CCGTTTCTTGGTTCGGTTTA
Hsac_DN36333 F (SPRYSEC)	TGTCAAAGGAGCGAGGGATT
Hsac_DN36333 R (SPRYSEC)	ATGAGGCTTCCACAGTTCCA
Hsac_DN43534 F (SPRYSEC)	AAATTCCCAGCTGCGATCAC
Hsac_DN43534 R (SPRYSEC)	GTCGGAAGCAAATCGGTCTC
<b>Primers for qPCR</b>	
Hsac_q cellulase F	CTGGACCGACGAAACAGTGA
Hsac_q cellulase R	TCGATCGCTTTGTCCGAAA
Hsac_q Chor Mutase F	AGCATTGACGACTTTGTGCG
Hsac_q Chor Mutase R	ATTCGTCGGTTGCCTCTGT
Hsac_DN36333 qF (SPRYSEC)	GGGGCCATTCTGTCAAAGGA
Hsac_DN36333 qR (SPRYSEC)	TGAGGCTTCCACAGTTCCAC
Hsac_DN43534 qF (SPRYSEC)	AGAAACGTCGACAGTGGCAT
Hsac_DN43534 qR (SPRYSEC)	GACAAGCGCAAACCTCAGCAA
Hsac_DN34918F Reference gene for qPCR (EIF 1A)	ATACGCTTGGGTGCTTGACA
Hsac_DN34918R Reference gene for qPCR (EIF 1A)	CCAGCCTCAAATTCTCCCGT
<b>Primers used for gene cloning</b>	
Hsac_DN36333 CF (SPRYSEC)	ACCATGCGTATTGAAGACAACAAT
Hsac_DN36333 R (SPRYSEC)	TCAGATCGACTCATCGGA
Hsac_DN43534 CF (SPRYSEC)	ACCATGGTCGCGGAAGATGT
Hsac_DN43534 R (SPRYSEC)	TCAATCACCAAGAAGTGGCTTG
Hsac_DN37996F (CLE)	TATCAAATCATGTTTGCCTCCA
Hsac_DN37996R (CLE)	CGTTATTCACITTTAATGGTGCTG

**Table S1:** Primers used in this study.








Figure 2: Analysis of spatio-temporal expression patterns of cellulase (left panels) and chorismate mutase (right panels) identified in the *H. sacchari* transcriptome. In situ hybridisation analysis showed that both genes are expressed in the subventral gland cells of second stage juveniles (J2s) of *H. sacchari* (purple staining; upper panels). Analysis using qPCR shows that the cellulase is expressed specifically at the J2 life stage while the chorismate mutase is expressed at all life stages tested (lower panels). Error bars represent standard errors across the biological replicates.


Best match of *G. rostochiensis* genes in *H. sacchari*

Figure 3: Histograms showing distributions of percentage identity between *H. sacchari* effectors (left panel) and non-effectors (right panel) with their homologues in *G. rostochiensis*. This analysis suggests that effector-like transcripts have accumulated more mutations than other transcripts since the divergence of these two species.




Figure 4: Analysis of spatio-temporal expression patterns of two putative SPRYSEC-encoding transcripts of *H. sacchari*. In situ hybridisation analysis showed that both genes are expressed in the dorsal gland cell of *H. sacchari* (purple staining, arrows; upper panels). Analysis by qPCR shows that both sequences are upregulated in later parasitic stage nematodes (lower panels). Error bars represent standard errors across the biological replicates.

### The HYP gene family




Figure 5: Overview of the HYP gene family in cyst nematodes. Left, A schematic representation of the *H. sacchari* HYP-like sequences shows the position of a novel structural variant outside the hyper-variable domain. The faded region of the hyper-variable domain of HYP-1 and -3 represents the region of variable sequence. Right, a midpoint re-rooted phylogenetic construction of HYP sequences from *G. pallida* and *H. sacchari* position those of *H. sacchari* as a distinct sub-family (names named here sub-family-0).




Figure 6: CLE-like effectors of *H. sacchari*. A) A midpoint re-rooted phylogenetic construction of the seven six complete CLE-like sequences in the *H. sacchari* transcriptome. Six of these sequences encode a canonical CLE domain at the C terminus, while the seventh encodes tandem repeats of an unknown peptide. The CLE domain containing effectors form two groups, each of which has a distinct CLE domain (red box). B) A similarity network of CLE domains from plants and *H. sacchari*. CLEs from selected species are highlighted: Rice CLEs are highlighted in green; *H. sacchari* CLE-like sequences in orange; *H. glycines* in purple; *G. rostochiensis* in red; *H. avenae* in dark blue; *R. reniformis* in light blue. Nodes in the network are scaled by connectivity. *H. sacchari* are the only nematode CLEs with connections to rice CLEs.




Figure 7: Functional adaptation of *H. sacchari* CLE effectors to monocot hosts. Exogenous application of synthetic *H. sacchari* CLE peptides causes a short root phenotype in rice when compared to a scrambled version of this peptide, a CLE peptide from the related dicot parasitic *H. glycines*, or no peptide control (n=24 per condition,  $p < 0.001$ , Tukey's HSD).



Figure S1: Negative control in situ hybridisation reactions for chorismate mutase (left panel) and DN36333 (right panel). No staining is seen in these nematodes.

254x190mm (96 x 96 DPI)




Figure S21: Alignment of CLE-effector-like sequences assembled from the of *H. sacchari* transcriptome. Highlighted the position of the predicted signal peptide (green), the unknown repeat of DN37996\_c0\_g2\_i1 (not a CLE encoding peptide, purple), and the canonical CLE domain of the other six sequences (red). Arrow indicates polymorphism in CLE domain.



**Significance statement:** This is the first study that tackles the issue of host adaptation in plant parasitic nematodes. This is a topic of high scientific interest as "host-jumps" (adaptation of a pathogen to a new host) are of fundamental importance to plant sciences in general, and global food security in particular