



Expansion Phase 1 Study of Pegargiminase Plus Pemetrexed and Cisplatin in Patients With Argininosuccinate Synthetase 1-Deficient Mesothelioma: Safety, Efficacy, and Resistance Mechanisms

Peter W. Szlosarek, MD, PhD,^{a,*} Melissa M. Phillips, MD, PhD,^{a,b} Iuliia Pavlyk, PhD,^a Jeremy Steele, MD,^b Jonathan Shamash, MD,^b James Spicer, MD, PhD,^c Sanjeev Kumar, MD,^d Simon Pacey, MD, PhD,^d Xiaoxing Feng, PhD,^e Amanda Johnston, PhD,^e John Bomalaski, MD,^e Graeme Moir, MD,^f Kelvin Lau, MD,^g Stephen Ellis, MD,^h Michael Sheaff, MDⁱ

^aCenter for Cancer Biomarkers and Biotherapeutics, Barts Cancer Institute (BCI)—A Cancer Research UK Center of Excellence, Queen Mary University of London, London, United Kingdom

^bDepartment of Medical Oncology, Barts Health NHS Trust, St Bartholomew's Hospital, London, United Kingdom

^cSchool of Cancer and Pharmaceutical Sciences, King's College London, Guy's Hospital, London, United Kingdom

^dDepartment of Oncology, University of Cambridge, Cambridge, United Kingdom

^ePolaris Pharmaceuticals, Inc., San Diego, California

^fDepartment of Plastic Surgery, Barts Health NHS Trust, Royal London Hospital, London, United Kingdom

^gDepartment of Cardiothoracic Surgery, Barts Health NHS Trust, St Bartholomew's Hospital, London, United Kingdom

^hDepartment of Diagnostic Imaging, Barts Health NHS Trust, St Bartholomew's Hospital, London, United Kingdom

ⁱDepartment of Histopathology, Pathology and Pharmacy Building, Barts Health NHS Trust, Royal London Hospital, London, United Kingdom

Received 26 August 2020; accepted 27 August 2020

Available online - 03 September 2020

ABSTRACT

Introduction: Pegargiminase (ADI-PEG 20; ADI) degrades arginine and potentiates pemetrexed (Pem) cytotoxicity in argininosuccinate synthetase 1 (ASS1)-deficient malignant pleural mesothelioma (MPM). We conducted a phase 1 dose-expansion study at the recommended phase 2 dose of ADI-PEG 20 with Pem and cisplatin (ADIPemCis), to further evaluate arginine-lowering therapy in ASS1-deficient MPM and explore the mechanisms of resistance.

Methods: A total of 32 patients with ASS1-deficient MPM (11 epithelioid; 10 biphasic; 11 sarcomatoid) who were chemonaïve received weekly intramuscular pegargiminase (36 mg/m²) with Pem (500 mg/m²) and cisplatin (75 mg/m²) intravenously, every 3 weeks (six cycles maximum). Maintenance pegargiminase was permitted until disease progression or withdrawal. Safety, pharmacodynamics, immunogenicity, and efficacy were determined. Biopsies were performed in progressing patients to explore the mechanisms of resistance to pegargiminase.

Results: The treatment was well tolerated. Most adverse events were of grade 1/2, whereas four nonhematologic

grade 3/4 adverse events related to pegargiminase were reversible. Plasma arginine decreased whereas citrulline increased; this was maintained by 18 weeks of ADIPemCis

*Corresponding author.

Disclosure: Dr. Szlosarek has received support from the Higher Education Funding Council for England and research support from the Polaris Group. Drs. Bomalaski, Johnson, and Feng are paid employees of Polaris Pharmaceuticals. The remaining authors declare no conflict of interest.

Address for correspondence: Peter W. Szlosarek, MD, PhD, Center for Cancer Biomarkers and Biotherapeutics, Barts Cancer Institute—A Cancer Research UK Center of Excellence, Queen Mary University of London, Barts and The London Medical School, John Vane Science Center, Charterhouse Square, London EC1M 6BQ, United Kingdom. E-mail: p.w.szlosarek@qmul.ac.uk

Cite this article as: Szlosarek PW, et al. Expansion Phase 1 Study of Pegargiminase Plus Pemetrexed and Cisplatin in Patients With Argininosuccinate Synthetase 1-Deficient Mesothelioma: Safety, Efficacy, and Resistance Mechanisms. *JTO Clin Res Rep* 1:100093

© 2020 The Authors. Published by Elsevier Inc. on behalf of the International Association for the Study of Lung Cancer. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

ISSN: 2666-3643

<https://doi.org/10.1016/j.jtocrr.2020.100093>

therapy. The disease control rate in 31 assessed patients was 93.5% ($n = 29$ of 31; 95% confidence interval [CI]: 78.6%–99.2%), with a partial response rate of 35.5% ($n = 11$ of 31; 95% CI: 19.2%–54.6%). The median progression-free and overall survivals were 5.6 (95% CI: 4.0–6.0) and 10.1 (95% CI: 6.1–11.1) months, respectively. Progression biopsies on pegargiminase revealed a statistically significant influx of macrophages ($n = 6$; $p = 0.0255$) and patchy tumoral ASS1 reexpression ($n = 2$ of 6). In addition, we observed increased tumoral programmed death-ligand 1—an ADI-PEG 20 inducible gene—and the formation of CD3-positive T lymphocyte aggregates on disease progression ($n = 2$ of 5).

Conclusions: The dose expansion of ADIPemCis confirmed the high clinical activity and good tolerability in ASS1-deficient poor-prognosis mesothelioma, underpinning an ongoing phase 3 study ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02709512) NCT02709512). Notably, resistance to pegargiminase correlated with marked macrophage recruitment and—along with the tumor immune microenvironment—warrants further study to optimize arginine deprivation for the treatment of mesothelioma.

© 2020 The Authors. Published by Elsevier Inc. on behalf of the International Association for the Study of Lung Cancer. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Arginine; ASS1; ADIPemCis; Mesothelioma; Macrophages

Introduction

Malignant pleural mesothelioma (MPM) is predominantly an asbestos-driven thoracic tumor notable for its chemoresistance and poor prognosis. Median survivals range from 3.5 to 6.6 months for the nonepithelioid, sarcomatoid, and biphasic variants and up to 18 months for the epithelioid subtype.^{1,2} No new frontline therapies for mesothelioma have been licensed since the antifolate pemetrexed with cisplatin in 2004.³

In preclinical studies, we identified arginine depletion as a rational antimetabolite strategy that targets mesothelioma cells displaying epigenetic inactivation of the urea cycle enzyme argininosuccinate synthetase 1 (ASS1).⁴ Arginine deprivation affects multiple biosynthetic pathways, including proteins, polyamines, nucleotides, and nitric oxide, emphasizing an essential role for the amino acid in the growth or auxotrophy of mesothelioma and other cancers.^{5–7} Consequently, bacteria-derived polyethylene glycol (PEG)ylated arginine deiminase (ADI-PEG 20, ADI, or pegargiminase) or bioengineered forms of human arginase are currently in development for patients with a range of advanced malignancies.^{8–10}

Clinically, pegargiminase, which degrades arginine into citrulline and ammonia, improved progression-free survival (PFS) in patients with ASS1-deficient MPM in the ADAM (Arginine Deiminase And Mesothelioma) study, representing the first biomarker-driven randomized trial of arginine deprivation versus best supportive care in cancer.¹¹ Moreover, ASS1 was prognostic, with the ASS1-deficient disease conferring a worse survival compared with ASS1-proficient disease, consistent with data linking dysregulation of urea cycle enzymes to accelerated tumorigenesis.^{5,11} In addition, when ADI-PEG 20 was combined with pemetrexed and cisplatin chemotherapy (ADIPemCis) in the phase I dose-escalation Tumors Requiring Arginine to Assess ADI-PEG 20, Pemetrexed and cisplatin (TRAP) study, a 100% disease control (78% partial response) rate was observed in nine patients with thoracic cancers (lung adenocarcinoma and MPM), including four of five patients with nonepithelioid MPM.¹² Nevertheless, despite prolonged suppression of plasma arginine and a reciprocal increase in citrulline, patients progressed on ADIPemCis therapy, thereby implicating tumoral, rather than drug-innate, mechanisms of resistance to arginine deprivation.

First, reexpression of ASS1, and thus the recycling of citrulline to arginine, after long-term culture of tumor cell lines, including MPM cells in ADI-PEG 20, has been identified as a viable resistance mechanism with confirmatory studies in patients with melanoma.^{13–15} Second, autophagy (degradation and recycling of cellular components) is known to protect ASS1-negative MPM cells from arginine depletion.¹⁶ Third, the tumor microenvironment may also mediate cancer cell resistance to arginine withdrawal; however, this has not been addressed specifically in the context of pegargiminase. Tumor-associated macrophages (TAMs), in particular, constitute up to 30% of the total cell population of mesothelioma and play a key role in asbestos-mediated tumorigenesis.^{17–20} As such, TAMs might also play a role in resistance to arginine deprivation therapy.

We treated a dose-expansion cohort of 32 patients with ASS1-deficient MPM at the recommended phase 2 dose (RP2D) of ADI-PEG 20 (36 mg/m²) in combination with standard doses of pemetrexed and cisplatin. The main aims of this phase 1 dose-expansion study were to define further the safety and preliminary activity of the ADIPemCis triplet in patients with MPM and to elucidate mechanisms of resistance to arginine deprivation by analyzing patients' tumors at progression.

Materials and Methods

Patient Eligibility

The patients were 18 years or over and chemonaive with histologically proven ASS1-deficient advanced

MPM (see Beddowes et al.¹² for the methods). Additional eligibility included an Eastern Cooperative Oncology Group performance status of 0 or 1, no major comorbidities, a minimum expected survival of 3 months, and a measurable disease by modified Response Evaluation Criteria in Solid Tumors criteria for MPM. Exclusion criteria included recent major operation, history of another active primary cancer, and previous therapy with pegargiminase. All patients signed written informed consent.

Study Design and Treatment

This dose-expansion multicenter phase 1 study evaluated the RP2D of 36 mg/m² weekly intramuscular ADI-PEG 20 plus 3-weekly 75 mg/m² cisplatin and 500 mg/m² pemetrexed derived from the previous dose-escalation TRAP study.¹¹ Standard premedication was administered, including oral dexamethasone, daily folic acid, and 1000 µg intramuscular hydroxycobalamin every 9 weeks. The initial dose of intramuscular ADI-PEG 20 was administered 48 hours before the first dose of cytotoxic drugs. The patients received up to six cycles (18 wk) of ADIPemCis chemotherapy and could continue on maintenance pegargiminase until disease progression or withdrawal. Blood samples were taken at baseline, during ADIPemCis chemotherapy, and on disease progression or withdrawal from the study. Tumor biopsies were required at baseline and were optional at disease progression.

The primary objective of the dose-expansion study was to determine the safety and tolerability and to estimate the preliminary efficacy of ADIPemCis in patients with ASS1-deficient MPM. Secondary objectives included measuring pharmacodynamics, immunogenicity, and exploration of resistance mechanisms to pegargiminase.

Safety

Evaluation of safety was based on the National Cancer Institute Common Terminology Criteria for Adverse Events version 4.03, vital signs, physical examination, electrocardiogram, and laboratory blood analyses.

Pharmacodynamic and Efficacy Evaluations

Blood samples were analyzed by Polaris Pharmaceuticals, Inc. (San Diego, CA) for arginine and citrulline levels and anti-ADI-PEG 20 antibody titers, as described previously.^{11,12} Efficacy was assessed by computed tomography imaging using the modified Response Evaluation Criteria in Solid Tumors criteria every 6 weeks while on ADIPemCis and then every 2 months on maintenance pegargiminase.

Patient Tumor Immunohistochemistry

Tumor biopsy results were assessed for ASS1 expression using the monoclonal antibody 195-21-1 from Polaris Pharmaceuticals, Inc., San Diego, California. Infiltrating CD68^{pos} macrophages were identified using a murine antihuman antibody (KP-1) and quantified as a percentage of the number of malignant cells, taking an average from five high-power fields at ×400 magnification. The expression of programmed death-ligand 1 (PD-L1) (Cell Signaling Technology E1L3N and Ventana-Roche SP-263 antibodies) and CD3 (Ventana-Roche 2GV6 antibody) was performed subsequent to the CD68 staining using residual tissue. PD-L1 was scored as a percentage of positive tumor cells and CD3 summarized descriptively.

Statistical Analyses

No formal sample size calculation was made for the dose-expansion TRAP study in patients with MPM, which aimed to recruit up to 30 patients as per protocol. Adverse events (AEs) were collated, and response rates, PFS, and overall survival (OS) were characterized according to the MPM subtype. The results of the patients' tumor biopsies were analyzed using a paired *t* test in GraphPad Prism version 8.3.1 (GraphPad Software, San Diego, CA). A *p* value of less than 0.05 was considered to be statistically significant. This trial is registered with [ClinicalTrials.gov](https://clinicaltrials.gov), number NCT02029690.

Ethical Considerations

The clinical protocol ([ClinicalTrials.gov](https://clinicaltrials.gov) identifier NCT02029690) was approved by the Leeds East Research Ethics Committee (14/YH/0090) and was sponsored by Polaris Pharmaceuticals, Inc.

Results

Patient Demographics

Patient enrollment in the dose-expansion study began in February 2015 and was completed in May 2016. A total of 93 patients were screened for recruiting 32 patients with ASS1-deficient MPM treated with ADIPemCis, including the following: 11 with epithelioid, 10 with biphasic, and 11 with the sarcomatoid subtype (Fig. 1). The protocol amendment for the dose-expansion cohort specified the enrollment of 30 patients at the RP2D; one patient was deemed ineligible and replaced owing to occult malignant melena, and an additional patient consented as the study recruitment was closing. All subjects were included for the safety analysis and 31 for the efficacy analysis (Table 1).




Figure 1. CONSORT diagram. AE, adverse event; ASS1, argininosuccinate synthetase 1; MPM, malignant pleural mesothelioma; SARC, sarcomatoid.

Safety

Consistent with the previous dose-escalation study, the ADIPemCis treatment was well tolerated ([Supplementary Tables 1 and 2](#)). AEs were reported in 24 of 32 patients (75%), most of which were related to cisplatin and pemetrexed (22 of 32; 68.8%), and to pegargiminase (12 of 32; 37.5%) patients. Most of the patients had grade 1 or 2 (116 of 137; 84.7%), particularly nausea and vomiting and decreased blood counts, with the remainder having grade 3 or 4 only (21 of 137; 15.3%). There were four nonhematologic grade 3 or 4 AEs related to pegargiminase, which include the following: increased alkaline phosphatase, hyperuricemia, skin rash, and posterior reversible encephalopathy

syndrome. The last AE was unexpected and occurred in a patient with a sarcomatoid mesothelioma presenting with agitation and characteristic magnetic resonance imaging features during maintenance pegargiminase. He recovered completely after anxiolytics, steroids, and pegargiminase discontinuation ([Supplementary Fig. 1](#)).

Pharmacodynamics

Pegargiminase decreased plasma arginine with a reciprocal increase in plasma citrulline levels in the patients ([Fig. 2A](#)). As reported in the dose-escalation cohort, plasma levels of the amino acids remained differentially altered compared with pretreatment levels

Table 1. Demographics

Characteristic	Epithelioid (n = 11)	Nonepithelioid (n = 21)	
		Biphasic (n = 10)	Sarcomatoid (n = 11)*
Age (y), median (range)	67 (61-77)	66 (49-82)	68 (58-79)
Sex			
Male	10	8	11
Female	1	2	0
Performance status			
0	1	1	0
1	10	9	11
Previous operation			
Yes	3	2	1
No	8	8	10
Disease stage ^a	1A (n = 1); 1B (n = 4)	1A (n = 1)	1B (n = 6)
	II (n = 1)	1B (n = 6)	II (n = 1)
	IIIA (n = 2)	IIIA (n = 1)	IIIA (n = 1)
	IV (n = 3)	IV (n = 2)	IV (n = 3)
Time on study treatment (mo), median (range)	4.6 (0.5-7.0)	6.1 (1.9-18.0)	4.1 (1.2-5.9)

^aEighth TNM classification for mesothelioma.




Figure 2. Pharmacodynamics and response. (A) Pharmacodynamics of arginine and citrulline in patients treated with ADI-PemCis. Serum arginine and citrulline are revealed by week of treatment (mean \pm SEM). (B) Serum levels of anti-ADI-PEG 20 antibodies in all patients by week of ADIPemCis (mean \pm SEM). (C) Waterfall plot of response by modified RECIST to ADI-PemCis. (D) Spider plots revealing response duration to ADIPemCis. Ab, antibody; ADI, arginine deiminase; ADIPemCis, ADI-PEG 20 with pemetrexed and cisplatin; B, biphasic; E, epithelioid; PEG, PEGylated; RECIST, Response Evaluation Criteria in Solid Tumors; S, sarcomatoid.

by 18 weeks, despite a concomitant increase in anti-ADI-PEG 20 antibodies (Fig. 2B).

Efficacy

The ADIPemCis treatment induced a high disease control rate of 93.5% ($n = 29$ of 31; 95% confidence interval [CI]: 78.6%–99.2%), with a partial response rate of 35.5% ($n = 11$ of 31; 95% CI: 19.2%–54.6%) in a cohort of patients enriched by ASS1 loss for nonepithelioid MPM (Fig. 2C and D). The median PFS and OS were 5.6 (95% CI: 4.0–6.0) and 10.1 (95% CI: 6.1–11.1) months, respectively (Fig. 3A and B). Subsequently, 11 of 31 patients (35.5%) received antiprogrammed cell death protein 1 (PD-1) therapy with pembrolizumab, achieving stable disease in one patient (9.1%), whereas nine patients had progressive disease (81.8%), and one patient was nonassessable (9.1%). PD-L1 expression before treatment was available in nine of 11 patients ranging from 0% ($n = 4$) to 15% to 30% ($n = 5$). Owing to rapidly progressive disease, a few patients received second-line and subsequent therapies (vinorelbine and gemcitabine).

Exploratory Tumor Biopsies

To understand drug resistance, six patients on pegargiminas therapy consented to a tumor rebiopsy at

progression allowing a comparison with the pretreatment biopsies. The ASS1 levels increased in a subpopulation of MPM cells in two of six patients with epithelioid and sarcomatoid disease during cycles 5 and 6 of ADI-PemCis (Fig. 4A). There was a significant increase of CD68^{pos} ASS1^{pos} macrophages at disease progression in ASS1^{neg} tumor areas, which included four patients receiving maintenance pegargiminas for up to 18 months ($p = 0.0255$; $n = 6$; Fig. 4B and C). Owing to patchy ASS1 tumoral reexpression found in two patients only, we were unable to quantify the amount of macrophage infiltration specifically in ASS1^{pos} tumor areas. We also noted an increase in tumoral PD-L1 expression and clustering of CD3^{pos} T lymphocytes within MPM tumor cell islands in two of five patients with available tissue for immunohistochemistry (Fig. 5). In the remaining three patients, we detected variable effects on PD-L1 expression and T cell localization at disease progression (Supplementary Table 3).

Discussion

In this ASS1 biomarker-led study, we observed good tolerability and a high rate of disease control in patients with poor-prognosis MPM treated with the RP2D of ADIPemCis, expanding on the preliminary signal in the




Figure 3. Survival outcomes. (A) PFS by MPM histologic subtype. (B) Kaplan-Meier survival estimates by MPM histologic subtype. CI, confidence interval; MPM, malignant pleural mesothelioma; OS, overall survival; PFS, progression-free survival.

dose-escalation trial. Nonetheless, tumor progression on pegargiminase was universal, and instead of widespread ASS1 tumoral reexpression, correlated with robust macrophage infiltration on rebiopsy, pointing to a stromal-mediated resistance pathway that may be leveraged to optimize arginine-depleting cancer therapeutics. We also describe induction of tumoral PD-L1 expression and modulation of T lymphocytes, which segues into the developing area of mesothelioma immunotherapy.

The toxicities were mostly grade 1 or 2 nausea and vomiting and hematologic and injection skin reactions, whereas grade 3 or 4 events were manageable and

reversible. There was only one serious grade 3 toxicity attributed to pegargiminase maintenance therapy, namely posterior reversible encephalopathy syndrome, a known complication of several biochemotherapies, including bevacizumab and the enzyme therapeutic, asparaginase, but described here for the first time with arginine deprivation.²¹

The median OS and PFS of 10.1 and 5.6 months, respectively, are encouraging in a patient cohort enriched for poor-prognosis ASS1-deficient disease. Biomarker screening selected two to three times as many patients with nonepithelioid compared with epithelioid disease, consistent with previous data sets for ASS1 loss in MPM,




Figure 4. Baseline and progression biopsies analyzed for ASS1 and CD68. (A) Tumoral ASS1 reexpression at progression noted in two patients ($\times 200$ magnification; epithelioid and sarcomatoid). (B) Increase in CD68^{pos} macrophages at disease progression in ASS1-deficient tumoral regions (n = 6; $p = 0.0255$; paired t test); two epithelioid, one sarcomatoid, and three biphasic tumors (N.B. one epithelioid tumor was reclassified as biphasic on operative rebiopsy). (C) Representative serial sections of epithelioid, biphasic, and sarcomatoid MPM at baseline and progression stained for ASS1 and CD68, revealing the increase in ASS1^{pos}CD68^{pos} macrophages at progression ($\times 200$ magnification). ASS1, argininosuccinate synthetase 1; MPM, malignant pleural mesothelioma; N.B., nota bene; pos, positive.

and accounting for the unusually high rate of patients enrolled with the nonepithelioid disease (65.6%).^{4,12,22} The median OS for epithelioid disease was 11.1 months, lower than that reported in recent phase 3 studies with a

median OS of 16.1 months for patients with predominantly epithelioid disease (81%–97%) in the standard chemotherapy groups (LUME-meso [Nintedanib in combination with pemetrexed and cisplatin in mesothelioma]




Figure 5. Baseline and progression biopsies analyzed for PD-L1 and CD3 modulation of PD-L1 expression and CD3^{pos} lymphocytes in two patients at progression ($\times 200$ magnification; epithelioid and biphasic). PD-L1 increased from 10% to 30% (in epithelioid disease) and 0% to 5% (in biphasic disease) with clustering of CD3^{pos} T cells in both patients at progression. PD-L1, programmed death-ligand 1; pos, positive.

and MAPS [Mesothelioma Avastin Cisplatin Pemetrexed Study]].^{23,24} Moreover, twice as many patients were alive at 15 months with biphasic compared with the epithelioid disease (40% versus 20%), indicating that the latter subgroup is at the aggressive end of the spectrum and concurring with the poor-prognosis epithelioid disease defined by nuclear grading and p16 loss on multivariate analysis.^{25,26}

Notably, the 8.2 month median OS for nonepithelioid disease compares favorably with the recent SWOG S0905 trial reporting a median OS of 6.3 months for PemCis plus placebo or 6.5 months for PemCis plus the vascular endothelial growth factor receptor antagonist, cediranib ($n = 23$; nonepithelioid).²⁷ In addition, we observed a doubling of the median survival (6.5 versus 3.5 mo) and a three-fold increase in survival at 12 months (30% versus 10%), compared with historical controls for sarcomatoid mesothelioma.^{1,26} Although response assessment in mesothelioma is challenging, and reported infrequently in trials for the nonepithelioid disease, the 93.5% disease control rate is encouraging and consistent with the earlier dose-escalation study findings.¹² Collectively, these data benchmarked the design of the ATOMIC-meso (ADI-PEG 20 Targeting Of Malignancy Induces Cytotoxicity-Mesothelioma) study, which transitioned from phase 2 to phase 3 earlier this year after successful recruitment of 176 patients with nonepithelioid mesothelioma; a further 210 patients are being enrolled to report on the primary end point of OS ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02709512) identifier NCT02709512).

A key exploratory aim of the dose-expansion study was to understand resistance to the ADI-PEG 20-based therapy by sampling patients' tumors on progression. Six patients consented to repeat biopsies, which were incorporated into patient management, most often for control of a recurrent pleural effusion. Owing to limited

baseline tissue, we analyzed the ASS1 status followed by CD68 expression on macrophages, and lastly, PD-L1 expression and CD3 expression on lymphocytes. Patchy induction of tumoral ASS1 was identified in two patients, supporting a limited role for ASS1 reexpression as a mechanism of resistance to pegargiminas as identified in the long-term MPM cell line culture under ADI-PEG 20.¹³ This contrasted with a robust and statistically significant influx of CD68^{pos} ASS1^{pos} macrophages in ASS1-deficient tumoral regions, which is of particular interest because myeloid cells are known to express abundant ASS1 under proinflammatory cytokine control.²⁸ Moreover, arginine metabolism is a critical component of macrophage function, including nitric oxide synthesis for pathogen recognition and polyamine synthesis for wound healing.²⁹ Interestingly, we also observed an influx of CD68^{pos} ASS1^{pos} macrophages in ASS1-deficient tumoral regions in the rebiopsy of two patients in a separate expansion study of ADIPemCis in patients with NSCLC, in which p equals 0.0079 for the entire thoracic patient cohort of ADIPemCis ([Supplementary Fig. 2](#)).³⁰ Separately, we have identified a novel mechanism whereby ADI-PEG 20 induces several chemoattractant proinflammatory cytokines from MPM cells, triggering resistance to arginine deprivation by macrophage-derived argininosuccinate, the immediate precursor for arginine synthesis (manuscript under review).

Our analysis of resistance was limited by the availability of patient tissue, especially the polarization of the infiltrating CD68^{pos} ASS1^{pos} macrophages (i.e., M1 and M2 macrophage subtypes) and the potential role of autophagy which will require further study.^{31,32} Indeed, autophagy was inferred in a separate expansion cohort study of ADIPemCis in glioblastoma multiforme with a patient exhibiting prolonged remission on maintenance ADI-PEG 20 with quinine sulfate, an antimalarial

autophagy inhibitor, and on a background of durable arginine depletion (20.8 mo).³³ Indeed, autophagy as a contributory resistance mechanism has been described preclinically in various cancer cell lines, including MPM cells and is similarly abrogated with chloroquine.^{16,34,35} Finally, pharmacologic resistance owing to neutralizing antibodies to ADI-PEG 20 cannot be excluded entirely, as arginine levels, although persistently low compared with pretreatment levels, increased at the end of the 18-week sampling period (with reciprocal changes in citrulline). Nonetheless, the pharmacodynamic changes were durable in the dose-escalation ADIPemCis study, which reported a higher median OS of 13.9 months in patients with thoracic cancers; these interstudy differences may be explained in part by a variation in the amount of blood sampling performed at each time point owing to earlier subject withdrawal in this study (Supplementary Table 4).¹² It is also relevant that blood draws were performed weekly and just before ADI-PEG 20 dosing, reflecting static rather than dynamic changes in amino acid levels.

The limited analysis of tumoral PD-L1 expression and CD3 lymphocytes at progression in the remaining five biopsies was insufficient to draw firm conclusions. However, the up-regulation of PD-L1 and CD3 lymphocytes in two of the five patients on rebiopsy is consistent with earlier preclinical work on ADI-PEG 20 inducing PD-L1 in tumor cell lines and T cell infiltration in syngeneic tumor mouse models.³⁶ Recently, a phase 1 study of pegargiminase and pembrolizumab in solid tumors completed accrual with on-treatment biopsies that evaluate the effects of pegargiminase specifically on T cell markers in the tumor microenvironment before PD-1 blockade (ClinicalTrials.gov identifier NCT03254732).³⁷ Although a third of the patients received pembrolizumab on progression (6 of 11 epithelioid and 5 of 11 nonepithelioid), the disease control rate of 11.1% ($n = 1$ of 11; biphasic disease) was disappointing and lower than that reported in larger patient studies of PD-1/PD-L1 blockade in mesothelioma (47%–72%).^{38–41} However, four of the patients treated with pembrolizumab expressed PD-L1 less than 1% ($n = 4$ of 9 or 44.4), which is known to correlate with lower responses to PD-1 blockade compared with greater than 1% PD-L1 MPM expression (Supplementary Table 5).⁴¹ Furthermore, the influx of TAMs reported previously may have contributed potentially to a more immunosuppressive environment constraining PD1-based immune checkpoint therapy.⁴²

Clearly, further dissection of the complex effects of arginine deprivation on the mesothelioma microenvironment will be needed to understand the role of ADI-PEG 20 in the context of mesothelioma immunotherapy.^{43–45} Moreover, studies in urea cycle-

dysregulated cancers suggest that biomarker analysis will be of increasing importance in guiding prognosis and therapeutic response to arginine-based therapeutics.⁴⁶ We propose that the macrophage influx may be exploited pharmacologically to optimize arginine deprivation as a novel antimetabolite therapy for mesothelioma and other treatment-resistant cancers. Indeed, several approaches are under clinical evaluation, including CSF-1R, CXCR2, CD47 (don't eat me), and PD-1 antagonists, to retarget TAMs for tumor cell killing.^{47–52}

In summary, ADIPemCis is safe and active in an expansion cohort of patients with aggressive ASS1-deficient MPM, and a phase 3 trial is underway. Our data also provide novel insights into resistance pathways to arginine deprivation, going beyond tumoral ASS1 reexpression, namely macrophage trafficking. Validation of this innate-immunometabolic relationship by targeting macrophages alongside tumor cells with pegargiminase therapy has the potential to improve patient outcomes with mesothelioma and other arginine-auxotrophic cancers.

Acknowledgments

This work was funded and sponsored by Polaris Pharmaceuticals, Inc., and supported by Barts (Queen Mary University of London), King's College (University of London), and Cambridge Experimental Cancer Medicine Centers and the National Institute for Health Research Cambridge Clinical Research Facility. The Cambridge Human Research Tissue Bank is supported by the National Institute for Health Research Cambridge Biomedical Research Centre. The authors are indebted to all the patients and their families who took part in the dose-expansion Tumors Requiring Arginine to Assess ADI-PEG 20, Pemetrexed and cisplatin (Thoracic cohort) study. The authors also thank Suzanne Jordan for her clinical pathology services in Barts.

Supplementary Data

Note: To access the supplementary material accompanying this article, visit the online version of the *JTO Clinical and Research Reports* at www.jtocrr.org and at <https://doi.org/10.1016/j.jtocrr.2020.100093>.

References

1. Klebe S, Brownlee NA, Mahar A, et al. Sarcomatoid mesothelioma: a clinical-pathologic correlation of 326 cases. *Mod Pathol*. 2010;23:470-479.
2. Vigneswaran WT, Kircheva DY, Ananthanarayanan V, et al. Amount of epithelioid differentiation is a predictor of survival in malignant pleural mesothelioma. *Ann Thorac Surg*. 2017;103:962-966.

3. Vogelzang NJ, Rusthoven JJ, Symanowski J, et al. Phase III study of pemetrexed in combination with cisplatin versus cisplatin alone in patients with malignant pleural mesothelioma. *J Clin Oncol*. 2003;21:2636-2644.
4. Szlosarek PW, Klabatsa A, Pallaska A, et al. In vivo loss of expression of argininosuccinate synthetase in malignant pleural mesothelioma is a biomarker for susceptibility to arginine depletion. *Clin Cancer Res*. 2006;12:7126-7131.
5. Allen MD, Luong P, Hudson C, et al. Prognostic and therapeutic impact of argininosuccinate synthetase 1 control in bladder cancer as monitored longitudinally by PET imaging. *Cancer Res*. 2014;74:896-907.
6. Rabinovich S, Adler L, Yizhak K, et al. Diversion of aspartate in ASS1-deficient tumors fosters de novo pyrimidine synthesis. *Nature*. 2015;527:379-383.
7. Keshet R, Szlosarek P, Carracedo A, Erez A. Rewiring urea cycle metabolism in cancer to support anabolism. *Nat Rev Cancer*. 2018;18:634-645.
8. Ensor CM, Holtsberg FW, Bomalaski JS, Clark MA. PEGylated arginine deiminase (ADI-SS PEG20,000 mw) inhibits human melanomas and hepatocellular carcinomas in vitro and in vivo. *Cancer Res*. 2002;62:5443-5450.
9. Cheng PN, Lam TL, Lam WM, et al. PEGylated recombinant human arginase (rhArg-peg5,000mw) inhibits the in vitro and in vivo proliferation of human hepatocellular carcinoma through arginine depletion. *Cancer Res*. 2007;67:309-317.
10. Phillips MM, Sheaff MT, Szlosarek PW. Targeting arginine-dependent cancers with arginine-degrading enzymes: opportunities and challenges. *Cancer Res Treat*. 2013;45:251-262.
11. Szlosarek PW, Steele JP, Nolan L, et al. Arginine deprivation with PEGylated arginine deiminase in patients with argininosuccinate synthetase 1-deficient malignant pleural mesothelioma: a randomized clinical trial. *JAMA Oncol*. 2017;3:58-66.
12. Beddowes E, Spicer J, Chan PY, et al. Phase 1 dose-escalation study of PEGylated arginine deiminase, cisplatin, and pemetrexed in patients with argininosuccinate synthetase 1-deficient thoracic cancers. *J Clin Oncol*. 2017;35:1778-1785.
13. Locke M, Ghazaly E, Freitas MO, et al. Inhibition of the polyamine synthesis pathway is synthetically lethal with loss of argininosuccinate synthase 1. *Cell Rep*. 2016;16:1604-1613.
14. Tsai WB, Aiba I, Lee SY, Feun L, Savaraj N, Kuo MT. Resistance to arginine deiminase treatment in melanoma cells is associated with induced argininosuccinate synthetase expression involving c-Myc/HIF-1 α /Sp4. *Mol Cancer Ther*. 2009;8:3223-3233.
15. Feun LG, Marini A, Walker G, et al. Negative argininosuccinate synthetase expression in melanoma tumors may predict clinical benefit from arginine-depleting therapy with PEGylated arginine deiminase. *Br J Cancer*. 2012;106:1481-1485.
16. Battisti S, Valente D, Albonici L, Bei R, Modesti A, Palumbo C. Nutritional stress and arginine auxotrophy confer high sensitivity to chloroquine toxicity in mesothelioma cells. *Am J Respir Cell Mol Biol*. 2012;46:498-506.
17. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144:646-674.
18. Balkwill FR, Mantovani A. Cancer-related inflammation: common themes and therapeutic opportunities. *Semin Cancer Biol*. 2012;22:33-40.
19. Burt BM, Rodig SJ, Tillemann TR, Elbardissi AW, Bueno R, Sugarbaker DJ. Circulating and tumor-infiltrating myeloid cells predict survival in human pleural mesothelioma. *Cancer*. 2011;117:5234-5244.
20. Miselis NR, Wu ZJ, Van Rooijen N, Kane AB. Targeting tumor-associated macrophages in an orthotopic murine model of diffuse malignant mesothelioma. *Mol Cancer Ther*. 2008;7:788-799.
21. Peddi PF, Peddi S, Santos ES, Morgensztern D. Central nervous system toxicities of chemotherapeutic agents. *Expert Rev Anticancer Ther*. 2014;14:857-863.
22. Bueno R, Stawiski EW, Goldstein LD, et al. Comprehensive genomic analysis of malignant pleural mesothelioma identifies recurrent mutations, gene fusions and splicing alterations. *Nat Genet*. 2016;48:407-416.
23. Scagliotti GV, Gaafar R, Nowak AK, et al. Nintedanib in combination with pemetrexed and cisplatin for chemotherapy-naïve patients with advanced malignant pleural mesothelioma (LUME-Meso): a double-blind, randomised, placebo-controlled phase 3 trial. *Lancet Respir Med*. 2019;7:569-580.
24. Zalcman G, Mazieres J, Margery J, et al. Bevacizumab for newly diagnosed pleural mesothelioma in the Mesothelioma Avastin Cisplatin Pemetrexed Study (MAPS): a randomised, controlled, open-label, phase 3 trial [published correction appears in *Lancet*. 2016;387:e24]. *Lancet*. 2016;387:1405-1414.
25. Forest F, Patoir A, Dal Col P, et al. Nuclear grading, BAP1, mesothelin and PD-L1 expression in malignant pleural mesothelioma: prognostic implications. *Pathology*. 2018;50:635-641.
26. Chou A, Toon CW, Clarkson A, Sheen A, Sioson L, Gill AJ. The epithelioid BAP1-negative and p16-positive phenotype predicts prolonged survival in pleural mesothelioma. *Histopathology*. 2018;72:509-515.
27. Tsao AS, Miao J, Wistuba II, et al. Phase II trial of cediranib in combination with cisplatin and pemetrexed in chemotherapy-naïve patients with unresectable malignant pleural mesothelioma (SWOG S0905). *J Clin Oncol*. 2019;37:2537-2547.
28. Nussler AK, Billiar TR, Liu ZZ, Morris SM Jr. Coinduction of nitric oxide synthase and argininosuccinate synthetase in a murine macrophage cell line. Implications for regulation of nitric oxide production. *J Biol Chem*. 1994;269:1257-1261.
29. Phillips M, Szyszko T, Hall P, et al. Abstract B33: expansion study of PEGylated arginine deiminase (ADI-PEG 20), pemetrexed, and cisplatin in patients with ASS1-deficient non-squamous non-small cell lung cancer (TRAP). *Clin Cancer Res*. 2018;24(suppl).
30. Phillips M, Szlosarek PW. Arginine metabolism and tumour associated macrophages. In: Lawrence T, Hagemann T, eds. *Tumour-Associated Macrophages*. New York, NY: Springer; 2012:77-90.
31. Jayasingam SD, Citartan M, Thang TH, Mat Zin AA, Ang KC, Ch'ng ES. Evaluating the polarization of tumor-

- associated macrophages into M1 and M2 phenotypes in human cancer tissue: technicalities and challenges in routine clinical practice. *Front Oncol.* 2020;9:1512.
32. Klionsky DJ, Abdelmohsen K, Abe A, et al. Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition) [published correction appears in *Autophagy*. 2016;12:443. Selliez, Iban [corrected to Seilliez, Iban]]. *Autophagy*. 2016;12:1-222.
 33. Hall PE, Lewis R, Syed N, et al. A phase I study of PEGylated arginine deiminase (pegarginase), cisplatin, and pemetrexed in argininosuccinate synthetase 1-deficient recurrent high-grade glioma. *Clin Cancer Res.* 2019;25:2708-2716.
 34. Kim RH, Coates JM, Bowles TL, et al. Arginine deiminase as a novel therapy for prostate cancer induces autophagy and caspase-independent apoptosis. *Cancer Res.* 2009;69:700-708.
 35. Delage B, Luong P, Maharaj L, et al. Promoter methylation of argininosuccinate synthetase-1 sensitises lymphomas to arginine deiminase treatment, autophagy and caspase-dependent apoptosis. *Cell Death Dis.* 2012;3:e342.
 36. Brin E, Wu K, Lu HT, He Y, Dai Z, He W. PEGylated arginine deiminase can modulate tumor immune microenvironment by affecting immune checkpoint expression, decreasing regulatory T cell accumulation and inducing tumor T cell infiltration. *Oncotarget.* 2017;8:58948-58963.
 37. Chang KY, Chiang NJ, Yen CJ, et al. A phase Ib study of ADI-PEG 20 plus pembrolizumab in advanced solid cancers. *J Clin Oncol.* 2018;36(supple15), 2556-2556.
 38. Alley EW, Lopez J, Santoro A, et al. Clinical safety and activity of pembrolizumab in patients with malignant pleural mesothelioma (KEYNOTE-028): preliminary results from a non-randomised, open-label, phase 1b trial. *Lancet Oncol.* 2017;18:623-630.
 39. Quispel-Janssen J, van der Noort V, de Vries JF, et al. Programmed death 1 blockade with nivolumab in patients with recurrent malignant pleural mesothelioma. *J Thorac Oncol.* 2018;13:1569-1576.
 40. Hassan R, Thomas A, Nemunaitis JJ, et al. Efficacy and safety of avelumab treatment in patients with advanced unresectable mesothelioma: phase 1b results from the JAVELIN solid tumor trial. *JAMA Oncol.* 2019;5:351-357.
 41. Okada M, Kijima T, Aoe K, et al. Clinical efficacy and safety of nivolumab: results of a multicenter, open-label, single-arm, Japanese phase II study in malignant pleural mesothelioma (MERIT). *Clin Cancer Res.* 2019;25:5485-5492.
 42. Awad MM, Jones RE, Liu H, et al. Cytotoxic T cells in PD-L1-positive malignant pleural mesotheliomas are counterbalanced by distinct immunosuppressive factors. *Cancer Immunol Res.* 2016;4:1038-1048.
 43. Rodriguez PC, Ochoa AC, Al-Khami AA. Arginine metabolism in myeloid cells shapes innate and adaptive immunity. *Front Immunol.* 2017;8:93.
 44. Murray PJ. Amino acid auxotrophy as a system of immunological control nodes. *Nat Immunol.* 2016;17:132-139.
 45. Lemos H, Huang L, Prendergast GC, Mellor AL. Immune control by amino acid catabolism during tumorigenesis and therapy. *Nat Rev Cancer.* 2019;19:162-175.
 46. Lee JS, Adler L, Karathia H, et al. Urea cycle dysregulation generates clinically relevant genomic and biochemical signatures. *Cell.* 2018;174:1559-1570.
 47. Ries CH, Cannarile MA, Hoves S, et al. Targeting tumor-associated macrophages with anti-CSF-1R antibody reveals a strategy for cancer therapy. *Cancer Cell.* 2014;25:846-859.
 48. Sharma B, Nawandar DM, Nannuru KC, Varney ML, Singh RK. Targeting CXCR2 enhances chemotherapeutic response, inhibits mammary tumor growth, angiogenesis, and lung metastasis. *Mol Cancer Ther.* 2013;12:799-808.
 49. Matlung HL, Szilagyi K, Barclay NA, van den Berg TK. The CD47-SIRP α signaling axis as an innate immune checkpoint in cancer. *Immunol Rev.* 2017;276:145-164.
 50. Schürch CM, Forster S, Brühl F, Yang SH, Felley-Bosco E, Hewer E. The “don’t eat me” signal CD47 is a novel diagnostic biomarker and potential therapeutic target for diffuse malignant mesothelioma. *Oncoimmunology.* 2017;7:e1373235.
 51. Gordon SR, Maute RL, Dulken BW, et al. PD-1 expression by tumor-associated macrophages inhibits phagocytosis and tumor immunity. *Nature.* 2017;545:495-499.
 52. Vitale I, Manic G, Coussens LM, Kroemer G, Galluzzi L. Macrophages and metabolism in the tumor microenvironment. *Cell Metab.* 2019;30:36-50.