Risk factors for the development of neonatal sepsis in a neonatal intensive care unit of a tertiary care hospital of Nepal

Sulochana Manandhar ^{1,2}, Puja Amatya ³, Imran Ansari ³, Niva Joshi ¹, Nhukesh Maharjan ¹, Sabina Dongol ¹, Buddha Basnyat ¹, Sameer M. Dixit ⁴, Stephen Baker ⁵ and Abhilasha Karkey ^{1*}

¹Oxford University Clinical Research Unit, Patan Academy of Health Sciences, Kathmandu, Nepal

²Centre for Tropical Medicine and Global Health, Medical sciences division, Nuffield

Department of Medicine, University of Oxford, Linacre College, Oxford, UK

³ Department of Pediatrics, Patan Academy of Health Sciences, Patan Hospital, Kathmandu, Nepal

⁴Center for Molecular Dynamics Nepal, Kathmandu, Nepal

⁵ Cambridge Institute of Therapeutic Immunology & Infectious Disease (CITIID) Department of Medicine, University of Cambridge, Cambridge, UK

*Correspondence

Dr Abhilasha Karkey akarkey@oucru.org

File name: Additional file 3

File format: .pdf

Title of data: Overall workflow for sepsis diagnosis

Description of data: The figure depicts details on the workflow used for sepsis diagnosis and the diagnostic outcomes of the neonates that were enrolled in this study

