Validation of lymphocyte density determined by computational pathology as a predictor of pathological complete response in the ARTemis trial

H. Raza Ali, MB BCh, PhD^{1,2*}; Aliakbar Dariush, PhD^{3*}; Jeremy Thomas, MBBS⁴; Elena
Provenzano, MD, PhD⁵⁻⁷; Janet Dunn, PhD⁸; Louise Hiller, PhD⁸; Anne-Laure Vallier, MSc^{5,7}; Jean Abraham, MBBS, PhD^{5,7}; Tammy Piper MSc⁴; John M. S. Bartlett, PhD^{4,9}; David A. Cameron, MD⁴; Larry Hayward, MB ChB, PhD⁴; James D. Brenton, MD, PhD^{1,5,7}; Paul D. P. Pharoah, MB BCh, PhD^{5,7}; Mike Irwin, PhD³; Nicholas A. Walton, PhD³; Helena M. Earl, MBBS, PhD^{5,7†}; Carlos Caldas, MD^{1,5,7†}

10

15

Affiliations

1. Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre, Cambridge, UK.

2. Department of Pathology, University of Cambridge, Cambridge, UK

- 3. Institute of Astronomy, University of Cambridge, Cambridge, UK
- 4. Edinburgh Cancer Research Centre, Western General Hospital, Edinburgh, UK

5. Department of Oncology, University of Cambridge, Addenbrooke's Hospital, Cambridge, UK.

6. Department of Histopathology, Addenbrooke's Hospital, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK.

²⁰ 7. Cambridge Experimental Cancer Medicine Centre and NIHR Cambridge Biomedical Research Centre, Cambridge, UK.

8. Warwick Clinical Trials Unit, University of Warwick, Coventry, UK

9. Ontario Institute for Cancer Research, Toronto, ON, Canada

25

^{*}These authors contributed equally.

[†]Co-senior, corresponding authors:

Prof. Helena M. Earl: hme22@cam.ac.uk

Professor of Clinical Cancer Medicine
 University of Cambridge
 Cambridge Biomedical Campus
 Cambridge, U.K.

Prof. Carlos Caldas: carlos.caldas@cruk.cam.ac.uk Professor of Cancer Medicine Senior Group Leader, CRUK Cambridge Institute Director, Breast Cancer Programme, Cambridge Cancer Centre University of Cambridge, Cambridge, UK

40

Word count: 1,338

Key points

50

55

45 **Question**: Is lymphocyte density, as quantified by computational pathology, associated with response to neoadjuvant chemotherapy in breast cancer?

Findings: This analysis of 609 patients enrolled in the randomised phase III ARTemis trial showed that pre-treatment lymphocyte density was significantly associated with pathological complete response (pCR), with 15% of patients experiencing pCR in the lower quartile of density compared to 35% in the top quartile. An increase in pre-to-post treatment lymphocyte density was inversely associated with pCR (53% experiencing pCR in the lower quartile compared to 18% in the first).

Meaning: These findings independently validate those of our previous computational pathology study, confirming lymphocyte density as a clinically valid predictor of chemotherapy response for breast cancer patients.

Abstract

Importance

60

65

We have previously shown lymphocyte density, measured using computational pathology, is associated with pathological complete response (pCR) in breast cancer. The clinical validity of this finding in independent studies, among patients receiving different chemotherapy, is unknown.

Objective

To determine whether lymphocyte density and change in lymphocyte density are associated with pCR among breast cancer patients treated with neoadjuvant chemotherapy.

Design, setting and participants

The ARTemis trial randomly assigned 800 women with early stage breast cancer between May 2009 and January 2013 to 3 cycles of docetaxel, followed by 3 cycles of fluorouracil, epirubicin and cyclophosphamide once every 21 days with or without 4 cycles of bevacizumab. The primary

⁷⁰ endpoint was pCR (absence of invasive cancer in the breast and lymph nodes). We quantified lymphocyte density within haematoxylin and eosin (H&E) whole slide images using our previously described computational pathology approach: for every detected lymphocyte the average distance to the nearest fifty lymphocytes was calculated and the density derived from this statistic. We analysed both pre-treatment biopsies and post-treatment surgical samples of the tumour bed.

75 Main outcomes and measures

Lymphocyte density was tested for associations with pCR.

Results

Of the 781 patients originally included in the primary endpoint analysis of the trial, 609 (78%) were included for baseline lymphocyte density analyses and a subset of 383 (49% of 781) for analyses of

change in lymphocyte density. The main reason for loss of patients was the availability of digitised whole slide images. Pre-treatment lymphocyte density modelled as a continuous variable was associated with pCR on univariate analysis (odds ratio [OR], 2.92; 95% CI, 1.78-4.85; p<0.001) and after adjustment for clinical covariates (OR, 2.13; 95% CI, 1.24-3.67; p=0.006). Increased pre-to-post treatment lymphocyte density showed an independent inverse association with pCR (adjusted OR,

⁸⁵ 0.1; 95% CI, 0.033-0.31; p<0.001).

Conclusions and relevance

Lymphocyte density in pre-treatment biopsies was validated as an independent predictor of pCR in breast cancer. Computational pathology is emerging as a viable and objective means of identifying predictive biomarkers for cancer patients.

90 ClinicalTrials.gov: NCT01093235

Funding: Cancer Research UK (CRUK/08/037), Roche, Sanofi-Aventis

Introduction

Tumour-infiltrating lymphocytes (TILs) have been widely investigated as a prognostic and predictive biomarker in breast cancer¹. However, routine assessment of TILs in the clinical setting is hindered by poor reproducibility of their manual pathological evaluation. We previously conducted a systematic analysis of quantitative pathology metrics in the Neo-tAnGo trial² and found that pretreatment tumour lymphocyte density was independently associated with pathological complete response (pCR)³. Our computational pathology approach is automated, objective and quantitative and may, therefore, facilitate clinical implementation. In addition, we found that a relative increase in lymphocyte density after treatment was inversely associated with pCR and that this relationship significantly differed by taxane sequencing³, suggesting that in a subset of patients chemotherapy modulates the post-treatment immune microenvironment.

105

95

100

The ARTemis trial showed that the addition of bevacizumab to standard neoadjuvant chemotherapy significantly increased the proportion of patients with a pCR⁴, but this has not impacted on disease free and overall survival (Earl HM et al.⁵). Here, we tested whether our original findings could be validated in this independent study, and have also conducted exploratory analyses of associations with disease-free and overall survival.

Methods

115

Study design

ARTemis was a multicentre phase III randomised controlled trial conducted to test whether the addition of bevacizumab to 3 cycles of docetaxel, followed by 3 cycles of fluorouracil, epirubicin and cyclophosphamide increased the proportion of patients with a pCR⁴. Women with human epidermal growth factor receptor 2 (HER2)-negative early breast cancer were recruited from May 2009 until January 2013. Of the 800 patients randomised, 781 were available for the primary endpoint analysis. The primary endpoint was pCR (absence of invasive cancer in both the breast and lymph nodes). Here, whether a pCR had occurred was either determined based on central pathology review or, where central review was not possible, on histopathology reports⁶. Details of eligibility and follow-up procedures are provided in the main trial report⁴. The trial was approved by the multicentre and local research ethics committees. All patients provided written, informed consent. The trial was registered at ClinicalTrials.gov (NCT01093235). Supplementary Table 1 details characteristics of patients included in this analysis.

125 **Computational pathology**

Digital whole slide images of haematoxylin and eosin (H&E) stained tissue sections both before and after treatment, were captured using a Hamamatsu Nanozoomer (Hamamatsu City, Shizuoka Pref., Japan). Blinded to all pathological and clinical parameters, we used our computational pathology analysis pipeline to compute cellular metrics from these images. Supplementary Figure 1 summarises the computational pathology workflow. Briefly, the algorithm segments cell nuclei and, based on a training set of approximately 1000 objects per category, uses machine learning (support-vector-machine) to classify cells into three categories: cancer, stromal and lymphocyte. Finally, based on these classes descriptive cellular metrics are computed, including cellular density. Here lymphocyte density is calculated as follows: for every detected lymphocyte in a section, the average distance R to the fifty nearest lymphocytes (N=50) is calculated using a K-nearest-neighbour approach. For each lymphocyte, density is estimated as N/(π R²) and the median of this value for all detected lymphocytes is taken as the summary statistic for a given section. The computational pathology approach has been described in detail previously³ and the analysis code is available at http://www.ast.cam.ac.uk/~adariush/files/codes/.

140

130

135

Statistical analyses

We tested for associations between lymphocyte density and pCR using logistic regression, reporting odds ratios (OR) and 95% confidence intervals (95% CI). Lymphocyte density and change in lymphocyte density were modelled as continuous variables. Multivariable models were adjusted for age, randomisation arm, histological grade, estrogen receptor (ER) status, tumour size and lymph

node status at randomisation. Age and histological grade were modelled as continuous variables. Tumour size (<51mm versus >50mm) and lymph node status (negative versus positive) were modelled as categorical variables. Associations with categorical clinical variables were tested using Kruskal-Wallis tests. Associations with overall survival (OS) defined as all-cause mortality, and disease-free survival (DFS) were tested using Cox proportional-hazards models, where follow-up commenced from day of surgery. DFS was calculated to date of first relapse (loco-regional or distant, not including DCIS); to date of death in women dying without invasive relapse; or to date of censoring in women alive and disease free. Survival analyses were conducted separately by ER-status to account for known violations of the proportional-hazards assumption⁷. Statistical analyses were conducted using Stata SE version 14.2 (Stata Corp, College Station, TX, USA).

150

Results

160

165

170

Of the 781 patients included in the ARTemis primary analysis, 609 (78%) had computational pathology and baseline outcome data (Figure 1), where 109 (18%) experienced pCR, a similar proportion to the entire group of 781 patients where 20% experienced pCR. Of these 609, 383 patients had matched pre- and post-treatment samples to calculate change in lymphocyte density; of which 17 (4%) achieved pCR (Supplementary Table 1). Median time at risk for OS was 3.1 years (range 0.07-6.3 years). Among the 609 patients, there were 140 DFS events and 98 OS events.

Pre-treatment lymphocyte density was associated with ER status (p<0.001), tumour size (p=0.003), and histological grade (p<0.001) (Supplementary Figure 2).

Higher pre-treatment lymphocyte density was associated with a greater chance of pCR in unadjusted (OR, 2.93; 95% CI, 1.77-4.85; p<0.001) and adjusted (OR, 2.13; 95% CI, 1.24-3.67; p=0.006) analyses (Table 1 and Figure 2). However, there was no association between pre-treatment lymphocyte density and survival (OS or DFS) in either ER-positive or ER-negative disease (Supplementary Table 2). Consistent with our previous observations³, an increase in lymphocyte density between pre- and post-treatment was associated with residual disease (adjusted OR for pCR, 0.1; 95% CI, 0.033-0.31; p<0.001; Figure 2 and Supplementary Table 3). Change in lymphocyte

density was not associated with OS or DFS in either ER-positive or ER-negative disease

175

(Supplementary Table 2).

Discussion

180

185

In this computational pathology analysis of the ARTemis trial, we have validated our previous observation that higher pre-treatment lymphocyte density is associated with pCR and that an increase in lymphocyte density after treatment is seen in a subset of surgical resection samples with residual disease.

Pre-treatment lymphocyte density, while predicting pCR independent of clinical variables, was not associated with survival. Although this contrasts with the findings of past studies⁸⁻¹⁰, it should be noted that in these published reports lymphocyte density was not quantified using the approach described here. Our finding should also be interpreted cautiously since analyses were modestly powered due to small sample sizes and limited follow-up time.

Our analyses were limited to tissue morphology in H&E slides. While this is a pragmatic and therefore clinically feasible approach, it overlooks functional differences in infiltrating lymphocytes, which have been shown to influence clinical outcome¹¹⁻¹³. A second limitation was the incomplete representation of post-treatment specimens. A possible explanation for this, and for the lower proportion of patients with pCR in this subset, is that slides from surgical samples in which a pCR is observed are less likely to be digitised since they do not contain cancer cells. Similarly, we were not able to include all patients recruited to the trial because some slides were not available for digitization. Importantly, the findings validate those of our previous independent study and therefore are more likely to be generalizable.

Our findings validate pre-treatment lymphocyte density - a computational pathology metric - as a predictor of pCR. This highlights the potential of automated quantitative pathology methods to improve the standard histopathological evaluation of tumour samples. Such approaches have the additional advantage of being objective and reproducible. Moreover, our finding that an increase in pre-to-post treatment lymphocyte density is associated with residual disease again highlights perturbations in the immune microenvironment secondary to, and presumably caused by, treatment. We speculate that such a comparative metric could serve as a biomarker to identify patients likely to respond to post-neoadjuvant immunotherapy.

205

200

Higher pre-treatment lymphocyte density is validated as a predictor of pCR among women with early stage breast cancer. In addition, an increase in lymphocyte density following chemotherapy is again observed to be associated with residual disease. Patients with low pre-treatment lymphocyte density may benefit from more aggressive therapies or enrolment into clinical trials. In addition,

immunotherapies may prove more effective following an increase in lymphocyte density following neoadjuvant chemotherapy.

References

230

235

245

- 15 1. Ingold Heppner B, Loibl S, Denkert C. Tumor-Infiltrating Lymphocytes: A Promising Biomarker in Breast Cancer. *Breast Care (Basel)*. 2016;11(2):96-100.
 - 2. Earl HM, Vallier AL, Hiller L, et al. Effects of the addition of gemcitabine, and paclitaxelfirst sequencing, in neoadjuvant sequential epirubicin, cyclophosphamide, and paclitaxel for women with high-risk early breast cancer (Neo-tAnGo): an open-label, 2x2 factorial randomised phase 3 trial. *Lancet Oncol.* 2014;15(2):201-212.
 - 3. Ali HR, Dariush A, Provenzano E, et al. Computational pathology of pre-treatment biopsies identifies lymphocyte density as a predictor of response to neoadjuvant chemotherapy in breast cancer. *Breast Cancer Res.* 2016;18(1):21.
 - 4. Earl HM, Hiller L, Dunn JA, et al. Efficacy of neoadjuvant bevacizumab added to docetaxel followed by fluorouracil, epirubicin, and cyclophosphamide, for women with HER2-negative early breast cancer (ARTemis): an open-label, randomised, phase 3 trial. *Lancet Oncol.* 2015;16(6):656-666.
 - 5. Earl HM, Hiller L, Dunn D, et al. Disease-free and overall survival at 3.5 years for neoadjuvant bevacizumab added to docetaxel followed by fluorouracil, epirubicin and cyclophosphamide, for women with HER2 negative early breast cancer: ARTemis Trial. *Annals of Oncology (In press).* 2017.
 - 6. Thomas J, Provenzano E, Hiller L, et al. Central pathology review with two-stage quality assurance for pathological response after neoadjuvant chemotherapy in the ARTemis Trial. *Modern Pathology (In press).* 2017.
 - 7. Blows F, Driver K, Schmidt M, et al. Subtyping of breast cancer by immunohistochemistry to investigate a relationship between subtype and short and long term survival: a collaborative analysis of data for 10,159 cases from 12 studies. *PLoS Med.* 2010;7(5):e1000279.
 - 8. Ali HR, Provenzano E, Dawson SJ, et al. Association between CD8+ T-cell infiltration and breast cancer survival in 12 439 patients. *Ann Oncol.* 2014;25(8):1536-1543.
- 9. Loi S, Michiels S, Salgado R, et al. Tumor infiltrating lymphocytes are prognostic in triple negative breast cancer and predictive for trastuzumab benefit in early breast cancer: results from the FinHER trial. *Ann Oncol.* 2014;25(8):1544-1550.
 - 10. Adams S, Gray RJ, Demaria S, et al. Prognostic value of tumor-infiltrating lymphocytes in triple-negative breast cancers from two phase III randomized adjuvant breast cancer trials: ECOG 2197 and ECOG 1199. *J Clin Oncol.* 2014;32(27):2959-2966.
 - 11. Ali HR, Chlon L, Pharoah PD, Markowetz F, Caldas C. Patterns of Immune Infiltration in Breast Cancer and Their Clinical Implications: A Gene-Expression-Based Retrospective Study. *PLoS Med.* 2016;13(12):e1002194.
 - 12. Gentles AJ, Newman AM, Liu CL, et al. The prognostic landscape of genes and infiltrating immune cells across human cancers. *Nature medicine*. 2015;21(8):938-945.
 - 13. Bense RD, Sotiriou C, Piccart-Gebhart MJ, et al. Relevance of Tumor-Infiltrating Immune Cell Composition and Functionality for Disease Outcome in Breast Cancer. *J Natl Cancer Inst.* 2017;109(1).

255 **Tables**

Table 1. Univariable and multivariable logistic regression of lymphocyte density and clinical covariates against pCR.

		Univariate					Multivariate				
Variable	Categories	Odds ratio	95% CI	P-value	Observations	Odds ratio	95% CI	P-value	Observations		
Median lymphocyte density	Continuous	2.93	1.77-4.85	0.00003	609	2.13	1.24-3.67	0.006			
Grade	1,2,3	4.82	2.80-8.29	< 0.00001	557	2.80	1.58-4.96	0.0004			
ER status	Negative, Positive	0.19	0.12-0.30	< 0.00001	609	0.29	0.18-0.47	< 0.00001			
Age	Continuous	0.97	0.94-0.99	0.007	609	0.98	0.95-1.00	0.06	557		
Node status	Negative, Positive	0.69	0.45-1.04	0.08	609	0.65	0.41-1.05	0.08			
Chemotherapy	BEV+D FEC, D FEC	0.72	0.48-1.10	0.13	609	0.60	0.38-0.97	0.04			
Tumour size	<51mm, >50mm	0.73	0.42-1.26	0.25	609	1.05	0.56-1.97	0.87			

Abbreviations: a.u., arbitrary units, FEC, fluorouracil, epirubicin and cyclophosphamide; BEV, bevacizumab; pCR,

260 pathological complete response

Figures

265

Figure 1. Flowchart of patients and samples through analytic stages.

Figure 2. Association between lymphocyte density, change in lymphocyte density, cellular proportions and chemotherapy response.

Observations are ranked by pre-treatment lymphocyte density scores. Lymphocyte density has been rescaled to between zero and one for illustration (a.u., arbitrary units; pCR, pathological complete response; RD residual disease)

Supplementary Material

270 Supplementary Figures

275

<u>Supplementary Figure 1. Computational pathology workflow.</u> A. Digital whole slide images were segmented into subimages (B) and cellular nuclei segmented from these images. C. Support-vector-machine classification was used to classify nuclei into cancer, stromal and lymphocyte categories. D. Spatial statistics per slide, including lymphocyte density, were computed based on cell classes.

Supplementary Figure 2. Strip plots depicting the distribution of lymphocyte density by clinical variables.

P-values derived from Kruskal-Wallis tests.

280 Supplementary Tables

Supplementary Table 1. Summary of patient characteristics.

<u>Supplementary Table 2. Univariate Cox proportional hazards models for disease-free and overall</u> <u>survival, separately for ER-positive and ER-negative disease</u> <u>Supplementary Table 3. Univariate and multivariate logistic regression analyses for change in</u>

²⁸⁵ lymphocyte density and clinical covariates against pCR.





Supplementary Material

Supplementary Figures

Supplementary Figure 1. Computational pathology workflow.

A. Digital whole slide images were segmented into subimages (B) and cellular nuclei segmented from these images. C. Support-vector-machine classification was used to classify nuclei into cancer, stromal and lymphocyte categories. D. Spatial statistics per slide, including lymphocyte density, were computed based on cell classes.

Supplementary Figure 2. Strip plots depicting the distribution of lymphocyte density by clinical variables.

P-values derived from Kruskal-Wallis tests.

Supplementary Tables

Supplementary Table 1. Summary of patient characteristics.

Supplementary Table 2. Univariate Cox proportional hazards models for disease-free and overall

survival, separately for ER-positive and ER-negative disease

Supplementary Table 3. Univariate and multivariate logistic regression analyses for change in

lymphocyte density and clinical covariates against pCR.

Supplementary Figures



Supplementary Figure 1. Computational pathology workflow.

A. Digital whole slide images were segmented into subimages (B) and cellular nuclei segmented from these images. C. Support-vector-machine classification was used to classify nuclei into cancer, stromal and lymphocyte categories. D. Spatial statistics per slide, including lymphocyte density, were computed based on cell classes.



Supplementary Figure 2. Strip plots depicting the distribution of lymphocyte density by clinical variables.

P-values derived from Kruskal-Wallis tests.

Supplementary Tables

			Pathological co	omplete	response		
	No pCR		pCR			Total	
	No.	%	No.	%	No.	%	
Tumour size							
<51mm	393	78.6	91	83.5	484	79.5	
>50mm	107	21.4	18	16.5	125	20.5	
Total	500	100	109	100	609	100	
Node status							
Negative	233	46.6	61	56	294	48.3	
Positive	267	53.4	48	44	315	51.7	
Total	500	100	109	100	609	100	
Grade							
1	12	2.4	1	0.9	13	2.1	
2	212	42.4	15	13.8	227	37.3	
3	229	45.8	88	80.7	317	52.1	
Missing	47	9.4	5	4.6	52	8.5	
Total	500	100	109	100	609	100	
Chemotherapy							
BEV+D FEC	235	47	60	55	295	48.4	
D FEC	265	53	49	45	314	51.6	
Total	500	100	109	100	609	100	
ER status							
Negative	117	23.4	67	61.5	184	30.2	
Positive	383	76.6	42	38.5	425	69.8	
Total	500	100	109	100	609	100	
Diagnostic biopsies							
Analysed	500	100	109	100	609	100	
Total	500	100	109	100	609	100	
Surgical samples							
Missing	134	26.8	92	84.4	226	37.1	
Analysed	366	73.2	17	15.6	383	62.9	
Total	500	100	109	100	609	100	
All-cause mortality							
No	417	83.4	93	85.3	510	83.7	
Yes	83	16.6	16	14.7	99	16.3	
Total	500	100	109	100	609	100	
Disease recurrence							
No	378	75.6	89	81.7	467	76.7	
Yes	122	24.4	20	18.3	142	23.3	
Total	500	100	109	100	609	100	
		No	pCR		p	CR	
	N	Median	IQR	N	Median	IQR	
Median lymphocyte density*	500	-4.324	-4.755 to -4.065	109	-4.145	-4.482 to -3.910	
Change in lymphocyte density [†]	366	-0.154	-0.431 to 0.185	17	-0.473	-1.207 to -0.22	

Supplementary Table 1. Summary of patient characteristics.

*Log-transformed values

[†]Pre-post treatment value

Abbreviations: FEC, fluorouracil, epirubicin and cyclophosphamide; BEV, bevacizumab; pCR, pathological complete response

<u>Supplementary Table 2.</u> Univariate Cox proportional hazards models for disease-free and overall <u>survival</u>, separately for ER-positive and ER-negative disease

		ER-positive				ER-negative					
Variable	Categories	HR	95% CI	P-value	Observations	Events	HR	95% CI	P-value	Observations	Events
Overall survival											
Median lymphocyte density*	Continuous	1.34	0.79-2.28	0.28	418	49	1.02	0.58-1.78	0.95	178	49
Change in lymphocyte density*	Continuous	0.68	0.39-1.21	0.19	277	31	1.12	0.69-1.82	0.64	97	34
Disease-free survival							•				
Median lymphocyte density*	Continuous	1.27	0.85-1.92	0.24	418	80	0.92	0.55-1.52	0.74	178	60
Change in lymphocyte density*	Continuous	0.73	0.47-1.14	0.17	277	50	1.34	0.84-2.16	0.22	97	38

*Arbitrary units

Supplementary Table 3. Univariate and multivariate logistic regression analyses for change in

lymphoc	yte densit	y and cl	linical	covariates	against	pCR.
	-				-	

			Univ	variate		Multivariate				
Variable	Categories	Odds ratio	95% CI	P-value	Observations	Odds ratio	95% CI	P-value	Observations	
Median lymphocyte density*	Continuous	2.93	1.77-4.85	0.00003	609	0.49	0.093-2.61	0.41		
Change in lymphocyte density*	Continuous	0.24	0.11-0.50	0.0001	383	0.10	0.033-0.31	0.00007		
Age	Continuous	0.97	0.94-0.99	0.007	609	1.03	0.96-1.10	0.44		
Tumour size	<51mm, >50mm	0.73	0.42-1.26	0.25	609	0.70	0.13-3.69	0.67	240	
Grade	1,2,3	4.82	2.80-8.29	< 0.00001	557	6.06	1.16-31.7	0.03	349	
ER status	Negative, Positive	0.19	0.12-0.30	< 0.00001	609	0.14	0.031-0.59	0.008		
Chemotherapy	BEV+D FEC, D FEC	0.72	0.48-1.10	0.13	609	4.49	1.01-20.0	0.05		
Node status	Negative, Positive	0.69	0.45-1.04	0.08	609	0.69	0.20-2.36	0.56		

*Arbitrary units

Abbreviations: FEC, fluorouracil, epirubicin and cyclophosphamide; BEV, bevacizumab; pCR, pathological complete

response

_