A NOTE ON THE FIBRES OF MORI FIBRE SPACES

GIULIO CODOGNI, ANDREA FANELLI, ROBERTO SVALDI, AND LUCA TASIN

ABSTRACT. In this note we consider the problem of determining which Fano manifolds can be realised as fibres of a Mori fibre space. In particular, we study the case of toric varieties, Fano manifolds with high index and some Fano manifolds with high Picard rank.

Contents

Introduction	1
1. Toric Fano varieties	4
2. Fano manifolds of high index	12
3. Fano manifolds with high Picard number	15
4. Open questions	16
References	17

INTRODUCTION

In algebraic geometry, one of the main goals is to classify algebraic varieties. Rather than distinguishing varieties based on their isomorphism type, one can look at their birational type, i.e. the structure of a non-empty Zariski open set of the variety, which allows more flexibility in the choice of preferred model for the object under scrutiny.

From this point of view, the study of the canonical bundle of a smooth projective variety, that is, the determinant of the cotangent bundle, plays a central role in the classification. There is a stark dichotomy between algebraic varieties that admit sections of powers of the canonical bundle and those that do not.

In fact, while it is expected that the former are birational to a fibration in Calabi-Yau varieties over a base of (log-)general type, using a suitable realization of the litaka fibration, the latter are instead expected to be birational to a fibration in Fano varieties over a smaller dimensional base.

The following definition gives the precise notion needed in the latter case. Let us remind the reader that a normal projective variety is said to be Fano if the anticanonical divisor is ample (and in particular it is Q-Cartier).

Definition 0.1. Let $f: X \to Y$ be a dominant projective morphism of normal varieties. Then X is a *Mori fibre space* (or *MFS*) if

- $f_*\mathcal{O}_X = \mathcal{O}_Y$ and dim $Y < \dim X$;
- X is \mathbb{Q} -factorial with klt singularities;
- the general fibre is a Fano variety and $\rho(X/Y) := \rho(X) \rho(Y) = 1$.

²⁰¹⁰ Mathematics Subject Classification. 14J45, 14E30, 14M25, 14J35, 14J40.

While it is a difficult problem to prove that a variety having no non-zero pluricanonical forms is birational to a Mori fibre space, in [BCHM10] the authors show, among other things, that this is actually the case if instead the variety is assumed to have a non-pseudoeffective canonical bundle – the two conditions being expected to be equivalent as predicted by the existence of minimal models and the celebrated Abundance Conjecture, cf. [KM98, Conjecture 3.12]. In view of these considerations, it is natural to wonder about the following matter.

Question 0.2. What type of Fano varieties appear as fibers in a Mori fibre space?

Part of the difficulty in answering the above question lays in the lack of clarity as to what fibers of a MFS one should actually consider. To this end, the notion of fibre-like Fano variety was introduced in [CFST16]. For the sake of simplicity, we only consider the case of smooth fibers.

Definition 0.3. A Fano manifold F is *fibre-like* if it can be realised as a fibre of a Mori fibre space $f: X \to Y$ over the smooth locus of f.

Any Fano manifold with Picard number $\rho = 1$ is fibre-like, via the constant map to a point. When $\rho \geq 2$ the problem of determining whether a variety is fibre-like or not is highly non-trivial. Mori showed in [Mor82, Theorem 3.5] which Fano surfaces are fibre-like. In [CFST16], we systematically study fibre-like Fano varieties by analyzing the action of the monodromy of the MFS on the Néron-Severi group of a general fibre. Moreover, we fully characterize those threefolds that are fibre like, cf. [CFST16, Theorem 1.4] and give the following sufficient condition for fibre-likeness in any dimension.

Theorem 0.4 ([CFST16], Theorem 3.1). A smooth Fano variety F is fibre-like if

(1)
$$\operatorname{NS}(F)^{\operatorname{Aut}(F)}_{\mathbb{O}} = \mathbb{Q}K_F.$$

When F is rigid, then property (1) is equivalent to F being fibre-like.

In [CFST16], we also establish that the fibre-likeness of a Fano manifold F implies an analogous necessary condition: namely, F fibre-like implies

$$\mathrm{NS}(F)^{\mathrm{Mon}(F)}_{\mathbb{O}} = \mathbb{Q}K_F,$$

where Mon(F) is the maximal subgroup of $GL(NS(F), \mathbb{Z})$ which preserves the birational data of F.

Below we recall some of the consequences of our analysis. First, we construct a large class of examples of fibre-like Fano manifolds.

Corollary 0.5 ([CFST16], Corollary 4.6). Let r, k, d be integers with $n \ge 2$ and kd < n+1. Then any smooth complete intersection of k divisors of degree (d, \ldots, d) in $(\mathbb{P}^n)^r$ is fibre-like.

Moreover, since the notion of fibre-likeness is strictly intertwined with the monodromy action, we show that it forces a high degree of symmetry on facets of the nef cone of the variety. Recall that a facet of a polyhedral cone is just a maximal dimensional face. As Fano varieties are Mori dream spaces, any facet \mathcal{G} of the nef cone F corresponds to a contraction $\pi: F \to G$ such that $\mathcal{G} = \pi^* \operatorname{Nef}(G)$.

Corollary 0.6 ([CFST16], Corollary 3.9). Let F be a smooth fibre-like Fano variety. Let \mathcal{G} be a facet of the nef cone of F corresponding to a contraction $F \to G$.

Then for any other facet \mathcal{H} with contraction $F \to H$ we have that G and H are deformation equivalent.

Let us point out that the definition of fibre-like can be extended to singular varieties (cf. [CFST16], Definition 2.14) and all results illustrated so far still hold in that setting.

In this note we show that various natural and interesting classes of Fano manifolds are fibre-like.

We first focus on smooth toric Fano varieties. If $F(\Delta)$ is a smooth toric Fano variety with associated polytope Δ , then F is said to be vertex-transitive if the automorphism group of the polytope acts transitively on the set of vertices of Δ . A vertex-transitive toric Fano manifold is fibre-like (see Lemma 1.2) and all the known examples of toric fibre-like Fano varieties seem to be of this kind (cf. Question 4.1).

Among these, projective spaces and t-del Pezzo manifolds (see Definition 1.3) are classical examples of vertex-transitive varieties. A generalisation of t-del Pezzo manifolds has been introduced in [Kly84] and studied in [VK84]. We call those Klyachko varities (see Definition 1.17). We show that Klyachko varities are vertextransitive and we show that they constitute a fundamental building block in the theory of fibre-like toric Fano manifolds.

Theorem 0.7 (Propositions 1.13 and 1.23). Let $F = F(\Delta)$ be a d-dimensional vertex-transitive Fano manifold.

(i) If there are two vertices of Δ that are not in the same face (i.e. Δ is not 2-neighbourly), then F is a power of \mathbb{P}^1 or a power of t-del Pezzo manifolds. (ii) If d < 7, then F is a power of projective spaces or Klyachko varieties.

Next we look at Fano manifolds with high index. As mentioned above, fibrelikeness is completely understood for surfaces and threefolds. Hence, here we focus on Fano varieties of dimension at least 4.

For a Fano manifold F, the index i_F is defined as the largest integer that divides $-K_F$ in Pic(F). The index is one of the most basic numerical invariants for Fano varieties: there is a complete classification of Fano manifolds with Picard number at least 2 and index $i \ge n-2$ due to Kobayashi-Ochai [KO73], Fujita [Fuj82a, Fuj82b] and Mukai [Muk89]. Moreover, Wiśniewski [Wiś91] classified Fano manifolds with index $i \ge (n+1)/2$. Notice that the only case for which i < (n+1)/2 and $i \ge n-2$ is i = 2 and n = 4.

When the Fano index is greater then the above bounds, we show that fibre-like Fano varieties can be explicitly classified.

Theorem 0.8 (Propositions 2.2 and 2.3). Let F be a fibre-like Fano manifold with Picard number $\rho \geq 2$ and dimension $n \geq 4$.

- If $i_F \ge \frac{n+1}{2}$, then $F \cong \mathbb{P}^{n/2} \times \mathbb{P}^{n/2}$ (n even) or $F \cong \mathbb{P}\left(T_{\mathbb{P}^{\frac{n+1}{2}}}\right)$ (n odd). If n = 4 and $i_F = 2$ (i.e. F is a Mukai fourfold), then F is isomorphic to one of the following:
 - (i) a double cover of $\mathbb{P}^2 \times \mathbb{P}^2$ branched along a degree-(2,2) divisor;
 - (ii) an intersection of two degree–(1,1) divisors in $\mathbb{P}^3 \times \mathbb{P}^3$; (iii) $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$.

Finally, we discuss families of Fano manifolds with high Picard number. As there only exist finitely many families of Fano manifolds in any given dimension, there must be an upper bound on the Picard number in fixed dimension. It is a natural question to ask whether it is possible to compute such bound.

The only known examples of Fano manifolds – other than products – of dimension $n \ge 4$ and Picard number at least n + 5 are birational modifications of the blow-up of \mathbb{P}^4 at 8 points in general position. These manifolds have been studied in [CCF17]; among other results, the authors showed that they are fibre-like. Here, we study the only known examples of Fano manifolds – again, other than products – that have dimension $n \ge 4$ and Picard number at least n+4. These are birational modification of \mathbb{P}^n blown up at n + 3 general points, with n even. They are isomorphic to the space of (m-1)-planes in the intersection of two quadrics in \mathbb{P}^{n+2} , where n = 2m. Building on [AC17], we can prove the following result.

Theorem 0.9 (Theorem 3.2). Let $n \ge 4$ be an even integer. An *n* dimensional Fano manifold birational to the blow-up of \mathbb{P}^n at n+3 general points is fibre-like.

The structure of the paper is as follows. Section 1 is dedicated to toric Fano manifolds. In Section 2 we discuss the fibre-likeness of Fano manifolds with high index. Section 3 is devoted to the study of certain families of Fano manifolds with high Picard number. Finally, in Section 4 we present new questions and research directions.

Acknowledgement. We like to thank Cinzia Casagrande for interesting comments on this work. The second-named author has been supported by the DFG grant "Gromov-Witten Theorie, Geometrie und Darstellungen" (PE 2165/1-2) and by the Swiss National Science Foundation Grant "Algebraic subgroups of the Cremona groups" (200021–159921). The fourth-named author is supported by the DFG grant "Birationale Methoden in Topologie und Hyperkähler Geometrie".

Notation. The term *variety* stays for separated, integral and proper scheme of finite type over \mathbb{C} . A *manifold* is a smooth variety. For the definitions of singularities in the context of the minimal model program, see [Kol13, Section 2].

1. TORIC FANO VARIETIES

The first class of examples we want to discuss is given by toric varieties. We will recall here only some notions and refer to [CLS11] for an exhaustive treatment of the topic. Toric varieties can be described in terms of combinatorial data and this makes them particularly suitable to test general conjectures on Fano varieties.

Let N be a free abelian group of rank n and set $N_{\mathbb{Q}} := N \otimes_{\mathbb{Z}} \mathbb{Q}$. Let $\Sigma \subset N_{\mathbb{Q}}$ be a fan of a d-dimensional toric Fano variety F and let Δ be the polytope associated to the anti-canonical polarisation. Furthermore, M will denote the dual of N.

The vertices of Δ , denoted by $V(\Delta)$, are the generators of Σ . We denote by $O(\sigma)$ the closure of the orbit corresponding to $\sigma \in \Sigma$, which is an irreducible invariant subvariety.

Let A_1 be the group of 1-cycles on F modulo numerical equivalence and set $N_1 = A_1 \otimes \mathbb{Q}$. Inside N_1 , we consider the Kleiman-Mori cone $\overline{\text{NE}}(F)$ generated by the effective 1-cycles. We have the following standard exact sequence (cf. [CLS11, Chapter 4, Theorem 1.3])

(2)
$$0 \to A_1(F) \to \mathbb{Z}^{V(\Delta)} \to N \to 0,$$

which by duality yields the following one:

(3)
$$0 \to M \to \mathbb{Z}^{V(\Delta)} \to \mathrm{NS}(F) \to 0$$

1.1. Vertex-transitive Polytopes. The following definition of vertex-transitive for polytopes is classical, although some authors refer to them as isogonal polytopes (cf. [GO04, 19.5, Enumeration]).

Definition 1.1. A polytope Δ is *vertex-transitive* if Aut(Δ) acts transitively on the vertices of Δ . If Δ is associated to a toric Fano variety F, then F is vertex-transitive.

This class of varieties is interesting from our prospective for the following reason.

Lemma 1.2. Vertex-transitive Fano manifolds are fibre-like.

Proof. Let $F(\Delta)$ be a Fano toric variety and let $G = \operatorname{Aut}(\Delta)$. As explained in the proof of [CFST16, Theorem 5.7], the exact sequence in (3) yields the following sequence

$$0 \to M^G_{\mathbb{O}} \to (\mathbb{Q}^{V(\Delta)})^G \to \mathrm{NS}(F)^G_{\mathbb{O}} \to 0.$$

Then, F is fibre-like if and only if t - k = 1, where t is the number of orbits of the action of G on $V(\Delta)$ and $k = \dim M_{\mathbb{Q}}^G$.

If F is vertex-transitive, then t = 1 and so [CFST16, Lemma 5.10] implies that k = 0, which means that F is fibre-like.

Denote by d the dimension of the toric Fano variety F and by m the number of vertices of Δ .

The first non-trivial class of vertex-transitive Fano varieties are *t*-del Pezzo manifolds.

Definition 1.3. The *d*-dimensional *t*-del Pezzo manifold V_d (with *d* even) is the smooth toric Fano variety whose associated polytope has vertices

$$V(\Delta) = \{e_1, \dots, e_d, -e_1, \dots, -e_d, (e_1 + \dots + e_d), -(e_1 + \dots + e_d)\},\$$

where e_1, \ldots, e_d is the standard basis of $N_{\mathbb{O}}$.

Remark 1.4. In the literature on toric geometry, these manifolds are simply named del Pezzo varieties. We added the prefix "t-" in order to distinguish them from the del Pezzo manifolds appearing in Section 2.

T-del Pezzo polytopes are symmetric with respect to the origin, i.e., $-\Delta = \Delta$. Polytopes satisfying this condition are also said to be *centrally symmetric*, cf. [VK84]. A classic result by Voskresenkii and Klyachko shows that del Pezzo varieties are essentially the only centrally symmetric toric varieties.

Theorem 1.5 ([VK84, Theorem 6]). Let F be a toric Fano manifold such that Δ is centrally symmetric. Then F is isomorphic to a product of projective lines and t-del Pezzo varieties.

Coming back to vertex-transitive varieties, we can prove the following structural result.

Lemma 1.6. Let $F = F(\Delta)$ be a toric Fano manifold which is vertex-transitive. Then there exists a unique vertex-transitive Fano toric manifold F_{min} and a positive integer n such that $F \cong (F_{min})^n$. *Proof.* Let $\Delta = \Delta_1^{n_1} \times \ldots \times \Delta_r^{n_r}$ be a prime decomposition of Δ . The automorphism group of Δ is given by (see for example [GH16, Theorem A])

$$\operatorname{Aut}(\Delta) = \prod_{i=1}^{r} (\operatorname{Aut}(\Delta_i) \rtimes S_{n_i})$$

with its natural action on Δ .

Since $\operatorname{Aut}(\Delta)$ acts transitively on the vertices of Δ , the lemma follows immediately.

1.2. **Primitive collections.** Primitive collections are an essential tool to study the birational geometry of Fano toric varieties. We refer to [Rei83] and [Cas03] for further details.

Definition 1.7. Let $F = F(\Delta)$ be a toric Fano variety. A subset $P \subset V(\Delta)$ is called a *primitive collection* if the cone generated by P is not in Σ , but for any $x \in P$ the elements of $P \setminus \{x\}$ generate a cone in Σ .

For a primitive collection $P = \{x_1, \ldots, x_k\}$ denote by $\sigma(P)$ the (unique) minimal cone in Σ such that $(x_1 + \ldots + x_k) \in \sigma(P)$. Let y_1, \ldots, y_h be generators of $\sigma(P)$, then

(4)
$$r(P): x_1 + \ldots + x_k = b_1 y_1 + \ldots + b_h y_h$$

where b_i is a positive integer for all $1 \le i \le h$: we have simply written the element $x_1 + \ldots + x_k$ in terms of the generators y_1, \ldots, y_h (the coefficients are positive since $(x_1 + \ldots + x_k)$ is in the cone $\sigma(P)$).

The linear relation (4) is called the primitive relation of P and the cone $\sigma(P)$ is called the focus of P. The integer k is called the *length of* r(P) and the *degree of* P is defined as deg $P = k - \sum b_i$.

Here it is convenient to write down explicitly the group of 1-cylces A_1 of F as:

$$A_1(F) \cong \left\{ (b_x)_{x \in V(\Delta)} \in \operatorname{Hom}(\mathbb{Z}^m, \mathbb{Z}) \ \middle| \ \sum_{x \in V(\Delta)} b_x x = 0 \right\}.$$

The previous isomorphism is clear looking at the exact sequence (2). So it is natural to identify primitive relations with the associated cycles. Moreover, we work on Fano varieties, so deg $P = -(K_F \cdot r(P)) > 0$ for all primitive relations.

Consider now a primitive collection P on F for which the relation r(P) is *extremal*, meaning that it generates an extremal ray in $\overline{\text{NE}}(F)$. One sees that the exceptional locus of the associated contraction is given by $O(\sigma(P))$ and moreover, according to the dimension of $\sigma(P)$, one recovers:

- divisorial contraction when $\sigma(P)$ is a one-dimensional cone and the contracted divisor is precisely the one associated to the ray;
- Mori fibration, when $\sigma(P)$ coincides with the origin;
- *flipping contraction* otherwise.

Let us recall some useful results.

Proposition 1.8 ([Cas03, Prop. 4.3]). Let $\gamma \in \overline{NE}(F) \cap A_1(F)$ be a 1-cycle of F for which $(K_F \cdot \gamma) = -1$. Then γ is extremal.

Theorem 1.9 ([Rei83, Theorem 2.4], [Cas03, Theorem 1.5]). Let $R \subset \overline{NE}(F)$ be an extremal ray and let $\gamma \in R \cap A_1(F)$ be a primitive cycle. Then there exists a primitive collection $P = \{x_1, \ldots, x_k\}$ such that

 $\gamma = r(P) \colon x_1 + \ldots + x_k = b_1 y_1 + \ldots + b_h y_h.$

Moreover, for any cone $\nu = \langle z_1, \ldots, z_t \rangle$ which verifies

• $\{z_1, \ldots, z_t\} \cap \{x_1, \ldots, x_k, y_1, \ldots, y_h\} = \emptyset$; and

• $\langle y_1, \ldots, y_h \rangle + \nu \in \Sigma;$

the following holds for all i = 1, ..., h:

$$\langle x_1,\ldots,\check{x_i},\ldots,x_k,y_1\ldots,y_h\rangle+\nu\in\Sigma.$$

Proposition 1.10 ([Cas03, Prop. 3.4]). Let P a primitive extremal collection for F and write $\sigma(P) = \langle y_1, \ldots, y_h \rangle$. Then for any other primitive collection $Q \neq P$ for which $P \cap Q \neq \emptyset$, the set $(Q \setminus P) \cup \{y_1, \ldots, y_h\}$ contains a primitive collection.

The following observation is easy but useful for our analysis.

Remark 1.11. Consider a relation

$$_1x_1 + \dots + a_kx_k = b_1y_1 + \dots + b_hy_h$$

among the vertices of Δ , with $a_i, b_j > 0$ for all i, j. If $\sum a_i \geq \sum b_j$, then [Cas03, Lemma 1.4] implies that $\langle x_1 \cdots, x_k \rangle \notin \Sigma$.

We recall now the following definition.

Definition 1.12 (k-neighbourly polytope). A polytope is k-neighbourly if every set of k vertices lies on one of its face.

A Fano variety $F(\Delta)$ is k-neighbourly if the corresponding polytope Δ is.

We want now to understand the structure of vertex-transitive polytopes: the following result is the first step towards a classification of vertex-transitive those.

Proposition 1.13. Let $F = F(\Delta)$ be a vertex-transitive toric Fano manifold. Then either

- (1) $F = (\mathbb{P}^1)^d$ or $F = (V_k)^r$ for some r and k or
- (2) Δ is 2-neighbourly.

Proof. Let us assume that Δ is not 2-neighbourly, which implies the existence of a primitive collection with two elements. We claim that this primitive relation can be assumed to be of the form

To show this, we assume such a relation does not exist and seek for contradiction. Take a primitive collection $P_1 = \{x_1, x_2\}$ verifying the relation $R_1: x_1 + x_2 = y_1$. Let $\operatorname{Aut}(\Delta)$ act on P_1 to obtain a family of primitive collections $\mathcal{P} = \{P_i\}_{1 \leq i \leq r}$ with relations $\mathcal{R} = \{R_i\}_{1 \leq i \leq r}$. Since the action is transitive by hypothesis, any vertex of Δ appears the same number of times as right hand side of these relations and so the number of vertices $m := \#\{V(\Delta)\}$ divides r. This implies that the P_i 's cannot be all disjoint, otherwise 2r = m. Hence we may assume that $P_2 = \{x_1, x_3\}, x_2 \neq x_3$, with relation $R_2: x_1 + x_3 = y_2$.

The two relations R_1 and R_2 give $x_2 + y_2 = x_3 + y_1$, which implies, by Remark 1.11, that $\{x_2, y_2\}$ is also a primitive collection. The two relations

$$R': x_3 + y_1 = z_1$$
 and $R'': x_2 + y_2 = z_1$

are extremal, so Proposition 1.8 and Theorem 1.9 imply that $\langle y_1, y_2 \rangle \in \Sigma$. This is a contradiction, since $y_1 + y_2 = x_1 + z_1$. We proved the existence of (5).

Now act with $\operatorname{Aut}(\Delta)$ to get exactly m/2 relations of the same form. One can verify that those are disjoint. Using the vertex-transitivity of Δ , we deduce that for any vertex \bar{x} there is a vertex \bar{y} for which $\bar{x} + \bar{y} = 0$, i.e. Δ is centrally symmetric. Theorem 1.5 concludes the proof.

We study now the extremal contractions of 2-neighbourly vertex-transitive toric Fano manifolds.

Lemma 1.14. Let $F = F(\Delta)$ be a vertex-transitive, 2-neighbourly toric Fano manifold. Then there exist an integer $k \geq 3$ and a set of primitive collections

$$\mathcal{P} = \{P_i\}_{i=1,\dots,r}$$

such that r = m/k, $|P_i| = k$, $\sigma(P_i) = 0$ and $P_i \cap P_j = \emptyset$ for any $i \neq j$. Moreover, these are the only primitive relations with focus equal to zero.

Proof. The result in [Bat91, Proposition 3.2] implies that there exists a primitive collection P_1 with $\sigma(P_1) = 0$. Define $k := |P_1|$. Since Δ is 2-neighbourly, we have $k \geq 3$.

Act with $\operatorname{Aut}(\Delta)$ to get a set of primitive collections $\mathcal{P} = \{P_i\}_{1 \leq i \leq r}$ verifying $\sigma(P_i) = 0$ and for which $\bigcup_{i=1}^r P_i = V(\Delta)$. Let us prove they are disjoint, assuming that $P_i \cap P_j \neq \emptyset$ for some i, j and seeking for contradiction. Write $P_i = \{x_1, \ldots, x_k\}$ and $P_j = \{x_1, \ldots, x_h, y_{h+1}, \ldots, y_k\}$ with $y_s \neq x_t$ for any s, t. Then

$$x_{h+1} + \ldots + x_k = y_{h+1} + \ldots + y_k.$$

Remark 1.11 gives the required contradiction, since x_{h+1}, \ldots, x_k generate a cone in Σ . Moreover one sees that there are no other primitive relations with focus equal to zero.

Proposition 1.15. In the notation of Lemma 1.14, assume that one of the relations P_i is extremal. Then $F(\Delta) = (\mathbb{P}^{k-1})^r$. On the other hand, if any of these relations is not extremal then F does not admit any extremal contraction of fibre type.

Proof. Up to reordering, assume that P_1 is extremal. Acting with $\operatorname{Aut}(\Delta)$, we deduce that all P_i 's are extremal. We claim these are the only primitive collections. In fact, let \tilde{P} be a primitive collection such that $\tilde{P} \notin \mathcal{P}$ and \tilde{P} has minimal cardinality among the primitive collections which are not in \mathcal{P} . We may assume that $P_1 \cap \tilde{P} \neq \emptyset$. Using Proposition 1.10, we deduce that the set $(\tilde{P} \setminus P_1)$ contains a primitive collection. Contradiction, since $|\tilde{P}|$ is minimal.

Note that k is the index of K_F , dim F = d = (k - 1)r and $\rho(F) = r$ by [Bat91, Corollary 4.4]. So apply [Cas06, Theorem 1] (Mukai's conjecture) to obtain the first part of the statement.

For the last part, just observe that an extremal contraction of fibre type would provide a primitive collection P with trivial focus $\sigma(P) = 0$.

Lemma 1.16. Let $F = F(\Delta)$ be a vertex-transitive, 2-neighbourly toric Fano manifold. Then there are no extremal relations of the form

$$(6) x_1 + \ldots + x_k = by_1.$$

In particular F does not admit any extremal divisorial contraction.

Proof. Let us assume that an extremal relation of the form (6) exists and seek for contradiction. Let $\mathcal{R} = \{R_i\}_{1 \leq i \leq r}$ be the set of extremal relations obtained acting with $\operatorname{Aut}(\Delta)$ and denote with P_i the associated collections. Assume that x_1 appears only in one P_i . Since by transitivity any vertex appears the same number of times, we get that the P_i 's are disjoint. In particular m = kr, where $m = \#\{V(\Delta)\}$. On the other hand, we have that m divides r, because any vertex appears the same number of times as right hand side. This implies k = 1, which is a contradiction.

Hence there is an extremal primitive relation different from (6) of the form

$$x_1 + z_2 \ldots + z_k = by_2.$$

Assume $y_2 \notin \{x_2, \ldots, x_k, y_1\}$ (the other case is analogous).

We have $b_1y_2 + x_2 + \ldots + x_k = by_1 + z_2 + \ldots + z_k$, and, since Δ is 2-neighbourly, we know that $\langle y_1, y_2 \rangle$ is a cone of Σ . Theorem 1.9 implies that $\langle y_2, x_2, \ldots, x_k \rangle \in \Sigma$, but this contradicts Remark 1.11.

1.3. Klyachko varieties. Looking for interesting examples of vertex-transitive toric varieties, we found a generalisation of t-del Pezzo varieties, which were introduced in [Kly84] and studied in [VK84].

Let us remark that our notation is not the same as Klyachko (cf. Remark 1.18). Fix a basis e_1, \ldots, e_d of a lattice $N \cong \mathbb{Z}^d$, with $d \ge 2$ and let k be a positive integers such that (k-1)|d.

Definition 1.17. The Klyachko variety of order k and dimension d is the toric Fano variety W_d^k with polytope $\Delta_d^k \subset N$ having vertices

$$V(\Delta_d^k) = \{e_1, e_2, \dots, e_d, e_1 + \dots + e_d, \\ - (e_1 + \dots + e_{k-1}), -(e_k + \dots + e_{2k-2}), \dots, -(e_{d-k+2} + \dots + e_d), \\ - (e_1 + e_k + \dots + e_{d-k+2}), -(e_2 + e_{k+1} + \dots + e_{d-k+3}), \dots, \\ - (e_{k-1} + e_{2k-2} + \dots + e_d)\}.$$

Remark 1.18. When d is even, W_d^2 is the t-del Pezzo manifold V_d . In [VK84], the varieties W_d^k are introduced as $P_{m,n}$. The dictionary between the indices is:

$$d = (m-1)(n-1), \quad k = m \ (or \ n).$$

As we will see in Lemma 1.19, our definition of k is consistent.

If W_d^k is smooth (cf. Proposition 1.21), we can describe some birational geometry of Klyachko varieties.

The 1-dimensional cones of the fan of W_d^k coincide with the 1-dimensional cones of the fan of the blow-up Z_d^k of $(\mathbb{P}^{k-1})^{\frac{d}{k-1}}$ in k invariant points. This implies that W_d^k and Z_d^k are isomorphic in codimension one and W_d^k is a Fano model of Z_d^k (cf. Section 3 for other examples of fibre-like Fano manifolds obtained as small modifications of blow-ups of projective spaces).

Lemma 1.19. For any integers d and m, $W_{md}^{d+1} = W_{md}^{m+1}$.

Proof. Assume $m \leq d$ and consider the vertices of W_{md}^{d+1} :

 $\{ e_1, e_2, \dots, e_{md}, e_1 + \dots + e_{md}, \\ - (e_1 + \dots + e_d), -(e_{d+1} + \dots + e_{2d}), \dots, -(e_{m(d-1)+1} + \dots + e_{md}), \\ - (e_1 + e_{d+1} + \dots + e_{m(d-1)+1}), -(e_2 + e_{d+2} + \dots + e_{m(d-1)+2}), \dots, \\ - (e_d + e_{2d} + \dots + e_{md}) \}.$

The following transformation

$$e'_{mi+j} := e_{d(j-1)+i+1}$$

where $i \in \{0, ..., d\}$ and $j \in \{1, ..., m - 1\}$ gives the identification.

We study now symmetries and singularities of Klyachko varieties.

Lemma 1.20. The Fano varieties W_d^k are vertex-transitive, reflexive and have terminal singularities, for all d, k.

Proof. Let us fix k and observe that Lemma 1.19 provides the following identification: $W_{k-1}^k \cong W_{k-1}^2$.

Vertex-transitivity is proved by induction on d: assume that for any (k-1)|d'and d' < d, the variety $W_{d'}^k$ is vertex-transitive. We write the projections

$$\pi_i \colon \Delta_d^k \longrightarrow \langle e_{i+1}, e_{i+2}, \dots, e_{i+k-1} \rangle^\perp,$$

with $i = 0, \ldots, d - k + 1$. By inductive hypothesis, the images via the π_i 's of Δ_d^k are vertex-transitive and they are all isomorphic to Δ_{d-k+1}^k . To prove the transitivity for the whole polytope, we act with $\operatorname{GL}(N_{\mathbb{Q}})$ to exchange the subspaces $\langle e_{i+1}, e_{i+2}, \ldots, e_{i+k-1} \rangle$.

We write now $W = W_d^k$ and $\Delta = \Delta_d^k$ to simplify the notation and prove reflexivity. Look at the dual polytope $\Delta^* \subset M_{\mathbb{Q}}$: we claim that no lattice point lies between the affine hyperplane spanned by the facets of Δ^* and its parallel through the origin. The claim holds for the hyperplane $\{x_1 = -1\} \subset M_{\mathbb{Q}}$, so acting with $\operatorname{Aut}(\Delta)$ on Δ^* we conclude.

Terminality can be translated on polytopes with the condition

$$\Delta \cap N = V(\Delta) \cup \{0\}.$$

We assume there exists a non-zero $v \in \Delta \cap N$ which verifies $v \notin V(\Delta)$ and seek for contradiction. Without loss of generality, assume that v is not in the subspace Hgenerated by e_1, \ldots, e_{k-1} and let π_H be the projection from H. Then the image $\Delta_H := \pi_H(\Delta)$ is a Klyachko polytope, $\pi_H(v) \in \Delta_H$ and $\pi_H(v) \notin V(\Delta_H) \cup \{0\}$. Since W_d^2 is terminal for $d \geq 2$, we obtain terminality by induction. \Box

We analyse smoothness, together with \mathbb{Q} -factoriality, for Klyachko varieties. It turns out that these properties depend on some divisibility conditions on the indices d and k (cf. [VK84]).

Let us fix some notation. For any positive d and k let \overline{d}_k be the smallest nonnegative integer r which verifies $d \equiv r \mod k$.

Fix integers $k \ge 2$ and $h \ge 1$ and let $\{x_1, \ldots, x_d\}$ be coordinates on $N_{\mathbb{Q}}$. Then define the following linear form on $N_{\mathbb{Q}}$:

$$L_{k,h} := \sum_{i=0}^{k-3} (x_{h+ik} + x_{h+ik+1} + \ldots + x_{h+ik+k-2} - (k-1)x_{h+ik+(k-1)}).$$

10

The following proposition already appeared in [VK84], in a different notation.

Proposition 1.21. The Klyachko variety W_d^k is smooth if gcd(d-1,k) = 1. If $gcd(d-1,k) \neq 1$, then W_d^k is not \mathbb{Q} -factorial.

Proof. The polytope $\Delta := \Delta_d^k$ is not simplicial for $d = (k-1)^2$, since the hyperplane $\{L_{k,1} + x_d = 1\}$ supports a facet of Δ with k(k-1) vertices.

On the other hand, we claim that the polytope Δ is smooth for d = k(k-1). To show this, one can see that any facet of Δ containing the vertex $(1, 1, \ldots, 1)$ also contains at least (k-1)(k-2) + 1 elements of the standard basis. This implies that the hyperplane

$$\{a_1x_1+\ldots+a_dx_d=1\}$$

supporting the facet has (exactly as for the hyperplane $\{L_{k,1} + x_d = 1\}$):

- (k-1)(k-2) + 1 coefficients equal to 1;
- k-2 coefficients equal to -(k-1);
- k-1 coefficients equal to 0.

One can verify that the vertices of all these facets give a basis of $N_{\mathbb{Q}}$. Using the transitivity of $\operatorname{Aut}(\Delta)$ we obtain the claim.

The general result on smoothness is proved via induction on k and d. Two cases are easy:

- (1) k = 2 and any d;
- (2) d = 2.

Take Δ_d^k with $k, d \ge 3$: if $d < (k-1)^2$ then $\Delta_d^k \cong \Delta_d^{l+1}$, where $l := k - \overline{d}_k$ and d = l(k-1). Since $\gcd(d-1,k) = \gcd(l+1,k) = \gcd(l+1,d-1)$, we conclude by induction of k.

Assume now d > k(k-1). Define h := d - k(k-1) and take the plane Hgenerated by $\{e_{h+1}, e_{h+2}, \ldots, e_d\}$ with projection π_H . Let define $\Delta_H := \pi_H(\Delta_d^k)$; then $\Delta_H = \Delta_h^k$ and $\gcd(d-1,k) = \gcd(h-1,k)$. For any facet \mathcal{F} of Δ_h^k supported on the hyperplane $\{P(x_1, \ldots, x_h) = 1\}$ we get a facet \mathcal{F}' of Δ_d^k supported on $\{P + L_{k,(h+1)} + (x_{d-k+1} + \ldots + x_{d-1} - (k-1)x_d) = 1\}$.

Observe that $|V(\mathcal{F}')| = |V(\mathcal{F})| + k(k-1)$. So if Δ_h^k is not simplicial, neither Δ_d^k is so. Analogously, one checks that Δ_d^k is smooth if and only if Δ_h^k is so. We conclude via induction on d.

Remark 1.22. As a consequence of the previous proposition, if k is a prime number then W_d^k is smooth, unless $d \equiv 1 \mod k$.

1.4. Low dimension. The results and the methods of the previous subsections are enough to classify all vertex-transitive Fano manifolds up to dimension 7. The result is confirmed by Table 1, which collects the Fano toric manifolds up to dimension 8 which are fibre-like¹.

Proposition 1.23. Let $F = F(\Delta)$ be a d-dimensional vertex-transitive Fano manifold. If $d \leq 7$, then F is a power of projective spaces or Klyachko manifolds.

Proof. The result can be proven using the software MAGMA together with the classification of smooth toric Fano varieties from the Graded Ring Database $[BK^+]$ (cf. Table 1): giving as input a list of smooth Fano polytopes, MAGMA can check

¹The table appeared in [CFST16] and has been obtained using the software MAGMA together with the Graded Ring Database $[BK^+]$ (for further details on the classification, cf. $[\emptyset b07]$)

in which cases $\operatorname{Aut}(\Delta)$ acts transitively on the vertexes².

When the dimension is at most 4, we are able to provide the following short argument, which does not require computer computations.

Let us start with d = 2. If Δ is not 2-neighbourly, then Proposition 1.13 implies that $F \cong \mathbb{P}^1 \times \mathbb{P}^1$ or $F \cong V_2$. If Δ is 2-neighbourly, then there is an extremal collection $P = \{x_1, x_2, x_3\}$ for which $\sigma(P) = 0$ and so, by Proposition 1.15, we have $F \cong \mathbb{P}^2$.

Assume now d = 3. If Δ is not 2-neighbourly, then Proposition 1.13 implies that $F \cong (\mathbb{P}^1)^3$. If Δ is 2-neighbourly, then the extremal relations could only be of the form $x_1 + x_2 + x_3 = 0$ (contradiction by Proposition 1.15), $x_1 + x_2 + x_3 = y_1$ (contradiction by Lemma 1.16) or $x_1 + x_2 + x_3 + x_4 = 0$. In this last case, $F \cong \mathbb{P}^3$ by Proposition 1.15.

Assume finally that d = 4. If Δ is not 2-neighbourly, then by Proposition 1.13 we get $X \cong (\mathbb{P}^1)^4$, $X \cong (V_2)^2$ or $X \cong V_4$. If there is an extremal relation of the form $x_1 + x_2 + x_3 = 0$, then by Proposition 1.15 we have $X \cong \mathbb{P}^2 \times \mathbb{P}^2$.

Hence assume that Δ is 2-neighbourly and let P be an extremal primitive collection. By Theorem 1.9 we have $|P| + |\sigma(P)| \leq 5$. By Lemma 1.16 we conclude that there is an extremal relation of the form $x_1 + x_2 + x_3 = y_1 + y_2$ or $x_1 + \ldots + x_5 = 0$. In the second case $X \cong \mathbb{P}^4$ and so we can assume to have $x_1 + x_2 + x_3 = y_1 + y_2$. From here it is not difficult to see that one should have $|V(\Delta)| \geq 12$, which is impossible by [Cas06, Theorem 1].

Remark 1.24. The 8-dimensional polytope denoted by \tilde{W} in Table 1 is not a Klyachko variety and we do not have a classic description of it.

2. Fano manifolds of high index

An important invariant of a Fano manifold F is its index, i_F , defined as the largest integer that divides $-K_F$ in Pic(F). There is a complete classification of Fano manifolds with index $i_F \ge n-2$ and $i_F \ge (n+1)/2$. In this section we investigate the fibre-likeness of these varieties, assuming $n \ge 4$ (fibre-like Fano 3-folds have been classified in [CFST16]).

In [Wiś91], Wiśniewski classified Fano manifolds with index $i_F \ge (n+1)/2$. Let us denote by $Q^j \subset \mathbb{P}^{j+1}$ the *j*-dimensional smooth projective quadric and by $T_{\mathbb{P}^l}$ the tangent bundle of \mathbb{P}^l .

Theorem 2.1. [Wiś91] Let F be a Fano manifold of dimension n and index $i_F \ge (n+1)/2$. Then F verifies one of the following:

```
(i) \rho(F) = 1;
```

(ii) *n* is even and $F \simeq \mathbb{P}^{\frac{n}{2}} \times \mathbb{P}^{\frac{n}{2}}$;

²We briefly describe the MAGMA code. Given an integer i, the function PolytopeSmoothFano(i) gives the polytope of the *i*-th toric Fano in the Graded Ring Database; we denote by N the number of Fano polytopes in the database; at the end of a run of the following code, the variable Pol will contain the list of vertex-transitive toric Fano polytopes in the Graded Ring Database.

 $[\]operatorname{Pol}:=[**];$

for i := 1 to N do if $\#\{\{v * G : G \text{ in AutomorphismGroup}(\text{PolytopeSmoothFano}(i))\}$:

 $v \text{ in Vertices}(\texttt{PolytopeSmoothFano}(i))\}$

 $^{- \}mbox{Dimension}(\mbox{FixedSubspaceToPolyhedron}(\mbox{AutomorphismGroup}(\mbox{PolytopeSmoothFano}(i)))) \ \mbox{eq 1 then Pol} = \mbox{Append}(\mbox{Pol,PolytopeSmoothFano}(i));$

 $\begin{array}{ll} (iii) \ n \ is \ odd \ and \ F \simeq \mathbb{P}^{\frac{n-1}{2}} \times Q^{\frac{n+1}{2}};\\ (iv) \ n \ is \ odd \ and \ F \simeq \mathbb{P}\Big(T_{\mathbb{P}^{\frac{n+1}{2}}}\Big);\\ (v) \ n \ is \ odd, \ \mathcal{O} = \mathcal{O}_{\mathbb{P}^{\frac{n+1}{2}}} \ and \ F \simeq \mathbb{P}\Big(\mathcal{O}(1) \oplus \mathcal{O}^{\frac{n-1}{2}}\Big). \end{array}$

Let us note that in case (iv) (resp. (v)) of the above theorem F can be alternatively described as a smooth divisor of degree (1,1) in $\mathbb{P}^{\frac{n+1}{2}} \times \mathbb{P}^{\frac{n+1}{2}}$ (resp. as the blow-up of \mathbb{P}^n along a linear $\mathbb{P}^{\frac{n-3}{2}}$).

Looking at the above list, we are able to classify fibre-like Fano manifolds with high index.

Dimension	# Vertices	Description	ID
2	6	V_2	2
2	4	$\mathbb{P}^1 \times \mathbb{P}^1$	4
2	3	\mathbb{P}^2	5
3	6	$(\mathbb{P}^1)^3$	21
3	4	\mathbb{P}^3	23
4	10	V_4	63
4	12	$V_2 \times V_2$	100
4	8	$(\mathbb{P}^1)^4$	142
4	6	$\mathbb{P}^2 \times \mathbb{P}^2$	146
4	5	\mathbb{P}^4	147
5	10	$(\mathbb{P}^1)^5$	1003
5	6	\mathbb{P}^5	1013
6	14	V_6	1930
6	12	W_{6}^{3}	5817
6	18	$(V_2)^3$	7568
6	12	$(\mathbb{P}^1)^6$	8611
6	9	$(\mathbb{P}^2)^3$	8631
6	8	$(\mathbb{P}^3)^2$	8634
6	7	\mathbb{P}^6	8635
7	14	$(\mathbb{P}^1)^7$	80835
7	8	\mathbb{P}^7	80891
8	18	V_8	106303
8	15	W_{8}^{3}	277415
8	20	$(V_4)^2$	442179
8	24	$(V_2)^4$	790981
8	12	\tilde{W}	830429
8	16	$(\mathbb{P}^1)^8$	830635
8	12	$(\mathbb{P}^2)^4$	830767
8	10	$(\mathbb{P}^4)^2$	830782
8	9	\mathbb{P}^8	830783

TABLE 1. Toric Fano manifolds of dimension at most 8 that are fibre-like (the entry of the last column is the ID number in the Graded Ring Database $[BK^+]$).

Proposition 2.2. Let F be a fibre-like Fano manifold of dimension $n \ge 4$, index $i_F \ge (n+1)/2$ and $\rho(F) > 1$. If n is even then F is isomorphic to $F \simeq \mathbb{P}^{\frac{n}{2}} \times \mathbb{P}^{\frac{n}{2}}$.

If n is odd then F is isomorphic to $F \simeq \mathbb{P}\left(T_{\mathbb{P}^{\frac{n+1}{2}}}\right)$.

Proof. Use Corollary 0.6 to show that cases (iii) and (v) are not fibre-like: (iii) is clear, while (v) comes with a divisorial contraction to \mathbb{P}^n and a fibration to $\mathbb{P}^{\frac{n+1}{2}}$. Fibre-likeness of (ii) is a consequence of Theorem 0.4, where $G = \mathbb{Z}/2\mathbb{Z}$ exchanges the two factors, and case (iv) follows by Corollary 0.5.

In [KO73] Kobayashi and Ochai proved that $i_F \leq n+1$, where $n = \dim X$ and equality holds if and only if $F \cong \mathbb{P}^n$. They also showed that $i_F = n$ if and only if F is a quadric hypersurface. Fano manifolds with index n-1 are called *del Pezzo manifolds* and they have been classified by Fujita [Fuj82a] and [Fuj82b], while Fano manifolds with index n-2 are called *Mukai manifolds* and their classification appeared in [Muk89]. In dimension $n \geq 3$, del Pezzo manifolds have index $i_F \geq (n+1)/2$ and they have already been studied in Proposition 2.2. For $n \geq 5$, also Mukai manifolds are included in Wiśniewski's list, so we only need to study the case n = 4, $i_F = 2$ (see [Sha99, Table 12.7] for the complete list).

Proposition 2.3. Let F be a 4-dimensional fibre-like Fano manifold of index $i_F = 2$ and $\rho(F) > 1$. Then F is isomorphic to one of the following:

- (i) a double cover of $\mathbb{P}^2 \times \mathbb{P}^2$ branched along a degree-(2,2) divisor;
- (ii) an intersection of two degree-(1,1) divisors in $\mathbb{P}^3 \times \mathbb{P}^3$;
- (*iii*) $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$.

Proof. We follow the enumeration in [Sha99, Table 12.7] of the 18 families. Applying Corollary 0.6 to the cases

$$(1), (2), (3), (5), (6), (8), (9), (14), (16) and (17)$$

we immediately see that they are not fibre-like. Cases

(10) and (12)

are also not fibre-like, since they are obtained as a blow-up of Q^4 but come with a 2-dimensional fibration over \mathbb{P}^{n-2} . Cases

$$(13)$$
 and (15)

are \mathbb{P}^1 -bundles over \mathbb{P}^3 or Q^3 and the other ray of the nef cone corresponds to the contraction of the section. The case which requires more care is (11), in which case F is isomorphic to the projectivisation of the null-correlation bundle over \mathbb{P}^3 . Although the two extremal rays of the nef cone of F both yield fibrations, the image of the fibration associated to the ray not inducing the bundle structure is the quadric Q^3 , see [SW90, Proposition 3.4]. Hence, F is not fibre-like.

We now prove fibre-likeness for the remaining cases. Case (18) (corresponding to case (*iii*) in our list), has an action of S_4 and is clearly fibre-like because of Theorem 0.4. For Case (7) (corresponding to case (*ii*)), we can directly apply Corollary 0.5. Let us analyse now the case (4) (corresponding to case (*i*) in our list): it is obtained as a member of the linear system $|2H_1 + 2H_2|$ in the toric variety Z with weight data

x_0	x_1	x_2	y_0	y_1	y_2	z	
1	1	1	0	0	0	1	H_1 .
0	0	0	1	1	1	1	H_2

Since Z comes with a $\mathbb{Z}/2\mathbb{Z}$ -action exchanging the divisors H_1 and H_2 we have $\dim \operatorname{NS}(Z)_{\mathbb{Q}}^{\mathbb{Z}/2\mathbb{Z}} = 1$ and we can apply [CFST16, Theorem 4.5] to conclude that case (i) is fibre-like.

3. FANO MANIFOLDS WITH HIGH PICARD NUMBER

Fano manifolds of a given dimension form a bounded family, so their Picard number is bounded. In spite of this boundeness result, their classification in dimension at least four is an open and rather difficult problem. A first step towards such classification would be to identify an effective bound on the Picard number of those Fano manifolds that are not a product of lower dimensional manifolds. Already this simpler problem is actually quite difficult. So far, the only known examples of families of Fano manifolds of dimension $n \ge 4$, which are not product and have Picard number at least n + 4 are

- (i) a birational model of the blow up of \mathbb{P}^n in n+3 points in general position, with $n \ge 4$ and even;
- (ii) a birational model of the blow up of \mathbb{P}^4 in 8 points in general position.

The first family appears in any even dimension, the second example is sporadic. In this section we are going to show that all these examples are fibre-like. The first family with n odd gives non-Q-factorial Fano varieties of Picard rank 1, see [AC17, page 3029]. We do not know if there exists a general connection between having high Picard number and being fibre-like.

The sporadic example is discussed in [CCF17]. There the authors prove, following [Muk05], that the Fano manifold under investigation is isomorphic to the moduli space of rank two vector bundles on a del Pezzo surface of degree one. Varying the stability conditions, the authors can explicitly describe the birational geometry of the Fano manifolds. Thanks to this analysis, it is possible to describe the automorphism group of the manifold, and to show the following result.

Theorem 3.1. [CCF17, Proposition 6.22] The Fano model of \mathbb{P}^4 blown-up in 8 points in general position is fibre-like.

We now focus on the first example. We use the results of [AC17] and [Rei72, Section 3]. Take an even integer $n = 2m \ge 2$ and consider a smooth complete intersection Z of two quadrics in \mathbb{P}^{n+2} . Let $\mathcal{F}_{m-1} = \mathcal{F}_{m-1}(Z)$ be the variety of (m-1)-planes in Z. This is a smooth Fano variety of dimension n. It can be seen as a higher dimensional generalisation of the quartic del Pezzo surface and has been extensively studied in the recent work [AC17].

The geometry of \mathcal{F}_{m-1} can be studied from another point of view, which we briefly recall here (see the survey [Cas17] for the notation about Mori Dream Spaces). Let $X := X_r^n$ be the blow up of \mathbb{P}^n at r points in general position. Then it follows from [Muk05] and [CT06, Theorem 1.3] that X is a Mori Dream Space if and only if n and r verify the inequality

(7)
$$\frac{1}{n+1} + \frac{1}{r-n-1} > \frac{1}{2},$$

cf. [Cas17, Example 3.6].

The manifolds appearing in Theorem 3.1 are, in this notation, obtained as Fano models of X_8^4 .

Look at X_{n+3}^n , with $n \ge 2$ even. For this class, inequality (7) holds, so X_{n+3}^n is a Mori Dream Space and we can consider its Fano model F_{n+3}^n . Bauer in [Bau91] proved that X_{n+3}^n and \mathcal{F}_{m-1} are isomorphic in codimension one – see also [AC17, Theorem 1.4]. By [AC17, Remark 4.10] it follows that \mathcal{F}_{m-1} is actually isomorphic to F_{n+3}^n . In [AC17, Proposition 7.1] the authors describe the automorphism groups of \mathcal{F}_{m-1} showing that

$$\left(\left(\mathbb{Z}/2\mathbb{Z}\right)^{n+2}\subseteq\right)\operatorname{Aut}(\mathcal{F}_{m-1})\subseteq W(D_{n+3})\left(=\left(\mathbb{Z}/2\mathbb{Z}\right)^{n+2}\rtimes S_{n+2}\right),$$

where $W(D_{n+3})$ is the Weyl group of automorphism of a D_{n+3} -lattice. The inclusion $(\mathbb{Z}/2\mathbb{Z})^{n+2} \subset \operatorname{Aut}(\mathcal{F}_{m-1})$ is an actual equality for a general choice of \mathcal{F}_{m-1} . The action of $(\mathbb{Z}/2\mathbb{Z})^{n+2}$ can be described by presenting Z as the locus

$$\sum_{i=0}^{n+3} x_i^2 = \sum_{i=0}^{n+3} \lambda_i x_i^2 = 0.$$

Then the group acts by changing the signs of the coordinates. We can use this to prove the following.

Theorem 3.2. Let $n = 2m \ge 4$ be an integer. Then the smooth n-dimensional Fano variety $\mathcal{F}_{m-1}(Z)$ of (m-1)-planes in the intersection of two quadrics $Z \subset \mathbb{P}^{n+2}$ is fibre-like.

Proof. Consider the isomorphism $H^2(\mathcal{F}_{m-1},\mathbb{Z}) \simeq \mathrm{NS}(\mathcal{F}_{m-1})$. The action of $G := (\mathbb{Z}/2\mathbb{Z})^{n+2}$ via pseudo-automorphisms of X_{n+3}^n is explicitly described in [Dol04, Sections 4.4-4.6] (see also [AC17, Remark 7.2]). Let $x_0, \ldots, x_{n+2} \in \mathbb{P}^n$ be blown-up points; we can assume that the first n+1 are the coordinate points and

- $x_{n+1} = [1 : \ldots : 1];$
- $x_{n+2} = [c_0 : \ldots : c_n].$

The pseudo-automorphism $\phi_{n+2,n+3} \colon X_{n+3}^n \to X_{n+3}^n$ is defined on \mathbb{P}^n as $\rho \circ \iota$, where $\iota \colon \mathbb{P}^n \dashrightarrow \mathbb{P}^n$ is the standard Cremona involution and ρ is the diagonal projective transformation $[t_0 \colon \ldots \colon t_n] \mapsto [c_0 t_0 \colon \ldots \colon c_n t_n]$. Analogously, one defines $\phi_{i,j}$ with i < j, which exchanges the exceptional divisors of X_{n+3}^n and fixes $K_{X_{n+3}^n}$. This implies that $\mathrm{NS}(X_{n+3}^n)_{\mathbb{Q}}^G = \mathbb{Q}K_{X_{n+3}^n}$. Moreover, the analysis in [AC17, Proposition 5.4] implies that $\mathrm{NS}(X_{n+3}^n)^G \simeq \mathrm{NS}(\mathcal{F}_{m-1})^G$. So we apply Theorem 0.4 to conclude. \Box

4. Open questions

We conclude this note with some questions regarding fibre-like varieties that are still open.

Since all the examples of smooth toric fibre-like Fano varieties are vertex transitive, we ask the following:

Question 4.1. Is any fibre-like toric Fano manifold vertex-transitive?

Answering this question affermatively would give a complete classification of fibre-like toric Fano manifolds.

On a different note, the study of fibre-like Fano varieties in positive characteristic seems to be still very far from being satisfactory.

16

After the recent developments for the MMP in positive characteristic for threefods (cf. [HX15], [Bir16], [BW17]), the picture that we delineated in the Introduction holds almost in the same way in characteristic > 5 as long as we only focus in dimension 2 and 3. Hence, it is natural to try to extend the results in [CFST16] to positive characteristic. If k is any algebraically closed field and F is a smooth Fano variety over k, the definition of fibre-likeness still makes sense and one can in particular ask the following:

Question 4.2. If char k = p > 0, are there sufficient or necessary conditions that determine whether a smooth Fano F is fibre-like?

At present time, the situation appears to be quite obscure: we do not even know if a del Pezzo surface of degree 8 is fibre-like in positive characteristic. The approach outlined in [CFST16] relaying on the study of a suitable monodromy action on the Néron-Severi does not generalize directly to this case.

References

- [AC17] Carolina Araujo and Cinzia Casagrande. On the Fano variety of linear spaces contained in two odd-dimensional quadrics. *Geom. Topol.*, 21(5):3009–3045, 2017.
- [Bat91] Victor V. Batyrev. On the classification of smooth projective toric varieties. Tohoku Math. J. (2), 43(4):569–585, 1991.
- [Bau91] Stefan Bauer. Parabolic bundles, elliptic surfaces and SU(2)-representation spaces of genus zero Fuchsian groups. Math. Ann., 290(3):509–526, 1991.
- [BCHM10] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James McKernan. Existence of minimal models for varieties of log general type. J. Amer. Math. Soc., 23(2):405– 468, 2010.
- [Bir16] Caucher Birkar. Existence of flips and minimal models for 3-folds in char p. Ann. Sci. Éc. Norm. Supér. (4), 49(1):169–212, 2016.
- [BK⁺] Gavin Brown, Alexander M. Kasprzyk, et al. Graded ring database. Online. Available at http://grdb.lboro.ac.uk.
- [BW17] Caucher Birkar and Joe Waldron. Existence of Mori fibre spaces for 3-folds in char p. Adv. Math., 313:62–101, 2017.
- [Cas03] Cinzia Casagrande. Contractible classes in toric varieties. Math. Z., 243(1):99–126, 2003.
- [Cas06] Cinzia Casagrande. The number of vertices of a Fano polytope. Ann. Inst. Fourier (Grenoble), 56(1):121–130, 2006.
- [Cas17] Ana-Maria Castravet. Mori Dream Spaces and Blow-Ups. ArXiv e-prints, to appear on the Proceedings of the AMS Summer Institute in Algebraic Geometry (University of Utah, July 2015), January 2017.
- [CCF17] Cinzia Casagrande, Giulio Codogni, and Andrea Fanelli. The blow-up of \mathbb{P}^4 at 8 points and its Fano model, via vector bundles on a degree 1 del Pezzo surface. ArXiv e-prints, July 2017.
- [CFST16] Giulio Codogni, Andrea Fanelli, Roberto Svaldi, and Luca Tasin. Fano varieties in Mori fibre spaces. Int. Math. Res. Not. IMRN, (7):2026–2067, 2016.
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck. Toric varieties, volume 124 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2011.
- [CT06] Ana-Maria Castravet and Jenia Tevelev. Hilbert's 14th problem and Cox rings. Compos. Math., 142(6):1479–1498, 2006.
- [Dol04] Igor V. Dolgachev. On certain families of elliptic curves in projective space. Ann. Mat. Pura Appl., 183:317–331, 2004.
- [Fuj82a] Takao Fujita. Classification of projective varieties of Δ-genus one. Proc. Japan Acad. Ser. A Math. Sci., 58(3):113–116, 1982.
- [Fuj82b] Takao Fujita. On polarized varieties of small Δ-genera. Tohoku Math. J. (2), 34(3):319– 341, 1982.
- [GH16] Ian Gleason and Isabel Hubard. Products of abstract polytopes. ArXiv e-prints, March 2016.

- [GO04] Jacob E. Goodman and Joseph O'Rourke, editors. Handbook of discrete and computational geometry. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, second edition, 2004.
- [HX15] Christopher D. Hacon and Chenyang Xu. On the three dimensional minimal model program in positive characteristic. J. Amer. Math. Soc., 28:711–744, 2015.
- [Kly84] A. A. Klyachko. Demazure models for a special class of tori. Selecta Math. Soviet., 3(1):57– 61, 1983/84. Selected translations.
- [KM98] János Kollár and Shigefumi Mori. Birational geometry of algebraic varieties, volume 134 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [KO73] Shoshichi Kobayashi and Takushiro Ochiai. Characterizations of complex projective spaces and hyperquadrics. J. Math. Kyoto Univ., 13:31–47, 1973.
- [Kol13] János Kollár. Singularities of the minimal model program, volume 200 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2013. With a collaboration of Sándor Kovács.
- [Mor82] Shigefumi Mori. Threefolds whose canonical bundles are not numerically effective. Ann. of Math. (2), 116(1):133–176, 1982.
- [Muk89] Shigeru Mukai. Birational classification of Fano 3-folds and Fano manifolds of coindex 3. Proc. Nat. Acad. Sci. U.S.A., 86(9):3000–3002, 1989.
- [Muk05] Shigeru Mukai. Finite generation of the Nagata invariant rings in A–D–E cases. RIMS preprint, Kyoto, (1502), 2005.
- [Øb07] Mikkel Øbro. An algorithm for the classification of smooth Fano polytopes. ArXiv e-prints, April 2007.
- [Rei72] Miles Reid. The complete intersection of two or more quadrics. Ph.D. thesis, 1972.
- [Rei83] Miles Reid. Decomposition of toric morphisms. In Arithmetic and geometry, Vol. II, volume 36 of Progr. Math., pages 395–418. Birkhäuser Boston, Boston, MA, 1983.
- [Sha99] I. R. Shafarevich, editor. Algebraic geometry. V, volume 47 of Encyclopaedia of Mathematical Sciences. Springer-Verlag, Berlin, 1999. Fano varieties, A translation of it Algebraic geometry. 5 (Russian), Ross. Akad. Nauk, Vseross. Inst. Nauchn. i Tekhn. Inform., Moscow, Translation edited by A. N. Parshin and I. R. Shafarevich.
- [SW90] Michał Szurek and Jarosław A. Wiśniewski. Fano bundles over \mathbf{P}^3 and Q_3 . Pacific J. Math., 141(1):197–208, 1990.
- [VK84] V. E. Voskresenskiĭ and A. A. Klyachko. Toric Fano varieties and systems of roots. Izv. Akad. Nauk SSSR Ser. Mat., 48(2):237–263, 1984.
- [Wiś91] Jarosław A. Wiśniewski. On Fano manifolds of large index. Manuscripta Math., 70(2):145–152, 1991.

EPFL, SB MATHGEOM CAG, MA B3 635 (Bâtiment MA), Station 8, CH-1015 Lausanne, Switzerland.

E-mail address: giulio.codogni@epfl.ch

Mathematisches Institut, Heinrich-Heine-Universität Düsseldorf, Universitätsstr. 1, 40204 Düsseldorf, Germany.

E-mail address: fanelli@hhu.de

DPMMS - University of Cambridge Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WB, UK.

E-mail address: rs872@cam.ac.uk

MATHEMATICAL INSTITUTE OF THE UNIVERSITY OF BONN, ENDENICHER ALLEE 60, D-53115 BONN, GERMANY.

E-mail address: tasin@math.uni-bonn.de