

The spontaneous regeneration of myelin: an unexpected central role of the androgen receptor

Bartosz Bielecki^{a,b}, Claudia Mattern^c, Abdel Ghomari^a, Sumaira Javaid^{a,d}, Kaja Smietanka^{a,b}, Charly Abi Ghanem^a, Sakina Mhaouty-Kodja^e, Said Ghandour^f, Etienne-Emile Baulieu^{a,1}, Robin J.M. Franklin^g, Michael Schumacher^{a,1,2}, and Elisabeth Traiffort^{a,2}

^aU1195 Inserm and University Paris-Sud and University Paris-Saclay, Kremlin-Bicêtre, France; ^bDepartment of Neurology and Stroke, Medical University of Lodz, Poland; ^cMattern Foundation, Vaduz, Liechtenstein; ^d H.E.J. Research Institute of Chemistry, ICCBS, University of Karachi, Pakistan;; ^eU1130 Inserm, UMR 8246 CNRS, University Pierre and Marie Curie, Paris, France; ^fUMR 7357 CNRS, University of Strasbourg, France and Department of Anatomy and Neurobiology, VCU, Richmond, Virginia, USA; ^gWellcome Trust-MRC Cambridge Stem Cell Institute and Department of Clinical Neurosciences, University of Cambridge, UK.

Author contributions: BB, MS, and ET designed research; BB, AG, SJ, KS, and CAG performed research; CM, SMK, SG, EEB, and RJMF analyzed and interpreted data; and BB, EEB, RJMF, MS and ET wrote the paper.

¹To whom correspondence may be addressed. E-mail: etienne.baulieu@inserm.fr (EEB) and michael.schumacher@inserm.fr (MS).

²These authors contributed equally to this work.

Classification: Biological Sciences - Neuroscience

Short title: Androgens and Remyelination

Abstract

Lost myelin can be replaced after injury or during demyelinating diseases in a regenerative process called remyelination. In the central nervous system (CNS), the myelin sheaths, which protect axons and allow the fast propagation of electrical impulses, are produced by oligodendrocytes. The abundance and widespread distribution of oligodendrocyte progenitors (OPs) within the adult CNS account for this remarkable regenerative potential. Here we report a key role for the male gonad, testosterone and the androgen receptor (AR) in CNS remyelination. After lysolecithin-induced demyelination of the male mouse ventral spinal cord white matter, the recruitment of glial fibrillary acidic protein-expressing astrocytes was compromised in the absence of testes and testosterone signaling via AR. Concomitantly, the differentiation of OPs into oligodendrocytes forming myelin basic protein (MBP)⁺ and proteolipid protein (PLP)⁺ myelin was impaired. Instead, in the absence of astrocytes, axons were remyelinated by protein zero (P0)⁺ and peripheral myelin protein 22-kDa (PMP22)⁺ myelin, normally only produced by Schwann cells in the peripheral nervous system (PNS). Thus, testosterone favours astrocyte recruitment and spontaneous oligodendrocyte-mediated remyelination. This finding may have important implications for demyelinating diseases, psychiatric disorders and cognitive aging. The testosterone-dependency of CNS oligodendrocyte remyelination may have roots in the evolutionary history of the AR, as the receptor has evolved from an ancestral 3-ketosteroid receptor through gene duplication at the time when myelin appeared in jawed vertebrates.

Key words: Myelin, oligodendrocytes, astrocytes, Schwann cells, testosterone, androgen receptor.

Significance

Myelin allows the rapid conduction of electrical signals and provides axons with metabolic support. It appeared relatively late in evolution in hinge-jawed vertebrates and concomitantly with the emergence of the receptor of androgen hormones (AR), hinting at a specific role for androgens in myelination. Here we show that after demyelination of the central nervous system (CNS), the male gonad, testosterone and AR favor astrocyte recruitment and myelin regeneration by oligodendrocytes. In their absence, astrocytes remain sparse and peripheral-type remyelination, classically associated with Schwann cells, is predominantly detected. These findings reveal a novel role of androgen signaling in CNS myelin formation and glial cell communication, with possible implications for demyelinating, psychiatric and age-related disorders.

\body

Introduction

The remyelination of axons is a complex process, involving interactions between different types of neural cells. It shows a peculiarity that has been observed in both experimental models and demyelinating diseases such as multiple sclerosis (1). Although most of the remyelination is normally accomplished by oligodendrocytes derived from oligodendrocyte progenitors (OPs), a variable proportion of central nervous system (CNS) axons can be remyelinated by cells with the immunophenotypic and ultrastructural characteristics of Schwann cells, the presence of which is normally limited to the peripheral nervous system (PNS). These cells indeed express P0, a sensitive and specific marker of Schwann cells (2). Moreover, analysis by electron microscopy demonstrated that they have the typical morphology of myelinating Schwann cells (1-4). These ultrastructural findings indicate that the expression of P0 by part of the remyelinating cells is not merely ectopic, but that these cells may correspond to authentic Schwann cells. This was further supported by their expression of Schwann cell-specific transcription factors (5).

Importantly, there is strong evidence that large numbers of the Schwann cells present in demyelinating CNS lesions are derived from OPs. The first evidence came from the transplantation of purified OPs into demyelinating CNS lesions, in which endogenous remyelination was prevented by X-irradiation (4, 6, 7). Unequivocal confirmation that most Schwann cells contributing to CNS remyelination are indeed derived from OPs has then been provided by genetic fate-mapping in transgenic mice (5). This plasticity of OPs with a neuroepithelial origin may come as a surprise, because Schwann cells normally arise from the neural crest, and this problem still presents a matter of controversy. However, under the influence of specific morphogenetic factors, neuroepithelial cells can differentiate into neural crest stem cells and give rise to neural cell types of both CNS and PNS (4, 8).

Astrocytes play an important role in determining the balance between oligodendrocyte and Schwann cell remyelination in the CNS. Thus, within a demyelinated lesion, oligodendrocyte remyelination occurs in regions where astrocytes are present, whereas Schwann cell remyelination preferentially occurs in the absence of astrocytes (7, 9, 10). Moreover, reducing astrocyte activation by a conditional genetic deletion promoted Schwann cell remyelination (1).

Like other members of the steroid receptor family, the AR functions as a ligand-activated transcription factor and also directly interacts with components of extranuclear signaling pathways (11, 12). Here, by investigating the role of AR signaling in the spontaneous regeneration of myelin within the adult CNS, we show that the recruitment of glial fibrillary acidic protein (GFAP)⁺ astrocytes into a demyelinated area of the ventral funiculus of male mice is compromised in the absence of testes, testosterone or AR. Consistent with a key role of astrocytes in CNS remyelination, in their absence the newly formed myelin was P0⁺ and PMP22⁺, most likely reflecting Schwann cell remyelination.

Results

The Role of Testes and Testosterone in the Recruitment of Astrocytes and Spontaneous Remyelination

We used an experimental model in which acute demyelination is followed by spontaneous remyelination. Axons of the right ventrolateral white matter tract of the adult male mouse spinal cord (ventral funiculus) were locally demyelinated by stereotaxic microinjection of the demyelinating toxin lysolecithin. Within 3 days, the injected area was depleted of astrocytes, oligodendrocytes and MBP, a major component and established marker of CNS myelin (13).

In gonadally intact males, the area of demyelination was rapidly replenished by GFAP⁺ astrocytes (Fig. 1A, left). Concomitantly, axons were remyelinated by MBP⁺ CNS myelin (Fig. 1A, right). However, after surgical castration, the recruitment of astrocytes was severely impaired (Fig. 1B, left), and axons were preferentially remyelinated by P0⁺ PNS-like myelin (Fig. 1B, right). P0 immunostaining appeared sparse, consistent with previous results showing that Schwann cell remyelinated areas contain more demyelinated axons than oligodendrocyte remyelinated ones (1). Similar results were obtained when tissue sections were analyzed with PMP22 antibody, another specific marker of peripheral myelin (14). Immunostaining of MBP remained very sparse within the lesion, and perfectly matched immunostaining of PLP, another major CNS myelin protein (Fig. S1). Despite the presence of low levels of MBP isoforms in peripheral myelin (14), these are likely not detected since the monoclonal anti-MBP antibodies used in the present study only weakly stain PNS myelin.

We then showed that testosterone was the testicular factor necessary for the recruitment of astrocytes and the remyelination of axons by oligodendrocytes. When castrated males received for 4 weeks a subcutaneous Silastic implant filled with testosterone, GFAP⁺ astrocytes and MBP immunostaining were fully restored within the lesion area (Fig. 1C), as in uncastrated males (Fig. 1A). The implant produced physiological plasma and brain levels of testosterone (10 to 15 nM) as determined by gas chromatography/mass spectrometry (15).

The role of testosterone treatment in the concomitant replenishment of astrocytes and CNS remyelination is also illustrated in Fig. S2. Although axons were not counted, the staining of large-caliber axons on the sagittal spinal cord sections with an antibody against neurofilament 200 kD (NF-200) revealed neither a substantial loss of axons after lysolecithin-induced demyelination, nor a marked increase in axonal density by testosterone treatment (Fig. S3). These observations point to a crucial role of the male gonads and testosterone in the recruitment of astrocytes and in the balance between CNS and PNS-like remyelination.

The Androgen Receptor is Required for the Recruitment of Astrocytes and the Spontaneous Regeneration of CNS Myelin

In the CNS, testosterone exerts its effects either after binding to intracellular AR or via its metabolites estradiol, 3 α - or 3 β -androstenediol (Fig. S4). Testosterone signaling via its conversion to estradiol may correspond to a phylogenetically ancient mechanism, as the ancestral intracellular steroid receptor was a receptor for estrogens (16). In addition, signaling of steroid metabolites via membrane proteins may have been an early evolutionary feature (17). This led us to investigate whether the testosterone-dependent recruitment of astrocytes and spontaneous regeneration of MBP⁺ myelin involved AR. We first used AR^{Tfm} mice, which display a frame shift mutation in exon 1 of the AR gene producing a non-functional receptor (18). Despite testosterone supplementation during 4 weeks, the area of demyelination remained largely depleted of astrocytes (Fig. 1D, left) in AR^{Tfm} mice, and axons were mainly remyelinated by P0⁺ (Fig. 1D, right) and PMP22⁺ myelin.

We then tested whether AR expression within the CNS is necessary for the stimulation of remyelination by testosterone. Transgenic AR^{NesCre} mice, displaying selective ablation of the AR in neurons, astrocytes and oligodendrocytes, but with microglial cells being spared (19), were castrated and received a testosterone-filled subcutaneous Silastic implant during 4 weeks following lysolecithin-induced demyelination. As in AR^{Tfm} mice and in contrast to uncastrated wild-type animals, testosterone therapy failed to restore astrocytes and MBP-immunoreactive myelin within the lesion. Instead, PNS-type P0⁺ remyelination prevailed (Fig. 1E). Thus, testosterone signaling via AR stimulates the recovery of astrocytes and shifts the balance towards the regeneration of CNS myelin.

Testosterone Increased the Number of GFAP⁺ Astrocytes at the Border of the Demyelinated Area and in Glial Cell Culture

In the vicinity of a demyelinating lesion, new astrocytes are mainly generated from preexisting ones expressing the fibroblast growth factor receptor 3 (5). Therefore, we investigated the effects of testosterone on the rapid astrocytic response after lysolecithin administration. At the top border of the lysolecithin-demyelinated area, the GFAP-immunoreactive surface was slightly increased in castrated males as early as 4 days after lysolecithin injection. However, the extent of GFAP labeling was markedly enhanced following testosterone treatment (Fig. 2 A,B). At 14 days post-lysolecithin, the demyelinated lesion was nearly entirely occupied by GFAP⁺ astrocytes, whereas the remyelination by oligodendrocytes was still incomplete (1), indicating that astrocyte proliferation and recruitment precede the formation of new MBP⁺ myelin.

In mixed cultures of glial cells prepared from postnatal day 1 to 3 male mice, the number of reactive GFAP⁺ astrocytes was significantly increased 72 hours after a 12-hour exposure to lysolecithin. When the culture medium was supplemented with testosterone (1 μ M) for 72 hours after lysolecithin removal, a further increase in GFAP immunostaining was observed (Fig. 2 C,D). Thus, testosterone increased the astrocytic response both *in vivo* at the border of the lesion and under *in vitro* conditions in mixed cultures of glial cells.

The Regeneration of MBP-Immunoreactive Myelin is Accompanied by the Recovery of Mature Oligodendrocytes

We confirmed that the spontaneous regeneration of MBP-immunoreactive CNS myelin 4 weeks after a lysolecithin-induced demyelination was also dependent on the presence of testes, testosterone and the AR. Indeed, in castrated males and in the absence of a functional AR (AR^{Tfm} and AR^{NesCre} mice), the recovery of MBP⁺ myelin was markedly

impaired (Fig. 3). Importantly, the regeneration of MBP⁺ myelin was always accompanied by full replenishment of the demyelinated lesion by oligodendrocytes expressing carbonic anhydrase II (CA II). In contrast, few CA II⁺ oligodendrocytes were observed after castration or in the absence of AR (Fig. 3). Thus, testosterone-dependent CNS remyelination goes together with the parallel recruitment of both oligodendrocytes and astrocytes.

In the Presence of Astrocytes, Testosterone Increases the Proliferation and Differentiation of Oligodendrocyte Progenitors

Following oligodendrocyte loss, CNS remyelination requires the recruitment of OPs and their differentiation into mature oligodendrocytes (20). To provide further insight into the mechanisms by which testosterone promotes CNS myelin regeneration in the presence of astrocytes, we first investigated its effect on neonatal OPs in mixed primary glial cell cultures prepared from the brains of postnatal day 1 to 3 transgenic male mice selectively expressing the enhanced green fluorescent protein (EGFP) in the oligodendrocyte lineage under the control of the mouse *Plp* gene promoter (*Plp*-EGFP mice) (21). Confluent cultures, composed of OPs and oligodendrocytes (green fluorescent) and astrocytes, were exposed for 12 hours to lysolecithin damage and were then treated with vehicle or testosterone. Three days later, cells were fixed and immunostained with an antibody against the chondroitin sulfate proteoglycan (NG2), a proven marker of OPs (Fig. S5A-D).

The number of EGFP⁺ cells was reduced by half after exposure to lysolecithin and remained low in the absence of testosterone. Culture medium supplementation with testosterone (1 μ M) significantly increased the number of both EGFP⁺ and NG2⁺ cells, which exhibited a differentiated branched morphology (Fig. S5A-D). Thus, adding testosterone to mixed glial cell cultures increases proliferation and differentiation of neonatal OPs.

Compared with their neonatal counterparts, adult OPs are more quiescent and show other distinct features (22, 23). We thus also examined the impact of testosterone on Olig2-expressing cells in adult castrated male mice. This transcription factor is confined to the CNS and is critically involved in OPs differentiation (24). Four weeks after injection of lysolecithin into the myelinated ventrolateral white matter of the castrated male spinal cord, Olig2⁺ cells were significantly more abundant in males receiving a testosterone-filled Silastic implant when compared to males receiving an empty implant (Fig. S5E). Testosterone replacement indeed significantly increased the number of Olig2⁺ cells, which even reached a higher density when compared to unlesioned control animals. Moreover, double immunolabeling with proliferating cell nuclear antigen (PCNA) showed that testosterone stimulated OPs proliferation.

Discussion

Our data establish the importance of the male gonad, testosterone and the AR in the efficient recruitment of astrocytes into a demyelinated lesion, with implications for the balance of oligodendrocyte versus peripheral-type remyelination. In gonadally intact male mice or in castrated animals treated with testosterone, the area of lysolecithin-induced demyelination was replenished with astrocytes after 4 weeks. Consequently, axons were mainly remyelinated by oligodendrocytes. On the contrary, in the absence of testosterone or a functional AR, astrocytes remained sparse and P0⁺ myelin was preferentially detected within the lesion. This observation is consistent with previous studies showing that Schwann cell remyelination occurs in areas of the CNS where astrocytes are absent (9, 10).

Testosterone treatment has recently been shown to stimulate the regeneration of myelin in a mouse model of severe chronic demyelination, where no spontaneous remyelination was observed. Moreover, the strong remyelinating effect of testosterone was

AR-dependent (15). These findings already qualified the brain AR as a promising drug target for remyelination therapy. Nevertheless, it came as a surprise in the present study that spontaneous remyelination by oligodendrocytes was markedly inhibited in the absence of testes, testosterone or AR.

Encoded by a single gene, AR only arose lately in jawed vertebrates, after the cyclostome-gnathostome divergence (25, 26). This has allowed androgens to acquire new signaling functions related to reproduction and sexual differentiation (27, 28). In parallel, testosterone and its metabolite 5 α -dihydrotestosterone (5 α -DHT), both ligands of the AR, may also have acquired new functions in the CNS, as suggested by the widespread distribution of AR within the brain and spinal cord (29, 30).

Intriguingly, the evolutionary acquisition of AR was concomitant with the advent of the myelin sheath, constituting one of the most recent structural innovations of the vertebrate nervous system (Fig. 4). Like the AR, myelin first appeared in vertebrates with a hinged jaw, most likely in placoderms, facilitating predatory and escape behaviors and improving information processing by increasing the conduction speed of axons (31). Both myelin and AR are present in extant cartilaginous fishes (sharks, rays), but absent in jawless fishes (hagfishes and lampreys) (28, 32). The parallel evolution of AR and myelin, together with the widespread distribution of AR in the CNS, suggest that androgen signaling may have become involved in the regulation of myelin formation and play a role in the regenerative capacity of myelin, which is preserved in the adult CNS.

The recruitment of astrocytes is likely to play a key role in the remyelinating effect of testosterone, as their presence was associated with oligodendrocyte remyelination, whereas in their absence, P0⁺/PMP22⁺ peripheral-type remyelination was observed. How astrocytes participate in the remyelinating actions of testosterone remains to be clarified. The quality of presently available AR antibodies unfortunately did not allow us to study the cellular

distribution of the receptor by dual labelling. In previous studies, AR-immunoreactivity has been mainly localized to neurons and reactive microglia and to subpopulations of astrocytes and oligodendrocytes (30, 33). In our conditional AR^{NesCre} mice, the AR was selectively inactivated in neurons, astrocytes and oligodendrocytes, whereas microglial cells were spared (19). Thus, the effect of testosterone observed here does not involve microglial AR. We gave advantage to the use of the Nestin-Cre mouse strain for two major reasons. First, the high efficiency of the Cre-mediated recombination well characterized for this strain results in the almost complete ablation of the floxed genes in neurons and macroglial cells (15, 19, 34). Then, this choice allowed us to analyze the role of testosterone without any preconceived idea about its neural targets. Nevertheless, cell-specific conditional AR knockout will be further needed to identify the direct target cells of androgens. It is conceivable that an androgen-dependent neuronal signal may attract astrocytes to the demyelinating lesion or alternatively testosterone may directly act on astrocytes or oligodendrocytes.

Under the present experimental conditions, the magnitude of the effects of castration, testosterone therapy and AR inactivation on the replenishment of astrocytes and on the balance of oligodendrocyte versus peripheral-type remyelination was noticeable. However, after a demyelinating lesion induced by the injection of ethidium bromide into the caudal cerebellar peduncle of male or female rats, remyelination by oligodendrocytes or Schwann cells was previously reported not to be affected by gonadectomy (35). The use of different species and different experimental conditions may explain this apparent discrepancy. Alternatively, the effect of castration may also have been masked by the local synthesis of androgens in the brain. Thus, an exercise-dependent increase in androgen biosynthesis within the hippocampus has been shown to stimulate neurogenesis (36). Other "neurosteroids" synthesized within the CNS, such as progesterone, may also positively influence the regeneration of CNS myelin (37, 38). However, in our system, the effect of AR signaling was

very robust and AR inactivation was not compensated by other pro-myelinating signaling mechanisms.

The multipotency of OPs and their capacity to differentiate into Schwann cells remain amazing (5). How the male gonad-dependent recruitment of astrocytes and its influence on the balance of central- versus peripheral-type remyelination affect functional recovery after a demyelinating event remains to be explored. Moreover, the presence of testes has been shown also to influence the thickness of the myelin sheaths as well as the number and turnover of oligodendrocytes, which therefore differ between sexes (39).

Neuroprotective and immunosuppressive actions of testosterone have previously been documented in experimental autoimmune encephalomyelitis (EAE), a widely used animal model of MS (40). These studies have provided the support for a small clinical pilot study, suggesting that prolonged testosterone therapy of men with MS may have neuroprotective and anti-inflammatory effects and also improve cognitive performance (41). The present study may also gain significance from clinical observations showing that: 1) the incidence of MS and the relapse rate throughout the disease are lower in men (42); 2) low testosterone levels are associated with an increased risk of developing MS (43); 3) levels of testosterone are decreased in men with MS and reduced levels of testosterone have been associated with disability (44, 45); 4) men reach disability milestones more rapidly than women (46).

Testosterone levels progressively decrease with age, but the hormonal decline occurs more rapidly in as many as 20% of aging men (47). Whether age-dependent changes in testosterone are a contributing factor to age-related white matter abnormalities (48) or the reduced regenerative capacity of myelin (49) remains to be explored. Importantly, the influence of androgens on the plasticity and regeneration of myelin may not only have consequence for demyelinating diseases, as recent studies have linked myelin abnormalities to a wide range of psychiatric disorders and cognitive aging (50, 51).

Material and Methods

Mice. Wild-type gonadally intact or castrated C57/Bl6 male mice were purchased at the age of 8 to 12 weeks from Janvier Breeding Center (France). AR^{NesCre} male mice were obtained by crossing floxed AR mice (AR^{fl/Y}) (52) with transgenic mice expressing the Cre recombinase driven by the promoter and the CNS-specific enhancer of rat nestin (Nes), which is selectively active in neuronal and glial precursor cells (34). Both strains were on a C57/BL6 background. The AR^{NesCre} mice have been extensively characterized (19) and were castrated at 2–3 months of age under ketamine/xylazine anesthesia. AR^{Tfm} mice were obtained from the French Atomic Energy Commission (CEA, Dr. René Habert) (18). For *in vitro* experiments, *Plp*-EGFP mice were obtained from Dr. Wendy Macklin (University of Colorado, USA) (21). All animals were housed in standard conditions: 12 hours light-dark cycle with food and water *ad libitum*. All procedures were performed according to the European Communities Council Directive (86/806/EEC) for the care and use of laboratory animals.

Surgical Procedures. Lysolecithin (Sigma-Aldrich) was microinjected into right ventral funiculus of the male mouse spinal cord under ketamine/xylazine anesthesia. After shaving the skin at thoracic level T5, a midline incision was made caudally in order to show thoracolumbar fascia and musculature. Two incisions were performed on the fascia and external edges of dorsal muscles on both sides of the vertebral column that allowed immobilization of the vertebral column on the stereotaxic device (Kopf Instrument). The space between T12-T13 was identified and carefully dissected until the spinal cord was exposed. The central vein was localized and the dura mater was carefully pierced using a 32G dental needle. The injection was performed with a fine tip glass capillary, elongated until reaching a diameter not exceeding 50 μm (Harvard Apparatus) and mounted on a stereotaxic device after connection to a Hamilton syringe with a plastic tube. By using an infusion pump,

1 μ l of 1% lysolecithin solution was unilaterally injected. After suturing the muscles, a Silastic® tube either empty (control) or filled with testosterone (Fluka) was subcutaneously implanted.

Mixed Glial Cell Cultures. Male *Plp*-EGFP mouse pups were used between postnatal day 0 (P0) and P3 (53). Briefly, immediately after decapitation, the brain was dissected and placed into Dulbecco's Modified Eagle Medium (DMEM) supplemented with antibiotics. After the removal of meninges, the brain including the hemispheres, cerebellum and cerebral trunk was placed in fresh DMEM supplemented with antibiotics and 10% fetal bovine serum (FBS) and then passed through a sterile 100 μ m-nylon mesh. A total of 8 ml of cell suspension was transferred to poly-L-Lysine-coated 100 mm Petri dish. The cells were allowed to grow in DMEM supplemented with 10% FBS and antibiotics at 37 °C in humidified 5% CO₂. The medium was changed after 4 days and later twice a week. Cells were used for experiments after 14 days in culture. Prior to the experimental procedure, cells were transferred to a 24-well plate containing sterilized and poly-L-lysine-coated glass coverslips.

***In vitro* Induction of Oligodendrocyte Damage and Hormonal Treatment.** Images of the mixed glial cell cultures were taken under a fluorescent microscope prior to any treatment and then daily to check viability and morphology of the cells. Lysolecithin (0.1 mg/ml for 12 hrs) was used for inducing cell damage. After lysolecithin removal, testosterone (1 μ M) or vehicle alone (0.1% ethanol) was added to the culture medium. Immunohistochemical analysis was carried out after 4% paraformaldehyde fixation of the cells for 10 min, PBS washing, incubation for 1 hr in 10% normal goat serum (Vector) and overnight incubation with the primary antibodies at 4 °C.

Histological Procedures, Antibodies Used, Image Acquisition and Analysis are specified in *SI Materials and Methods*.

Statistical Analysis. Statistical analyses was performed using the Statistica 12 software (Statsoft Inc.). All data were given as arithmetic means \pm standard error of the mean (S.E.M.). For the *in vivo* experiments, each animal group comprised at least 5 mice. For the *in vitro* experiments, the groups consisted of at least 3 wells (average values obtained from 3 fields of vision per well). Comparisons of means were done by using one-way ANOVA followed by LSD post-hoc test.

Acknowledgments

We thank Wendy B. Macklin (University of Colorado) for sharing *Plp*-EGFP mice and René Habert for providing Tfm mice (University Paris-Diderot and CEA). B.B. was successively supported by the European Leukodystrophy Association (ELA) Foundation (France) and by the Mattern Foundation. The fellowship of S.J. was funded by the Higher Education Commission (HEC) of Pakistan and French Embassy in Pakistan. This work was supported by grants of ELA, the French Multiple Sclerosis Foundation (ARSEP) and The UK Multiple Sclerosis Society.

Figure Legends

Figure 1. Testes, testosterone and AR are required for the recruitment of astrocytes into a demyelinating lesion, thus determining the balance of central versus peripheral types of remyelination. (A-E) (Left panels) Visualization of astrocytes stained with an antibody to GFAP (red). (Right panels) Immunohistochemical staining of MBP⁺ CNS myelin (red) and P0⁺ PNS myelin (green) 4 weeks after lysolecithin-induced demyelination. Cell nuclei were blue-counterstained with DAPI. Dotted lines delineate the boundary between the ventral white matter (VWM) and top grey matter. (A) Recruitment of astrocytes and recovery of MBP⁺ CNS myelin in the lesion area of gonadally intact males. Scale bars: 100 μ m. (B) After testes removal and treatment with an empty implant (+V), astrocytes remained sparse within the remyelinated area and remyelination was mediated by Schwann cells (P0⁺ myelin). (C) Recruitment of GFAP⁺ astrocytes and regeneration of MBP⁺ myelin in castrated males treated with testosterone (+T). (D, E) Despite testosterone treatment, astrocytes were almost absent within the remyelinated area in AR^{Tfm} or AR^{NesCre} mice, and axons were remyelinated by Schwann cells. (F) (Left) Mean area of GFAP immunolabeling (\pm SEM) ($F_{5,28} = 5.67$, $P \leq 0.001$). (Right) Mean area of P0 immunolabeling (\pm SEM) ($F_{5,46} = 3.78$, $P \leq 0.01$). Letters on top of columns indicate least significant differences (Left: $P \leq 0.001$, Right: $P \leq 0.01$, post-hoc LSD tests, n = 5-6).

Figure 2. Testosterone increases the reactivity and number of astrocytes *in vivo* and in mixed cultures of glial cells. (A) (Top) Immunohistochemical staining of reactive GFAP⁺ astrocytes (green) and MBP⁺ myelin (red) in the ventrolateral part of the unlesioned male mouse spinal cord (CTRL). Cell nuclei were blue-counterstained with DAPI. Scale bar: 50 μ m. (Center and bottom) GFAP⁺ astrocytes and MBP⁺ myelin at the top border of the demyelinated area (dotted line) in castrated male mice which received lysolecithin injection and were treated for 4 days

with an empty (+ V) or testosterone filled (+T) subcutaneous Silastic implant. (B) Mean area covered by GFAP⁺ astrocytes (\pm SEM) ($F_{2,21} = 14.19$, $P \leq 0.001$). Letters on top of columns indicate least significant differences ($P \leq 0.05$, post-hoc LSD tests, $n = 5-6$). (C) (Top) Immunohistochemical staining of GFAP⁺ astrocytes (red) in control (CTRL) mixed cultures of glial cells. (Center) Exposing the cultures for 12 hours to lysolecithin and treating them for 3 days with vehicle (+V) increased the density of astrocytes (see legend of Fig. S5 for details). (Bottom) Further increase in the density of astrocytes was observed when cultures were treated for 3 days with testosterone (1 μ M) (+ T). (D) Mean area covered by GFAP⁺ astrocytes (\pm SEM) ($F_{2,43} = 21.3$, $P < 0.001$). Letters on top of columns indicate least significant differences (Left: $P \leq 0.05$, post-hoc LSD tests, $n = 3-4$).

Figure 3. Testes, testosterone and AR are required for the spontaneous regeneration of CNS myelin by oligodendrocytes. (A-F) (Left images) MBP-immunoreactive CNS myelin (green) in sagittal sections at 4 weeks after lysolecithin microinjection into the right ventrolateral white matter tract of the spinal cord. Cell nuclei were counterstained with DAPI. (Right images) Immunohistochemical staining of CA II⁺ oligodendrocytes (red). (A) Recovery of MBP and mature oligodendrocytes in gonadally intact wild-type (Wt) mice. Dotted lines delineate the boundary between the ventral white matter (VWM) and the top grey matter. Scale bars: 100 μ m. (B) Absence of recovery of MBP and oligodendrocytes after the removal of testes and treatment with an empty subcutaneous implant (+ V). Castration was performed at the age of 4-6 weeks, 4 weeks prior to lysolecithin-induced demyelination. (C) Regeneration of MBP- immunoreactive myelin and replenishment of oligodendrocytes in castrated males treated with a testosterone-filled implant (+ T). In (D) AR^{Tfm} mice with a non-functional AR, or in (E) AR^{NesCre} mice with CNS-selective ablation of AR, testosterone failed to stimulate CNS remyelination and the replenishment of oligodendrocytes. (F) (Left) Mean

area devoid of MBP immunostaining ($F_{5,30} = 44.8$, $P \leq 0.001$). Letters on top of columns indicate least significant differences (Left: $P \leq 0.01$, Right: $P \leq 0.05$, post-hoc LSD tests, $n = 5-6$). (Right) Mean number of CA II⁺ oligodendrocytes (\pm SEM) within the area of demyelination ($F_{5,34} = 9.2$, $P \leq 0.001$).

Figure 4. Coordinated acquisition of a hinged jaw and myelin, and parallel diversification of steroid receptors during vertebrate evolution. The surrounding of axons with myelin was a prerequisite for the emergence of predation and escape behavior in vertebrates. The evolutionary appearance of myelin indeed parallels the acquisition of a hinged jaw (gnathostomata) (31). The oldest contemporary vertebrates with myelin are cartilaginous fishes (rays, sharks), whereas myelin is absent in jawless fishes (cyclostomata: hagfishes, lampreys) (32). Notably, at the time when jaw and myelin appeared during vertebrate evolution, the 6 nuclear steroid hormone receptors diverged from 3 ancestral receptors during a second round of whole-genome duplication, namely, the androgen (AR), progesterone (PR), glucocorticosteroid (GR), mineralocorticosteroid (MR) and the two estrogen (ER) receptors (25, 26). In extant cyclostomata, only ancestral progesterone (Anc PR), corticoid (Anc CR) and estrogen (Anc ER) receptors are present, themselves derived from 2 ancestral cephalochordata 3-ketosteroid (AncSR) and estrogen (AncER) receptors (16). Cartilaginous fishes, the earliest group of living jawed vertebrates with myelinated axons, contain the most ancient type of AR activated by testosterone (27).

References

1. Monteiro de Castro G, Deja NA, Ma D, Zhao C, & Franklin RJ (2015) Astrocyte Activation via Stat3 Signaling Determines the Balance of Oligodendrocyte versus Schwann Cell Remyelination. *Am. J. Pathol.* 185(9):2431-2440.
2. Keirstead HS, *et al.* (1999) Polysialylated neural cell adhesion molecule-positive CNS precursors generate both oligodendrocytes and Schwann cells to remyelinate the CNS after transplantation. *J. Neurosci.* 19:7529-7536.
3. Crang AJ, Gilson JM, Li WW, & Blakemore WF (2004) The remyelinating potential and in vitro differentiation of MOG-expressing oligodendrocyte precursors isolated from the adult rat CNS. *Eur.J Neurosci.* 20:1445-1460.
4. Talbott JF, *et al.* (2006) Schwann cell-like differentiation by adult oligodendrocyte precursor cells following engraftment into the demyelinated spinal cord is BMP-dependent. *Glia* 54:147-159.
5. Zawadzka M, *et al.* (2010) CNS-resident glial progenitor/stem cells produce Schwann cells as well as oligodendrocytes during repair of CNS demyelination. *Cell Stem Cell* 6:578-590.
6. Keirstead HS, Morgan SV, Wilby MJ, & Fawcett JW (1999) Enhanced axonal regeneration following combined demyelination plus schwann cell transplantation therapy in the injured adult spinal cord. *Exp.Neurol.* 159:225-236.
7. Blakemore WF, Gilson JM, & Crang AJ (2003) The presence of astrocytes in areas of demyelination influences remyelination following transplantation of oligodendrocyte progenitors. *Exp.Neurol.* 184:955-963.
8. Mujtaba T, Mayer-Proschel M, & Rao MS (1998) A common neural progenitor for the CNS and PNS. *Dev. Biol.* 200(1):1-15.
9. Woodruff RH & Franklin RJ (1999) Demyelination and remyelination of the caudal cerebellar peduncle of adult rats following stereotaxic injections of lysolecithin, ethidium bromide, and complement/anti-galactocerebroside: a comparative study. *Glia* 25(3):216-228.
10. Talbott JF, *et al.* (2005) Endogenous Nkx2.2+/Olig2+ oligodendrocyte precursor cells fail to remyelinate the demyelinated adult rat spinal cord in the absence of astrocytes. *Exp.Neurol.* 192:11-24.
11. Matsumoto T, *et al.* (2013) The androgen receptor in health and disease. *Annu.Rev Physiol* 75:201-224.
12. Levin ER (2015) Extranuclear steroid receptors are essential for steroid hormone actions. *Annu. Rev. Med.* 66:271-280.
13. Arnett HA, *et al.* (2004) bHLH transcription factor Olig1 is required to repair demyelinated lesions in the CNS. *Science* 306:2111-2115.
14. Hagedorn L, Suter U, & Sommer L (1999) P0 and PMP22 mark a multipotent neural crest-derived cell type that displays community effects in response to TGF-beta family factors. *Development* 126(17):3781-3794.
15. Hussain R, *et al.* (2013) The neural androgen receptor: a therapeutic target for myelin repair in chronic demyelination. *Brain* 136:132-146.
16. Thornton JW (2001) Evolution of vertebrate steroid receptors from an ancestral estrogen receptor by ligand exploitation and serial genome expansions. *Proc. Natl. Acad. Sci. U. S. A.* 98(10):5671-5676.
17. Padiaditakis I, *et al.* (2015) Dehydroepiandrosterone: an ancestral ligand of neurotrophin receptors. *Endocrinology* 156(1):16-23.
18. Merlet J, Racine C, Moreau E, Moreno SG, & Habert R (2007) Male fetal germ cells are targets for androgens that physiologically inhibit their proliferation. *Proc.Natl.Acad.Sci.U.S.A* 104:3615-3620.

19. Raskin K, *et al.* (2009) Conditional inactivation of androgen receptor gene in the nervous system: effects on male behavioral and neuroendocrine responses. *J.Neurosci.* 29:4461-4470.
20. Crawford AH, Stockley JH, Tripathi RB, Richardson WD, & Franklin RJ (2014) Oligodendrocyte progenitors: adult stem cells of the central nervous system? *Exp. Neurol.* 260:50-55.
21. Mallon BS, Shick HE, Kidd GJ, & Macklin WB (2002) Proteolipid promoter activity distinguishes two populations of NG2-positive cells throughout neonatal cortical development. *J.Neurosci.* 22:876-885.
22. Shi J, Marinovich A, & Barres BA (1998) Purification and characterization of adult oligodendrocyte precursor cells from the rat optic nerve. *J. Neurosci.* 18(12):4627-4636.
23. Psachoulia K, Jamen F, Young KM, & Richardson WD (2009) Cell cycle dynamics of NG2 cells in the postnatal and ageing brain. *Neuron glia biology* 5(3-4):57-67.
24. Fancy SP, Chan JR, Baranzini SE, Franklin RJ, & Rowitch DH (2011) Myelin regeneration: a recapitulation of development? *Annu.Rev Neurosci.* 34:21-43.
25. Eick GN & Thornton JW (2011) Evolution of steroid receptors from an estrogen-sensitive ancestral receptor. *Mol. Cell. Endocrinol.* 334(1-2):31-38.
26. Baker ME, Nelson DR, & Studer RA (2015) Origin of the response to adrenal and sex steroids: Roles of promiscuity and co-evolution of enzymes and steroid receptors. *J. Steroid Biochem. Mol. Biol.* 151:12-24.
27. Ogino Y, Katoh H, Kuraku S, & Yamada G (2009) Evolutionary history and functional characterization of androgen receptor genes in jawed vertebrates. *Endocrinology* 150(12):5415-5427.
28. Ogino Y, *et al.* (2016) Neofunctionalization of Androgen Receptor by Gain-of-Function Mutations in Teleost Fish Lineage. *Mol. Biol. Evol.* 33(1):228-244.
29. Simerly RB, Chang C, Muramatsu M, & Swanson LW (1990) Distribution of androgen and estrogen receptor mRNA-containing cells in the rat brain: an in situ hybridization study. *J. Comp. Neurol.* 294:76-95.
30. DonCarlos LL, *et al.* (2006) Novel cellular phenotypes and subcellular sites for androgen action in the forebrain. *Neuroscience* 138:801-807.
31. Zalc B, Goujet D, & Colman D (2008) The origin of the myelination program in vertebrates. *Curr. Biol.* 18(12):R511-512.
32. Bullock TH, Moore JK, & Fields RD (1984) Evolution of myelin sheaths: both lamprey and hagfish lack myelin. *Neurosci. Lett.* 48(2):145-148.
33. Finley SK & Kritzer MF (1999) Immunoreactivity for intracellular androgen receptors in identified subpopulations of neurons, astrocytes and oligodendrocytes in primate prefrontal cortex. *J.Neurobiol.* 40:446-457.
34. Tronche F, *et al.* (1999) Disruption of the glucocorticoid receptor gene in the nervous system results in reduced anxiety. *Nat Genet.* 23:99-103.
35. Li WW, Penderis J, Zhao C, Schumacher M, & Franklin RJM (2006) Females remyelinate more efficiently than males following demyelination in the aged but not young adult CNS. *Exp.Neurol.* 202:250-254.
36. Okamoto M, *et al.* (2012) Mild exercise increases dihydrotestosterone in hippocampus providing evidence for androgenic mediation of neurogenesis. *Proc. Natl. Acad. Sci. U. S. A.* 109(32):13100-13105.
37. Hussain R, *et al.* (2011) Progesterone and nestorone facilitate axon remyelination: a role for progesterone receptors. *Endocrinology* 152:3820-3831.
38. El-Etr M, *et al.* (2015) Progesterone and nestorone promote myelin regeneration in chronic demyelinating lesions of corpus callosum and cerebral cortex. *Glia* 63(1):104-117.
39. Cerghet M, *et al.* (2006) Proliferation and death of oligodendrocytes and myelin proteins are differentially regulated in male and female rodents. *J.Neurosci.* 26:1439-1447.

40. Palaszynski KM, Loo KK, Ashouri JF, Liu HB, & Voskuhl RR (2004) Androgens are protective in experimental autoimmune encephalomyelitis: implications for multiple sclerosis. *J.Neuroimmunol.* 146:144-152.
41. Sicotte NL, *et al.* (2007) Testosterone treatment in multiple sclerosis: a pilot study. *Arch.Neurol* 64:683-688.
42. Kalincik T, *et al.* (2013) Sex as a determinant of relapse incidence and progressive course of multiple sclerosis. *Brain* 136(Pt 12):3609-3617.
43. Pakpoor J, Goldacre R, Schmierer K, Giovannoni G, & Goldacre MJ (2014) Testicular hypofunction and multiple sclerosis risk: a record-linkage study. *Ann. Neurol.* 76(4):625-628.
44. Bove R, *et al.* (2014) Low testosterone is associated with disability in men with multiple sclerosis. *Mult. Scler.* 20(12):1584-1592.
45. Safarinejad MR (2008) Evaluation of endocrine profile, hypothalamic-pituitary-testis axis and semen quality in multiple sclerosis. *J.Neuroendocrinol.* 20:1368-1375.
46. Confavreux C & Vukusic S (2006) Age at disability milestones in multiple sclerosis. *Brain* 129:595-605.
47. Harman SM, *et al.* (2001) Longitudinal effects of aging on serum total and free testosterone levels in healthy men. Baltimore Longitudinal Study of Aging. *J. Clin. Endocrinol. Metab.* 86(2):724-731.
48. Peters A & Sethares C (2002) Aging and the myelinated fibers in prefrontal cortex and corpus callosum of the monkey. *J. Comp. Neurol.* 442(3):277-291.
49. Ruckh JM, *et al.* (2012) Rejuvenation of regeneration in the aging central nervous system. *Cell Stem Cell* 10:96-103.
50. Fields RD (2008) White matter in learning, cognition and psychiatric disorders. *Trends Neurosci.* 31:361-370.
51. Haroutunian V, *et al.* (2014) Myelination, oligodendrocytes, and serious mental illness. *Glia* 62(11):1856-1877.
52. De Gendt K, *et al.* (2004) A Sertoli cell-selective knockout of the androgen receptor causes spermatogenic arrest in meiosis. *Proc.Natl.Acad.Sci.U.S.A* 101:1327-1332.
53. Feutz AC, Pham-Dinh D, Allinquant B, Miehe M, & Ghandour MS (2001) An immortalized jimpy oligodendrocyte cell line: defects in cell cycle and cAMP pathway. *Glia* 34:241-252.
54. Arevalo MA, Azcoitia I, & Garcia-Segura LM (2015) The neuroprotective actions of oestradiol and oestrogen receptors. *Nat Rev Neurosci* 16(1):17-29.
55. Boon WC, Chow JD, & Simpson ER (2010) The multiple roles of estrogens and the enzyme aromatase. *Prog. Brain Res.* 181:209-232.
56. van de Wijngaart DJ, Dubbink HJ, van Royen ME, Trapman J, & Jenster G (2012) Androgen receptor coregulators: recruitment via the coactivator binding groove. *Mol. Cell. Endocrinol.* 352(1-2):57-69.
57. Belelli D & Lambert JJ (2005) Neurosteroids: endogenous regulators of the GABA(A) receptor. *Nat.Rev.Neurosci.* 6:565-575.
58. Reddy DS (2010) Neurosteroids: endogenous role in the human brain and therapeutic potentials. *Prog. Brain Res.* 186:113-137.
59. Lund TD, Hinds LR, & Handa RJ (2006) The androgen 5alpha-dihydrotestosterone and its metabolite 5alpha-androstan-3beta, 17beta-diol inhibit the hypothalamo-pituitary-adrenal response to stress by acting through estrogen receptor beta-expressing neurons in the hypothalamus. *J. Neurosci.* 26(5):1448-1456.




Figure 1




Figure 2




Figure 3




Figure 4