



A revised catalogue of 294 Galactic supernova remnants

D. A. GREEN 

Astrophysics Group, Cavendish Laboratory, 19 J. J. Thomson Avenue, Cambridge CB3 0HE, UK.
E-mail: D.A.Green@mrao.cam.ac.uk

MS received 11 June 2019; accepted 4 July 2019

Abstract. A revised catalogue of Galactic supernova remnants (SNRs) is presented, along with some simple statistics of their properties. Six new SNRs have been added to the catalogue since the previous published version from 2014, and six entries have been removed, as they have been identified as H II regions, leaving the number of entries in the catalogue at 294. Some simple statistics of the remnants in the catalogue, and the selection effects that apply, are discussed, along with some recently proposed Galactic SNR candidates.

Keywords. Supernova remnants—catalogues—ISM: general.

1. Introduction

This paper presents the latest version of a catalogue of Galactic supernova remnants (SNRs) which I have compiled for several decades. Previous versions have been published in Green (1984, 1988, 1991, 2004, 2009, 2014) and Stephenson & Green (2002). In addition, more detailed web-based versions of the catalogue have been produced since 1995 – most recently in 2017 – which either correspond to one of the published catalogues, or are an intermediate revision. This version of the catalogue contains 294 entries. Section 2 gives the details of the entries in the catalogue, and Section 3 discusses the entries added or removed from the catalogue since the last published version (Green 2014). Section 4 discusses some simple statistics of the remnants in the current catalogue, the selection effects that apply to the identification of Galactic SNRs, and some recently proposed candidate SNRs.

2. The catalogue format

This catalogue is based on the literature published up to the end of 2018, and contains 294 entries. For each SNR in the catalogue, the following parameters are given:

- *Galactic coordinates of the remnant.* These are quoted to a tenth of a degree, as is conventional. In this catalogue additional leading zeros are not

used. These are generally taken from the Galactic coordinate based name used for the remnant in the literature. It should be noted that when these names were first defined, they may not have followed the IAU recommendation¹ that coordinates should be truncated, not rounded to construct such names.

- *Right ascension and declination.* The right ascension and declination of J2000.0 equatorial coordinates of the source centroid, for which an accuracy of the quoted values depends on the size of the remnant. For small remnants, they are to the nearest few seconds of time and the nearest minute of arc respectively, whereas for larger remnants they are rounded to coarser values, but are in every case sufficient to specify a point within the boundary of the remnant. These coordinates are usually deduced from radio images rather than from X-ray or optical observations.
- *Angular size.* The angular size of the remnant in arcminutes. This is usually taken from the highest resolution radio image available. The boundary of most remnants approximates reasonably well to either a circle or to an ellipse. A single value is quoted for the angular size of the more nearly circular remnants, which is the diameter of a circle with an area equal to that of the remnant. For more elongated remnants, the product of two values is

¹See, <http://cdsweb.u-strasbg.fr/Dic/iau-spec.htx>.

given, which are the major and minor diameters of the remnant boundary modelled as an ellipse. In a small number of cases an ellipse is not a good description of the boundary of the object (which will be noted in the description of the object given in its catalogue entry), although an angular size is still quoted for information. For ‘filled-centre’ type remnants (see below), the size quoted is for the largest extent of the observed emission, not, as at times has been used by others, the half-width of the centrally brightened peak.

- *Type of the SNR.* This is ‘S’ or ‘F’ if the remnant shows a ‘shell’ or ‘filled-centre’ structure, or ‘C’ if it shows ‘composite’ (or ‘combination’) radio structure, with a combination of shell and filled-centre characteristics. If there is some uncertainty, the type is given as ‘S?’, ‘F?’ or ‘C?’, and as ‘?’ in several cases where an object is conventionally regarded as an SNR even though its nature is poorly known or it is not well-understood. (Note: the term ‘composite’ has been used, by some authors, in a different sense, to describe remnants with radio shell and centrally-brightened X-ray emission. An alternative term used to describe such remnants is ‘mixed morphology’, e.g. see [Rho & Petre 1998](#).)
- *Flux density.* The flux density of the remnant at a frequency of 1 GHz, in jansky. This is *not* a measured value, but is instead derived from the observed radio spectrum of the source. The frequency of 1 GHz is chosen because flux density measurements are usually available at both higher and lower frequencies. Some young remnants – notably G111.7–2.1 (=Cassiopeia A) and G184.6–5.8 (=Crab Nebula), but also G130.7+3.1 (=3C58) and G120.1+1.4 (=Tycho) – show secular variations in their radio flux density. In this revision of the catalogue, the 1-GHz flux densities for G111.7–2.1 and G184.6–5.8 have been taken from [Perley and Butler \(2017\)](#), for an epoch of 2016. Results from the primary literature should be used for any detailed quantitative studies of the radio spectra of these and other remnants.
- *Spectral index.* The spectral index of the integrated radio emission from the remnant, α (here defined in the sense, $S \propto \nu^{-\alpha}$, where S is the flux density at frequency ν). This is either a value that is quoted in the literature, or one deduced from the available integrated flux densities of the remnant. For several SNRs a simple power law is not adequate to describe their radio spectra, either

because there is evidence that the integrated spectrum is curved or the spectral index varies across the face of the remnant. In these cases the spectral index is given as ‘varies’ (refer to the description of the remnant and appropriate references in the detailed catalogue entry for more information). In some cases, for example where the remnant is highly confused with thermal emission, the spectral index is given as ‘?’ since no value can be deduced with any confidence. These spectral indices have a very wide range of quality, and the primary literature should be consulted for any detailed study of the radio spectral indices of these remnants.

- *Other names that are commonly used for the remnant.* Note that these are given in parentheses if the remnant is only a part of the source. For some well-known remnants – e.g. G184.6–5.8, the Crab Nebula – not all common names are given.

A summary of the data available for all 294 remnants in the catalogue is given in Table 1.

A more detailed version of the catalogue is available at <http://www.mrao.cam.ac.uk/surveys/snrs/>. In addition to the basic parameters which are given in Table 1, the detailed catalogue contains the following additional information.

- (i) Notes on the remnant. For example, if other Galactic coordinates have at times been used to label it (usually before good observations have revealed the full extent of the object, but sometimes in error); if the SNR is thought to be the remnant of a historical SN.
- (ii) Short descriptions of the observed structure/properties of the remnant at radio, optical and X-ray wavelengths, as appropriate from available observations.
- (iii) Comments on distance determinations, and any point sources or pulsars in or near the object (although they may not necessarily be related to the remnant).
- (iv) References to observations are given for each remnant, complete with journal, volume, page, and a short description of what information each paper contains (e.g. for radio observations these generally include the telescopes used, the observing frequencies and resolutions, together with any flux density determinations). These references are *not* complete, but cover recent and representative observations of the remnant that

Table 1. 294 Galactic supernova remnants: summary data.

<i>l</i> /°	<i>b</i> /°	RA (J2000) Dec		Size	Type	Flux at 1 GHz/Jy	Spectral index	Other name(s)
		/(h m s)	/(° ′)	/arcmin				
0.0	+0.0	17 45 44	−29 00	3.5×2.5	S	100?	0.8?	Sgr A East
0.3	+0.0	17 46 15	−28 38	15×8	S	22	0.6	
0.9	+0.1	17 47 21	−28 09	8	C	18?	varies	
1.0	−0.1	17 48 30	−28 09	8	S	15	0.6?	
1.4	−0.1	17 49 39	−27 46	10	S	2?	?	
1.9	+0.3	17 48 45	−27 10	1.5	S	0.6	0.6	
3.7	−0.2	17 55 26	−25 50	14×11	S	2.3	0.65	
3.8	+0.3	17 52 55	−25 28	18	S?	3?	0.6	
4.2	−3.5	18 08 55	−27 03	28	S	3.2?	0.6?	
4.5	+6.8	17 30 42	−21 29	3	S	19	0.64	Kepler, SN1604, 3C358
4.8	+6.2	17 33 25	−21 34	18	S	3	0.6	
5.2	−2.6	18 07 30	−25 45	18	S	2.6?	0.6?	
5.4	−1.2	18 02 10	−24 54	35	C?	35?	0.2?	Milne 56
5.5	+0.3	17 57 04	−24 00	15×12	S	5.5	0.7	
5.9	+3.1	17 47 20	−22 16	20	S	3.3?	0.4?	
6.1	+0.5	17 57 29	−23 25	18×12	S	4.5	0.9	
6.1	+1.2	17 54 55	−23 05	30×26	F	4.0?	0.3?	
6.4	−0.1	18 00 30	−23 26	48	C	310	varies	W28
6.4	+4.0	17 45 10	−21 22	31	S	1.3?	0.4?	
6.5	−0.4	18 02 11	−23 34	18	S	27	0.6	
7.0	−0.1	18 01 50	−22 54	15	S	2.5?	0.5?	
7.2	+0.2	18 01 07	−22 38	12	S	2.8	0.6	
7.7	−3.7	18 17 25	−24 04	22	S	11	0.32	1814–24
8.3	−0.0	18 04 34	−21 49	5×4	S	1.2	0.6	
8.7	−5.0	18 24 10	−23 48	26	S	4.4	0.3	
8.7	−0.1	18 05 30	−21 26	45	S?	80	0.5	(W30)
8.9	+0.4	18 03 58	−21 03	24	S	9	0.6	
9.7	−0.0	18 07 22	−20 35	15×11	S	3.7	0.6	
9.8	+0.6	18 05 08	−20 14	12	S	3.9	0.5	
9.9	−0.8	18 10 41	−20 43	12	S	6.7	0.4	
10.5	−0.0	18 09 08	−19 47	6	S	0.9	0.6	
11.0	−0.0	18 10 04	−19 25	11×9	S	1.3	0.6	
11.1	−1.0	18 14 03	−19 46	18×12	S	5.8	0.5	
11.1	−0.7	18 12 46	−19 38	11×7	S	1.0	0.7	
11.1	+0.1	18 09 47	−19 12	12×10	S	2.3	0.4	
11.2	−0.3	18 11 27	−19 25	4	C	22	0.5	
11.4	−0.1	18 10 47	−19 05	8	S?	6	0.5	
11.8	−0.2	18 12 25	−18 44	4	S	0.7	0.3	
12.0	−0.1	18 12 11	−18 37	7?	?	3.5	0.7	
12.2	+0.3	18 11 17	−18 10	6×5	S	0.8	0.7	
12.5	+0.2	18 12 14	−17 55	6×5	C?	0.6	0.4	
12.7	−0.0	18 13 19	−17 54	6	S	0.8	0.8	
12.8	−0.0	18 13 37	−17 49	3	C?	0.8	0.5	
13.3	−1.3	18 19 20	−18 00	70×40	S?	?	?	
13.5	+0.2	18 14 14	−17 12	5×4	S	3.5?	1.0?	

Table 1. *Continued.*

l / $^{\circ}$	b / $^{\circ}$	RA (J2000) /(h m s)	Dec / $(^{\circ}')$	Size /arcmin	Type	Flux at 1 GHz/Jy	Spectral index	Other name(s)
14.1	-0.1	18 16 40	-16 41	6×5	S	0.5	0.6	
14.3	+0.1	18 15 58	-16 27	5×4	S	0.6	0.4	
15.1	-1.6	18 24 00	-16 34	30×24	S?	5.5?	0.0?	
15.4	+0.1	18 18 02	-15 27	15×14	C?	5.6	0.62	
15.9	+0.2	18 18 52	-15 02	7×5	S?	5.0	0.63	
16.0	-0.5	18 21 56	-15 14	15×10	S	2.7	0.6	
16.2	-2.7	18 29 40	-16 08	17	S	2.5	0.4	
16.4	-0.5	18 22 38	-14 55	13	S	4.6	0.3?	
16.7	+0.1	18 20 56	-14 20	4	C	3.0	0.6	
17.0	-0.0	18 21 57	-14 08	5	S	0.5	0.5	
17.4	-2.3	18 30 55	-14 52	24?	S	5	0.5?	
17.4	-0.1	18 23 08	-13 46	6	S	0.4	0.7	
17.8	-2.6	18 32 50	-14 39	24	S	5	0.5	
18.1	-0.1	18 24 34	-13 11	8	S	4.6	0.5	
18.6	-0.2	18 25 55	-12 50	6	S	1.4	0.4	
18.8	+0.3	18 23 58	-12 23	17×11	S	33	0.46	Kes 67
18.9	-1.1	18 29 50	-12 58	33	C?	37	0.39	
19.1	+0.2	18 24 56	-12 07	27	S	10	0.5	
20.0	-0.2	18 28 07	-11 35	10	F	10	0.1	
21.0	-0.4	18 31 12	-10 47	9×7	S	1.1	0.6	
21.5	-0.9	18 33 33	-10 35	5	C	7	varies	
21.6	-0.8	18 33 40	-10 25	13	S	1.4	0.5?	
21.8	-0.6	18 32 45	-10 08	20	S	65	0.56	Kes 69
22.7	-0.2	18 33 15	-09 13	26	S?	33	0.6	
23.3	-0.3	18 34 45	-08 48	27	S	70	0.5	W41
24.7	-0.6	18 38 43	-07 32	15?	S?	8	0.5	
24.7	+0.6	18 34 10	-07 05	30×15	C?	20?	0.2?	
25.1	-2.3	18 45 10	-08 00	80×30?	S	8	0.5?	
27.4	+0.0	18 41 19	-04 56	4	S	6	0.68	4C-04.71
27.8	+0.6	18 39 50	-04 24	50×30	F	30	varies	
28.6	-0.1	18 43 55	-03 53	13×9	S	3?	?	
28.8	+1.5	18 39 00	-02 55	100?	S?	?	0.4?	
29.6	+0.1	18 44 52	-02 57	5	S	1.5?	0.5?	
29.7	-0.3	18 46 25	-02 59	3	C	10	0.63	Kes 75
30.7	-2.0	18 54 25	-02 54	16	?	0.5?	0.7?	
30.7	+1.0	18 44 00	-01 32	24×18	S?	6	0.4	
31.5	-0.6	18 51 10	-01 31	18?	S?	2?	?	
31.9	+0.0	18 49 25	-00 55	7×5	S	25	varies	3C391
32.0	-4.9	19 06 00	-03 00	60?	S?	22?	0.5?	3C396.1
32.1	-0.9	18 53 10	-01 08	40?	C?	?	?	
32.4	+0.1	18 50 05	-00 25	6	S	0.25?	?	
32.8	-0.1	18 51 25	-00 08	22×15	S?	11?	0.2?	Kes 78
33.2	-0.6	18 53 50	-00 02	18	S	3.5	varies	
33.6	+0.1	18 52 48	+00 41	10	S	20	0.51	Kes 79, 4C00.70, HC13
34.7	-0.4	18 56 00	+01 22	35×27	C	240	0.37	W44, 3C392

Table 1. *Continued.*

<i>l</i> /°	<i>b</i> /°	RA (J2000) /(h m s)	Dec /(° ′)	Size /arcmin	Type	Flux at 1 GHz/Jy	Spectral index	Other name(s)
35.6	−0.4	18 57 55	+02 13	15×11	S?	9	0.5	
36.6	−0.7	19 00 35	+02 56	25?	S?	1.0	0.7?	
36.6	+2.6	18 48 49	+04 26	17×13?	S	0.7?	0.5?	
38.7	−1.3	19 06 40	+04 28	32×19?	S	?	?	
39.2	−0.3	19 04 08	+05 28	8×6	C	18	0.34	3C396, HC24, NRAO 593
39.7	−2.0	19 12 20	+04 55	120×60	?	85?	0.7?	W50, SS433
40.5	−0.5	19 07 10	+06 31	22	S	11	0.4	
41.1	−0.3	19 07 34	+07 08	4.5×2.5	S	25	0.50	3C397
41.5	+0.4	19 05 50	+07 46	10	S?	1?	?	
42.0	−0.1	19 08 10	+08 00	8	S?	0.5?	?	
42.8	+0.6	19 07 20	+09 05	24	S	3?	0.5?	
43.3	−0.2	19 11 08	+09 06	4×3	S	38	0.46	W49B
43.9	+1.6	19 05 50	+10 30	60?	S?	9.0	0.5	
45.7	−0.4	19 16 25	+11 09	22	S	4.2?	0.4?	
46.8	−0.3	19 18 10	+12 09	15	S	17	0.54	(HC30)
49.2	−0.7	19 23 50	+14 06	30	S?	160?	0.3?	(W51)
53.4	+0.0	19 29 57	+18 10	10?	S	1.5	0.6?	
53.6	−2.2	19 38 50	+17 14	33×28	S	8	0.50	3C400.2, NRAO 611
54.1	+0.3	19 30 31	+18 52	12?	C?	0.5	0.1	
54.4	−0.3	19 33 20	+18 56	40	S	28	0.5	(HC40)
55.0	+0.3	19 32 00	+19 50	20×15?	S	0.5?	0.5?	
55.7	+3.4	19 21 20	+21 44	23	S	1?	0.3?	
57.2	+0.8	19 34 59	+21 57	12?	S?	1.8	0.35	(4C21.53)
59.5	+0.1	19 42 33	+23 35	15	S	3?	?	
63.7	+1.1	19 47 52	+27 45	8	F	1.8	0.24	
64.5	+0.9	19 50 25	+28 16	8	S?	0.15?	0.5	
65.1	+0.6	19 54 40	+28 35	90×50	S	5.5	0.61	
65.3	+5.7	19 33 00	+31 10	310×240	S?	42	0.6	
65.7	+1.2	19 52 10	+29 26	22	F	5.1	varies	DA 495
66.0	−0.0	19 57 50	+29 03	31×25?	S	?	?	
67.6	+0.9	19 57 45	+30 53	50×45?	S	?	?	
67.7	+1.8	19 54 32	+31 29	15×12	S	1.0	0.61	
67.8	+0.5	20 00 00	+30 51	7×5	?	?	?	
68.6	−1.2	20 08 40	+30 37	23	?	1.1	0.2	
69.0	+2.7	19 53 20	+32 55	80?	?	120?	varies	CTB 80
69.7	+1.0	20 02 40	+32 43	16×14	S	2.0	0.7	
70.0	−21.5	21 24 00	+19 23	330×240	S	?	?	
73.9	+0.9	20 14 15	+36 12	27	S?	9	0.23	
74.0	−8.5	20 51 00	+30 40	230×160	S	210	varies	Cygnus Loop
74.9	+1.2	20 16 02	+37 12	8×6	F	9	varies	CTB 87
76.9	+1.0	20 22 20	+38 43	9	C	2?	?	
78.2	+2.1	20 20 50	+40 26	60	S	320	0.51	DR4, γ Cygni SNR
82.2	+5.3	20 19 00	+45 30	95×65	S	120?	0.5?	W63
83.0	−0.3	20 46 55	+42 52	9×7	S	1	0.4	
84.2	−0.8	20 53 20	+43 27	20×16	S	11	0.5	

Table 1. *Continued.*

l /°	b /°	RA (J2000) /(h m s)	Dec /(° ′)	Size /arcmin	Type	Flux at 1 GHz/Jy	Spectral index	Other name(s)
85.4	+0.7	20 50 40	+45 22	24?	S	?	0.2	
85.9	-0.6	20 58 40	+44 53	24	S	?	0.2	
89.0	+4.7	20 45 00	+50 35	120×90	S	220	0.38	HB21
93.3	+6.9	20 52 25	+55 21	27×20	C?	9	0.45	DA 530, 4C(T)55.38.1
93.7	-0.2	21 29 20	+50 50	80	S	65	0.65	CTB 104A, DA 551
94.0	+1.0	21 24 50	+51 53	30×25	S	13	0.45	3C434.1
96.0	+2.0	21 30 30	+53 59	26	S	0.35	0.6	
106.3	+2.7	22 27 30	+60 50	60×24	C?	6	0.6	
108.2	-0.6	22 53 40	+58 50	70×54	S	8	0.5	
109.1	-1.0	23 01 35	+58 53	28	S	20	0.45	CTB 109
111.7	-2.1	23 23 26	+58 48	5	S	2300	0.77	Cassiopeia A, 3C461
113.0	+0.2	23 26 50	+61 26	40×17?	?	4	0.5?	
114.3	+0.3	23 37 00	+61 55	90×55	S	5.5	0.5	
116.5	+1.1	23 53 40	+63 15	80×60	S	10	0.5	
116.9	+0.2	23 59 10	+62 26	34	S	8	0.57	CTB 1
119.5	+10.2	00 06 40	+72 45	90?	S	36	0.6	CTA 1
120.1	+1.4	00 25 18	+64 09	8	S	50	0.58	Tycho, 3C10, SN1572
126.2	+1.6	01 22 00	+64 15	70	S?	6	0.5	
127.1	+0.5	01 28 20	+63 10	45	S	12	0.45	R5
130.7	+3.1	02 05 41	+64 49	9×5	F	33	0.07	3C58, SN1181
132.7	+1.3	02 17 40	+62 45	80	S	45	0.6	HB3
150.3	+4.5	04 27 00	+55 28	180×150	S	?	?	
152.4	-2.1	04 07 50	+49 11	100×95	S	3.5?	0.7?	
156.2	+5.7	04 58 40	+51 50	110	S	5	0.5	
159.6	+7.3	05 20 00	+50 00	240×180?	S	?	?	
160.9	+2.6	05 01 00	+46 40	140×120	S	110	0.64	HB9
166.0	+4.3	05 26 30	+42 56	55×35	S	7	0.37	VRO 42.05.01
178.2	-4.2	05 25 05	+28 11	72×62	S	2	0.5	
179.0	+2.6	05 53 40	+31 05	70	S?	7	0.4	
180.0	-1.7	05 39 00	+27 50	180	S	65	varies	S147
181.1	+9.5	06 26 40	+32 30	74	S	?	0.45?	
182.4	+4.3	06 08 10	+29 00	50	S	0.5	0.4	
184.6	-5.8	05 34 31	+22 01	7×5	F	900	0.30	Crab Nebula, 3C144, SN1054
189.1	+3.0	06 17 00	+22 34	45	C	165	0.36	IC443, 3C157
190.9	-2.2	06 01 55	+18 24	70×60	S	1.3?	0.7?	
205.5	+0.5	06 39 00	+06 30	220	S	140	0.4	Monoceros Nebula
206.9	+2.3	06 48 40	+06 26	60×40	S?	6	0.5	PKS 0646+06
213.0	-0.6	06 50 50	-00 30	160×140?	S	21	0.4	
260.4	-3.4	08 22 10	-43 00	60×50	S	130	0.5	Puppis A, MSH 08-44
261.9	+5.5	09 04 20	-38 42	40×30	S	10?	0.4?	
263.9	-3.3	08 34 00	-45 50	255	C	1750	varies	Vela (XYZ)
266.2	-1.2	08 52 00	-46 20	120	S	50?	0.3?	RX J0852.0-4622
272.2	-3.2	09 06 50	-52 07	15?	S?	0.4	0.6	
279.0	+1.1	09 57 40	-53 15	95	S	30?	0.6?	
284.3	-1.8	10 18 15	-59 00	24?	S	11?	0.3?	MSH 10-53

Table 1. *Continued.*

<i>l</i> /°	<i>b</i> /°	RA (J2000) /(h m s)	Dec /(° ′)	Size /arcmin	Type	Flux at 1 GHz/Jy	Spectral index	Other name(s)
286.5	-1.2	10 35 40	-59 42	26×6	S?	1.4?	?	
289.7	-0.3	11 01 15	-60 18	18×14	S	6.2	0.2?	
290.1	-0.8	11 03 05	-60 56	19×14	S	42	0.4	MSH 11-61A
291.0	-0.1	11 11 54	-60 38	15×13	C	16	0.29	(MSH 11-62)
292.0	+1.8	11 24 36	-59 16	12×8	C	15	0.4	MSH 11-54
292.2	-0.5	11 19 20	-61 28	20×15	S	7	0.5	
293.8	+0.6	11 35 00	-60 54	20	C	5?	0.6?	
294.1	-0.0	11 36 10	-61 38	40	S	>2?	?	
296.1	-0.5	11 51 10	-62 34	37×25	S	8?	0.6?	
296.5	+10.0	12 09 40	-52 25	90×65	S	48	0.5	PKS 1209-51/52
296.7	-0.9	11 55 30	-63 08	15×8	S	3	0.5	
296.8	-0.3	11 58 30	-62 35	20×14	S	9	0.6	1156-62
298.5	-0.3	12 12 40	-62 52	5?	?	5?	0.4?	
298.6	-0.0	12 13 41	-62 37	12×9	S	5?	0.3	
299.2	-2.9	12 15 13	-65 30	18×11	S	0.5?	?	
299.6	-0.5	12 21 45	-63 09	13	S	1.0?	?	
301.4	-1.0	12 37 55	-63 49	37×23	S	2.1?	?	
302.3	+0.7	12 45 55	-62 08	17	S	5?	0.4?	
304.6	+0.1	13 05 59	-62 42	8	S	14	0.5	Kes 17
306.3	-0.9	13 21 50	-63 34	4	S?	0.16?	0.5?	
308.1	-0.7	13 37 37	-63 04	13	S	1.2?	?	
308.4	-1.4	13 41 30	-63 44	12×6?	S?	0.4?	?	
308.8	-0.1	13 42 30	-62 23	30×20?	C?	15?	0.4?	
309.2	-0.6	13 46 31	-62 54	15×12	S	7?	0.4?	
309.8	+0.0	13 50 30	-62 05	25×19	S	17	0.5	
310.6	-1.6	14 00 45	-63 26	2.5	C?	?	?	
310.6	-0.3	13 58 00	-62 09	8	S	5?	?	Kes 20B
310.8	-0.4	14 00 00	-62 17	12	S	6?	?	Kes 20A
311.5	-0.3	14 05 38	-61 58	5	S	3?	0.5	
312.4	-0.4	14 13 00	-61 44	38	S	45	0.36	
312.5	-3.0	14 21 00	-64 12	20×18	S	3.5?	?	
315.1	+2.7	14 24 30	-57 50	190×150	S	?	?	
315.4	-2.3	14 43 00	-62 30	42	S	49	0.6	RCW 86, MSH 14-63
315.4	-0.3	14 35 55	-60 36	24×13	?	8	0.4	
315.9	-0.0	14 38 25	-60 11	25×14	S	0.8?	?	
316.3	-0.0	14 41 30	-60 00	29×14	S	20?	0.4	(MSH 14-57)
317.3	-0.2	14 49 40	-59 46	11	S	4.7?	?	
318.2	+0.1	14 54 50	-59 04	40×35	S	>3.9?	?	
318.9	+0.4	14 58 30	-58 29	30×14	C	4?	0.2?	
320.4	-1.2	15 14 30	-59 08	35	C	60?	0.4	MSH 15-52, RCW 89
320.6	-1.6	15 17 50	-59 16	60×30	S	?	?	
321.9	-1.1	15 23 45	-58 13	28	S	>3.4?	?	
321.9	-0.3	15 20 40	-57 34	31×23	S	13	0.3	
322.1	+0.0	15 20 49	-57 10	8×4.5?	S?	?	?	
322.5	-0.1	15 23 23	-57 06	15	C	1.5	0.4	

Table 1. *Continued.*

l /°	b /°	RA (J2000) /(h m s)	Dec /(° ′)	Size /arcmin	Type	Flux at 1 GHz/Jy	Spectral index	Other name(s)
323.5	+0.1	15 28 42	−56 21	13	S	3?	0.4?	
323.7	−1.0	15 34 30	−57 12	51×38	S	?	?	
326.3	−1.8	15 53 00	−56 10	38	C	145	varies	MSH 15–56
327.1	−1.1	15 54 25	−55 09	18	C	7?	?	
327.2	−0.1	15 50 55	−54 18	5	S	0.4	?	
327.4	+0.4	15 48 20	−53 49	21	S	30?	0.6	Kes 27
327.4	+1.0	15 46 48	−53 20	14	S	1.9?	?	
327.6	+14.6	15 02 50	−41 56	30	S	19	0.6	SN1006, PKS 1459–41
328.4	+0.2	15 55 30	−53 17	5	F	15	0.0	(MSH 15–57)
329.7	+0.4	16 01 20	−52 18	40×33	S	>34?	?	
330.0	+15.0	15 10 00	−40 00	180?	S	350?	0.5?	Lupus Loop
330.2	+1.0	16 01 06	−51 34	11	S?	5?	0.3	
332.0	+0.2	16 13 17	−50 53	12	S	8?	0.5	
332.4	−0.4	16 17 33	−51 02	10	S	28	0.5	RCW 103
332.4	+0.1	16 15 20	−50 42	15	S	26	0.5	MSH 16–51, Kes 32
332.5	−5.6	16 43 20	−54 30	35	S	2?	0.7?	
335.2	+0.1	16 27 45	−48 47	21	S	16	0.5	
336.7	+0.5	16 32 11	−47 19	14×10	S	6	0.5	
337.0	−0.1	16 35 57	−47 36	1.5	S	1.5	0.6?	(CTB 33)
337.2	−0.7	16 39 28	−47 51	6	S	1.5	0.4	
337.2	+0.1	16 35 55	−47 20	3×2	?	1.5?	?	
337.3	+1.0	16 32 39	−46 36	15×12	S	16	0.55	Kes 40
337.8	−0.1	16 39 01	−46 59	9×6	S	15	0.5	Kes 41
338.1	+0.4	16 37 59	−46 24	15?	S	4?	0.4	
338.3	−0.0	16 41 00	−46 34	8	C?	7?	?	
338.5	+0.1	16 41 09	−46 19	9	?	12?	?	
340.4	+0.4	16 46 31	−44 39	10×7	S	5	0.4	
340.6	+0.3	16 47 41	−44 34	6	S	5?	0.4?	
341.2	+0.9	16 47 35	−43 47	22×16	C	1.5?	0.6?	
341.9	−0.3	16 55 01	−44 01	7	S	2.5	0.5	
342.0	−0.2	16 54 50	−43 53	12×9	S	3.5?	0.4?	
342.1	+0.9	16 50 43	−43 04	10×9	S	0.5?	?	
343.0	−6.0	17 25 00	−46 30	250	S	?	?	RCW 114
343.1	−2.3	17 08 00	−44 16	32?	C?	8?	0.5?	
343.1	−0.7	17 00 25	−43 14	27×21	S	7.8	0.55	
344.7	−0.1	17 03 51	−41 42	8	C?	2.5?	0.3?	
345.7	−0.2	17 07 20	−40 53	6	S	0.6?	?	
346.6	−0.2	17 10 19	−40 11	8	S	8?	0.5?	
347.3	−0.5	17 13 50	−39 45	65×55	S?	30?	?	RX J1713.7–3946
348.5	−0.0	17 15 26	−38 28	10?	S?	10?	0.4?	
348.5	+0.1	17 14 06	−38 32	15	S	72	0.3	CTB 37A
348.7	+0.3	17 13 55	−38 11	17?	S	26	0.3	CTB 37B
349.2	−0.1	17 17 15	−38 04	9×6	S	1.4?	?	
349.7	+0.2	17 17 59	−37 26	2.5×2	S	20	0.5	
350.0	−2.0	17 27 50	−38 32	45	S	26	0.4	

Table 1. *Continued.*

<i>l</i> /°	<i>b</i> /°	RA (J2000) /(h m s)	Dec /(° ′)	Size /arcmin	Type	Flux at 1 GHz/Jy	Spectral index	Other name(s)
350.1	−0.3	17 21 05	−37 27	4?	?	6?	0.8?	
351.0	−5.4	17 46 00	−39 25	30	S	?	?	
351.2	+0.1	17 22 27	−36 11	7	C?	5?	0.4	
351.7	+0.8	17 21 00	−35 27	18×14	S	10	0.5?	
351.9	−0.9	17 28 52	−36 16	12×9	S	1.8?	?	
352.7	−0.1	17 27 40	−35 07	8×6	S	4	0.6	
353.6	−0.7	17 32 00	−34 44	30	S	2.5?	?	
353.9	−2.0	17 38 55	−35 11	13	S	1?	0.5?	
354.1	+0.1	17 30 28	−33 46	15×3?	C?	?	varies	
354.8	−0.8	17 36 00	−33 42	19	S	2.8?	?	
355.4	+0.7	17 31 20	−32 26	25	S	5?	?	
355.6	−0.0	17 35 16	−32 38	8×6	S	3?	?	
355.9	−2.5	17 45 53	−33 43	13	S	8	0.5	
356.2	+4.5	17 19 00	−29 40	25	S	4	0.7	
356.3	−1.5	17 42 35	−32 52	20×15	S	3?	?	
356.3	−0.3	17 37 56	−32 16	11×7	S	3?	?	
357.7	−0.1	17 40 29	−30 58	8×3?	?	37	0.4	MSH 17–39
357.7	+0.3	17 38 35	−30 44	24	S	10	0.4?	
358.0	+3.8	17 26 00	−28 36	38	S	1.5?	?	
358.1	+1.0	17 37 00	−29 59	20	S	2?	?	
358.5	−0.9	17 46 10	−30 40	17	S	4?	?	
359.0	−0.9	17 46 50	−30 16	23	S	23	0.5	
359.1	−0.5	17 45 30	−29 57	24	S	14	0.4?	
359.1	+0.9	17 39 36	−29 11	12×11	S	2?	?	

are available, and should themselves include references to earlier work. These references are from the published literature up to the end of 2018.

The detailed version of the catalogue is available in pdf format for downloading and printing, or as web pages, including a page for each individual remnant. The web pages for each remnant include links to the ‘NASA Astrophysics Data System’ for each of the over three thousand references that are included in the detailed listings for individual SNRs.

Some of the parameters included in the catalogue are themselves of variable quality. For example, the radio flux density of each remnant at 1 GHz is generally obtained from several radio observations over a range of frequencies, both above and below 1 GHz, so it is of good quality. However, there are 21 remnants – often those which have been identified at other than radio wavelengths – for which no reliable radio flux density is available yet, because they have either not

been detected or well observed in the radio. Although the detailed version of the catalogue contains notes on distances for many remnants reported in the literature, these are highly variable in terms of reliability and accuracy. Consequently, the distances given within the detailed catalogue should be used with caution in any statistical studies, and reference should be made to the primary literature cited in the detailed catalogue.

The detailed version of the catalogue also contains notes both on those objects no longer thought to be SNRs, and on the many possible and probable remnants that have been reported in the literature (including possible large, old remnants, seen from radio continuum, X-ray or HI observations). See Section 4.3 below for discussion of some recently proposed remnants.

It should be noted that the catalogue is far from homogeneous. Although many remnants, or possible remnants, were first identified from wide-area radio surveys, there are many others that have been observed with diverse observational parameters, making uniform

criteria for inclusion in the main catalogue difficult. For an alternative, high-energy catalogue of SNRs, see [Ferrand and Safi-Harb \(2012\)](#).

3. SNRs added to, objects removed from the catalogue

Since the last published version ([Green 2014](#)), the following supernova remnants have been added to the catalogue.

- G351.0–5.4, which was identified by [de Gasperin et al. \(2014\)](#) from radio and other observations.
- A very high-latitude remnant, G70.0–21.5, identified primarily from optical observations by [Fesen et al. \(2015\)](#). Previously, [Boumis et al. \(2002\)](#) had noted optical filaments in this region, which they suggested were indicative of one or more SNRs. As noted by both [Boumis et al. \(2002\)](#) and [Fesen et al. \(2015\)](#), there is also faint X-ray emission from this remnant.
- G181.1+9.5, another high-latitude remnant, identified from radio observations by [Kothes et al. \(2017\)](#).
- G323.7–1.0, which was one of the several candidate remnant given by [Green et al. \(2014\)](#), was confirmed as a SNR from γ -ray observations, see [Araya \(2017\)](#) and [H.E.S.S. Collaboration et al. \(2018\)](#).
- The possible faint radio SNRs near $l = 150^\circ 5$, $b = 4^\circ 0$ have been reported by [Gerbrandt et al. \(2014\)](#) and [Gao & Han \(2014\)](#). [Gao & Han \(2014\)](#) proposed a large ($180 \times 150 \text{ arcmin}^2$) remnant, G150.3+4.5, whereas [Gerbrandt et al. \(2014\)](#) proposed part of this as a smaller ($61 \times 18 \text{ arcmin}^2$) remnant, G150.8+3.8. Recently [Ackermann et al. \(2018\)](#) showed the extended γ -ray emission from much of G150.3+4.5, confirming it as a SNR.
- G53.4+0.0 was confirmed as a SNR by [Driessen et al. \(2018\)](#), from radio and X-ray observations. This is one of the several candidate SNRs in this region (e.g. [Anderson et al. 2017](#)). See also [Dokara et al. \(2018\)](#).

In this version of the catalogue, five objects previously listed as SNRs have been removed, namely (G20.4+0.1, G21.5–0.1, G23.6+0.3, G59.8+1.2 and G65.8–0.5), as they have been identified as H II regions by [Anderson et al. \(2017\)](#). Also, G192.8–1.1 has been removed, as [Gao et al. \(2011\)](#) showed that this is not a SNR (see also, [Kang et al. 2014](#)). Erroneously it was not removed from the 2014 version of the catalogue. Note that G358.1+1.0 was erroneously labelled G358.1+0.1 in the 2009 and

2014 versions of the catalogue, which has now been corrected.

4. Discussion

4.1 Some simple statistics

There are 21 Galactic SNRs which do not have a flux density at 1 GHz in the catalogue. This is because either the remnant has not been detected at radio wavelengths, or it is poorly defined by current radio observations, so that their flux density at 1 GHz cannot be determined with any confidence: i.e. 93% of the remnants do have a flux density at 1 GHz in the catalogue. Of the catalogued remnants, $\approx 42\%$ are detected in X-rays, and $\approx 31\%$ in the optical. The smaller proportion of SNR identified in the optical and X-ray wavebands is due to Galactic absorption, which hampers the detection of distant remnants.

In this version of the catalogue, 80% of remnants are classified as shell (or possible shell) remnants, 13% are composite (or possible composite) remnants, and just 3% are filled-centre (or possible filled centre) remnants. The types of the remaining remnants are not clear from current observations (or else they are objects which are conventionally regarded as SNRs although they do not fit well into any of the conventional types, e.g. CTB80 (=G69.0+2.7), MSH 17–39 (=G357.7–0.1).

4.2 Selection effects

In previous papers (e.g. [Green 1991, 2005](#)), the selection effects that apply to the identification of Galactic SNRs were discussed. Although some SNRs are identified first at other than radio wavelengths, most SNRs have been identified first in the radio. The selection effects for the SNR catalogue are therefore dominated by those that apply at radio wavelengths. These are: (i) the difficulty in finding low surface brightness remnants, and (ii) the difficulty in finding small angular size remnants.

In [Green \(2005\)](#), a surface brightness completeness limit of $\Sigma \approx 10^{-20} \text{ W m}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$ at 1 GHz was derived. This nominal completeness limit is supported by various searches for SNRs, as no remnants with a surface brightness above this limit have been added to the published versions of the catalogue since [Green \(2009\)](#). [Xu et al. \(2013\)](#) used multi-frequency radio observations to separate thermal and non-thermal radio emissions in a large region around Cygnus X ($66^\circ \leq l \leq 90^\circ$, $|b| < 4^\circ$). They did not find any new large SNRs with $\Sigma > 0.37 \times 10^{-20} \text{ W m}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$,

consistent with the previously quoted completeness limit. More recently, [Anderson et al. \(2017\)](#) have identified many candidate SNRs in the region $17^{\circ}5 < l < 67^{\circ}4$, $|b| \leq 1^{\circ}25$, from THOR ([Beuther et al. 2016](#)) and VGPS ([Stil et al. 2006](#)) radio continuum observations at 1.4 GHz and mid-IR surveys. This covers a large fraction of the inner Galaxy – where most Galactic SNRs are expected to be – but only 2 of these candidates appear to have a surface brightness at 1 GHz above $10^{-20} \text{ W m}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$ (assuming a spectral index of 0.5 to scale the observed 1.4 GHz flux density to 1 GHz).

This surface brightness completeness limit of $10^{-20} \text{ W m}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$ was used in [Green \(2015\)](#) to select a sample of 69 SNRs from the 2014 version of the catalogue. This sample was then used to derive constraints on the distribution of remnants with Galactocentric radius. Of the six new remnants added to the catalogue since the 2014 version, only one (G53.4+0.0) has an integrated flux density at 1 GHz, and it is fainter than $\Sigma \approx 10^{-20} \text{ W m}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$. The other five are either not detected in the radio (G70.0–21.5) or do not currently have integrated radio flux densities as they are faint. Of the objects removed from the catalogue since 2014, two had a surface brightness of 1.2×10^{-20} and $2.1 \times 10^{-20} \text{ W m}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$ (for G20.4+0.1 and G23.6+0.3 respectively). So, the current catalogue contains 67 remnants with a surface brightness above $10^{-20} \text{ W m}^{-2} \text{ Hz}^{-1} \text{ sr}^{-1}$. This is two less than in the sample used in [Green \(2015\)](#), so that the results derived there will not be significantly changed.

Small angular size remnants – which will be the young but distant SNRs in the Galaxy – need to be resolved, for their structure to be recognized. Most wide-field radio surveys have not had small enough resolutions to easily identify such small angular size remnants. There are only 9 SNRs with angular diameters ≤ 3 arcmin in the catalogue, and none of these have been added to the catalogue since the 2009 version of the catalogue.

4.3 Recently proposed SNRs

As noted above, the detailed version of the catalogue includes notes on many objects that have been reported in the literature as possible or probable SNRs. Some of the recently proposed SNR candidates are discussed here in more detail:

- [Demetroullas et al. \(2015\)](#) suggested that a region of radio emission, which they label NGC 6334D, might be a SNR. This region, near $l = 351^{\circ}6$, $b = 0^{\circ}2$ was identified from their 31-GHz




Figure 1. MAGPIS ([Helfand et al. 2006](#)) 1.4 GHz image, in Galactic coordinates, of the candidate SNR G38.83–0.01 reported in [Anderson et al. \(2017\)](#). Contour levels are $\pm 0.8, 1.6, 2.4, 3.2, 4.0 \text{ mJy beam}^{-1}$ (with the negative contours dashed). This image is observations made with the Very Large Array (VLA), in B, C and D configurations, with a resolution of approximately 6 arcsec.

observations (with a resolution of ~ 4.5 arcmin), apparently with a non-thermal radio spectrum. However, other available observations of this region do not support a SNR identification for NGC 6334D. [Demetroullas et al. \(2015\)](#) noted that there are two sources in the NRAO VLA Sky Survey (NVSS, [Condon et al. 1998](#), at 1.4 GHz with a resolution of 45 arcsec) in the region of NGC 6334D, with peaks at 2.1 and 2.0 Jy beam^{-1} . Each of these sources have integrated flux densities of about 3.8 Jy in the NVSS catalogue, and other observations (e.g. [Murphy et al. 2007](#)) show they have relatively flat radio spectra. They are each associated with one or more compact H II regions identified by [Giveon et al. \(2005\)](#), from higher resolution 5-GHz and IR observations. The NVSS sources are separated by about 4 arcmin, and – with flat radio spectra – explain the emission of NGC 6334D seen in [Demetroullas et al. \(2015\)](#)’s lower resolution 31-GHz image. Higher quality 1.4-GHz observations from the SGPS ([Haverkorn et al. 2006](#)) do not show any obvious emission in this region – apart from that of the NVSS sources – that might indicate a SNR.

- A sample of ‘giant radio sources’ identified in the NVSS from pattern recognition techniques

is presented by Proctor (2016). One of these sources, NVGRC J205051.1+312728 – which is annotated with ‘SNR?’ as one of the several possibilities – is actually part of the Cygnus Loop (=G74.0–8.5, e.g. Green 1990). Several other of these sources also correspond to known SNRs, including other parts of the Cygnus Loop.

- As noted above, Anderson *et al.* (2017) identified many SNR candidates, which include several very small objects, 6 having a radius of ≤ 1 arcmin. If these were SNRs, then they would have to be physically small, even if in the distant Galaxy, and so would be scientifically interesting. Higher resolution radio images are available from the Multi-Array Galactic Plane Imaging Survey (MAGPIS, Helfand *et al.* 2006)² for several of these, but none of these look like young SNRs. In particular, G38.83–0.01 – which was reported as having a radius of 0.6 arcmin and a flux density at 1.4 GHz of 10 mJy – is clearly resolved into two compact sources, see Figure 1, so is not a young SNR.
- Dzib *et al.* (2018) presented observations of a small (only about 15 arcsec in extent) radio shell, which they suggested may be a SNR. If this was a SNR, it would have to be very young, given its small angular size, even younger than the youngest known Galactic SNR G1.9+0.3 (e.g. Green *et al.* 2008; Reynolds *et al.* 2008). However, this source has already been identified as a candidate PN by Froebrich *et al.* (2015).

Acknowledgements

The author is grateful to colleagues for numerous comments and corrections to the various versions of the Galactic SNR catalogue. This research has made use of NASA’s Astrophysics Data System Bibliographic Services, and the SIMBAD database, operated at CDS, Strasbourg, France.

References

- Ackermann M. *et al.* 2018, ApJS, 237, 32
 Anderson L. D. *et al.* 2017, A&A, 605, A58
 Araya M. 2017, ApJ, 843, 12
 Beuther H. *et al.* 2016, A&A, 595, A32
 Boumis P., Mavromatakis F., Paleologou E. V., Becker W. 2002, A&A, 396, 225
- Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J. 1998, AJ, 115, 1693
 de Gasperin F., Evoli C., Brüggem M., Hektor A., Cardillo M., Thorman P., Dawson W. A., Morrison C. B. 2014, A&A, 568, A107
 Demetroullas C. *et al.* 2015, MNRAS, 453, 2082
 Dokara R. *et al.* 2018, ApJ, 866, 61
 Driessen L. N., Domček V., Vink J., Hessels J. W. T., Arias M., Gelfand J. D. 2018, ApJ, 860, 133
 Dzib S. A., Rodríguez L. F., Karuppusamy R., Loinard L., Medina S.-N. X. 2018, ApJ, 866, 100
 Ferrand G., Safi-Harb S. 2012, AdSpR, 49, 1313
 Fesen R. A., Neustadt J. M. M., Black C. S., Koepfel A. H. D. 2015, ApJ, 812, 37
 Froebrich D. *et al.* 2015, MNRAS, 454, 2586
 Gao X. Y., Han J. L. 2014, A&A, 567, A59
 Gao X. Y., Han J. L., Reich W., Reich P., Sun X. H., Xiao L. 2011, A&A, 529, A159
 Gerbrandt S., Foster T. J., Kothes R., Geisbüsch J., Tung A. 2014, A&A, 566, A76
 Giveon U., Becker R. H., Helfand D. J., White R. L. 2005, AJ, 129, 348
 Green D. A. 1984, MNRAS, 209, 449
 Green D. A. 1988, Ap&SS, 148, 3
 Green D. A. 1990, AJ, 100, 192
 Green D. A. 1991, PASP, 103, 209
 Green D. A. 2004, BASI, 32, 335
 Green D. A. 2005, MmSAI, 76, 534
 Green D. A. 2009, BASI, 37, 45
 Green D. A. 2014, BASI, 42, 47
 Green D. A. 2015, MNRAS, 454, 1517
 Green D. A., Reynolds S. P., Borkowski K. J., Hwang U., Harrus I., Petre R. 2008, MNRAS, 387, L54
 Green A. J., Reeves S. N., Murphy T. 2014, PASA, 31, e042
 Haverkorn M., Gaensler B. M., McClure-Griffiths N. M., Dickey J. M., Green A. J. 2006, ApJS, 167, 230
 HESS Collaboration: Abdalla H. *et al.* 2018, A&A, 612, A8
 Helfand D. J., Becker R. H., White R. L., Fallon A., Tuttle S. 2006, AJ, 131, 2525
 Kang J.-H., Koo B.-C., Byun D.-Y. 2014, JKAS, 47, 259
 Kothes R., Reich P., Foster T. J., Reich W. 2017, A&A, 597, A116
 Murphy T., Mauch T., Green A., Hunstead R. W., Piestrzynska B., Kels A. P., Sztajer P. 2007, MNRAS, 382, 382
 Perley R. A., Butler B. J. 2017, ApJS, 230, 7
 Proctor D. D. 2016, ApJS, 224, 18
 Reynolds S. P., Borkowski K. J., Green D. A., Hwang U., Harrus I., Petre R. 2008, ApJ, 680, L41
 Rho J., Petre R. 1998, ApJ, 503, L167
 Stephenson F. R., Green D. A. 2002, Historical supernovae and their remnants, Oxford University Press
 Stil J. M. *et al.* 2006, AJ, 132, 1158
 Xu W. F., Gao X. Y., Han J. L., Liu F. S. 2013, A&A, 559, A81

²See also: <https://third.ucllnl.org/gps/>.