

SCIENTIFIC DATA

OPEN

SUBJECT CATEGORIES

- » Population dynamics
- » Ecological epidemiology
- » Genetic variation

Data Descriptor: Bat trait, genetic and pathogen data from large-scale investigations of African fruit bats, *Eidolon helvum*

Alison J. Peel^{1,2,3}, Kate S. Baker^{1,2,4,5}, David T.S. Hayman^{1,2,6}, Richard Suu-Ire^{7,8}, Andrew C. Breed⁹, Guy-Crispin Gembu¹⁰, Tiziana Lembo¹¹, Andrés Fernández-Loras^{2,12}, David R. Sargan¹, Anthony R. Fooks⁹, Andrew A. Cunningham² & James L.N. Wood¹

Received: 10 May 2016

Accepted: 19 May 2016

Published: 1 August 2016

Bats, including African straw-coloured fruit bats (*Eidolon helvum*), have been highlighted as reservoirs of many recently emerged zoonotic viruses. This common, widespread and ecologically important species was the focus of longitudinal and continent-wide studies of the epidemiological and ecology of Lagos bat virus, henipaviruses and Achimota viruses. Here we present a spatial, morphological, demographic, genetic and serological dataset encompassing 2827 bats from nine countries over an 8-year period. Genetic data comprises cytochrome *b* mitochondrial sequences ($n = 608$) and microsatellite genotypes from 18 loci ($n = 544$). Tooth-cementum analyses ($n = 316$) allowed derivation of rare age-specific serologic data for a lyssavirus, a henipavirus and two rubulaviruses. This dataset contributes a substantial volume of data on the ecology of *E. helvum* and its viruses and will be valuable for a wide range of studies, including viral transmission dynamic modelling in age-structured populations, investigation of seasonal reproductive asynchrony in wide-ranging species, ecological niche modelling, inference of island colonisation history, exploration of relationships between island and body size, and various spatial analyses of demographic, morphometric or serological data.

Design Type(s)	epidemiological study • genotyping design • population genetics analysis objective • individual genetic characteristics comparison design
Measurement Type(s)	genetic characteristics information • viral infection
Technology Type(s)	genotyping assay • serum neutralization of viral infectivity assay • DNA sequencing
Factor Type(s)	
Sample Characteristic(s)	<i>Eidolon helvum</i> • plagiopatagium • blood • Ilha de Sao Tome • Accra • Kisumu District • Bondo District • Mandra District • Kumasi • Techiman Municipal District • City of Kisangani • Kasanka National Park • Blantyre City • Dar es Salaam Region • Morogoro • Municipality of Jinja • City of Kampala • Ilha da Principe • Annobon Island • Municipality of Malabo • Bioko Island • Rio Muni

¹Department of Veterinary Medicine, University of Cambridge, Cambridge CB3 0ES, UK. ²Institute of Zoology, Zoological Society of London, Regent's Park, London NW1 4RY, UK. ³Environmental Futures Research Institute, Griffith University, Brisbane, Queensland 4111 Australia. ⁴Wellcome Trust Sanger Institute, Hinxton CB10 1SA, UK. ⁵Institute for Integrative Biology, University of Liverpool, Liverpool L69 7ZB, UK. ⁶Molecular Epidemiology and Public Health Laboratory, Hopkirk Research Institute, Massey University, Private Bag, 11 222, Palmerston North 4442, New Zealand. ⁷Wildlife Division, Ghana Forestry Commission, Accra, Ghana. ⁸University of Ghana, Faculty of Animal Biology and Conservation Science, Box LG 571, Legon, Accra, Ghana. ⁹Animal and Plant Health Agency (APHA), Addlestone, Surrey KT15 3NB, UK. ¹⁰Faculté des Sciences, Université de Kisangani, 4, Avenue Kithima, commune Makiso, BP 2012, Kisangani, République Démocratique du Congo. ¹¹Institute of Biodiversity, Animal Health and Comparative Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow, Graham Kerr Building, Glasgow, G12 8QQ, Scotland. ¹²Museo Nacional de Ciencias Naturales, CSIC, José Gutiérrez Abascal 2, Madrid 28006, Spain. Correspondence and requests for materials should be addressed to A.J.P. (email: alisonpeel@gmail.com).

Background & Summary

The straw-coloured fruit bat (*Eidolon helvum*) is a common, widely distributed, migratory species, occurring across sub-Saharan Africa and some offshore islands (Fig. 1)^{1,2}. Since 2007, investigations into the epidemiology and ecology of zoonotic viral infections in *E. helvum* have been undertaken via longitudinal sampling of wild populations in Ghana. Complementing this, between 2008–2011 and in 2014, cross-sectional sampling events were undertaken to determine the genetic population structure of *E. helvum*, and to assess whether the serological findings in Ghana were representative across the species' range (Fig. 1).

Four viruses were the focus of our serological surveys in *E. helvum* bats: Lagos bat virus (LBV), African henipaviruses, Achimota virus 1 (AchPV1) and Achimota virus 2 (AchPV2). Lagos bat virus is one of at least 15 known species in the *Lyssavirus* genus³ and has been isolated from *E. helvum* on multiple occasions^{4,5}. An African henipavirus is still yet to be isolated, however a full genome sequence has been obtained (putative name: African bat henipavirus Eid_hel/GH-M74a/GHA/2009 (M74))⁶. Achimota viruses 1 and 2 are closely related rubulaviruses for which serological evidence suggestive of spillover to humans in Africa exists⁷.




Figure 1. Map showing location of *E. helvum* sampling locations for genetic and serological analyses. Shading represents the distribution range of *E. helvum*. Sampling locations are numbered as in Table 1 (available online only). Adapted with permission from Mickleburgh *et al.*⁴³ and Peel *et al.*¹².

The specific aims of the data collection were to:

- Investigate whether antibodies to LBV, henipaviruses and Achimota viruses are present in *E. helvum* across its continental and island range, and to explore the antibody dynamics where possible.
- Describe the genetic metapopulation structure of *E. helvum* using a combination of mitochondrial (mtDNA) and microsatellite markers.
- Gather information on *E. helvum* distribution and seasonal patterns of reproduction.
- Combine results from these multidisciplinary studies to make inferences about virus transmission dynamics, and ultimately make inferences on the spillover risk to human populations.

Samples in this dataset are from 2827 bats from nine countries over an 8-year period (Fig. 2, Table 1 (available online only)). Raw data comprises spatial (roost location), seasonal (timing of sampling and seasonal birth pulses), morphological (forearm length, body weight), demographic (age, sex, reproductive status, mother-offspring relationships) and identification (individually numbered thumb-band) components. Data generated includes genetic characterisation (mtDNA sequencing and microsatellite genotyping) and serological assay results (for LBV, henipaviruses and Achimota viruses).

Multiple publications have arisen from these data, however many aspects remain unexplored. Demographic analyses have estimated birth and survival rates^{8,9}, and explored the effect of hunting on the latter⁹. Variations in roost composition have suggested a fission-fusion social structure⁹. Serological analyses have identified: the presence of antibodies against LBV, henipaviruses and Achimota viruses in *E. helvum* in Ghana^{7,10,11} and more broadly across the species' range, including isolated off-shore islands^{7,12,13}; that these viruses circulate endemically in *E. helvum* in Ghana, with evidence of horizontal transmission^{7,14,15}; and that *E. helvum* bats previously infected with LBV can have long-survival post infection¹⁶. The henipavirus dataset was used to develop a Bayesian method to determine appropriate cutoffs for serological assays¹⁷. Population genetic analyses identified that *E. helvum* are panmictic across their continental range, but that genetically isolated populations exist on isolated islands¹².

Other publications arising from these samples, but based on analyses not included here, include the development of a universal real-time assay and a pseudotype neutralisation assay for Lyssaviruses^{18,19}, microsatellite loci characterisation²⁰, estimation of divergence times between *Eidolon* sister species²¹, inference of movement ecology based on stable isotope ratios²², demonstration of Ebola antibodies^{16,23}, identification of multiple novel viruses²⁴, and novel Bartonella species in bat flies collected from *E. helvum*^{25,26}.




Figure 2. Sampling intensity per month, by country. Red represents high sampling intensity and the numbers of samples collected per month is recorded within each grid cell. Records in the database with unknown collection date are not represented here (nine from Annobón, seven from Bioko, 14 from Príncipe, 10 from Rio Muni and 13 from São Tomé).

This dataset contributes a substantial volume of data on the ecology of *E. helvum* and its viruses and will be valuable for a wide range of studies. In particular, an age-specific dataset such as this is rare and valuable for wildlife, especially bats. Further analyses could include viral transmission dynamic modelling in age-structured populations, including the use of cutting-edge Bayesian approaches to address complex epidemiological questions²⁷; time-series analyses on 5 years of wild henipavirus serological data from the same study site in Ghana ($n=1486$ data points), investigation of seasonal reproductive asynchrony in wide-ranging species; ecological niche modelling; inference of island colonisation history, exploration of relationships between island and body size; and various spatial analyses of demographic, morphometric or serological data. Field samples (e.g. serum, blood cells, urine, skin samples) and extracted DNA from individual bats in this dataset exist in storage and the authors are open to collaborative requests to undertake further analyses.

Methods

These methods are expanded and modified versions of descriptions in our previous publications, as cited in each section below. All associated data can be found in ‘Eidolon helvum data 2007–2014.csv’ [Data Citation 1].

Capture and Data Collection

Capture and sampling information has been described previously e.g. (refs 10,12). Sampling locations comprised 13 *E. helvum* roosting sites in continental Africa, and 14 in the four main islands in the Gulf of Guinea (Fig. 1, further detail in Table 1 (available online only)). In the majority of locations, data are from a single sampling event (sometimes comprising multiple sampling sessions within a one month period). Repeated sampling was conducted in Ghana (multiple sampling events per year over four years), Tanzania (one sampling event per year over two years) and Annobón (three sampling events over 4 years) as these locations were the focus of specific research studies. All fieldwork was undertaken under permits granted by national and local authorities (listed in Acknowledgements) and under ethics approval from the Zoological Society of London Ethics Committee (WLE/0489 and WLE/0467), using field protocols which followed ASM guidelines²⁸. Bats were captured at the roost with mist nets (6–18 m; 38 mm) as they departed the roost site at dusk, or returned at dawn. Except for a proportion of bats that were euthanased for virological studies ($n=238$), bats were released following sampling. Additional samples and data were obtained from other research groups ($n=152$) and in collaboration with local hunters in São Tomé ($n=102$), where bats are hunted for human consumption.

Personal protective equipment (long clothing, face masks, eye protection and gloves) was worn during sample collection. Morphometric and demographic details were recorded from bats under manual restraint. Female reproductive status was assigned as non-reproductive, pregnant, or lactating, according to the descriptions provided in Table 2. The phase in the reproductive cycle (i.e. the time in months between the sampling date and the beginning of the last birthing season) was estimated based on published data and the pregnancy status of females (foetal size, assuming a true gestation period of 4 months (Mutere 1965)) or degree of juvenile development during sampling.

Age was assessed by morphological characteristics (Table 3) and all individuals were placed into one of four age classes: Neonate (N; <2mths), Juvenile (J; 2–<6 months), Sexually Immature (SI; 6–<24 months) or Adult (A; ≥24 months). For a subset of samples, the timing of sampling in relation to the birthing season permitted further classification of SI individuals into 6-month age groups SI.1, SI.2 and SI.3 (6–<12, 12–<18, 18–<24 months, respectively). Additionally, for bats that were hunted or euthanased following capture, upper canine teeth were extracted, air dried and shipped to the USA (Matson’s laboratory, USA) for histological examination to assess the number of tooth cementum annuli present^{29,30}. Following previous studies³¹, it was assumed that each observed cementum layer represented

Phase	Reproductive status	Abbreviation	Description
1–2	Lactating	L	Neonate attached and suckling, or milk able to be expressed from mammary glands of females
3–8	Non-reproductive	NR	Not detectably pregnant on abdominal palpation
9	Very early pregnant	VEP	Uterine bulge palpable on abdominal palpation, up to 1cm diameter (detectable from ~0.5 cm diameter with careful palpation)
10	Early pregnant	EP	Uterine bulge palpable on abdominal palpation, 1–2 cm diameter
11	Mid pregnant	MP	Uterine bulge and foetus palpable on abdominal palpation, but no obvious abdominal distension of the female (2–~3.5 cm)
12	Late pregnant	LP	Foetus palpable on abdominal palpation, with distension of the female’s abdomen so that it is wider than the base of the ribs

Table 2. Reproductive status classifications for female *E. helvum*. Phase represents the estimated time in months since the beginning of the last birthing season.

Age class		Abbr.	Age (mths)	Timing of presence	Classification features
Standard age classes	Extended age classes				
Neonate	Neonate		< 2	Up to 2 months after birth pulse	Suckling
Juvenile	Juvenile	J	2– < 6	2–6 months after birth pulse	Small body size
Sexually Immature	SI	6 – < 24		Adult size (or nearly adult size), but undeveloped nipples and genitalia compared with adults	
	Sexually Immature 1	SI.1*	6 – < 12	6–12 months after birth pulse	Early in this period, distinguishable from SI.3 by smaller body size, and from Adults by lack of sexual development. In the last 3 months, females expected to be pregnant, but with undeveloped nipples.
	Sexually Immature 2	SI.2	12 – < 18	Up to 6 months after birth pulse	Clearly distinguishable from juveniles by size and adults by lack of sexual development
	Sexually Immature 3†	SI.3*	18 – < 24	6–12 months after birth pulse	Early in this period, distinguishable from SI.1 by larger body size, and from Adults by lack of sexual development. In the last 3 months this is less clear, however females expected to be pregnant with their first offspring (primiparous), but with undeveloped nipples.†
Adult	Adult	A	≥ 24	All year	Full body size within ranges expected for an adult. Females have developed nipples, males with developed testicles.

Table 3. Age classification system used for *E. helvum* *SI.1 and SI.3 stages represent two different birth cohorts, overlapping in time. Where distinction between SI.1 and SI.3 was unclear during sampling and analysis of measurements, individuals were simply classified as ‘SI’ and omitted from extended age class analyses. †SI individuals become sexually active during the last 4 months of their second year, however the highly synchronous and seasonal nature of the birthing period means that it is simpler for individuals to be recognised as sexually mature only after the beginning of the birthing pulse. While pregnant SI.3 (primiparous) individuals are therefore not technically sexually ‘immature’, by classifying them as such, they can be differentiated from adult females (>2 years of age).




Figure 3. Histological sections of upper canine teeth from *E. helvum* for cementum age analysis (Giemsa stain). Photographs and captions courtesy of Gary Matson, Matson’s Laboratory, MT, USA. Each age estimation was scored with a certainty code: A: highest certainty of reported age, B: histological evidence supported a given age result ± 0.5 –1.5 years, or C: tooth or section quality was too compromised to accurately age. (a) Bat ID 424. Cementum age 2, certainty code A. 100X. The tooth was in excellent histological condition, as indicated by the presence of periodontal membrane and good differential staining between annuli and light cementum. (b) Bat ID 62. Cementum age 6, certainty code A. 100X. Annuli are complex, with at least two components each year. A key feature of age analysis is resolving uncertainty about whether complex annuli or individual components are being used as age indicators. (c) Bat ID 44. Cementum age 13, certainty code B (13–15 yrs). 400X. The root tip of this tooth had been broken off during extraction. Missing cementum complicates age analysis, reducing the evidence available for evaluating whether annuli observed at one point may be clearly identifiable as components of complex annuli at another point.

one year. Each age estimation was scored with a certainty code: A: highest certainty of reported age (51% of samples, e.g. Fig. 3a,b), B: histological evidence supported a given age result ± 0.5 –1.5 years (46% of samples, e.g. Fig. 3c), or C: tooth or section quality was too compromised to accurately age (3% of samples).

Genetic and blood samples were collected under manual restraint. Wing membrane biopsies (4-mm) were placed into 70% alcohol. Up to 1 ml blood was collected from the propatagial vein using a citrated 1 ml syringe and placed into a plain 1.5 ml eppendorf tube.

Molecular methods

Molecular methods have been described previously^{12,20}. Genomic DNA was extracted from *E. helvum* tissues (predominantly wing membrane biopsies, but also liver and muscle samples, all stored in ethanol) using DNeasy Blood and Tissue Kits (QIAGEN Ltd., Crawley, West Sussex, UK). DNA was quantified using Quant-iT PicoGreen dsDNA kits (Molecular probes, UK), and later using a Nanodrop ND-1000 Spectrophotometer (Thermo Fisher Scientific, UK) and diluted to a standard concentration.

Twenty *E. helvum* loci developed in a previous study²⁰ were quality-checked using a subset of samples. Loci E and Ae were discarded due to difficulty in scoring or high error rates and data from locus Ag were re-binned and re-scored, correcting earlier issues with allelic dropout. In total, 170 continental and 385 island samples were run as multiplex PCRs at 18 loci (TSY, FWB, MNQX, AgPK, AcAfAi, AdAh) in 10 µl PCRs, containing 4ng template DNA, 0.2 µM of each primer, and 5 µl Type-it Multiplex PCR Master Mix (QIAGEN Ltd.). Positive and negative controls were included on each plate and amplification was performed using the following conditions: 5 min at 94 °C; 30 cycles of 30 s at 95 °C, 90 s at 57 °C, and 30 s at 72 °C; then 30 s at 60 °C. Genotyping was performed by capillary electrophoresis using a Beckman CEQ 8000 (Beckman, UK). Allele sizes were scored automatically prior to manual verification. Genotyping data from 18 loci are provided in 'Eidolon helvum data 2007–2014.csv' [Data Citation 1]. Loci B has previously been identified as being X-linked²⁰.

Fragments of the mitochondrial DNA cytochrome b (cytb) gene were amplified from continental samples by PCR using the generic primers L14722 (5'-CGA AGC TTG ATA TGA AAA ACC ATC GTT G)³² and H15149 (5'-AAA CTG CAG CCC CTC AGA ATG ATA TTT GTC CTC A)³³ in 20 µl reactions, containing 0.1–1 ng template DNA, 0.2 µM of each primer, 0.25 mM of each dNTP, 1.5 mM MgCl₂, 0.25 µl of *Taq* polymerase (Invitrogen), and 0.2 µl 10X reaction buffer and with the following conditions: 5 min at 94 °C; 40 cycles of 1 min at 93 °C, 1 min at 54 °C, and 2 min at 72 °C; then 7 min at 72 °C. Although these generic primers were adequate with continental samples (8% PCR failure), amplification from isolated Gulf of Guinea island samples was less successful (48% PCR failure). Shortened primers (EhM2814 (5'-GCT TGA TAT GAA AAA CCA TCG TTG) and EhM2815 (5'-CAG CCC CTC AGA ATG ATA TTT GT) resulted in successful amplification when using Microzone MegaMix-Gold reagent (Microzone Ltd, UK). PCRs were performed in 20 µl reactions, containing 2 ng template DNA, 0.25 µM of each primer, and 10 µl MegaMix-Gold, using the following conditions: 5 min at 95 °C; 33 cycles of 30 sec at 95 °C, 30 sec at 53 °C, and 45 sec at 72 °C. PCR products were checked by gel electrophoresis on 1% agarose gels, purified using Exosap-IT clean-up (USB Europe, Germany) and sequenced in both directions on an ABI 3730xl DNA Analyser, (Applied Biosystems). Paired sequences were edited and aligned using the STADEN Package v1.6 (ref. 34). Multiple sequence alignment was performed using default settings in T-COFFEE³⁵. Sequences were checked manually and trimmed to a standard length (397 bp) in JALVIEW v2 (ref. 36). No sequence differences were detected in 38 samples sequenced using both primer pairs, so data were combined.

Data from 608 and 544 individuals is available for cytb and microsatellite analyses (at 18 loci), respectively (Table 1 (available online only)).




Figure 4. Correlation between $\ln(\text{MFI})$ values pre- and post-repair of the Luminex machine used to run the assays, based on 293 samples. The linear regression line is in red. ($R^2 0.81$, F-statistic: 1306 (1, 296), $P < 2.2\text{e-}16$).

MEASURES ADOPTED TO REDUCE AND DETECT ERRORS
Sampling
• Standard protocols followed for collection and labelling samples • Sampling information logged on paper sampling sheets, and retained as backup • Sampling data transferred to a single sampling database
DNA extraction
• Inclusion of negative controls to monitor contamination • Consistency in extraction protocol • Extraction performed in a different room to downstream analyses
DNA amplification
• Inclusion of negative control to monitor contamination • Inclusion of positive control as a reference sample • Consistency in amplification protocol • Replicate amplifications - by main user, and also another user (blinded)
Sequencing
• Inclusion of negative control to monitor contamination • Inclusion of positive control as a reference sample • Consistency in sequencing protocol • Sequencing in both forward and reverse directions • Replicate sequencing - including multiple samples individually repeated, and a single sample repeated many times
Sequencing Analyses
• Alignment of forward and reverse sequences, with automated and manual checking of inconsistencies • Alignment against GenBank database • Multiple alignment - manual checking of all polymorphisms
Genotyping
• Perform pilot study to assess suitability of new loci for larger study • Automated scoring, with manual checking • Inclusion of positive control as a reference sample • Discard low-quality samples • Inclusion of a locus invariant in 99% of samples to monitor inter-assay variability of CEQ allele migration • Cross-reading of datasets, and repeat scoring of overall dataset if required
Genotyping Analyses
• Discard poor markers from pilot study • Quantify overall genotyping error rate and assess acceptability • Consider genotyping errors as a possible cause of Hardy-Weinberg or linkage disequilibrium

Table 5. Measures adopted to minimise and allow assessment of errors which may occur during the sampling, DNA extraction, amplification, sequencing, genotyping and data analysis processes (adapted from Bonin *et al.* 2004; Table 4).

Serological analyses

Serological methods have been described previously^{7,10–14}. A modified fluorescent antibody virus neutralization (mFAVN) assay using the LBNig56 isolate was used to detect neutralising antibodies against LBV^{10,37}. Samples were tested in duplicate using threefold serial dilutions (representing reciprocal titres of 9, 27, 81, and 243–19,683). Human rabies immunoglobulin, LBV-positive rabbit serum, and rabies-vaccinated mouse serum were used as positive controls and negative rabbit and mouse serum were negative controls. Titres were considered positive at IC100 endpoint reciprocal dilutions >1: 9 (100% neutralisation of virus). Henipavirus antibodies detected in African fruit bat samples using virus neutralisation assays, multiplexed microsphere assays and pseudotype assays developed to target other known henipaviruses (Hendra and Nipah viruses) and are presumed to represent cross-neutralisation or cross-reactivity¹². Here, Luminex multiplexed microsphere binding assays were used to detect antibodies against henipaviruses (HeV and NiV). In these assays, purified recombinant expressed henipavirus soluble G glycoproteins³⁸ are conjugated to internally coloured and distinguishable microspheres, allowing

	cytb original primers	cytb new primers
Number of samples	207	438
Number of samples repeat PCR and sequenced	28	1
Number of duplicates run per sample	1	70
Total number of repeats	28	70
% of sequences repeated	13.5%	16.0%
Number of base pairs in sequence	418	397
Number of repeated sequences	28	70
Number of base pairs compared	11704	27790
Number of sequence errors (mismatches)	1	0
% PCR/sequence error (No of mismatches/no bp compared)	0.01%	0%
% samples extracted in duplicate, cytb PCR and sequenced	2.4%	
Number of errors (mismatches)	0	
% error	0.0%	

Table 6. Mitochondrial DNA PCR repeat rates and sequencing error rates.

multiplexing. For African bat samples, stronger results were consistently observed in NiV binding assays and virus neutralisation tests¹³, so only NiV binding assay results are included in the dataset. Binding results are outputted as median fluorescence intensity (MFI) values of at least 100 microspheres for each virus type. In mid-December 2010, major repair work was undertaken on the Luminex machine being used for serological analysis. A subset of samples that had been analysed before the repairs were repeated to calibrate results ($n = 293$). MFI values pre- and post-repair work were significantly different, making the use of a single cutoff inappropriate¹⁷. Two approaches were taken to designate results as seropositive or seronegative. First, a Bayesian mixture model was applied as described in¹⁷. Cutoffs for pre- and post-repair work were determined so that samples above this cutoff were $\geq 99\%$ likely to be in the seropositive distribution (MFI = 156.1 and 127.5, respectively). Second, linear regression of pre- and post-maintenance $\ln(\text{MFI})$ values demonstrated a significant linear relationship (Fig. 4, $R^2 0.81$, F-statistic: 1306 (1, 296), $P < 2.2\text{e-}16$), and the variance decreases for higher MFI values (above the cutoff). Pre-maintenance MFI values were converted to post-maintenance values using the formula:

$$\text{NEW_MFI} = \exp(0.7795774 \cdot \ln(\text{OLD_MFI}) + 0.4392832).$$

The Bayesian mixture model was applied to this transformed and combined data using the same method. From this analysis, MFI values > 94.2 were $\geq 99\%$ likely to be in the seropositive distribution. Results from the two methods were compared and the second method resulted in the highest congruence between pre- and post-maintenance paired results (congruence in 266/298 samples versus 250/298 samples for the first method), and these data were therefore used in the final dataset. Raw MFI values are available on request.

Antibodies against Achimota viruses 1 and 2 were detected using virus neutralisation assays⁷, with all testing in duplicate. Samples were diluted to 1:20 and incubated with 200 TCID₅₀ of virus for 30 min at 37 °C prior to the addition of Vero cell suspension at an MOI equivalent to 0.01. Cell monolayers were assessed for evidence of virus neutralization 7 days post infection. Where sample volume permitted, positive samples were titrated in a 2-fold dilution series from 1: 20 to 1: 160 and retested using the same protocol.

Data Records

The data are contained in a single comma-separated file (.csv format), entitled ‘Eidolon helvum data 2007–2014’ (Data Citation 1). Each row below the header represents an individual bat ($n = 2,827$), and the columns ($n = 68$) contain sample identifier information, demographic and morphometric data, and results of genetic and serological assays. Full descriptions of the column titles are included in the Table 4 (available online only).

Technical Validation

Molecular analyses

Recommendations for minimisation and assessment of errors that may occur during the sampling, DNA extraction, amplification, sequencing, genotyping and data analysis processes were followed where possible³⁹ (Table 5).

As previously described²⁰, microsatellite loci were tested for evidence of departure from Hardy–Weinberg equilibrium (HWE) and genotypic disequilibrium using FSTAT 2.9 (ref. 40), with appropriate Bonferroni corrections for multiple testing. All loci were analysed in MICRO-

CHECKER⁴¹ to test for null alleles, stuttering and large allelic dropout as a cause of departure from HWE. Additionally, since Locus M displayed extremely low polymorphism (99.1% of individuals were homozygous for a particular allele), this locus was included in all PCR plates as a positive control and to determine inter-assay variability in allele fragment length. Error rates for microsatellite loci are reported in Peel *et al.*²⁰ Inter-assay genotyping variability, measured by the variation in fragment length of the dominant allele of locus M on each plate, was low (range 134.32 – 134.66) across 27 runs and two control samples. Loci Y proved difficult to confidently bin due to alleles of single nucleotide difference and was therefore not included in the dataset.

Error rates for cytb analyses were assessed by replicate extractions (performed on 2.4% of samples), replicate PCR and sequencing reactions (performed on 8–14% of extracted samples), and by inclusion of positive and negative controls for all extractions and PCRs. Poor quality mtDNA sequence traces were excluded. Background PCR and sequencing error rates of the new *E. helvum* cytb primers EhM2814 and EhM2815 were assessed by running 70 replicates of a single sample. PCR and sequencing error rates were calculated at the base-pair level. Sequencing error rate was negligible (0–0.01%) across samples repeated in duplicate, and no substitutions were observed in the 70 replicate sequences obtained from a single sample (Table 6).

Serological analyses

All serological assays included positive and negative controls. Samples were tested in duplicate (LBV and Achimota viruses) or with 100 replicates (henipaviruses). Further validation procedures for multiplexed microsphere binding assays are presented as part of the methods, above.

Usage Notes

Users of these data are advised that importing the.csv data file (Data Citation 1) into Microsoft Excel can result in formatting errors, particularly with the column ‘Teeth.Age.Range’. Rather than opening the file with Excel (by double-clicking, for example), it is suggested that users instead select ‘File>Import>csv file >Delimited’, then select the ‘Teeth.Age.Range’ column and set the column data format as ‘Text’. Alternatively, importing and processing the data into the software ‘R’⁴² may be preferable.

References

- DeFrees, S. L. & Wilson, D. E. *Eidolon helvum*. *Mammalian Species* **312**, 1–5 (1988).
- Bergmans, W. Taxonomy and biogeography of African fruit bats (Mammalia, Megachiroptera). 3. The genera *Scotonycteris* Matschie, 1894, *Casinycteris* Thomas, 1910, *Pteropus* Brisson, 1762, *Eidolon* Rafinesque, 1815. *Beaufortia* **40**, 111–177 (1990).
- Kuzmin, I. V. in *Current Laboratory Techniques in Rabies Diagnosis, Research and Prevention* (eds Rupprecht C. & Nagarajan T.1) 3–21 (Elsevier, 2014).
- Boulger, L. & Porterfield, J. Isolation of a virus from Nigerian fruit bats. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **52**, 421–424 (1958).
- Kuzmin, I. V. *et al.* Lagos bat virus in Kenya. *J Clin Microbiol* **46**, 1451–1461 (2008).
- Drexler, J. F., Corman, V. & Müller, M. Bats host major mammalian paramyxoviruses. *Nature Communications* **3**, 1–12 (2012).
- Baker, K. S. *et al.* Novel, Potentially Zoonotic Paramyxoviruses from the African Straw-Colored Fruit Bat *Eidolon helvum*. *Journal of Virology* **87**, 1348–1358 (2013).
- Hayman, D. T. S. *et al.* Demography of straw-colored fruit bats in Ghana. *J. Mammal* **93**, 1393–1404 (2012).
- Hayman, D. T. S. & Peel, A. J. Can survival analyses detect hunting pressure in a highly connected species? Lessons from straw-coloured fruit bats. *Biological Conservation* **200**, 131–139 (2016).
- Hayman, D. T. S. *et al.* Antibodies against Lagos bat virus in megachiroptera from West Africa. *Emerg Infect Dis* **14**, 926–928 (2008).
- Hayman, D. T. S. *et al.* Evidence of henipavirus infection in West African fruit bats. *PLoS ONE* **3**, e2739 (2008).
- Peel, A. J. *et al.* Continent-wide panmixia of an African fruit bat facilitates transmission of potentially zoonotic viruses. *Nature Communications* **4**, 3770 (2013).
- Peel, A. J. *et al.* Henipavirus neutralising antibodies in an isolated island population of African fruit bats. *PLoS ONE* **7**, e30346 (2012).
- Hayman, D. T. S. *et al.* Endemic Lagos bat virus infection in *Eidolon helvum*. *Epidemiol Infect* **140**, 2163–2171 (2012).
- Peel, A. J. The epidemiology of Lagos bat virus and henipaviruses in straw-coloured fruit bats (*Eidolon helvum*). 1–327 (University of Cambridge, 2012). at <http://www.dspace.cam.ac.uk/handle/1810/244571>.
- Hayman, D. T. S. *et al.* Long-term survival of an urban fruit bat seropositive for Ebola and Lagos bat viruses. *PLoS ONE* **5**, e11978 (2010).
- Peel, A. J. *et al.* Use of cross-reactive serological assays for detecting novel pathogens in wildlife: assessing an appropriate cutoff for henipavirus assays in African bats. *Journal of Virological Methods* **193**, 295–303 (2013).
- Hayman, D. T. S. *et al.* A universal real-time assay for the detection of Lyssaviruses. *Journal of Virological Methods* **177**, 87–93 (2011).
- Wright, E. *et al.* Virus neutralising activity of African fruit bat (*Eidolon helvum*) sera against emerging lyssaviruses. *Virology* **408**, 183–189 (2010).
- Peel, A. J., Rossiter, S. J., Wood, J. L. N., Cunningham, A. A. & Sargan, D. R. Characterization of microsatellite loci in the straw-colored fruit bat, *Eidolon helvum* (Pteropodidae). *Conserv Genet Resour* **2**, 279–282 (2010).
- Shi, J. J. *et al.* A deep divergence time between sister species of *Eidolon* (Pteropodidae) with evidence for widespread panmixia. *Acta Chiropterologica* **16**, 279–292 (2014).
- Ossa, G., Kramer-Schadt, S., Peel, A. J., Scharf, A. K. & Voigt, C. C. The movement ecology of the straw-colored fruit bat, *Eidolon helvum*, in sub-Saharan Africa assessed by stable isotope ratios. *PLoS ONE* **7**, e45729 (2012).
- Hayman, D. T. S. *et al.* Ebola Virus Antibodies in Fruit Bats, Ghana, West Africa. *Emerg Infect Dis* **18**, 1207–1209 (2012).
- Baker, K. S. *et al.* Metagenomic study of the viruses of African straw-coloured fruit bats: detection of a chiropteran poxvirus and isolation of a novel adenovirus. *Virology* **441**, 95–106 (2013).
- Billeter, S. A. *et al.* *Bartonella* species in bat flies (Diptera: Nycteribiidae) from western Africa. *Parasitology* **139**, 324–329 (2012).

26. Bai, Y., Hayman, D. T. S., McKee, C. D. & Kosoy, M. Y. Classification of Bartonella Strains Associated with Straw-Colored Fruit Bats (*Eidolon helvum*) across Africa Using a Multi-locus Sequence Typing Platform. *PLoS Negl Trop Dis* **9**, e0003478 (2015).
27. Viana, M. *et al.* Dynamics of a morbillivirus at the domestic-wildlife interface: Canine distemper virus in domestic dogs and lions. *PNAS* **112**, 1464–1469 (2015).
28. Sikes, R. S. & Gannon, W. L. The Animal Care and Use Committee of the American Society of Mammalogists. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *J Mammal* **92**, 235–253 (2011).
29. Matson, G. *et al.* A laboratory manual for cementum age determination of Alaska brown bear first premolar teeth (Matson's Laboratory, 1993).
30. Bodkin, J. L., Ames, J. A., Jameson, R. J., Johnson, A. M. & Matson, G. M. Estimating age of sea otters with cementum layers in the first premolar. *The Journal of Wildlife Management* **61**, 967–973 (1997).
31. Divljan, A., Parry-Jones, K. & Wardle, G. M. Age determination in the grey-headed flying fox. *J Wildlife Manage* **70**, 607–611 (2006).
32. Juste, J. *et al.* Phylogeography of African Fruitbats (Megachiroptera). *Mol Phylogenet Evol* **13**, 596–604 (1999).
33. Kocher, T. *et al.* Dynamics of Mitochondrial DNA Evolution in Animals: Amplification and Sequencing with Conserved Primers. *PNAS* **86**, 6196–6200 (1989).
34. Staden, R., Judge, D., Bonfield, J., Krawetz, S., Womble, D. in *Introduction to Bioinformatics. A Theoretical and Practical Approach* (eds Krawetz S. A. & Womble D. D.) 760 (Human Press, 2003).
35. Notredame, C., Higgins, D. & Heringa, J. T-coffee: a novel method for fast and accurate multiple sequence alignment. *Journal of Molecular Biology* **302**, 205–217 (2000).
36. Waterhouse, A., Procter, J., Martin, D., Clamp, M. & Barton, G. Jalview Version 2 - a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189–1191 (2009).
37. Brookes, S. M. *et al.* European bat lyssavirus in Scottish bats. *Emerg Infect Dis* **11**, 572–578 (2005).
38. Bossart, K. N. *et al.* Neutralization assays for differential henipavirus serology using Bio-Plex Protein Array Systems. *Journal of Virological Methods* **142**, 29–40 (2007).
39. Bonin, A. *et al.* How to track and assess genotyping errors in population genetics studies. *Mol Ecol* **13**, 3261–3273 (2004).
40. Goudet, J. FSTAT (Version 1.2): A computer program to calculate F-statistics. *J Hered* **86**, 485–486 (1995).
41. van Oosterhout, C., Hutchinson, W., Wills, D. & Shipley, P. Micro-Checker: software for identifying and correcting genotyping errors in microsatellite data. *Mol Ecol Notes* **4**, 535–538 (2004).
42. R Core Team. R: A language and environment for statistical computing. Version 3.0 (2014) at <http://www.R-project.org/>.
43. Mickleburgh, S., Hutson, A. M., Bergmans, W., Fahr, J. & Racey, P. *Eidolon helvum*, In: IUCN 2008. 2008 IUCN Red List of Threatened Species. Available from www.iucnredlist.org Downloaded on 6 February 2011. (2008).
44. Juste, J., Ibanez, C. & Machordom, A. Morphological and allozyme variation of *Eidolon helvum* (Mammalia: Megachiroptera) in the islands of the Gulf of Guinea. *Biol J Linn Soc* **71**, 359–378 (2000).

Data Citation

1. Peel, A. *et al.* Dryad Digital Repository. <http://dx.doi.org/10.5061/dryad.2fp34> (2016).

Acknowledgements

The authors thank the governments of Ghana, Tanzania, Malawi, Zambia, Uganda, Equatorial Guinea, and São Tomé and Príncipe for facilitating this research. For their invaluable support in planning and implementing the field work, we also thank the Wildlife Division of the Forestry Commission, and the Veterinary Services Directorate, Ghana; Zambian Wildlife Authority; Kasanka Trust; Malawi Ministry of Tourism, Wildlife and Culture, Malawi Ministry of Agriculture, Irrigation and Food Security; former Tanzania Ministry of Livestock and Fisheries Development (now Ministry of Agriculture, Livestock and Fisheries); Tanzania Wildlife Research Institute; Sokoine University of Agriculture, Tanzania; Makerere University, Uganda; Equatorial Guinea Ministerio de Agricultura y Bosques; Equatorial Guinea Instituto del Desarrollo Forestal y Gestión de las Áreas Protegidas; Universidad Nacional de Guinea Ecuatorial; São Tomé and Príncipe Ministério de Agricultura, Desenvolvimento Rural e Pesca; Ecosystèmes Forestiers d'Afrique Centrale; Associação Monte Pico, São Tomé and Príncipe; Iñaki Rodríguez-Prieto; Alex Torrance; Lucrecia Bilé Osa Ahara; Heidi Ruffler; Ricardo Castro César de Sá; Ricardo Faustino de Lima; Mariana Carvalho and Meyir Ziekah. Additional genetic samples from the Gulf of Guinea islands and Rio Muni were supplied by Javier Juste (Museo Nacional de Ciencias Naturales, CSIC) and DJ Long (Californian Academy of Sciences). Additional sera were supplied by Ivan Kuzmin (previously Centers for Disease Control and Prevention). Christopher Broder (Uniformed Services University of the Health Sciences) supplied HeV and NiV soluble G glycoprotein for the henipavirus serological assays. The authors are indebted to Linfa Wang, Gary Cramer, Jennifer Barr and others at the CSIRO Australian Animal Health Laboratory, Geelong, Australia and Daniel Horton and David Seldon at the Animal Health and Veterinary Laboratories Agency (now Animal and Plant Health Agency), UK for providing invaluable support, constructive discussions and technical assistance throughout the project.

Funding for this study was provided by the Cambridge Infectious Diseases Consortium (grant VT0105) (A.J.P., D.T.S.H., K.S.B.), The Charles Slater Trust (A.J.P.), Zebra Foundation for Veterinary Zoological Education (A.J.P.), Isaac Newton Trust (A.J.P.), the Wellcome Trust (D.T.S.H., K.S.B.), a David H. Smith postdoctoral fellowship (D.T.S.H.), the RAPIDD program of the Science and Technology Directorate, Department of Homeland Security, Fogarty International Center, National Institutes of Health (D.T.S.H., J.L.N.W., A.R.F.), The Alborada Trust (J.L.N.W.), a European Union FP7 project ANTIGONE (Anticipating Global Onset of Novel Epidemics 278976) (A.C.B., J.L.N.W., A.A.C., A.R.F.), Royal Society Wolfson Research Merit Award (A.A.C.), Lincoln Park Zoo, Chicago, US (T.L.) and a Wellcome Trust Value in People award through the University of Glasgow (T.L.). KSB is in receipt of a Wellcome Trust Postdoctoral Training fellowship for clinicians (106690/Z/14/Z) and is supported by Wellcome Trust grant number 098051.

Author Contributions

A.J.P. conceived the research, conducted fieldwork, conducted genetic analyses, conducted serological assays, curated data file, and wrote the manuscript. J.L.N.W. conceived the research. K.S.B. conceived the research, conducted fieldwork, conducted serological assays, and edited the manuscript. A.C.B. conceived the research, and conducted fieldwork. G.C.G. conducted fieldwork. T.L. conducted fieldwork and edited the manuscript. R.S. conducted fieldwork. A.A.C. conceived the research. D.T.S.H. conceived the research, conducted fieldwork, conducted serological assays. A.F.L. conducted fieldwork. D.R.S. conceived the research and edited the manuscript. A.R.F. conceived the research and conducted serological assays.

Additional Information

Tables 1 and 4 are only available in the online version of this paper.

Competing financial interests: The authors declare no competing financial interests.

How to cite: Peel, A. J. *et al.* Bat trait, genetic and pathogen data from large-scale investigations of African fruit bats, *Eidolon helvum*. *Sci. Data* 3:160049 doi: 10.1038/sdata.2016.49 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0>

Metadata associated with this Data Descriptor is available at <http://www.nature.com/sdata/> and is released under the CC0 waiver to maximize reuse.