

Childhood adversity, pubertal timing and self-harm: a longitudinal cohort study

Abigail Emma Russell¹, Carol Joinson², Elystan Roberts², Jon Heron², Tamsin Ford³, David Gunnell^{2,4}, Paul Moran^{2,4}, Caroline Relton⁵, Matthew Suderman⁶ and Becky Mars^{2,4}

¹ Children and Young People's Mental Health Research Collaboration, University of Exeter College of Medicine and Health

² Centre for Academic Mental Health, Population Health Sciences, University of Bristol Medical School

³ University of Cambridge Department of Psychiatry

⁴ NIHR Biomedical Research Centre at the University Hospitals Bristol NHS Foundation Trust and the University of Bristol.

⁵ MRC Integrative Epidemiology Unit, University of Bristol Medical School; Population Health Sciences, University of Bristol Medical School

Correspondence to: Abigail Emma Russell, South Cloisters, St Luke's Campus, University of Exeter
ae.russell@exeter.ac.uk - +44(0)1392 722985

Accepted version: Psychological Medicine 8/2/21

Abstract

Background: The occurrence of early childhood adversity is strongly linked to later self-harm, but there is poor understanding of how this distal risk factor might influence later behaviours. One possible mechanism is through an earlier onset of puberty in children exposed to adversity, since early puberty is associated with an increased risk of adolescent self-harm. We investigated whether early pubertal timing mediates the association between childhood adversity and later self-harm.

Methods: Participants were 6698 young people from a UK population-based birth cohort (ALSPAC). We measured exposure to nine types of adversity from 0-9 years old, and self-harm when participants were aged 16 and 21 years. Pubertal timing measures were age at peak height velocity (aPHV-males and females) and age at menarche (AAM). We used generalised structural equation modelling for analyses. **Results:** For every additional type of adversity; participants had an average 12-14% increased risk of self-harm by 16. Relative Risk estimates were stronger for direct effects when outcomes were self-harm with suicidal intent. There was no evidence that earlier pubertal timing mediated the association between adversity and self-harm (indirect effect RR 1.00, 95% CI 1.00, 1.00 for aPHV and RR 1.00 95% CI 1.00, 1.01 for AAM). **Conclusions:** A cumulative measure of exposure to multiple types of adversity does not confer an increased risk of self-harm via early pubertal timing, however both childhood adversity and early puberty are risk factors for later self-harm. Research identifying mechanisms underlying the link between childhood adversity and later self-harm is needed to inform interventions.

Funding: Medical Research Foundation and Medical Research Council (MR/R004889/1).

Keywords: ALSPAC, puberty, menarche, peak height velocity, adversity, ACEs, self-harm, suicide attempt, mediation

Introduction

Self-harm, defined in this study as intentionally hurting oneself in any way with or without suicidal intent, is a global public health concern. There is evidence that rates of self-harm are rising, and it is one of the strongest predictors of later suicide (Hawton et al., 2003; McManus et al., 2019). Understanding the causal antecedents of self-harm is important in reducing risk and developing effective treatments. One of the strongest risk factors for self-harm is experiencing adversity early in life (Björkenstam, Kosidou, and Björkenstam, 2017; Björkenstam, Kosidou, and Björkenstam, 2016; Cha et al., 2018; Hughes et al., 2017). Adverse childhood experiences (ACEs) are commonly conceptualised as indicators of abuse, child maltreatment or household dysfunction (Björkenstam et al., 2017; Björkenstam et al., 2016; Dube, Felitti, Dong, Giles, and Anda, 2003), including sexual, physical and emotional abuse as well as parental psychopathology and witnessing domestic violence. Studies exploring the role of ACEs in relation to health outcomes commonly consider the multiple types of adversity an individual has experienced, rather than focussing on individual adversities (Enns et al., 2006). Because adversities rarely occur in isolation and some, such as sexual abuse, are less common, it is difficult to disentangle the role of an individual adversity on child outcomes in population-based samples.

Understanding the pathways that link ACEs to an increased risk of later self-harm could improve the identification of those most at risk and may lead to novel intervention targets. However, the mechanisms linking ACEs to self-harm in adolescence and adulthood are unclear (Cha et al., 2018). One possible mechanism is through an earlier onset of puberty in children exposed to ACEs, since early puberty is associated with an increased risk of self-harm in adolescence and young adulthood in both sexes (Cha et al., 2018; Deng et al., 2011; Larsson and Sund, 2008; Michaud, Suris, and Deppen, 2006; Roberts, Fraser, Gunnell, Joinson, and Mars, 2020; Wichstrøm, 2000). There is some evidence that childhood adversity is associated with an earlier age at onset of puberty in both females and males (Henrichs et al., 2014; Lei, Beach, and Simons, 2018; Mendle, 2014) but empirical findings are mixed (Zhang, Zhang, and Sun, 2019). Adverse exposures that have been found to be associated with early pubertal timing in females include physical abuse, sexual abuse, parental mental illness, parent drinking, father absence, parental unemployment, frequent relocation and family violence (Arim, Tramonte, Shapka, Dahinten, and Willms, 2011; Barrios et al., 2015; Bogaert, 2005; Boynton-Jarrett and Harville, 2012; Boynton-Jarrett et al., 2013; Clutterbuck, Adams, and Nettle, 2015; Henrichs et al., 2014; Hulanicka, Gronkiewicz, and Koniarek, 2001; Kelly, Zilanawala, Sacker, Hiatt, and Viner, 2017; Li, Denholm, and Power, 2014; Magnus et al., 2018; Romans, Martin, Gendall, and Herbison, 2003). However, studies have also reported a later age at menarche for females affected by abuse, domestic violence, family conflict, and maternal alcohol abuse (Boynton-

Jarrett and Harville, 2012; Boynton-Jarrett et al., 2013; Li et al., 2014). In males, there is evidence that low parental education and father absence (Arim et al., 2011; Bogaert, 2005) are associated with earlier age at puberty.

Mechanisms and Intervention Potential

The developmental origins of health and disease (DoHAD) theoretical framework proposes that environmental stress impacts on the hypothalamic-pituitary-adrenal (HPA) axis, leading to elevated levels of stress hormones that are involved in the regulation of pubertal timing (Belsky and Shalev, 2016). HPA axis dysregulation and other stress-related physiological and epigenetic pathways have also been found to be associated with a greater risk of suicide and self-harm (Berardelli et al., 2020; Turecki and Brent, 2016).

Understanding if there are biological pathways that underpin the relationship between childhood adversity and self-harm will increase our understanding of factors that contribute to the aetiology of self-harm. Understanding this, and the relative contributions of biological and psychological mediators will highlight potential targets for psychosocial interventions. For example, early pubertal timing may lead to self-harm via adolescent social maladjustment, where those with early pubertal timing may associate with older peers, engaging in more risky behaviours such as alcohol use sooner than their same-age peers, increasing risk of self-harm (Patton et al., 2007). Interventions could promote adaptive coping skills, social negotiation skills and assertiveness and emotion regulation. The timing and quality of puberty education in schools (an ideal setting for such an intervention) could be improved to ensure those who have an earlier puberty are supported adequately; positive psychosocial adjustment at this point could reduce risk of later self-harm. Psychosocial interventions may also intervene on the pathways from ACEs to pubertal timing by using stress-tolerance techniques or by supporting families and children to maintain good mental health in the face of adversity and to reduce children's experiences of adversity.

Given these hypothesised common causal mechanisms from adversity to earlier onset of puberty and self-harm, we investigated whether pubertal timing lies on the causal pathway from ACEs to self-harm. Specifically, we aimed to assess whether pubertal timing mediated the association between the cumulative types of adversity a young person has experienced and risk of self-harm in adolescence and young adulthood, using data from a longitudinal birth cohort. We hypothesised that ACEs would be associated with earlier pubertal timing (measured as a continuous variable), and that pubertal timing would lie on the causal pathway between adversity and self-harm.

The current study

To our knowledge, there is no published research investigating whether early pubertal timing is a mediator of the relationship between ACEs and risk of self-harm. We used data from a large UK birth cohort to investigate the relationship between ACEs (assessed from birth to 9 years), pubertal timing and self-harm, and we specifically examined whether early pubertal timing is a mediator of the relationship between ACEs and self-harm at 16 and 21 years.

Methods

Sample

The Avon Longitudinal Study of Parents and Children (ALSPAC) is a birth cohort that originally recruited pregnant women in what was previously the county of Avon, UK with expected dates of delivery 1st April 1991 to 31st December 1992. We included individuals from the initial enrolment sample: the initial number of pregnancies was 14,541. Of these, there was a total of 14,676 fetuses, resulting in 14,062 live births and 13,988 children who were alive at 1 year of age (Boyd et al., 2013; Fraser et al., 2012; Northstone et al., 2019). Twins are usually highly similar due to growing up in the same environment and therefore their data are not independent. Including twins in our study sample may therefore inflate any population-level association found. To mitigate this risk, we excluded the second-born of each twin pair. The study sample comprised those who had data on any of the putative pubertal timing mediators (N=6689): multiple imputation using chained equations was utilised to impute missing data on exposure, outcome and covariates. Data were collected via questionnaires and research clinics. The ALSPAC study website contains details of all the data that is available through a fully searchable data dictionary and variable search tool (<http://www.bristol.ac.uk/alspac/researchers/our-data/>).

Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees.

Measures

Childhood adversity

Nine domains of childhood adversity from birth to 9 years old were assessed using parent and child reports (Houtepen, Heron, Suderman, Tilling, and Howe, 2018; Russell et al., 2019). We assessed adversity up to age 9 to ensure that our exposure variable preceded onset of menarche (only 12 participants reported onset of menarche prior to age 9 in the study sample). The domains of adversity have been widely used in other studies of childhood adversity (Finkelhor, Shattuck, Turner, and Hamby, 2013; Hughes et al., 2017) and include physical, sexual or emotional abuse, parental

mental health problems or suicide attempt, child being bullied, violence between parents, parental separation, parental criminal conviction and parental substance use (Supplementary Table 1). Exposure to adversity was conceptualised as a continuous score of the number of different adversities experienced (range 0-9), as existing evidence shows a dose-response relationship between the number of types of adversity experienced and a range of negative health outcomes in later life (Anda et al., 2006; Hughes et al., 2017).

Self-harm

At age 16, young people provided self-reports of their lifetime history of self-harm in response to the question “Have you ever hurt yourself on purpose in any way (e.g. by taking an overdose of pills, or by cutting yourself)?”

Mediators: pubertal timing

We used age at peak height velocity (aPHV) and age at menarche (AAM) as indicators of pubertal timing. aPHV is an objectively measured indicator of pubertal timing in both sexes that indicates the age at which the adolescent ‘growth spurt’ is at its most rapid (Demirjian, Buschang, Tanguay, and Patterson, 1985). The distribution of aPHV is normal in the general population and correlates well with other measures of pubertal timing (e.g. 0.79 with age at menarche in ALSPAC) (Demirjian et al., 1985; Marshall and Tanner, 1969, 1970). aPHV was calculated from multiple height measurements recorded at research clinics across childhood and adolescence using Superimposition by Translation and Rotation (SITAR), a mixed effects growth curve analysis described in detail elsewhere (Frysz, Howe, Tobias, and Paternoster, 2018).

AAM was assessed using data from nine postal questionnaires relating to pubertal development that were administered approximately annually from age 8–17. The questionnaires asked whether menstruation had started, and if so at what age (in years and months). The first-reported age at menarche was used to minimise recall error (Joinson, Heron, Araya, and Lewis, 2013).

Confounders and covariates

Participant sex, father absence during pregnancy, maternal education, housing tenure, material hardship and financial difficulties were treated as confounders of all paths in our models. We included body mass index (BMI) at age 9 as an intermediate confounder, because it could potentially be affected by childhood adversity and causally influence pubertal timing and self-harm.

Sensitivity analyses

We examined associations with four secondary dichotomous outcomes: lifetime history of self-harm with suicidal intent at age 16, multiple self-harm in past year (i.e. reported >1 episode) at age 16, self-harm at age 21 and lifetime history of self-harm with suicidal intent at age 21. We considered suicidal intent to be present if young people indicated that they “have ever seriously wanted to kill [themselves] on any occasion where [they] have hurt [themselves]” or the “last time [they] hurt themselves it was because [they] wanted to die”. Self-harm and suicidal intent at age 21 were measured with the same questions asked at age 16. We conducted sensitivity analyses to explore whether effects were independent of psychopathology by excluding those with a psychiatric disorder assessed using the Development and Wellbeing Assessment (DAWBA) at age 15. DSM-IV diagnoses were generated using information from parents and young people via a computer algorithm (Goodman, Heiervang, Collishaw, and Goodman, 2011; Goodman, Ford, Richards, Gatward, and Meltzer, 2000). The primary results presented are from the imputed sample (N=6698) with complete case results presented in the supplementary material.

Analysis

We treated aPHV and AAM (in months) as continuous variables. We used multiple imputation by chained equations (separately for males and females; 50 imputations) in order to mitigate the potential for bias due to missing data (N=6698), and compared this with results obtained from analyses in complete case data (n=2373). Results were largely consistent across the imputed and complete case samples.

We conducted mediation analyses using structural equation modelling to implement Poisson regression (because self-harm was not a rare outcome) with robust standard errors, partitioning effects into direct and indirect effects using the products of coefficients approach (conducted using the *gsem* package in Stata v15). In complete case data, we used bootstrapping (n reps=1000) to generate bias-corrected confidence intervals. In the aPHV model, there was no interaction between sex and aPHV ($p>0.1$), therefore we included males and females in a single model, with sex as a covariate. We ran sensitivity analyses to explore whether findings were consistent when excluding those with psychiatric disorder at age 15, and for the secondary outcomes (self-harm with suicidal intent at age 16, multiple self-harm in past year at age 16, self-harm at age 21, and self-harm with suicidal intent at age 21).


Results

Description of sample

Descriptive statistics for the imputed sample and complete cases are shown in Table 1. The study sample comprised young people from more socioeconomically advantaged families compared with those not included in the study sample (Supplementary Table 2). Of the nine adversities measured, parental mental health problems or suicide attempt were the most frequently experienced (39.3%), followed by parental separation and violence between parents (21.9% and 21.2% respectively; supplementary Tables 3 and 4). Figure 1 shows the number of adversities experienced across the sample, and supplementary Figures 1-4 show the distribution of pubertal timing measures by gender, as well as plots indicating the mean aPHV by number of ACEs and gender.

Table 1. Descriptive statistics: complete case and imputed sample

Figure 1. Number of adverse childhood experiences per child (complete case data N=2373)



Mediation results

There was robust evidence for direct effects of childhood adversity on risk of self-harm. For every additional type of adversity experienced; participants had on average a 12-14% increased risk of self-harm by age 16 (Table 2). The relative Risk (RR) for the direct effect was 1.14 (95% confidence interval (CI) 1.09, 1.20) in the model with aPHV as the mediator, whilst in the model with AAM as the mediator the RR was 1.12 (95% CI 1.08 - 1.17). RRs were stronger for direct effects when outcomes were restricted to those who self-harmed with suicidal intent at 16 or 21 years. There was no evidence that pubertal timing (assessed using aPHV or AAM) mediated the association between childhood adversity and self-harm (indirect effect RR 1.00, 95% CI 1.00, 1.00 for aPHV and RR 1.00 95% CI 1.00, 1.01 for AAM). On examining model results, we found evidence for associations between pubertal timing (aPHV and AAM) and self-harm, and between childhood adversity and self-harm. There was no evidence, however, of an association between childhood adversities and pubertal timing. RR estimates were marginally larger for complete case data, with wider confidence intervals (Supplementary Table 5).

Table 2 Results of mediation analyses assessing the relationship between number of adversities experienced by age 9, pubertal timing and later self-harm

Sensitivity analyses

In sensitivity analyses utilising outcomes of self-harm at age 21, self-harm with suicidal intent at ages 16 and 21, past-year multiple self-harm at age 16, and excluding those with psychiatric disorder at age 15, results for each mediator were consistent with the main findings (Tables 3 and 4). The direct effects were larger for self-harm with suicidal intent at 16 and 21 (aPHV model: RR at age 16= 1.28, 95% CI 1.18, 1.39; RR at age 21= 1.20, 95% CI 1.12, 1.28).

Table 3. Sensitivity analyses results of mediation analyses assessing the relationship between number of adversities experienced by age 9, age at peak height velocity and later self-harm

Table 4. Sensitivity analyses results of mediation analyses assessing the relationship between number of adversities experienced by age 9, age at menarche and later self-harm

Discussion

We found that childhood adversity and early timing of puberty are independent risk factors for later self-harm in both males and females. There was no evidence in this study for an association between childhood adversities and pubertal timing, and no mediating effect of early pubertal timing on the relationship between childhood adversity and self-harm. Our findings are consistent with a large study (n= 14,000 adolescents) that found no evidence for a mediating effect of pubertal timing on the relationship between childhood adversity and depressive symptoms in adolescence (Strong, Tsai, Lin, and Cheng, 2016). Just over one-in-five young people in our sample reported self-harm at age 16 (18.8% in complete case data and 22.7% in the imputed data). This is higher than the prevalence of self-harm in adolescents aged 13-18 in the community reported in a recent meta-analysis (16.9%, 95% CI 15.1-18.9%), although estimates vary across samples due to differences in population and methodology (Gillies et al., 2018). The prevalence of ACEs in our sample (58.7% reported at least one ACE) was consistent with estimates from a recent meta-analysis of over 250,000 participants and in a similar UK-based cohort assessing ACEs from age 0-5 years (57% and 50% respectively reported at least once ACE), whilst 5.0% in our sample reported four or more ACEs, compared with 13% with at least four ACEs in the meta-analysis and 1.4% in the Millennium Cohort (Hughes et al., 2017; Straatmann et al., 2020). We measured ACEs between the ages of 0 and 9 years whereas the meta-analysis largely includes data from adult samples. Differences in socioeconomic disadvantage across populations might also explain the difference in those accumulating more than four ACEs (Straatman et al., 2020). Our estimates of the distribution of age at menarche and peak height velocity are also in line with other studies (e.g. Granados, Gebremariam, and Lee, 2015; Parent et al., 2003).

There are currently fewer studies of adversity and pubertal timing in boys than girls, although research in this area is increasing (Hawton et al., 2003; Lei et al., 2018; Mendle, 2014; Zhang et al., 2019). Existing studies reporting null findings may also be subject to publication bias. Our study found no association between cumulative adversity and age at menarche. A study of the ALSPAC mothers' cohort found that only sexual abuse was associated with earlier menarche (Magnus et al., 2018). One study found evidence for differing dose-response relationships between several forms of abuse with age at menarche, with sexual and physical abuse predicting earlier AAM, and physical abuse also being associated with later AAM (Boynton-Jarrett et al., 2013). Exposure to specific types of adversity may also impact differentially on physiological development during

puberty, with findings for sexual abuse being particularly strong (Zabin, Emerson, and Rowland, 2005). Puberty and developmental maturational processes are largely controlled by biological systems, however there is evidence that environmental factors such as diet, nutrition, and the presence of a step-father in the household (Zabin et al., 2005) contribute to pubertal timing. We controlled for BMI and father absence during pregnancy, however there may be other factors that were not accounted for in our models.

Strengths and Limitations

We utilised data from a large sample of young people, who are broadly representative of the UK population although slightly more socioeconomically advantaged and less ethnically diverse. The most disadvantaged participants were more likely to drop-out (Wolke et al., 2009), but our findings were similar across imputed and complete case analyses. The longitudinal nature of the mediation analysis is a key strength of the study design, allowing us to draw inferences about the direction of effects with measures of adversity occurring before puberty, and pubertal timing prior to measures of self-harm. We used an objective measure of pubertal timing, aPHV, as our primary mediator. This was based on repeated measures of height taken during research clinics, rather than self-reported height, and allowed us to investigate associations for males as well as females. Males are underrepresented in research on puberty, and much of the existing studies that link adversity and puberty, and puberty and self-harm, are based on females. Our estimates of prevalence of ACEs and self-harm were in line with other studies, which gives confidence in the potential generalisability of our findings.

There are limitations to our study; there may be bias in reporting of adversity, as much of the information was collected from questionnaires to parents. The outcome data are now 10 years old, and rates of self-harm are now higher than 10 years ago (McManus et al., 2019; Sadler et al., 2018). Parents may under-report adversities due to perceived stigma or social desirability. However, estimates of adversities in ALSPAC are in line with similar studies. Similarly, self-harm was self-reported by individuals, and the prevalence is similar to other non-clinical samples. We did not account for the severity of adversities in our model, beyond including multiple forms of adversity. It is likely that exposure to severe and repeated adversity or multiple adversities will have differential impacts on pubertal timing, compared with brief exposure to one adversity. This was not adequately captured by our measures and future work could explore this further. We utilised a cumulative index of number of adversities experienced, which is common in the scientific literature as exposure to a range of ACEs has been shown to have associations with a range of health outcomes across the lifecourse (Hughes et al., 2017) and because ACEs do not often occur in isolation it is difficult to disentangle the impact of a single ACE on a single outcome.

Our sample was more socioeconomically advantaged and less ethnically diverse than the UK population. Socioeconomic disadvantage is associated with a higher burden of ACEs in studies from a range of countries (Straatmann et al., 2020; Walsh, McCartney, Smith, and Armour, 2019). In our sample this may mean that exposure to ACEs was underestimated and may explain why we have a smaller proportion of participants with four or more ACEs compared with other studies (Hughes et al, 2017) and therefore associations between adversities and self-harm would be attenuated towards the null. To some extent, this should be accounted for by our including several indicators of socioeconomic disadvantage as covariates in our models. Interestingly, studies have found that adversity confers additional risk of negative mental health outcomes after accounting for socioeconomic disadvantage and ethnicity of participants (Nurius, Logan-Greene, and Green, 2012), as seen in our findings also. There is also evidence that ethnic background or race may be differentially associated with likelihood of experiencing adversity. Most data on this comes from the US (e.g. Putnam-Hornstein, Needell, King, and Johnson-Motoyama, 2013) however studies based in UK populations have shown that ACEs were lower in those of Asian ethnicity, while those who were non-white and non-Asian had higher levels of adversity (Bellis, Hughes, Leckenby, Perkins, and Lowey, 2014). Our analysis may therefore either under or over-estimate associations between ACEs and self-harm in those with ethnic backgrounds that are not well represented in ALSPAC.

Implications

Further research should also explore the relationship between the severity and timing of exposure to adversity, and the role of specific types of adversity, on pubertal timing in both males and females in order to extend current understanding of how this relationship may impact on future health and wellbeing. Given the recent evidence for an increasing rate of self-harm in adolescence (Sadler et al., 2018), and the relationship with childhood adversity, research is needed to understand mechanisms that explain why adverse exposures in childhood increase the risk of self-harm. Identifying modifiable mediators could inform the development of prevention and intervention efforts. Adversity is known to have multiple impacts on development, and mediating mechanisms could include cognitive, behavioural, emotional, social and biological factors (Dube et al., 2003). Potential mediators that could be studied further include mental health problems (Enns et al., 2006) adolescent drinking and drug use (Dube et al., 2001; Turecki, Ernst, Jollant, Labonté, and Mechawar, 2012), family support (Cassels et al., 2018), immune functioning, and stress reactivity (Berens et al 2017). Biological mechanisms may also include dysregulated stress-response systems, or cognitive deficits that impact on ability to regulate behaviour and emotions (Turecki et al., 2012). As with many complex traits, it is likely that there are multiple interacting causal pathways from adversity to self-harm.

Conclusions

Our findings suggest that both childhood adversity and early pubertal timing are risk factors for later self-harm in both males and females. However, we found no evidence that the effect of adversity on self-harm is mediated through earlier pubertal timing. There is a need to identify mechanisms that explain why childhood adversity is associated with an increased risk of self-harm in order to inform the development of prevention and intervention efforts. School-based and psychosocial interventions targeting adolescent risky behaviours and social adjustment, as well as improved quality of puberty education in schools, are needed. Interventions should be targeted prior to and during pubertal onset, given that those who enter puberty earliest are at highest risk of later self-harm.

Tables

Table 1. Descriptive statistics: complete case and imputed sample

	Complete case (N=2373)		Imputed sample (N=6689, 50 imputed datasets)	
	mean	SD	mean	SE
Continuous variables				
Adversity score	1.08	1.23	1.38	0.0
Age at menarche (months)	153.1	13.9	151.5	0.2
Age at peak height velocity (PHV) (males and females, months)	150.6	14.1	151.3	0.2
Age at PHV (females, months)	141.9	9.81	141.5	0.0
Age at PHV (males, months)	161.8	10.3	162.3	0.2
BMI age 9	17.5	2.7	17.8	0.2
Birthweight (g)	3,369.0	655.0	3368.9	0.0
Maternal age at child birth (years)	29.93	4.26	29.0	8.0
Binary / categorical variables	n	%	%	SE
Psychiatric disorder at 15	87	4.1	6.59	0.4
Self-harm at 16	447	18.8	22.7	0.7
Self-harm at 21	372	21.8	23.7	0.7
Suicidal intent at 16	145	6.1	4.50	0.5
Suicidal intent at 21	202	8.5	6.53	0.6
Multiple self-harm at 16	242	10.5	14.6	0.6
Child sex (female)	1392	58.7	62.1	0.6
Housing tenure (not owned/mortgaged)	233	9.8	17.8	0.5
Maternal education				
<i>Degree</i>	590	24.9	16.5	0.5
<i>A-level</i>	731	30.8	26.7	0.5
<i>GCSE</i>	745	31.4	34.5	0.6
<i><GCSE</i>	307	12.9	22.3	0.5
Material hardship	411	17.3	27.1	0.5
Equivalent household income (quintiles)				
<i>Highest</i>	724	30.5	22.7	0.5
	606	25.5	21.8	0.5
	482	20.3	20.5	0.5
	351	14.8	18.7	0.5
<i>Lowest</i>	210	8.85	16.3	0.6
Father absence during pregnancy	<5	<0.2	1.7	0.5
Maternal smoking during pregnancy (ever)	295	12.4	19.3	0.5

Parity				
0	1,196	50.9	47.9	0.6
1	816	34.7	34.7	0.6
2	263	11.2	12.8	0.4
3+	75	3.2	4.6	0.3
White British ethnicity	2,335	98.8	98.0	0.2

Table 2 Results of mediation analyses assessing the relationship between number of adversities experienced by age 9, pubertal timing and later self-harm

Outcome	Mediator investigated	Direct effect			Indirect effect via mediator			Total effect		
		RR	95% CI	p value	RR	95% CI	p value	RR	95% CI	p value
Self-harm at 16	Age at peak height velocity (both sexes, main analysis N=6689)	1.14	1.09, 1.20	<0.001	1.00	1.00, 1.00	0.446	1.14	1.09, 1.20	<0.001
	Age at peak height velocity standardised by sex (both sexes, main analysis N=6689)	1.14	1.09, 1.19	<0.001	1.00	1.00, 1.00	0.454	1.14	1.08, 1.19	<0.001
Self-harm at 16	Age at menarche (females n=4040)	1.12	1.08, 1.17	<0.001	1.00	1.00, 1.01	0.224	1.12	1.08, 1.17	<0.001
Self-harm at 16	Age at peak height velocity (males only, n=2533)	1.14	1.04, 1.25	0.007	1.00	0.99, 1.01	0.878	1.14	1.04, 1.25	0.007
Self-harm at 16	Age at peak height velocity (females only, n=4156)	1.14	1.09, 1.20	<0.001	1.00	0.99, 1.00	0.273	1.14	1.09, 1.20	<0.001

Notes: RR relative risk 95% CI 95% confidence interval. RRs can be interpreted as the average percentage increase in risk (if >1) of the outcome for each additional type of adversity experienced prior to age 9 i.e. for row 1 direct effect, for every additional type of adversity experienced, participants were on average 14% more likely to report self-harm at age 16.

Table 3. Sensitivity analyses results of mediation analyses assessing the relationship between number of adversities experienced by age 9, age at peak height velocity and later self-harm

Outcome	Mediator	Direct effect			Indirect effect via mediator			Total effect		
		RR	95% CI	p value	RR	95% CI	p value	RR	95% CI	p value
Self-harm with suicidal intent at 16	Age at peak height velocity (both sexes N=6689)	1.28	1.18, 1.39	<0.001	1.00	1.00, 1.00	0.496	1.28	1.18, 1.39	<0.001
Multiple self-harm in past year at 16	Age at peak height velocity (both sexes N=6689)	1.12	1.05, 1.20	0.001	1.00	1.00, 1.00	0.504	1.12	1.04, 1.19	0.001
Self-harm at 16; no psychiatric disorder at 15	Age at peak height velocity (both sexes n=6535)	1.13	1.08, 1.19	<0.001	1.00	1.00, 1.00	0.442	1.13	1.08, 1.19	<0.001
Self-harm at 21	Age at peak height velocity (both sexes N=6689)	1.13	1.08, 1.18	<0.001	1.00	1.00, 1.00	0.572	1.13	1.08, 1.18	<0.001
Self-harm with suicidal intent at 21	Age at peak height velocity (both sexes N=6689)	1.20	1.12, 1.28	<0.001	1.00	1.00, 1.00	0.5	1.20	1.12, 1.28	<0.001

Notes: RR relative risk 95% CI 95% confidence interval. RRs can be interpreted as the average percentage increase in risk (if >1) of the outcome for each additional type of adversity experienced prior to age 9 i.e. for row 1 direct effect, for every additional type of adversity experienced, participants were on average 28% more likely to report self-harm with suicidal intent at age 16.

Table 4. Sensitivity analyses results of mediation analyses assessing the relationship between number of adversities experienced by age 9, age at menarche and later self-harm

Outcome	Mediator	Direct effect			Indirect effect via mediator			Total effect		
		RR	95% CI	p value	RR	95% CI	p value	RR	95% CI	p value
Self-harm with suicidal intent at 16	Age at menarche (females n=4040)	1.21	1.12, 1.29	<0.001	1.00	1.00, 1.01	0.229	1.21	1.13, 1.30	<0.001
Multiple self-harm in past year at 16	Age at menarche (females n=4040)	1.13	1.06, 1.20	<0.001	1.00	1.00, 1.01	0.221	1.13	1.07, 1.21	<0.001
Self-harm at 16; no psychiatric disorder at 15	Age at menarche (females n=3844)	1.12	1.07, 1.17	<0.001	1.00	1.00, 1.01	0.169	1.12	1.07, 1.17	<0.001
Self-harm at 21	Age at menarche (females n=4040)	1.09	1.05, 1.14	<0.001	1.00	1.00, 1.00	0.312	1.10	1.05, 1.14	<0.001
Self-harm with suicidal intent at 21	Age at menarche (females n=4040)	1.17	1.09, 1.24	<0.001	1.00	1.00, 1.01	0.225	1.17	1.10, 1.25	<0.001

Notes: RR relative risk 95% CI 95% confidence interval. RRs can be interpreted as the average percentage increase in risk (if >1) of the outcome for each additional type of adversity experienced prior to age 9 i.e. for row 1 direct effect, for every additional type of adversity experienced, participants were on average 21% more likely to report self-harm with suicidal intent at age 16.

Author contributions

Study conception and design, revising work critically for important intellectual content: all authors; Acquisition, analysis, and interpretation of data: AER, BM, JH, DJ, CJ; drafting manuscript: AER; final approval

Conflicts of interest

Conflicts of Interest: None

Ethical standards

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

Acknowledgements

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. We would like to thank Lotte Houtepen for her work on the ACE constructs in ALSPAC.

Financial support

This study was funded by the Medical Research Foundation and Medical Research Council (Grant ref: MR/R004889/1) Pathways to self-harm: Biological mechanisms and genetic contribution PI: Dr Becky Mars. The UK Medical Research Council and Wellcome (Grant ref: 217065/Z/19/Z) and the University of Bristol provide core support for ALSPAC. This publication is the work of the authors and Abigail Russell and Becky Mars serve as guarantors for the contents of this paper. A comprehensive list of grants funding is available on the ALSPAC website (<http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf>); this research was specifically funded by the Wellcome Trust (Grant ref: GR067797MA).

DG, PM, CR and BM are supported by the NIHR Biomedical Research Centre at University Hospitals Bristol NHS Foundation Trust and the University of Bristol, England. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health and Social Care.

Matthew Suderman and Caroline Relton work in the Medical Research Council Integrative Epidemiology Unit at the University of Bristol which is supported by the Medical Research Council and the University of Bristol [grant number MC_UU_00011/1, MC_UU_00011/4 and MC_UU_00011/5].

References

- Anda, R. F., Felitti, V. J., Bremner, J. D., Walker, J. D., Whitfield, C., Perry, B. D., . . . Giles, W. H. (2006). The enduring effects of abuse and related adverse experiences in childhood. *European archives of psychiatry and clinical neuroscience*, 256(3), 174-186. doi: [10.1007/s00406-005-0624-4](https://doi.org/10.1007/s00406-005-0624-4)
- Arim, R. G., Tramonte, L., Shapka, J. D., Dahinten, V. S., and Willms, J. D. (2011). The family antecedents and the subsequent outcomes of early puberty. *J Youth Adolesc*, 40(11), 1423-1435. doi:10.1007/s10964-011-9638-6
- Barrios, Y. V., Sanchez, S. E., Nicolaidis, C., Garcia, P. J., Gelaye, B., Zhong, Q. Y., and Williams, M. A. (2015). Childhood Abuse and Early Menarche Among Peruvian Women. *Journal of Adolescent Health*, 56(2), 197-202. doi:10.1016/j.jadohealth.2014.10.002
- Bellis, M. A., Hughes, K., Leckenby, N., Perkins, C., and Lowey, H. (2014). National household survey of adverse childhood experiences and their relationship with resilience to health-harming behaviors in England. *BMC medicine*, 12(1), 1-10. <https://doi.org/10.1186/1741-7015-12-72>
- Belsky, J., and Shalev, I. (2016). Contextual adversity, telomere erosion, pubertal development, and health: Two models of accelerated aging, or one? *Development and Psychopathology*, 28(4pt2), 1367-1383. doi: 10.1017/S0954579416000900
- Berardelli, I., Serafini, G., Cortese, N., Fiaschè, F., O'Connor, R. C., and Pompili, M. (2020). The Involvement of Hypothalamus–Pituitary–Adrenal (HPA) Axis in Suicide Risk. *Brain Sciences*, 10(9), 653. doi: 10.3390/brainsci10090653
- Björkenstam, C., Kosidou, K., and Björkenstam, E. (2017). Childhood adversity and risk of suicide: cohort study of 548 721 adolescents and young adults in Sweden. *BMJ*, 357, j1334. doi: <https://doi.org/10.1136/bmj.j1334>
- Björkenstam, E., Kosidou, K., and Björkenstam, C. (2016). Childhood household dysfunction and risk of self-harm: a cohort study of 107 518 young adults in Stockholm County. *International Journal of Epidemiology*, 45(2), 501-511. doi: 10.1093/ije/dyw012
- Bogaert, A. F. (2005). Age at puberty and father absence in a national probability sample. *Journal of Adolescence* 28(4), 541-546. doi: <https://doi.org/10.1016/j.adolescence.2004.10.008>
- Boyd, A., Golding, J., Macleod, J., Lawlor, D. A., Fraser, A., Henderson, J., . . . Davey Smith, G. (2013). Cohort profile: the 'children of the 90s'—the index offspring of the Avon Longitudinal Study of Parents and Children. *International Journal of Epidemiology*, 42(1), 111-127. doi: <https://doi.org/10.1093/ije/dys064>
- Boynton-Jarrett, R., and Harville, E. W. (2012). A prospective study of childhood social hardships and age at menarche. *Annals of Epidemiology*, 22(10), 731-737. doi:10.1016/j.annepidem.2012.08.005
- Boynton-Jarrett, R., Wright, R. J., Putnam, F. W., Hibert, E. L., Michels, K. B., Forman, M. R., and Rich-Edwards, J. (2013). Childhood Abuse and Age at Menarche. *Journal of Adolescent Health*, 52(2), 241-247. doi:10.1016/j.jadohealth.2012.06.006
- Cassels, M., van Harmelen, A. L., Neufeld, S., Goodyer, I., Jones, P. B., and Wilkinson, P. (2018). Poor family functioning mediates the link between childhood adversity and adolescent nonsuicidal self-injury. *Journal of Child Psychology and Psychiatry*, 59(8), 881-887. doi: <https://doi.org/10.1111/jcpp.12866>
- Cha, C. B., Franz, P. J., M. Guzmán, E., Glenn, C. R., Kleiman, E. M., and Nock, M. K. (2018). Annual Research Review: Suicide among youth—epidemiology,(potential) etiology, and treatment. *Journal of Child Psychology and Psychiatry*, 59(4), 460-482. doi: <https://doi.org/10.1111/jcpp.12831>
- Clutterbuck, S., Adams, J., and Nettle, D. (2015). Frequent Residential Relocations Cumulatively Accelerate Menarcheal Timing in a Sample of English Adolescent Girls. *Journal of Biosocial Science*, 47(2), 188-202. doi:10.1017/S0021932014000157

- Demirjian, A., Buschang, P. H., Tanguay, R., and Patterson, D. K. J. (1985). Interrelationships among measures of somatic, skeletal, dental, and sexual maturity. *American Journal of Orthodontics* 88(5), 433-438. doi: [https://doi.org/10.1016/0002-9416\(85\)90070-3](https://doi.org/10.1016/0002-9416(85)90070-3)
- Deng, F., Tao, F., Wan, Y., Hao, J., Su, P., and Cao, Y. (2011). Early menarche and psychopathological symptoms in young Chinese women. *Journal of Women's Health*, 20(2), 207-213. doi: <https://doi.org/10.1089/jwh.2010.2102>
- Dube, S. R., Anda, R. F., Felitti, V. J., Croft, J. B., Edwards, V. J., and Giles, W. H. (2001). Growing up with parental alcohol abuse: exposure to childhood abuse, neglect, and household dysfunction. *Child Abuse and Neglect*, 25(12), 1627-1640. doi: [https://doi.org/10.1016/S0145-2134\(01\)00293-9](https://doi.org/10.1016/S0145-2134(01)00293-9)
- Dube, S. R., Felitti, V. J., Dong, M., Giles, W. H., and Anda, R. F. (2003). The impact of adverse childhood experiences on health problems: evidence from four birth cohorts dating back to 1900. *Preventive Medicine*, 37(3), 268-277. doi: [https://doi.org/10.1016/S0091-7435\(03\)00123-3](https://doi.org/10.1016/S0091-7435(03)00123-3)
- Enns, M. W., Cox, B. J., Afifi, T. O., De Graaf, R., Ten Have, M., and Sareen, J. (2006). Childhood adversities and risk for suicidal ideation and attempts: a longitudinal population-based study. *Psychological Medicine*, 36(12), 1769-1778. doi: [doi:10.1017/S0033291706008646](https://doi.org/10.1017/S0033291706008646)
- Finkelhor, D., Shattuck, A., Turner, H., and Hamby, S. (2013). Improving the adverse childhood experiences study scale. *JAMA Pediatrics*, 167(1), 70-75. doi: [doi:10.1001/jamapediatrics.2013.420](https://doi.org/10.1001/jamapediatrics.2013.420)
- Fraser, A., Macdonald-Wallis, C., Tilling, K., Boyd, A., Golding, J., Davey Smith, G., . . . Lawlor, D. (2012). Cohort profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *International Journal of Epidemiology*, 42(1), 97-110. doi: <https://doi.org/10.1093/ije/dys066>
- Frysz, M., Howe, L. D., Tobias, J. H., and Paternoster, L. J. (2018). Using SITAR (SuperImposition by Translation and Rotation) to estimate age at peak height velocity in Avon Longitudinal Study of Parents and Children. *Wellcome Open Research* 3:90. doi: [10.12688/wellcomeopenres.14708.2](https://doi.org/10.12688/wellcomeopenres.14708.2)
- Gillies, D., Christou, M. A., Dixon, A. C., Featherston, O. J., Rapti, I., Garcia-Anguita, A., . . . Christou, P. (2018). Prevalence and characteristics of self-harm in adolescents: meta-analyses of community-based studies 1990–2015. *Journal of the American Academy of Child and Adolescent Psychiatry*, 57(10), 733-741. doi: <https://doi.org/10.1016/j.jaac.2018.06.018>
- Goodman, A., Heiervang, E., Collishaw, S., and Goodman, R. (2011). The 'DAWBA bands' as an ordered-categorical measure of child mental health: description and validation in British and Norwegian samples. *Social Psychiatry and Psychiatric Epidemiology*, 46(6), 521-532. doi: [10.1007/s00127-010-0219-x](https://doi.org/10.1007/s00127-010-0219-x)
- Goodman, R., Ford, T., Richards, H., Gatward, R., and Meltzer, H. (2000). The Development and Well-Being Assessment: description and initial validation of an integrated assessment of child and adolescent psychopathology. *The Journal of Child Psychology and Psychiatry and Allied Disciplines*, 41(5), 645-655. doi: <https://doi.org/10.1111/j.1469-7610.2000.tb02345.x>
- Granados, A., Gebremariam, A., and Lee, J. M. (2015). Relationship between timing of peak height velocity and pubertal staging in boys and girls. *Journal of Clinical Research in Pediatric Endocrinology*, 7(3), 235. doi: [10.4274/jcrpe.2007](https://doi.org/10.4274/jcrpe.2007)
- Hawton, K., Hall, S., Simkin, S., Bale, L., Bond, A., Codd, S., and Stewart, A. (2003). Deliberate self-harm in adolescents: A study of characteristics and trends in Oxford, 1990–2000. *Journal of Child Psychology and Psychiatry*, 44(8), 1191-1198. doi: <https://doi.org/10.1111/1469-7610.00200>
- Henrichs, K. L., McCauley, H. L., Miller, E., Styne, D. M., Saito, N., and Breslau, J. (2014). Early menarche and childhood adversities in a nationally representative sample. *International Journal of Pediatric Endocrinology*, 2014(1), 14. doi: <https://doi.org/10.1186/1687-9856-2014-14>

- Houtepen, L. C., Heron, J., Suderman, M. J., Tilling, K., and Howe, L. D. (2018). Adverse childhood experiences in the children of the Avon Longitudinal Study of Parents and Children (ALSPAC). *Wellcome Open Research*, 3:106. doi: 10.12688/wellcomeopenres.14716.1
- Hughes, K., Bellis, M. A., Hardcastle, K. A., Sethi, D., Butchart, A., Mikton, C., . . . Dunne, M. P. (2017). The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *The Lancet Public Health*, 2(8), e356-e366. doi:[https://doi.org/10.1016/S2468-2667\(17\)30118-4](https://doi.org/10.1016/S2468-2667(17)30118-4)
- Hulanicka, B., Gronkiewicz, L., and Koniarek, J. (2001). Effect of familial distress on growth and maturation of girls: a longitudinal study. *American Journal of Human Biology*, 13(6), 771-776. doi:10.1002/ajhb.1123
- Joinson, C., Heron, J., Araya, R., and Lewis, G. (2013). Early menarche and depressive symptoms from adolescence to young adulthood in a UK cohort. *Journal of the American Academy of Child and Adolescent Psychiatry*, 52(6), 591-598. e592. doi: <https://doi.org/10.1016/j.jaac.2013.03.018>
- Kelly, Y., Zilanawala, A., Sacker, A., Hiatt, R., and Viner, R. (2017). Early puberty in 11-year-old girls: Millennium Cohort Study findings. *Arch Dis Child*, 102(3), 232-237. doi:10.1136/archdischild-2016-310475
- Larsson, B., and Sund, A. M. (2008). Prevalence, course, incidence, and 1-year prediction of deliberate self-harm and suicide attempts in early Norwegian school adolescents. *Suicide and Life-Threatening Behavior*, 38(2), 152-165.
- Lei, M. K., Beach, S. R. H., and Simons, R. L. (2018). Childhood trauma, pubertal timing, and cardiovascular risk in adulthood. *Health Psychology*, 37(7), 613-617. doi:10.1037/hea0000609
- Li, L., Denholm, R., and Power, C. (2014). Child maltreatment and household dysfunction: associations with pubertal development in a British birth cohort. *International Journal of Epidemiology*, 43(4), 1163-1173. doi:10.1093/ije/dyu071
- Magnus, M. C., Anderson, E. L., Howe, L. D., Joinson, C. J., Penton-Voak, I. S., and Fraser, A. (2018). Childhood psychosocial adversity and female reproductive timing: a cohort study of the ALSPAC mothers. *Journal of Epidemiology and Community Health*, 72(1), 34-40. doi:10.1136/jech-2017-209488
- Marshall, W. A., and Tanner, J. M. (1969). Variations in pattern of pubertal changes in girls. *Archives of Disease in Childhood* (235), 44:291. doi: 10.1136/adc.44.235.291
- Marshall, W. A., and Tanner, J. M. (1970). Variations in the pattern of pubertal changes in boys. *Archives of Disease in Childhood* 45(239), 13-23. doi: <http://dx.doi.org/10.1136/adc.45.239.13>
- McManus, S., Gunnell, D., Cooper, C., Bebbington, P. E., Howard, L. M., Brugha, T., . . . Appleby, L. J. (2019). Prevalence of non-suicidal self-harm and service contact in England, 2000–14: repeated cross-sectional surveys of the general population. *The Lancet Psychiatry*, 6(7), 573-581. doi: [https://doi.org/10.1016/S2215-0366\(19\)30188-9](https://doi.org/10.1016/S2215-0366(19)30188-9)
- Mendle, J. (2014). Why puberty matters for psychopathology. *Child Development Perspectives*, 8(4), 218-222. doi: <https://doi.org/10.1111/cdep.12092>
- Michaud, P.-A., Suris, J.-C., and Deppen, A. (2006). Gender-related psychological and behavioural correlates of pubertal timing in a national sample of Swiss adolescents. *Molecular Cellular Endocrinology*, 254, 172-178. doi: <https://doi.org/10.1016/j.mce.2006.04.037>
- Northstone, K., Lewcock, M., Groom, A., Boyd, A., Macleod, J., Timpson, N., and Wells, N. J. (2019). The Avon Longitudinal Study of Parents and Children (ALSPAC): an update on the enrolled sample of index children in 2019. *Wellcome Open Research* 4:51. doi: 10.12688/wellcomeopenres.15132.1
- Nurius, P. S., Logan-Greene, P., and Green, S. (2012). ACEs within a social disadvantage framework: Distinguishing unique, cumulative, and moderated contributions to adult mental health.

- Journal of prevention and intervention in the community*, 40(4), 278. doi: 10.1080/10852352.2012.707443
- Parent, A.-S., Teilmann, G., Juul, A., Skakkebaek, N. E., Toppari, J., and Bourguignon, J.-P. (2003). The timing of normal puberty and the age limits of sexual precocity: variations around the world, secular trends, and changes after migration. *Endocrine reviews*, 24(5), 668-693. doi: <https://doi.org/10.1210/er.2002-0019>
- Patton, G. C., Hemphill, S. A., Beyers, J. M., Bond, L., Toumbourou, J. W., McMorris, B. J., and Catalano, R. F. (2007). Pubertal stage and deliberate self-harm in adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 46(4), 508-514. doi: <https://doi.org/10.1097/chi.0b013e31803065c7>
- Putnam-Hornstein, E., Needell, B., King, B., and Johnson-Motoyama, M. (2013). Racial and ethnic disparities: A population-based examination of risk factors for involvement with child protective services. *Child Abuse and Neglect*, 37(1), 33-46. doi:<https://doi.org/10.1016/j.chiabu.2012.08.005>
- Roberts, E., Fraser, A., Gunnell, D., Joinson, C., and Mars, B. (2020) Timing of menarche and self-harm in adolescence and adulthood: a population-based cohort study. *Psychological medicine*, 1-9. doi:10.1017/S0033291719002095
- Romans, S., Martin, J., Gendall, K., and Herbison, G. J. (2003). Age of menarche: The role of some psychosocial factors. *Psychological Medicine*: 33(5), 933-939. doi: 10.1017/S0033291703007530
- Russell, A. E., Heron, J., Gunnell, D., Ford, T., Hemani, G., Joinson, C., . . . Mars, B. (2019). Pathways between early-life adversity and adolescent self-harm: the mediating role of inflammation in the Avon Longitudinal Study of Parents and Children. *Journal of Child Psychology and Psychiatry*, 60(10) 1094 doi: <https://doi.org/10.1111/jcpp.13100>
- Sadler, K., Vizard, T., Ford, T., Goodman, A., Goodman, R., and McManus, S. (2020, 02 04). Mental Health of Children and Young People in England, 2017: Trends and characteristics. *Leeds, UK: NHS Digital*. Retrieved from <https://openaccess.city.ac.uk/id/eprint/23650/>
- Straatmann, V. S., Lai, E., Law, C., Whitehead, M., Strandberg-Larsen, K., and Taylor-Robinson, D. (2020). How do early-life adverse childhood experiences mediate the relationship between childhood socioeconomic conditions and adolescent health outcomes in the UK? *Journal of Epidemiology and Community Health*, 74(11), 969-975. doi:10.1136/jech-2020-213817
- Strong, C., Tsai, M. C., Lin, C. Y., and Cheng, C. P. (2016). Childhood Adversity, Timing of Puberty and Adolescent Depressive Symptoms: A Longitudinal Study in Taiwan. *Child Psychiatry Hum Dev*, 47(3), 347-357. doi:10.1007/s10578-015-0570-y
- Turecki, G., and Brent, D. A. (2016). Suicide and suicidal behaviour. *The Lancet*, 387(10024), 1227-1239. doi: [https://doi.org/10.1016/S0140-6736\(15\)00234-2](https://doi.org/10.1016/S0140-6736(15)00234-2)
- Turecki, G., Ernst, C., Jollant, F., Labonté, B., and Mechawar, N. (2012). The neurodevelopmental origins of suicidal behavior. *Trends in Neuroscience*, 35(1), 14-23. doi:10.1016/j.tins.2011.11.008
- Walsh, D., McCartney, G., Smith, M., and Armour, G. (2019). Relationship between childhood socioeconomic position and adverse childhood experiences (ACEs): a systematic review. *Journal of Epidemiology and Community Health*, 73(12), 1087-1093. doi: <http://dx.doi.org/10.1136/jech-2019-212738>
- Wichstrøm, L. (2000). Predictors of adolescent suicide attempts: a nationally representative longitudinal study of Norwegian adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 39(5), 603-610. doi: 10.1097/00004583-200005000-00014
- Wolke, D., Waylen, A., Samara, M., Steer, C., Goodman, R., Ford, T., and Lamberts, K. (2009). Selective drop-out in longitudinal studies and non-biased prediction of behaviour disorders. *The British Journal of Psychiatry*, 195(3), 249-256. doi: <https://doi.org/10.1192/bjp.bp.108.053751>

- Zabin, L. S., Emerson, M. R., and Rowland, D. L. (2005). Childhood sexual abuse and early menarche: the direction of their relationship and its implications. *Journal of Adolescent Health, 36*(5), 393-400. doi:10.1016/j.jadohealth.2004.07.013
- Zhang, L., Zhang, D., and Sun, Y. (2019). Adverse childhood experiences and early pubertal timing among girls: a meta-analysis. *International journal of Environmental Research and Public Health, 16*(16), 2887. doi: <https://doi.org/10.3390/ijerph16162887>

Childhood adversity, pubertal timing and self-harm: a longitudinal cohort study: supplementary material

Abigail Russell et al. Psychological Medicine 2021

Supplementary Table 1. Adverse Childhood Experiences (ACEs) definitions

ACE	Definition	Number of questions	Age range of questions asked	Respondents
Sexual abuse	Was the child sexually abused	7	18 months to age 9	Mother
Physical abuse	Whether physically cruel to child	31	8 weeks to 9 years	Mother and Partner
Emotional abuse	Whether or not mum/partner had been emotionally cruel to the child	32	8 months to 9 years	Mother and Partner
Parent substance use	Daily use of cannabis or any use of other drugs. Or, alcohol problem by self-reported problematic use, and saw a doctor because of it	62	8 weeks to 9 years	Mother and Partner
Parent mental health problems or suicide attempt	Depression scores (EPDS>12) and medication, presence of schizophrenia, bulimia, anorexia or attempted suicide.	57	8 weeks to 9 years	Mother and Partner
Violence between parents	Parent experienced physical cruelty from partner, or displayed (specific types) of violence towards partner	43	8 weeks to 9 years	Mother and Partner
Parental separation	Parents divorced or separated. Degree to which this impacted on the child.	32	8 weeks to 9 years	Mother and Partner
Bullying	Child bullied	6	8 years-8.5 years	Child
Parent convicted	Parent convicted off offence	18	8 weeks to 9 years	Mother and Partner

Notes: ACE adverse childhood experience. EPDS Edinburgh Postnatal Depression Scale. Several of the ACEs included questions about how much the ACE had impacted on the parent or child.

Supplementary Table 2. Descriptive statistics: sample for imputation and ALSPAC

Description	Sample for imputation (n=6689)		Rest of ALSPAC (n=7099)		
	mean	SD	mean	SD	P
Adversity score	1.15	1.28	1.00	1.32	0.036
Age at menarche (months)	151.5	14.10			
Age at peak height velocity (months)	151.3	14.43			
Age at PHV (females, months)	141.5	9.82			
Age at PHV (males, months)	162.3	10.26			
BMI age 9	17.7	2.87	17.8	3.05	0.2312
Birthweight (g)	3369	657.6	3351	673.38	0.1237
Maternal age at child birth	29.0	4.62	27.0	5.10	<0.001
	n	%	n	%	
Psychiatric disorder at 15	324	6.59	13.0	8.72	0.302
Self-harm at 16	837	19.7	68	12.19	<0.001
Self-harm at 21	766	21.9	77	14.10	<0.001
Suicidal intent at 16	301	7.09	25	4.48	0.021
Suicidal intent at 21	437	8.95	49	5.63	0.001
Multiple self-harm at 16	448	10.8	39	7.04	0.006
Child sex (female)	4156	62.1	2517	35.5	<0.001
Housing tenure (not owned/mortgaged)	1119	17.3	2312	36.2	<0.001
Maternal education					
<i>Degree</i>	1083	16.8	493	8.49	<0.001
<i>A-level</i>	1735	26.9	1022	17.6	
<i>GCSE</i>	2231	34.6	2007	34.54	
<i><GCSE</i>	1393	21.6	2288	39.38	
Material hardship	1812	27.1	3149	44.36	<0.001
Equivalent household income (quintiles)					
<i>Highest</i>	1380	23.4	611	15.59	<0.001
	1302	22.1	658	16.79	
	1216	20.7	734	18.73	
	1087	18.5	857	21.87	
<i>Lowest</i>	905	15.4	1058	27.00	
Father absence during pregnancy	102	1.6	245	3.81	<0.001
Maternal smoking during pregnancy (ever)	1268	19.1	2274	33.87	<0.001
Parity					
<i>0</i>	3087	47.8	2625	41.68	<0.001

1	2249	34.8	2217	35.20	
2	830	12.9	992	15.75	
3+	293	4.5	464	7.37	
White British ethnicity	6303	98.1	5543	96.55	<0.001


Supplementary Table 3. Adversities experienced by sample (sample for imputation and rest of ALSPAC)

	Sample for imputation (n=6689)		Rest of ALSPAC (n=7099)		p
	n	%	n	%	
Sexual abuse	35	0.58	16	0.46	0.453
Physical abuse	347	6.33	168	6.53	0.73
Emotional abuse	795	16.2	405	18.1	0.048
Child experiences bullying	592	11.3	194	13.9	0.006
Violence between parents	848	17.7	322	20.4	0.019
Parent substance use	459	8.65	237	10.3	0.021
Parent mental health problems or suicide	1967	36.5	965	40.9	<0.001
Parent criminal conviction	343	6.17	183	6.67	0.376
Parental separation	939	17.9	542	24.2	<0.001


Supplementary Table 4. Adversities experienced by sample (complete case and imputed data)

	COMPLETE CASE (n=2373)		Imputed sample (N=6689, 50 iterations)	
	n	%	Percentage	95% CI
Sexual abuse	9	0.38	0.82	0.55, 1.08
Physical abuse	153	6.45	6.98	6.23, 7.72
Emotional abuse	350	14.75	17.73	16.4, 19.0
Child experiences bullying	246	10.37	11.81	11.0, 12.7
Violence between parents	392	16.52	21.18	20.0, 22.4
Parent substance use	178	7.5	10.92	9.96, 11.9
Parent mental health problems or suicide	797	33.59	39.29	37.9, 40.7
Parent criminal conviction	137	5.77	6.89	6.19, 7.59
Parental separation	298	12.56	21.90	20.7, 23.1


Supplementary Figure 1. Age at peak height velocity in males in study sample




Supplementary Figure 2. Age at peak height velocity in females in study sample



Supplementary Figure 3. Age at menarche in females in study sample



Supplementary Figure 4. Descriptive chart showing mean age at peak height velocity for males and females by number of adversities experienced



Supplementary Table 5 Complete case mediation results

Outcome	Model	Direct effect			Indirect effect via mediator			Total effect		
		RR	95% CI	p value	RR	95% CI	p value	RR	95% CI	p value
Self-harm at 16	Age at peak height velocity (both sexes, main analysis N=2251)	1.16	1.09, 1.24	<0.001	1.00	0.99, 1.00	0.508	1.15	1.09, 1.24	<0.001
Self-harm at 16	Age at menarche (females n=1380)	1.16	1.09, 1.23	<0.001	1.00	0.99, 1.01	0.767	1.16	1.09, 1.23	<0.001
Self-harm at 16	Age at peak height velocity (males only, n=981)	1.18	1.04, 1.38	0.015	1.00	0.97, 1.01	0.765	1.18	1.04, 1.38	0.018
Self-harm at 16	Age at peak height velocity (females only, n=1270)	1.15	1.08, 1.23	<0.001	1.00	0.99, 1.00	0.611	1.15	1.08, 1.23	<0.001
Sensitivity analyses- age at peak height velocity										
Self-harm with suicidal intent at 16	Age at peak height velocity (both sexes N=3006)	1.28	1.13, 1.41	<0.001	1.00	0.99, 1.00	0.942	1.28	1.13, 1.41	<0.001
Multiple self-harm in past year at 16	Age at peak height velocity (both sexes N=2251)	1.10	1.01, 1.20	0.025	1.00	0.99, 1.00	0.662	1.10	1.01, 1.20	0.026
Self-harm at 16; no psychiatric disorder at 15	Age at peak height velocity (both sexes n=2873)	1.17	1.09, 1.23	<0.001	1.00	0.99, 1.00	0.661	1.17	1.09, 1.23	<0.001
Self-harm at 21	Age at peak height velocity (both sexes N=3006)	1.12	1.06, 1.20	<0.001	1.00	1.00, 1.00	0.957	1.12	1.06, 1.20	<0.001
Self-harm with suicidal intent at 21	Age at peak height velocity (both sexes, main analysis N=3705)	1.25	1.13, 1.35	<0.001	1.00	1.00, 1.00	0.931	1.25	1.14, 1.35	<0.001

Sensitivity analyses- age at menarche										
Self-harm with suicidal intent at 16	Age at menarche (females n=1762)	1.28	1.14, 1.44	<0.001	1.00	0.99, 1.02	0.629	1.28	1.14, 1.44	<0.001
Multiple self-harm in past year at 16	Age at menarche (females n=1380)	1.15	1.05, 1.27	0.002	1.00	0.99, 1.01	0.779	1.15	1.05, 1.27	0.002
Self-harm at 16; no psychiatric disorder at 15	Age at menarche (females n=1681)	1.17	1.10, 1.25	<0.001	1.00	1.00, 1.00	0.583	1.18	1.10, 1.26	<0.001
Self-harm at 21	Age at menarche (females n=1762)	1.14	1.07, 1.22	<0.001	1.00	1.00, 1.01	0.654	1.14	1.07, 1.22	<0.001
Self-harm with suicidal intent at 21	Age at menarche (females n=1762)	1.24	1.12, 1.36	<0.001	1.00	0.99, 1.01	0.617	1.24	1.12, 1.37	<0.001

